

# Unearthing the Impact of Diverse Nutrient Sources on Soil Quality in Transplanted Rice (Eastern Uttar Pradesh)

## ABSTRACT

A field experiment was conducted at Agronomy Research Farm, Acharya Narendra Deva University of Agriculture & Technology, Kumarganj, Ayodhya (U.P.) during Kharif season 2022 to assess the effect of different nutrients sources on physico-chemical properties of in randomized block design and replicated three times with seven treatment combinations in rice crop. The treatment combinations were **T<sub>1</sub>**:-Control, **T<sub>2</sub>**:-100% RDF, **T<sub>3</sub>**:-75% RDF + 25% N through Vermicompost+ Jeevamrit application at 30 DAT (Days after transplanting), **T<sub>4</sub>**:-75% RDF+25%N through FYM+ Jeevamrit application at 30 DAT, **T<sub>5</sub>**:-50% RDF+50% N through Vermicompost + Jeevamrit application at 45 DAT, **T<sub>6</sub>**:-50% RDF+50%N through FYM + Jeevamrit application at 45 DAT and **T<sub>7</sub>**:-50% RDF+25% N through Vermicompost +25% N through FYM + Jeevamrit application 30 and 45 DAT. Rice variety NDR- 2065 was taken as test crop. After harvesting soil was tested for physico-chemical parameters, according to which T<sub>5</sub> showed maximum available Nitrogen, phosphorus and potassium, in a similar way T<sub>5</sub> also showed improved E.C., p H, organic carbon as well as bulk density and particle density.

**Key Words: Rice, Jeevamrit, Vermicompost, FYM and Soil properties**

## INTRODUCTION

Intensive farming methods that rely heavily on agrochemicals have contaminated land, water, and the environment, negatively impacting human health. In response, alternative farming systems, particularly organic agriculture, are gaining traction due to concerns over environmental sustainability, safe food, and resource conservation. The overuse of chemical fertilizers poses risks, including high costs, depletion of non-renewable energy, and negative effects on soil health and human safety. Combining inorganic fertilizers with organic sources like farm yard manure (FYM), vermicompost (VC), and bio-inoculants can improve soil fertility and maintain crop yields while reducing chemical input dependency. Organic inputs, such as jeevamrit and liquid manures, enhance soil microbial activity and nutrient cycling, thus improving soil health and crop productivity without directly adding nutrients. Chemical fertilizer inputs used carelessly and unevenly have risky impacts on soil health and occasionally end up in the food chain, endangering human health (Karmakar *et al.*, 2013). Inorganic fertilizers can improve soil health and promote higher rice growth and productivity when used in conjunction with organic sources like FYM and vermicompost (Ram *et al.*, 2020). In order to maintain crop yields, organic manures must be applied since they help nutrients build up in the soil (Thirunavukkarasu and Vinoth 2013). In order to reduce the need for chemical inputs while maintaining yields, organic inputs and bio-inoculants are combined (Giraddi, 2000). The main ingredients used to make several fermented organic inputs, including panchagavya, jeevamrit, beejamrit, and vermiwash, include cow dung, urine, pulse flour, jaggery, living soil, and extracts of nearby plants (Kulkarni and Gargelwar, 2019).

Application of jeevamrit improves soil fertility while also enhancing microbial activity in the soil (Aulakh *et al.*, 2018). Utilizing liquid manures effectively promotes crop development and yields while improving the cycling of nutrients. The use of liquid manures increases microbial activity in the soil, improving soil fertility and supplying sufficient amounts of nutrients for crop growth and pest control (Joshi, 2012). Despite the fact that these manures may not directly add nutrients to the soil where they are applied, they do speed up the activity of soil microbes, which helps to preserve the soil's fertility (Yadav and Mowade, 2004).

## **MATERIALS AND METHODS:**

The experiments were conducted at Agronomy Farm of the Acharya Narendra Deva University of Agriculture and Technology in Kumarganj, Ayodhya, which is located in the subtropical Indo Gangetic Plains climate zone at 26.470 N latitude, 82.120 E longitude, and an elevation of 113 meters above mean sea level.

The treatment combinations were **T<sub>1</sub>** (Control), **T<sub>2</sub>** (100% Recommended dose of fertilizer), **T<sub>3</sub>** (75% RDF + 25% N through Vermicompost+ Jeevamrit application at 30 Days after transplanting(DAT)),**T<sub>4</sub>** (75% RDF+25%N through Farm yard manure(FYM)+ Jeevamrit application at 30 DAT, **T<sub>5</sub>** (50% RDF+50% N through Vermicompost + Jeevamrit application at 45 DAT, **T<sub>6</sub>** (50% RDF+50%N through FYM + Jeevamrit application at 45 DAT) and **T<sub>7</sub>** (50% RDF+25% N through Vermicompost +25% N through FYM + Jeevamrit application 30 and 45 DAT). The rate of 100 % Recommended dose of fertilizer (RDF) is 150:60:40 NPK kg ha<sup>-1</sup>.

Soil sampling randomly drawn from each replicated plot after, harvesting of crop. Collected samples were brought to the laboratory, air dried, grind and passed through 30 mesh sieve and representative sample (about half kg) were collected in polythene bag. The physico-chemical properties were analyzed.

Initial value of soil properties viz. Particle density, Bulk density, Soil pH, EC, organic C, available N, P, and K are 2.67 (Mg m<sup>-3</sup>), 1.35 (Mg m<sup>-3</sup>), 8.3, 0.23 (dsm<sup>-1</sup>), 3.65%, 183.14 (Kg ha<sup>-1</sup>), 12.15 (Kg ha<sup>-1</sup>), 240.78(Kg ha<sup>-1</sup>) respectively determined from the processed samples for each treatment (in triplicate) as per the standard methods. Physico-chemical study of soil before and after harvesting of crop. Initial values of of soil properties has been given in table no. 1.

Data collected were subjected to statistical analysis by using a computer program OPSTAT. Least Significant Difference test (LSD) at 5% probability level was applied to compare the differences among treatments means

## **RESULT AND DISCUSSION**

### **Physical analysis**

#### **Bulk Density (BD)**

The data for bulk density (Mg m<sup>-3</sup>) in table no. 2, showed that the treatment with the highest bulk density was T<sub>1</sub> (1.35) (control), which did not differ significantly from the other treatments. T<sub>5</sub> (1.31) had the lowest bulk density throughout the investigation. Due to the

stabilization of the soil structure brought about by the addition of organic manure in the form of vermicompost, changes in bulk density have been noticed. The findings are very similar to those published by Kumar *et al.* (2022) and Koushal *et al.* (2011).

### Particle Density (PD)

The data for particle density ( $\text{Mg m}^{-3}$ ) shown in table no. 2, showed that the rice field in  $T_1$  (control) had the maximum particle density after harvest (2.71), while  $T_5$  (2.62) and  $T_7$  (2.62) had the lowest particle density. The findings are very similar to those published by Koushal *et al.* (2011).

**Table 2: Effect of different nutrient sources on Bulk density and Particle density in soil after harvest.**

S. No.	Treatments	Bulk Density ( $\text{Mg m}^{-3}$ )	Particle density ( $\text{Mg m}^{-3}$ )
1	$T_1$ – Control	1.35	2.71
2	$T_2$ - 100% R.D. F	1.34	2.67
3	$T_3$ - 75% R.D.F + 25% N through Vermicompost+ Jeevamrit application at 30 DAT	1.32	2.64
4	$T_4$ - 75% R.D.F. +25%N through FYM+ Jeevamrit application at 30 DAT	1.32	2.64
5	$T_5$ - 50% R.D.F.+50% N through Vermicompost + Jeevamrit application at 45 DAT	1.31	2.62
6	$T_6$ - 50% R.D.F.+50%N through FYM + Jeevamrit application at 45 DAT	1.31	2.62
7	$T_7$ - 50% R.D.F. + 25% N through Vermicompost +25% N through FYM + Jeevamrit application 30 and 45 DAT	1.31	2.62
CD at 5%		NS	NS
SEM $\pm$		0.029	0.035

### Soil pH:

Table no 3, data on pH showed that the highest pH was found in the  $T_1$  (8.3) (Control) group and the lowest pH was found in the  $T_5$ (8.2) group. In comparison to applying chemical fertilizer alone, applying it along with manures marginally lowered the pH of the soil. The generation of organic acids during the breakdown of manures may be responsible for the pH drop. Sharma *et al.* (2022), Kaur *et al.* (2004), and Kumar *et al.* (2022) have all reported similar findings.

### Electrical Conductivity (EC):

According to the data on E.C. (Electrical Conductivity  $\text{dsm}^{-1}$ ) shown in table no. 3, the rice field with the greatest E.C. after harvesting was  $T_1$  (0.35) (Control), which was statistically equal to  $T_2$  (0.34) and significantly higher than  $T_3$  (0.25).  $T_5$  (0.21) had the lowest electrical conductivity throughout the investigation. Koushal *et al.* (2011), Kumar *et al.* (2022), and Sharma *et al.* (2022) have all found similar findings.

## Organic Carbon (%)

Examining the information in table no. 3, showed that the soil's organic carbon content rose from 0.24% to 0.35%. The treatments T<sub>5</sub> had the highest value of organic carbon (0.35%), which was statistically comparable to T<sub>7</sub>. However, compared to its initial value before 25% of the nitrogen in control and T<sub>2</sub> was replaced with manures, the content of organic carbon dropped. The soil's organic carbon content increased as the amount of nitrogen added by organic sources such FYM, vermicompost, and jeevamrit increased. It might be related to the organic source's sluggish degradation and the increased microbial activity in the soil. Vermicompost and FYM treatments differ in their organic carbon contents, which may be the cause of the variance. Additionally, Kaur *et al.* (2004), Kumar *et al.* (2022), Koushal *et al.* (2011), and Sharma *et al.* (2022) observed similar findings.

**Table 3: Effect of different nutrient sources on pH, EC (dsm<sup>-1</sup>) and organic carbon (%) in soil after harvest.**

S. No.	Treatments	pH	Electrical Conductivity (dsm <sup>-1</sup> )	Organic carbon (%)
1	T <sub>1</sub> – Control	8.30	0.35	0.24
2	T <sub>2</sub> - 100% R.D. F	8.24	0.34	0.28
3	T <sub>3</sub> - 75% R.D.F + 25% N through Vermicompost+ Jeevamrit application at 30 DAT	8.23	0.25	0.32
4	T <sub>4</sub> - 75% R.D.F. +25%N through FYM+ Jeevamrit application at 30 DAT	8.22	0.25	0.31
5	T <sub>5</sub> - 50% R.D.F.+50% N through Vermicompost + Jeevamrit application at 45 DAT	8.20	0.21	0.35
6	T <sub>6</sub> - 50% R.D.F.+50%N through FYM + Jeevamrit application at 45 DAT	8.20	0.23	0.32
7	T <sub>7</sub> - 50% R.D.F. + 25% N through Vermicompost +25% N through FYM + Jeevamrit application 30 and 45 DAT	8.20	0.22	0.35
	CD at 5%	NS	0.013	0.013
	SEm±	0.173	0.004	0.004

## Available N:

Analyzing the information in table no.4, showed that all of the treatments had significantly higher accessible N contents than the control. The treatment T<sub>5</sub> had the highest observed available N content (155.6 kg ha<sup>-1</sup>), which was statistically comparable to T<sub>7</sub>'s (153.1 kg ha<sup>-1</sup>), and much better than the other treatments. The control had the lowest level (135.56 kg ha<sup>-1</sup>). The treatment where nitrogen was applied by substituting vermicompost for 50% of the inorganic fertilizer showed the highest levels of accessible N. Vermicompost has the capacity to supply nutrients more quickly than FYM, and it aids in the transformation of unavailable forms of nitrogen into available forms. It may be explained by the fact that the

addition of organic manure increases the availability of organic matter to microorganisms, which leads to an increase in the activity of soil enzymes and microorganisms, increasing the amount of nitrogen (N) that is available in the soil. As more nitrogen was applied using organic sources, N became more readily available. It might be explained by the organic manure's sluggish breakdown and the plants' reduced absorption. Due of its gradual nutrient release pattern and lower nutritional content as compared to other organic manures, FYM may have a poor nitrogen availability. Kaur *et al.* (2004), Kumar *et al.* (2022), and Koushal *et al.* (2011) have reported similar findings.

**Available P:**

Critical observation of the data of table no. 4, revealed that there was a significant increase in the available P content in all the treatments as compared to the control. Except for the control, all of the treatments had higher available P content than their original values. The treatment T<sub>5</sub> had the highest available P content value (16.99 kg ha<sup>-1</sup>), which was statistically comparable to T<sub>7</sub>'s (16.45 kg ha<sup>-1</sup>), and the control had the lowest value (12.34 kg ha<sup>-1</sup>). It may be caused by the creation of organic acid under the simultaneous application of organic and inorganic sources, the mineralization of soil organic matter, and the solubilizing action of acids created during respirations. These factors may boost the availability of nutrients. Due to low phosphorus use efficiency, the majority of applied phosphorus is fixed in the soil, although over time, a little amount is made accessible for use by plants. By covering sesquioxide molecules and rendering them inactive, the organic matter created by the breakdown of organic manures lowers the soil's capacity to fix phosphate, which ultimately aids in the release of a significant amount of phosphorus. Native P is also made more soluble by the organic acids and CO<sub>2</sub> created during the breakdown of organic material. Both Kumar *et al.* (2022) and Koushal *et al.* (2011) observed similar findings.

**Available K:**

A further examination of the data (Table 4) showed that all of the treatments had significantly higher levels of accessible K than the control. The T<sub>1</sub> (control) group had the lowest amount of available K (250.54 kg ha<sup>-1</sup>). The treatment T<sub>5</sub> had the highest concentration of accessible K (280.35 kg ha<sup>-1</sup>), which was statistically comparable to treatment T<sub>7</sub>'s (275.4). By reducing potassium fixation, releasing potassium due to the interaction of organic matter with clay, and adding directly to the pool of accessible potassium in the soil, potassium availability increased after the application of organic manure. Koushal *et al.* (2011) and Kumar *et al.* (2022) also reported results that were similar.

**Table 4: Effect of different nutrient sources on Available Nitrogen, Phosphorus and potassium in soil after harvest.**

S. No.	Treatments	Nutrient available in soil (kg ha <sup>-1</sup> )		
		N	P	K
1	T <sub>1</sub> – Control	135.56	12.34	250.54
2	T <sub>2</sub> - 100% R.D.F	141.70	14.15	257.33

3	T <sub>3</sub> - 75% R.D.F + 25% N through Vermicompost + Jeevamrit application at 30 DAT	144.30	15.01	264.60
4	T <sub>4</sub> - 75% R.D.F. +25%N through FYM+ Jeevamrit application at 30 DAT	143.70	14.68	260.69
5	T <sub>5</sub> - 50% R.D.F.+50% N through Vermicompost + Jeevamrit application at 45 DAT	155.60	16.99	280.35
6	T <sub>6</sub> - 50% R.D.F.+50%N through FYM +Jeevamrit application at 45 DAT	145.90	15.37	265.90
7	T <sub>7</sub> - 50% R.D.F. + 25% N through Vermicompost +25% N through FYM + Jeevamrit application 30 and 45 DAT	153.10	16.45	275.40
CD at 5%		8.817	0.788	9.721
SEm±		2.83	0.253	3.120

### CONCLUSION:

On the basis of the current analysis, it can be deduced that organic sources had a significant impact on the soil's nutritional status after the crop was harvested. Vermicompost provided more residual major nutrients than FYM, and both organic sources improved the soil's physical characteristics, reducing the soil's bulk density and particle density. In addition to these advantages, organic sources also increased the soil's organic carbon content and biological activity. Therefore, it can be inferred that regular application of a combination of inorganic and organic nutrient sources enhances soil physical and biological features as well as soil nutrient status.

### REFERENCES:

- Aulakh, C. S., Singh, H., Walia, S. S., Phutela, R. P. and Singh, G. 2018. Effect of farmyard manure and jeevamrit in maize-wheat organic production system in Punjab. *Agric. Res. J.*, 55: 485-89.
- Giraddi, R. S. 2000. Influencing vermicomposting methods on the biodegradation of organic wastes. *Indian J. Agric. Sci.*, 70: 663-666.
- Jackson, M. L. 1973. *Soil Chemical Analysis*. Prentice Hall Private Limited, New Delhi. 498p.
- Joshi, M. 2012. *New Vistas of Organic Farming*. Scientific publishers, New Delhi. pp140.
- Karmakar, S., Brahmchari, K. and Gangopadhyay. 2013. Studies on agricultural waste management through preparation and utilization of organic manures for maintaining soil quality. *African J. Agric. Res.*, 8: 6351-58.
- Kaur, K., Kapoor, K. K., & Gupta, A. P. (2005). Impact of organic manures with and without mineral fertilizers on soil chemical and biological properties under tropical conditions. *Journal of Plant Nutrition and Soil Science*, 168(1), 117-122.

Koushal, S., Sharma, A. K., & SODHI, A. S. (2011). Yield performance, economics and soil fertility through direct and residual effects of organic and inorganic sources of nitrogen as substitute to chemical fertilizer in rice-wheat cropping system. *Research journal of agricultural science*, 43(3), 188-192.

Kulkarni, S. S. and Gargelwar, A. P. 2019. Production and microbial analysis of Jeevamrutham for Nitrogen fixers and Phosphate solubilizers in the rural area from Maharashtra. *J.Agric. Veterinary*, 12: 85-92.

Kumar, A., Singh, S. K., Kumar, R., Gautam, S. K., & Sharma, A. (2022). Evaluating the Effects of Organic Manures and Fertilizer N on Soil Properties in Rice Cultivation. *International Journal of Environment and Climate Change*, 12(12), 1140-1148.

Olsen, S. R., Cole, C.V., Watanable, F. S. and Dean L. A. 1954. Estimation of available phosphorus in soil by extraction with sodium carbonate. USDA Circular 939: 1-19.

Ram, M. S., Shankar, T., Maitra, S., & Duvvada, S. K. (2020). Effect of integrated nutrient management on growth, yield, nutrient content and economics of summer rice (*Oryza sativa* L.). *Indian J Pure Appl Biosci*, 8, 421-427.

Richards, L. A. (Ed.). (1954). *Diagnosis and improvement of saline and alkali soils* (No. 60). US Government Printing Office.

Sharma, K., Kaushal, R., Sharma, S., & Negi, M. (2022). Effect of organic and inorganic nutrient sources on soil physico-chemical and microbiological properties in cauliflower (*Brassica oleracea* var. botrytis L.) under mid hills of Himachal Pradesh. *Journal of Crop and Weed*, 18(1), 01-06.

Subbiah, H. V. and Asija H. L. 1956. A rapid procedure for the estimation of available nitrogen in soils. *Curr. Sci.*, 25: 259-260.

Thirunavukkarasu, M. and Vinoth, R. 2013. Influence of Vermicompost application along with nitrogen on growth, nutrients uptake, yield attributes and economics of rice (*Oryza sativa* L.). *Int. J. Agric. Environ. Biotechnol.*, 6: 599-604.

Walkely, A. and Black I. A. 1934. Examination of the method for determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Sci.*, 37: 29-38.

Yadav, A.K. and Mowade, S. M. 2004. Organic manures and compost. In: *Organic Farming- A Ray of Hope for Indian Farmer*. National Center of Organic farming, Ghaziabad, Uttar Pradesh