

Determination Of Better Acclimatization Process For *Channa Punctatus* , A Fresh Water Air Breathing Teleost In Laboratory Condition

Abstract

Acclimatization is a process in which an individual organism adjust to a change in its environment. The present experiment was designed to determine a better acclimatization process for *Channa punctatus* in different laboratory conditions. Maximum mortality of fishes was recorded in tank A (50%), then tank B (30%) and tank C (20%). Body pigmentation was found decreased among all the fishes in all the tanks (A-D). Less swimming activity, increase opercular activity and loss of equilibrium were also recorded among the experimental fishes. The maximum survival & adaptability were recorded among the fishes of tank D.

Keywords :- Acclimatization, *Channa punctatus*, Behavior anomalies, Mortality, Laboratory condition.

INTRODUCTION

Acclimatization is a process in which an individual organism adjust to a change in its environment (such as a change in altitude , [temperature](#)- humidity , photoperiod,hardness or pH) allowing it to maintain between a range of environmental conditions [1] .

Experimental Study is an important aspect of biological research. Experimental study using live intact animals (fishes, rats, mice, rabbit etc.) have continued to play essential role in development of new knowledge and better understanding of life process [2,3,4].

The use of intact animal in biological research for efficient and effective design is always required. In this context , different species of fishes (species of *Channa*, *Clarias*, *Heteropneustes*, *Anabas* etc) have been frequently used as biological model in the laboratory conditions [5,6,7,8,9,22,10,11,22].

Fishes form one of the important group of vertebrates influencing the human life in various ways. Fish is a cold blooded animal, which has the ability to adjust with the changed environment [12,3]. At the same time they are useful indicator of environmental quality and ecological integrity. [13,14].The unique adaptation and physiological specialization of fish made them especially suitable for use as physiological and biological model [15,4]. Fish occupies a peculiar position in the hierarchy of animal life because higher vertebrates evolved from them. Fishes have great importance as a secondary consumer in the food chain. This significance is magnified as it forms a major food for higher trophic levels [16].

Presently due to rapid industrialization and discharge of pollutant in ~~their~~ aquatic ecosystem, fishes and other aquatic animals are running under threat and ultimately causing higher mortality [17,18]. Many laboratory and field studies suggest that aquatic organism may acclimate to some toxicants . It has been hypothesized that animal exposed to sublethal level of a pollutant could become more tolerant or weakened depending upon the mode of action of the poison at which the animal has been acclimated [19]. Various acclimatization methods have been applied to overcome the fish mortality .

So far as *Channa* species are concerned it is not easy to keep them in laboratory for longer period as compared to other fishes like species of *Anabas* , *Clarias*, *Colisa* and *Heteropneustes*.

Keeping the above facts in mind , the present experiment was designed to study or to determine the better acclimatization process for *Channa punctatus* , a fresh water air breathing teleost fish in different laboratory conditions.

MATERIALS AND METHODS

Test Animal

The live healthy specimens of fresh water air breathing fish *Channa punctatus* (Total length 15.5 – 17.5cm and body weight 35-45 g) were collected from local fish market of Nathnagar & Bhagalpur, Bihar, and brought to the laboratory. *Channapunctatus*, locally known as Garai, is a snake headed air breathing fish belong to the family Channidae and order Channiformes. These fishes are mostly found in the fresh water pools & wetlands like ponds, ditches, chauras and swampy areas of Northern India. These fishes can survive in oxygen deficient water bodies [20,21,22].

Experiment Design

The experiment trials were conducted in four glass tanks (aquaria) arranged in a row on a wooden table in the departmental Research Laboratory. Fishes were categorized into four groups each containing 10 live specimens of *Channa punctatus*. These fishes were placed in four glass tanks (size 50cm×30cm×25cm) separately containing 25 liters of water in each tank. The tanks were named as A, B, C & D.

All tanks (A-D) containing 10 live test specimens, had different bottom habitat. Fishes were kept under observation for the period of 4 weeks for acclimatization to laboratory conditions. Tank A was with only glass bottom, tank B was covered with pure sand (1 inch thick) on the bottom, tank C was having equal proportion of sand and soil of the same thickness, whereas the tank D had mud at the bottom of same thickness.

All the four tanks (A, B, C & D) were kept in the laboratory and the photoperiod was maintained as per the normal day and night around 10 hours with proper optimal condition for fish. Around 50% of water from each tank was replaced with tap water after every week during the experiment period. No mortality occurred during this period. All the effort was made to provide natural environment to the fishes as far as possible in the laboratory. During this period fishes were fed boiled chicken eggs and chopped parts of small fishes @ 5% of body weight approximately on alternate days [4].

The experiment was conducted in Biodiversity laboratory of Univ. Dept. of Zoology , TMBU, Bhagalpur , Bihar during the month of october 2023. Fishes were exposed and frequent monitoring was made to observe the behavior pattern and survival of the fishes. A fish was considered dead when no response observed after touching/probing with glass rod. Dead fishes were immediately eliminated from the tank. Total mortality and abnormal behavior (if any) were monitored and recorded regularly between 8a.m. – 5p.m. Every observation and data were collected properly to draw better inferences.

RESULTS AND OBSERVATIONS

During the study of four weeks , The following observation were recorded :-

- No any mortality or significant changes in behavior was noticed among the fishes in the four days of experiment in all the tanks (A-D).
- At the beginning, fishes were found to be healthy and very active. But after four days of the experiment, fishes of tanks A,B&C expressed some structural and behavioural abnormalities.
- On fifth day of experiment some changes were noticed among the fishes of tanks A, B&C. Fishes were found less active. Swimming as well as body pigmentation were also found decreased in comparision to fishes of the tank D.Colour of the fishes changed from dark black to pale in colour.
- Feeding attempts and feeding activity were also found decreased in among the fishes in among all the tanks (A-D).
- On day eight , It was noticed that from some fishes of tanks A&B were showing random swimming movement and loss of equilibrium.
- Fishes of tank C were also found in stress, showed increased opercular activities .
- Fishes of tank D were expressing normal swimming movement . No sign of above mention abnormalities were noticed among the fishes of tank D till date.
- Mortality of fishes were recorded after eight days of experiment. 02 fishes from tank A&B and 01 fish from tank C was found dead in the

morning on day 9th of the experiment . These dead fishes floating on the water surface keeping their bodies upside down.

- Similarly, on day twelve again 02 fishes of tank A and 01 fishes each from tank B&C died.
- No mortality was recorded among the fishes of tank D. Fishes were normal and these were spending most of the time buried in the mud.They were coming to water surface for engulfing air at regular intervals.
- After 3 weeks(22nd day) of experiment 01 fish died from tank A&B in the afternoon.
- Before death it was noticed that fishes were assigned in vertical position with keeping its mouth near the water surface and tail in downward direction. Thereafter, they settled down passively at the bottom of the tank turning its bellies upward.
- At the completion of four weeks mortality rate was found maximum among the fishes of tank A (50%), 30% among the fishes of tank B, 20% among the fishes tank C and no mortality was recorded up to this date among the fish of tank D.
- Except colour changing and no sign of any further mortality behavioral anomalies were recorded among the fishes of tank D at the completion of this experiment.

Fish Tanks	No. of Fishes	Days of Mortality of the fishes					Total Mortalities	Behavior Observed	
		7 th	8 th	9 th	12 th	22 nd		Before 1 st week	After 1 st week
Tank A	10	-	02	-	02	01	05	L.A. L.F.A, L.P	L.E.
Tank B	10	-	02	-	01	01	04	S.F.A.,L.P.	L.E.
Tank C	10	-	-	01	01	-	02	F.O.A.&L.F	I.O.A
Tank D	10	-	-	-	-	-	-	N.	N.

Table 1 : Mortality and behavioral anomalies among the fishes during acclimatization indifferent laboratory conditions.

Formatted: Highlight

Abbreviations :-

L.A- Less Active, S.F.A - Slow Feeding Attempts, F.O.A - Fast Opercular Activity, N.- Normal, L. P- Less Pigmentation, L.S- Less swimming, L.F.A.-Less Feeding Attempts & I.O.A-Increase Opercular Activity. L.F.= ?




Fig 1. The number of mortality of fishes in different tanks (habitat) during acclimatization




Fig 2- The % of mortality of the fishes in tanks (A-D) in different habitat conditions.

DISCUSSION

The present study elucidate the better acclimatization of *Channa punctatus* in different laboratory conditions. The behavioral study gives direct responses of the fish to the ambient environment or chemical present in the water [4]. Warner *et al.*, 1966 and Radhaiah, *et al.*, 1987 [23,24] suggested that behavioral activities of an organism represent the final integrated result of diversified biochemical and physiological processes.

Most of the fishes under observation were found less active and expressed erratic swimming movement . It may be the effect of stress caused due to sudden change in their habitat condition of their new aquatic environment [25]. Roy *et al.*, 1989 [26] also reported that fishes need a stable environment during the acclimatization process in specially in terms of stable pH.

Body pigmentation were also decreasing gradually in almost all the fishes on all the tanks (A-D), may be due to dysfunction of the pituitary gland under stress causing changes in the number and distribution of chromatophores [27,28,29].

Behavioral changes as a result of stress are further accepted at the most sensitive indication of adverse effect of aquatic environment [30]. Koprucu, *et al.* [31] also observed lightening skin colour in the fingerlings of European cat fish exposed to contaminated water. There are few evidence that alteration of the chemical composition in the natural aquatic environment , usually effect behavioral and physiological system of inhabitants particularly fishes [24,32]. Bad water quality or changes in physico-chemical properties of water as well as the unfavourable habitat of the fish tank may also cause stress or may influence the behavioral pattern of fishes including loss of equilibrium, increase rate of opercular activity and less feeding attempts, may be the another reasons.

Faster opercular cavity and loss of equilibrium were observed among the fishes of tank A, B, & C. Hassanein & Okail [33] had also ~~observed~~ [observed](#) loss of buoyancy and balance with an initial increase in the opercular ventilation rate which then decreased significantly in *Ctenopharyngodon idella* (grass carp) after exposure to the biopesticide. According to Fulton & Key [34], the restlessness and hyperactivity in fish may occur due to the stress or inactivation of acetylcholinesterase, enzyme leading to accumulation of acetylcholine at synaptic junctions. Finally, fishes were almost paralysed and settled on the bottom of the tank and died. Mortality of fishes were recorded in tank A, B & C but maximum from tank A (50%) then tank B(30%) and tank C (20%) may be due to above mention reasons.

Survival rate was found maximum among the fishes of tank D (100%) may be due to favourable habitat and aquatic environment as *Channa punctatus* is an air-breathing fish and found in muddy and swampy areas with less oxygen content [2]. It was well acclimatized in mud water , and showed more adaptability in this habitat then other fishes of other tanks.

CONCLUSION

Study clearly revealed that the fishes in tank D expressed more adaptability. Survival rate was also found maximum among these fishes in comparision to fishes of other three tanks. Fishes expressed some behavioral anomalies due to stress caused by the sudden change in their natural habitat or aquatic environment. Mudwater and adequate size of the tank /aquarium containing sufficient volume of water is the best habitat for the better acclimatization of *Channa punctatus* in the different laboratory conditions.

REFERENCES

1. Mathur R.Effect of toxicants on movement of fish. Animal Behaviour. Fifth edition: 2019; 572.

2. Ghosh TK, Moitra A, Datta JS. Bimodal oxygen uptake in relation to body weight and seasonal temperature of an air-breathing murrel, *Channa punctatus*. J. Freshwater Biol.1990; 2(1):57-64.
3. Akter R, Ahmed K, Mollah RAM, Islam M. Effects of pre-exposure acclimatization to diazinon(60EC) on the mortality of freshwater murrel, *Channa punctatus* (Bloch, 1973). Bangladesh Journal of Environmental Science,2008; 13 (2):367-371.
4. Pandey AK, Nagpure NS, Trivedi SP, Kumar R, Kushwaha B, Lakra WS. Investigation on acute toxicity and behavioural changes in *Channa punctatus* due to organophosphate pesticide profenofos. Drug and Chemical Toxicology, 2011; 34 (4):424-428.
5. Hughes GM, Singh BN. Respiration in the air-breathing fish , the climbing perch, *Anabas testudineus* (Bloch). Oxygen and carbon dioxide release into air and water. J. Exp. Biol. 1970; 53 : 265-280.
6. Munshi JSD, Patra AK, Hughes GM. Oxygen consumption from air and in *Heteropneustes (=Saccobranhus) fossilis* (Bloch) in relation to body weight at three different seasons. Proc. Indian natn. Sci. Acad. B.1982; 48 (6) : 715-729.
7. Singh BR, Mishra AP, Singh RP. Development of the air-breathing organ in the snake-headed fish, *Channa punctatus* .Zool.Anz. 1982;208 : 428-439.
8. Patra AK, Munshi JSD, Hughes GM. Oxygen consumption of the freshwater air-breathing Indian siluroid fish, *Clarias batrachus* (Linn.) in relation to body size and seasons. Proc. Indian natn. Sci. Acad.B 1983; 49 (6):566-574.
9. Ghosh TK, Moitra A, Kunwar GK, Munshi JSD. Bimodal oxygen uptake in a freshwater air-breathing fish, *Notopterus chitala*. Japan J. Ichthyo.1986; 33 : 280-285.
10. Ojha J, Singh NK, Ghosh TK. Bimodal oxygen uptake in relation to body weight in a fresh-water air-breathing fish, *Colosa (=Trichogaster) fasciatus* (Bloch & Schneider). Arch. Hydrobiol. 1988; 111 (3) : 475-480.
11. Okpala COR, Sardo G, Vitale S. On the acclimatization of pelagic fish species juveniles in a closed environment: An atlantic horse mackerel case study. American Journal of Animal and Veterinary Sciencess, 2017.
12. Al-Arabi SAM, Alam MGM, Mazid MA. Toxic effects of Ripcord to the snake-headed, *Channa punctatus*. Bangladesh J. Training and Development.1993; 6(2): 97-101.

13. Roy PK, Munshi JSD. Toxicity of technical and commercial grade Malathion on *Cirrhinus mirgala*(Ham.), A major carp. Biol.Bull. of India 1987; 9(1) :50-56.
14. Khanna SS, Singh HR. Characters and Classification. A Textbook of Fish and Fisheries 3rd Edition: 2017; 1.
15. Lakra WS, Nagpure NS. Genotoxicology studies in fishes: a review. *Ind. J. Anim. Sci.*2009; 79:93-98.
16. Kumar S, Tembhre M. Preface.Fish and Fisheries.2014.
- 17.Ramona A, Biswas AK, Kundu S, Saha JK Yadav RBR. Efficacy of distillery effluent on seed germination and seedling rowth in mustard, cauliflower and radish. Proceedings of National Academy of Sciences, India. 1979.
- 18.Kumari AN, Ram Kumari S. Effect of polluted water on histochemical localization of carbohydrates in a freshwater teleost *C. punctatus* form Hussain Sugar Lake , Hyderabad, Andhra Pradesh. *Pollution Research* 1997;16:197-200.
- 19.Sprague JB, Elson PF, Saunders RL. Sublethal copper-zinc pollution in a salmon river – a field and laboratory study . *Int. J. Air Water Pollut.*1965; 9: 531-543.
- 20.Munshi JSD, Sinha AL, Ojha J. Oxygen uptake capacity of gills and skin in relation to body weight of the air-breathing siluroid fish, *Clarias batrachus* (Linn.). *Acta. Physiol.*1976; 84 (1): 23-33.
- 21.Ojha J,Patra AK, Biswas N, Munshi JSD. Interspecific variations in the bimodal oxygen uptake of the genus *Channa punctatus*. *Z. Tierphysiol.*1981; 46: 165-170.
- 22.Ghosh TK, Munshi JSD. Biomodal oxygen uptake in relation to body weight and seasonal temperature of an air-breathing climbing perch, *Anabas testudineus*(bloch).*Zool. Beitr. N.F.* 1987; 31 (3) :357-364.
- 23.Warner RE, Peterson KK, Burgman L. Behavioral pathology in fish. A quantitative study of sub-lethal pesticide toxication. *J. Appl. Ecolol.*1966; 3:223-242.
- 24.Radhaiah V, Girija M, Rao KJ. Changes in selected iochemical parameters in the kidney and blood of the fish, *Tilapia mossambica* (Peters), exposed to heptachlor. *Bull. Environ Contam Toxicol.*1987; 39: 1006-1011.
- 25.Srivastava AK,Mishra D, Shrivastava S,Srivastav SK, Srivasrav AK. Acute toxicity and behavioural responses of *Heteropneustes fossilis* to an

- organophosphate insecticide, Dimethoate. International Journal of Pharma and Bio Sciences.2010; 1 (4) : 359-363.
- 26.Roy PK, Munshi JSD. Effect of Saponin on oxygen uptake and haematology of air-breathing climbing perch, *Anabas testudineus* (Bloch). J. Freshwater Biol.1989; 1 (2), 167-172.
- 27.Pandey A, Kunwar GK, Munshi JSD. Integumentary chromatophores and mucus gland of fish as indicator of heavy metal pollution. J. Freshwater Biol.1990; 2: 117 – 121.
- 28.Yadav A, Neralia S, Gopesh A. Acute toxicity levels and ethological responses of *Channa punctatus* to fertilizer industrial wastewater. J. Environ Biol . 2007; 28: 159-162.
- 29.Ramesh M, Saravanan M. Haematological and biochemical responses in a freshwater fish *Cyprinus carpio* exposed to chlorpyrifos. I.J. Integ Biol. 2008; 3 ;80-83.
- 30.Nwani CD, Nagpure NS, Kumar R ,Kushwaha B,Kumar P, Lakra WS. Lethal concentration and toxicity stress of Carbosulfan, Glyphosate and Atrazine to freshwater air breathing fish *Channa punctatus*. Int. Aquat.Res.2010; 2: 105-111.
- 31.Koprucu SS, Koprucu K, Ural MS, Ispir U, Pala M. Acute toxicity of organophosphorus pesticide diazinon and its effects on behaviour and some haematological parameters of fingerling European catfish (*Silurus glanis* L.). Pesticide Biochemistry and Physiology. 2006; 86:99-105.
- 32.Khan MZ, Law FCP. Adverse effects of pesticides and related chemical on enzyme and hormone systems of fish, amphibians and reptiles: a review. Proc Pakistan Acad Sci.2005; 42: 315-323.
- 33.Hassanein HMA Okail HA. Toxicity determination and hypoglycaemic effect of neem biopesticide on the grass carp "*Ctenopharyngodon idella*". Egypt Acad. J. Biolog. Sci.2008; 1 (2): 37-49.
- 34.Fulton MH, Key PB. Acetylcholinesterase inhibition in estuarine fish and invertebrates as an indicator of organophosphorus insecticides exposure and effects. Environmental Toxicology Chem. 2001; 20: 37-45.