

# Influence of Antioxidants and Cytokinins on *In Vitro* Plantlet Regeneration of Banana cv. Ney Poovan (AB)

**Comment [s1]:** Research topic is good and novelty of research is important for research institute and industry

It can be accepted after correction of some minor correction which are given below

## ABSTRACT

**Aims:** Banana is one of the important fruit crop and food source for millions of people in many countries. In South India, many varieties of banana are cultivated. *In vitro* micropropagation of banana is nowadays pinned towards development of disease-free clones. An investigation was carried out in banana cv. Ney Poovan to study Influence of Antioxidants and cytokinins on in vitro plantlet regeneration.

**Study design:** The investigation was carried out in a completely randomized design with 25 treatments replicated thrice

**Place and Duration of Study:** Tissue culture laboratory, Bio Centre, Department of Horticulture, Shivamogga 2023-2024

**Methodology:** For controlling phenolic browning and to regenerate multiple plantlets production from the shoot tip, the explants were well sterilized and transferred to culture media containing MS media in combination with antioxidant (Ascorbic acid and Polyvinylpyrrolidone) and cytokinins (6-Benzylaminopurine and Kinetin)

**Results:** Ascorbic acid 200 mg/l showed less browning across all combinations of BAP and kinetin. On an average the minimum number of days taken for shoot initiation (8.13 days), maximum number of shoots per explant (5.02), maximum shoot length (5.20 cm) and maximum success rate (73.33 %) was recorded in explants treated with (T<sub>12</sub>) Ascorbic acid 200 mg/l + BAP 2.5 mg/l + Kinetin 2.0 mg/l.

**Conclusion:** Through this study, it is clear that shoot tip explant treated with MS basal medium + Activated charcoal 200 mg/l + Ascorbic acid 200 mg/l + BAP 2.5 mg/l + Kinetin 2.0 mg/l is best for controlling browning and *in vitro* shoot regeneration for banana cv. Ney poovan.

**Comment [s2]:** Aims is written only 1-2 line

**Keywords:** Micropropagation, Ney Poovan, Ascorbic acid, Polyvinylpyrrolidone (PVP), 6-Benzylaminopurine (BAP), Kinetin

## 1. INTRODUCTION

Banana (*Musa* spp. L.) is one of the most important cash crops and contributes immensely to global food security. It belongs to the family Musaceae and is distributed all around the world. Banana is considered as fourth most important food in the world after rice, wheat and maize. India occupies the third place in annual production of banana. It originated from the South East Asian region, where the greatest diversity of edible bananas is found [1]. Banana and plantains are traditionally propagated through sword suckers. The conventional propagules are not the ideal planting materials because they carry insect pest and disease causing pathogens. Micropropagation is an alternative and viable technology to produce and supply quality planting materials to the banana growing farmers without any pest and diseases. In terms of yield performance, tissue cultured plants have been reported to produce 39% higher yield than plants from sword suckers [2].

Different attempts have been made to alleviate browning problem, incorporation of antioxidants into the culture medium is one of them. Antioxidants play a vital role in the standardization of any micropropagation protocol to overcome browning of explant. As an antioxidant, potassium citrate-citrate, Activated charcoal, ascorbic acid, cystine, polyvinylpyrrolidone and silver nitrate can be used [3]. Cytokinins comprise a separate class of growth promoters. They stimulate the synthesis of proteins and actively take part in cell cycle control. They are known to overcome apical dominance and proclim lateral buds from dormant tissues. The cultivars have different proliferation rate which is significantly influenced by cytokinin type, concentration and ploidy level of the banana cultivars. Kinetin (kin) and BAP (6-benzylaminopurine) are major cytokinins used in tissue culture.

One of the best diploid banana cultivar is Ney Poovan (AB), which is widely grown commercially as mono cultivation, particularly in Tamil Nadu and Karnataka. Fruits are small, slender and attractive because of their ivory white pulp and bright yellow peel. Fruits are rich in carbohydrates, minerals, potassium and other nutritional elements. The cultivar is known for its taste, aroma and long keeping quality [4]. The plant is highly susceptible to Panama wilt and leaf spot diseases [5]. Cultivar also have browning problem under *in vitro* regeneration which is due to exudation of large amount of phenols. Therefore, in the present situation, requirement of large numbers of quality planting materials for local banana cultivar is needed. Therefore, to find out suitable ways for *in vitro* propagation of Ney Poovan, the objective of this study was to know the effect of different levels of antioxidants on browning of banana explant and to study the influence of Cytokinins on *in vitro* plantlet regeneration of Banana cv. Ney Poovan (AB).

## 2. EXPERIMENTAL DETAILS

The explants for the present experiment were collected from healthy and vigorously grown sword suckers of cv. 'Ney Poovan (AB group)' of 3 - 4 months old and in active growth phase, free from diseases and pests. The sword suckers were collected from a farmer's field, Mundre, Tarikere taluk, Chikmagalur, Karnataka. The study was conducted at the Tissue Culture Laboratory, Bio Centre, Department of Horticulture, Shivamogga from September 2023 to April 2024. The experiments were arranged in a completely randomized design (CRD) with 25 treatments each replicated thrice. For each replication, 5 explants were inoculated.

### 2.1 Sterilization of explant

After removing the outer sheaths of the suckers, they were cut into 5 to 10 cm of pieces. These suckers were kept under running tap water (for 30 mins.). The explants were soaked in Bavistin (2 g/l) solution overnight and then treated with 100 ml/l of sodium

**Comment [s3]:** In introduction section add more information about banana tissue culture

**Comment [s4]:** Add reference

**Comment [s5]:** Add more information about callus and rooting of *in vitro* culture of Banana Shukla, S., Rajbhar, Y., Sharma, A., Bhardwaj, P., Tiwari, V. K., Tiwari, H., ... & Kumar, A. (2020). Effect of auxins supplementation on callus induction and root regeneration in banana (*Musa paradisiaca* L.) cv. Udhayam under *in vitro* cultures.

hypochlorite solution for 10 minutes and washed with distilled water. These explants were next sterilized under aseptic conditions in the laminar airflow chamber by washing them with sterile water along with two to three drops of Tween 20 emulsifier solution followed by citric acid + Ascorbic acid (100 mg/l + 150 mg/l) respectively for 10 min, also with mercuric chloride solution (500 mg/l) for 10 min and then with Taxim solution 1 injection/l for 20 min, after each treatment explants were rinsed twice with sterile water so that no remaining chemical residue. After these treatments these explants were trimmed up to 3 to 5 cm before inoculation into MS media.

## 2.2 Culture media

For shoot initiation and control of explant browning, MS basal media along with activated charcoal 200 mg/l was supplemented with different combinations of ascorbic acid (150 and 200 mg/l) with BAP (2 and 2.5 mg/l) and kinetin (2 and 2.5 mg/l) (T<sub>1</sub>-T<sub>14</sub>) and polyvinylpyrrolidone (0.4 and 0.5 mg/l) with BAP (2 and 2.5 mg/l) and kinetin (2 and 2.5 mg/l) (T<sub>15</sub>-T<sub>25</sub>). For this study banana explants were subcultured after every twenty-one days into initiation stage (I<sub>1</sub> and I<sub>2</sub>), division stage (D<sub>1</sub> and D<sub>2</sub>) and multiplication stage (first multiplication cycle and second multiplication cycle).

## 2.3 Stock Solution and Media Preparation

The media namely MS [6] were prepared by dissolving the appropriate amount of macro and micro nutrient; and organic supplements. Similarly, growth regulators (BAP and Kinetin) stock solutions were prepared using the proportion of 20 mg: 20 ml and stored in the refrigerator at 4°C. The MS culture media were prepared from their respective stock solutions using the appropriate amount of sucrose, plant growth regulators, activated charcoal and agar (7 g/l). All individual and combinations of both growth regulators at two levels of BAP and two levels of kinetin were added separately to the media to study its effect on shoot regeneration. The bottles with media were then dispensed 35 ml each and autoclaved at 121°C for 25 minutes after adjusting the pH to 5.7 with 1N NaOH and/or 1N HCl.

## 2.4 Data Collection and Analysis

Five parameters *viz.*, Degree of explant browning, Number of days taken for shoot initiation, Number of shoots per explant, Length of the shoots (cm) and Success rate (%) were recorded after specific interval of time. Data were statistically analyzed using the analysis of variance (ANOVA) technique applicable to the completely randomized design. The significance of the treatment effect was determined using F-test; the means of the treatments are tested using the critical differences (CD) at the 1% probability level.

**Comment [s6]:** Mention the data analysis software

# 3. RESULTS AND DISCUSSION

## 3.1 Effect of antioxidants on the degree of explant browning

The effect of antioxidants at different concentrations on the degree of explant browning was observed visually during callus initiation and division stages of banana explant. Minimum phenolic browning was noticed when shoot tip explants were incubated onto (T<sub>6</sub> to T<sub>13</sub>) MS basal medium + Activated charcoal 200 mg/l + Ascorbic acid 200 mg/l across all combinations of BAP and kinetin (score +: minimum browning). Whereas, maximum phenolic browning was obtained when shoot tip explants were inoculated onto (T<sub>14</sub> to T<sub>17</sub>) MS basal medium + Activated charcoal 200 mg/l + PVP 0.4 mg/l and MS basal medium +

Activated charcoal 200 mg/l without antioxidants (control) (score + + +: maximum browning of the explants). Ascorbic acid gave better result in preventing browning due to its capacity to convert colorless o-quinones formed as a result of Polyphenol oxidases (PPO) activity back into diphenols. Ascorbic acid also have ability to directly inhibit the enzymatic browning process, scavenge free radicals and support overall cell metabolism makes it a more effective antioxidant in preventing browning in banana tissue culture compared to PVP. Similar results were observed in the findings of Ndakidemi et al. [7] Nguyen et al. [8] and Strosse et al. [9]

Comment [s7]: resentence the sentence

UNDER PEER REVIEW

**Table.1. The effect of antioxidant and cytokinins on browning of explant, shooting parameters and survival percentage in banana cv. Ney poovan**

Treatments	Treatment details (mg/l)	Degree of explant browning	Days taken for shoot initiation	Number of shoots per explant	Length of the shoots (cm)	Success rate (%)
			Mean	Mean	Mean	
T <sub>1</sub>	MS basal medium only (control)	+++	13.50	0.23	1.11	6.67 (8.86)
T <sub>2</sub>	Ascorbic acid 150 + BAP 2.0 + Kinetin 2.0	++	9.99	3.16	3.90	26.67 (30.79)
T <sub>3</sub>	Ascorbic acid 150 + BAP 2.0 + Kinetin 2.5	++	9.60	3.60	4.29	26.67 (30.79)
T <sub>4</sub>	Ascorbic acid 150 + BAP 2.5 + Kinetin 2.0	++	9.39	4.66	4.89	33.33 (35.01)
T <sub>5</sub>	Ascorbic acid 150 + BAP 2.5 + Kinetin 2.5	++	9.98	4.47	4.18	33.33 (35.01)
T <sub>6</sub>	Ascorbic acid 200 + BAP 2.0	+	11.93	2.56	3.71	46.67 (43.08)
T <sub>7</sub>	Ascorbic acid 200 + BAP 2.5	+	11.82	2.84	3.86	46.67 (42.70)
T <sub>8</sub>	Ascorbic acid 200 + Kinetin 2.0	+	12.15	1.89	3.51	46.67 (43.08)
T <sub>9</sub>	Ascorbic acid 200 + Kinetin 2.5	+	12.36	1.72	3.40	40.00 (39.23)
T <sub>10</sub>	Ascorbic acid 200 + BAP 2.0 + Kinetin 2.0	+	8.78	3.94	4.68	46.67 (43.08)
T <sub>11</sub>	Ascorbic acid 200 + BAP 2.0 + Kinetin 2.5	+	8.66	4.88	5.06	40.00 (39.23)
T <sub>12</sub>	Ascorbic acid 200 + BAP 2.5 + Kinetin 2.0	+	8.13	5.02	5.20	73.33(59.21)
T <sub>13</sub>	Ascorbic acid 200 + BAP 2.5 + Kinetin 2.5	+	9.04	4.79	4.92	46.67 (42.70)
T <sub>14</sub>	PVP 0.4 + BAP 2.0 + Kinetin 2.0	+++	10.21	3.32	3.43	26.67 (30.79)
T <sub>15</sub>	PVP 0.4 + BAP 2.0 + Kinetin 2.5	+++	10.00	3.55	3.61	20.00 (26.57)
T <sub>16</sub>	PVP 0.4 + BAP 2.5 + Kinetin 2.0	+++	9.57	3.89	3.93	20.00 (26.57)
T <sub>17</sub>	PVP 0.4 + BAP 2.5 + Kinetin 2.5	+++	10.38	3.92	3.29	26.67 (30.79)
T <sub>18</sub>	PVP 0.5 + BAP 2.0	++	12.16	2.78	3.23	46.67 (43.08)
T <sub>19</sub>	PVP 0.5 + BAP 2.5	++	11.82	2.94	3.35	33.33 (35.01)
T <sub>20</sub>	PVP 0.5 + Kinetin 2.0	++	12.27	1.78	3.00	33.33 (35.01)
T <sub>21</sub>	PVP 0.5 + Kinetin 2.5	++	12.24	1.34	2.93	26.67 (30.79)
T <sub>22</sub>	PVP 0.5 + BAP 2.0 + Kinetin 2.0	++	10.40	3.55	3.73	26.67 (30.79)
T <sub>23</sub>	PVP 0.5 + BAP 2.0 + Kinetin 2.5	++	10.17	3.50	3.81	33.33 (35.01)
T <sub>24</sub>	PVP 0.5 + BAP 2.5 + Kinetin 2.0	++	10.33	3.93	4.37	20.00 (26.57)
T <sub>25</sub>	PVP 0.5 + BAP 2.5 + Kinetin 2.5	++	10.60	3.51	3.75	33.33 (35.01)
	S.Em.±	-	0.17	0.06	0.06	4.47
	C.D. at 1%	-	0.65	0.22	0.21	16.93
	C.V	-	2.80	3.07	2.51	-

Note: MS basal medium + Activated charcoal 200 mg/l is common in all treatments, The values given in parenthesis are arc sine transformed values, BAP- 6- Benzylaminopurine, PVP- Polyvinylpyrrolidone

+ - minimum browning, + + - medium browning, + + + - maximum browning

### 3.2 Effect of antioxidants and cytokinins on shooting parameters of banana cv. Ney Poovan

#### Days taken for shoot initiation

After completion of second multiplication stage, on an average, the days taken for shoot initiation varied significantly (Table1). The minimum average number of days for shoot initiation (8.13 days) was recorded in (T<sub>12</sub>) MS basal medium + Activated charcoal 200 mg/l + Ascorbic acid 200 mg/l + BAP 2.5 mg/l + Kinetin 2.0 mg/l, which was on par with (T<sub>11</sub>) MS basal medium + Activated charcoal 200 mg/l + Ascorbic acid 200 mg/l + BAP 2.0 mg/l + Kinetin 2.5 mg/l (8.66 days) and (T<sub>10</sub>) MS basal medium + Activated charcoal 200 mg/l + Ascorbic acid 200 mg/l + BAP 2.0 mg/l + Kinetin 2.0 mg/l (8.78 days). The various combinations of BAP and kinetin were helpful for initiation and regeneration of plants, but the difference in the results may be due to the endogenous levels of cytokinin. These results also support the findings of Zaffari et al. [10], Shukla et al. [11] and Azam et al. [12]

**Comment [s8]:** What is the reason minimum was recorded in T-12 treatment?

#### The number of shoots per explant

The number of shoots per explant differed significantly among different treatments. On an average, after second multiplication stage the maximum number of shoots per explant (5.02) was observed in (T<sub>12</sub>) MS basal medium + Activated charcoal 200 mg/l + Ascorbic acid 200 mg/l + BAP 2.5 mg/l + Kinetin 2.0 mg/l, followed by (T<sub>11</sub>) MS basal medium + Activated charcoal 200 mg/l + Ascorbic acid 200 mg/l + BAP 2.0 mg/l + Kinetin 2.5 mg/l (4.88). The combination of BAP and kinetin gave better result in shoot proliferation than individual effect of both BAP and kinetin. BAP and kinetin in combination are generally known to reduce the apical dominance and induce both axillary and adventitious shoot formation from shoot tip explants in banana system Jafari et al. [13].

**Comment [s9]:** Add more discussion

#### Length of the shoots (cm)

Results on average length of the shoots varied significantly among different treatments. The maximum shoot length (5.20 cm) was observed in (T<sub>12</sub>) MS basal medium + Activated charcoal 200 mg/l + Ascorbic acid 200 mg/l + BAP 2.5 mg/l + Kinetin 2.0 mg/l, which was on par with (T<sub>11</sub>) MS basal medium + Activated charcoal 200 mg/l + Ascorbic acid 200 mg/l + BAP 2.0 mg/l + Kinetin 2.5 mg/l (5.06 cm). Adding ascorbic acid 200 mg/l to the culture media not only helped to reduce polyphenolic oxidation but also greatly increased the number of shoots produced and shoot length. Ascorbic acid can enhance the activity of cytokinins by maintaining a favorable redox state, which may improve the signaling pathways involved in shoot induction and multiplication Smirnov [14]. Correspondingly positive outcomes were noted in the studies of Habiba et al. [15] and Sunitha [16].

#### Success rate (%)

From the study, we observed that significantly maximum success rate in micropropagation of banana cv. Ney Poovan (73.33 %) was recorded in T<sub>12</sub> where explants were treated with MS basal medium + Activated charcoal 200 mg/l + Ascorbic acid 200 mg/l + BAP 2.5 mg/l + Kinetin 2.0 mg/l. This is because of overall reduction in contamination and browning of explants observed in this treatment.

**Fig. 1. Effect of antioxidants on explant browning of banana cv. Ney Poovan**



**T<sub>12</sub> - Minimum browning**




**T<sub>19</sub> - Medium browning**




**T<sub>1</sub> - Maximum browning**

**Fig. 2. Multiple shoots produced from explant cultured on MS medium supplemented with Ascorbic acid 200 mg/l + BAP 2.5 mg/l + Kinetin 2.0 mg/l**



**Fig. 3. Shoots produced from explant cultured on MS medium supplemented without any antioxidant and cytokinin**



#### 4. CONCLUSION

This study concludes that *in vitro* culture is the best method for the propagation of banana. Through, this study it is confirmed that basal medium (MS) supplemented with activated charcoal 200mg/l + Ascorbic acid 200 mg/l + BAP 2.5 mg/l + kinetin 2.0 mg/l is best for phenolic browning reduction, multiple shoot generation and growth. Thus, this research revealed important information related to producing disease-free plantlets by *In-vitro* micropropagation of banana cultivar Ney Poovan which is in demand.

**Comment [s10]:** Add 1-2 line for future research

#### REFERENCES

1. Stover RH, Simmonds NW, London DV, De Langhe EA. Feasibility of *in vitro* propagation of bananas and plantains. Trop. Agric. 1985;62:323-328.
2. Cronauer SS, Krikorian AD. Multiplication of Musa from excised stem tips. Ann. Bot. 1984;53(3):321-328.
3. Abdelwahd RN, Hakam M, Labhilli MU, Sripad S. Use of an adsorbent and antioxidants to reduce the effects of leached phenolics in *in vitro* plantlet regeneration of faba bean. Afr. J. Biotechnol. 2008;7(8):997-1002.
4. Prabhuling G, Rashmi H, Babu AG. Protocol for tissue culture propagation of banana cv. Ney poovan (AB). Int. J. Nat. Sci. 2017;8(4): 882-887.
5. Dita M, Barquero M, Heck D, Mizubut ESG, Staver CP. Fusarium wilt of banana: current knowledge on epidemiology and research needs toward sustainable disease management. Front. Plant Sci. 2018;9:1468.
6. Murashige T, Skoog F. A revised medium for rapid growth and bioassays with tobacco tissue culture. Physiol. Plant. 1962;15:473-497.
7. Ndakideml CF, Mneney E, Ndakidemi PA. Effects of ascorbic acid in controlling lethal browning in *in vitro* culture of *Brahylaenahuillensis* using nodal segments. Am. J. Plant. Sci. 2014;5(1): 2-7.
8. Nguyen QT, Vu DT, Chu HH. Optimization of *in vitro* propagation conditions for Cavendish banana (*Musa acuminata* L.) and elimination of browning by antioxidant additives. J. Biotechnol. 2020;18(2):265-272.
9. Strosse HI, Houwe VD, Panis B. Banana cell and tissue culture, banana improvement, cellular, molecular biology and induced mutations. Annal. Bot. 2004;3(6):1-12.
10. Zaffari GR, Kerbauy GB, Kraus JE, Romano EC. Hormonal and histological studies related to *in vitro* banana bud formation. Plant Cell Tissue Org. 2000;63:187-192.
11. ShuklaS, PrasadY, Yadav AK, Dixit P, PatelB, Yadav V. Effect of different auxins and cytokinins in callus induction and shoot regeneration in banana (*Musa paradisiaca* L.) variety Udhayam under *in vitro* condition. Int. J. Chem. Stud. 2018;6: 2041-2045.

**Comment [s11]:** Set the references according to journal guide line

12. Azam FMS, Islam S, Rahmatullah M, Zaman, A. Clonal propagation of banana (*Musa* spp.) cultivar 'BARI-1' (AAA genome). *Plant Cell Tissue Org.* 2008; **87**(9): 537-544.
13. Jafari N, Othman RY, Khalid, N. Effect of benzylaminopurine (BAP) pulsing on *in vitro* shoot multiplication of *Musa acuminata* (banana) cv. Berangan. *Afr. J. Biotechnol.* 2011; **10**(13):2446-2450.
14. Smirnoff N. The function and metabolism of ascorbic acid in plants. *Annal. Bot.* 1996; **78**: 661-669.
15. Habiba U, Reja S, Saha ML, Khan MR. Endogenous bacterial contamination during *in vitro* culture of table banana: Identification and prevention. *Plant Tissue Cult.* 2002; **12**(2): 117-124.
16. Sunitha SN. Regenerative Effect of L-ascorbic acid on the *in vitro* grown plants. *Br. Biotechnol. J.* 2014; **4**(12):1238-1252.

UNDER PEER REVIEW