

ECONOMIC BENEFITS OF APPLICATION OF NPK, FERMENTED ORGANIC MANURES AND MYCORRHIZA ON YIELD AND PRODUCTION OF GREEN GRAM (*Vigna radiata* L.) var. PDM-139.

Abstract

The research study titled "Economic benefits of application of NPK, fermented organic manures and mycorrhiza on yield and production of green gram (*Vigna radiata* L.) Var. PDM-139." was conducted during the Zaid season of 2023 at the Central Research Farm, Department of Soil Science and Agricultural Chemistry, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology, and Sciences, Prayagraj, Uttar Pradesh, India. The experimental layout followed a 3x3 Randomized Block Design (RBD), incorporating three levels of fermented organic manure (0%, 50%, and 100%), Mycorrhiza (0%, 50%, and 100%), and N, P, K at the Recommended Dose of Fertilizers (RDF). The variety chosen was PDM-139, which was released by the State Variety Release Committee (SVRC) of Uttar Pradesh in 1974. The recorded average yield ranged from 6 to 9 quintals per hectare. The response of total seed yield of green gram recorded as influenced by different levels NPK, fermented organic manure and mycorrhizal inoculation. The seed yield

of green gram was found to be increased significantly with the increase in levels of NPK, fermented organic manure and mycorrhizal inoculation. The maximum seed yield was recorded as 16.42 (q ha⁻¹) T₉ (NPK@RDF + M₂@ 10Kg ha⁻¹ 50% + F₂@ 50 l ha⁻¹ 50%) which was higher than any other treatment combination and the minimum seed yield was recorded as 8.76 (q ha⁻¹) in T₁ (control). However, as for the economy of different treatments is concerned, the treatment T₉ (NPK@RDF + M₂@ 10Kg ha⁻¹ 50% + F₂@ 50 l ha⁻¹ 100%) provide highest net profit of ₹ 68139.13 ha⁻¹ with highest cost benefit ratio is 1:1.605.

Keywords: NPK, Fermented organic manure, Mycorrhizal inoculation, seed yield, Cost-Benefit ratio, etc.

Introduction

The majority of the Indian population follows a vegetarian diet, and pulses are a cost-effective and excellent source of protein, providing 20-25% protein—more than twice the amount found in cereals. India imports around 3 million tonnes of pulses, with a projected demand of 28 million tonnes by 2017 (**Anonymous, 2011**). Currently, the availability of pulses in India is 36g per person per day, which is significantly lower than the minimum and optimal requirements of 80g and 104g per person per day, respectively. Due to limited opportunities for expanding the cultivation area for pulses, the potential for increasing production seems minimal. In such challenging circumstances, intensifying cropping practices, particularly through intercropping, offers a promising alternative (**Jat et al., 2014**).

Green gram (*Vigna radiata* L.), commonly known as moong, belongs to the Leguminosae family and is cultivated in countries such as India, Burma, Sri Lanka, Pakistan, China, Fiji, Queensland, and parts of Africa. (**Smartt, 2013**). India is the largest producer of green gram, grown in almost all states, covering an area of about 3.6 million hectares with a total production of approximately 1.7 million tonnes and an average yield of 500 kg per

hectare. Key green gram-producing states include Odisha, Maharashtra, Andhra Pradesh, Madhya Pradesh, Gujarat, Rajasthan, and Bihar.

Green gram is the third most important legume crop in India and holds particular significance as a Zaid season crop in Uttar Pradesh, typically sown with the onset of summer. However, many farmers do not apply the recommended dose of fertilizers or seed treatments due to the short growing season. (Nair *et al.*, 2020)

Green gram can thrive in a variety of soil types, from black cotton soils in North India to red laterite soils in the South and sandy soils in Uttar Pradesh. For optimal production, well-drained loamy to sandy loam soils are preferred. The seeds of moong beans contain 22-28% carbohydrates, 60-65% protein, 1-1.5% fats, 3-4.5% fibre, and 4.5-5.5% ash (USDA National database, 2023). This versatile crop is used for both seeds and forage, as it produces significant biomass and can regenerate after grazing to yield more seeds. As a protein-rich food, green gram is consumed as both whole and split pulses in India. Some prominent hybrids include RUM-I, PDM-139, PDM-4, K-851, and PDM-11.

Materials and Methods

The field trial was conducted during the Zaid season of 2023 at the Central Research Farm of the Department of Soil Science and Agricultural Chemistry at Sam Higginbottom University of Agriculture, Technology and Sciences (U.P.), located at 25°24'30'' N latitude, 81°51'10'' E longitude, and 98m above mean sea level. This location is part of the Agro-Ecological Sub Region [North Alluvium Plain Zone (0-1% slope)] and the Agro-Climatic Zone (Upper Gangetic Plain Region). Agro-climatically, Prayagraj District falls within the subtropical belt of Southeast Uttar Pradesh, experiencing extremely hot summers and fairly cold winters. The maximum temperature ranges up to 46°C, rarely dropping below 4-5°C, with relative humidity between 20-94%. The area receives an average annual rainfall of about 1100mm, primarily from July to the end of September, though occasional winter precipitation is not uncommon. During the crop season,

temperatures ranged from a minimum of 21.38°C to a maximum of 37.82°C, while humidity varied from 46.42% to 96.85%

Experimental design:

The experimental design was a 3x3 Randomized Block Design with three levels of fermented organic manure (0, 50, 100%), mycorrhiza (0, 50, 100%), and NPK at recommended doses (RDF). Each plot measured 2m x 2m, and the green gram variety used was PDM-139, aimed at evaluating soil properties and health.

Table 1. Treatment combination of green gram

Treatments	Treatment Combination	Symbol
T ₁	Absolute control	L ₀ M ₀ F ₀
T ₂	NPK@RDF + M ₀ + F ₁ @ 50%	LM ₀ F ₁
T ₃	NPK@RDF + M ₀ + F ₂ @ 100%	LM ₀ F ₂
T ₄	NPK@RDF + M ₁ @ 50% + F ₀	LM ₁ F ₀
T ₅	NPK@RDF + M ₁ @ 50% + F ₁ @ 50%	LM ₁ F ₁
T ₆	NPK@RDF + M ₁ @ 50% + F ₂ @ 100%	LM ₁ F ₂
T ₇	NPK@RDF + M ₂ @ 100% + F ₀	LM ₂ F ₀
T ₈	NPK@RDF + M ₂ @ 100% + F ₁ @ 50%	LM ₂ F ₁
T ₉	NPK@RDF + M ₂ @ 100% + F ₂ @ 100%	LM ₂ F ₂

Note: RDF = Recommended dose of fertilizer, M= Mycorrhiza and F= Fermented organic manure.

Methodology:

The grain yield obtained from each plot in each harvest was sum up which gave the total grain yield per plot and thereafter with the use of multiplication factor this yield was converted into per hectare for each treatment.

Economics of treatments

As per as existing market process, the input and output costs were computed treatment wise and different economics parameters *viz.* Net profit and Cost benefit ratio were calculated.

i) Cost of cultivation (₹ ha⁻¹)

The cost of cultivation of each treatment was calculated separately taken into consideration all the cultural practices followed in the cultivation of green gram.

ii) Gross return (₹ ha⁻¹)

The gross return from each treatment was calculated taking into consideration the cost of cultivation and the market price of the produce.

iii) Net Profit (₹ ha⁻¹)

The net profit from each treatment was calculated separately by using the formula given below.

Net profit (₹ ha⁻¹) = Gross return - Cost of cultivation.

iv) Cost Benefit Ratio (C: B)

The benefit ratio for each treatment was calculated by using following formula.

Cost Benefit Ratio = Gross Return / Cost of cultivation

STATISTICAL ANALYSIS

A randomized block design (RBD) is an experimental design where the experimental units are in groups called blocks. The treatments are randomly allocated to the experimental units inside each block. When all treatments appear at least once in each block, we have a completely randomized block design. Otherwise, we have an incomplete randomized block design. This kind of design is used to minimize the effects of systematic error. If the experimenter focuses exclusively on the differences between treatments, the effects due to variations between the different blocks should be eliminated.

The data recorded during the course of investigation subjected to statistical analysis by analysis of variance (ANOVA) technique. The significant and nonsignificant of treatment effect was judged with the help of 'F' (variance ratio) table (**Fisher 1960**). The significant different between the mean were tested against the critical difference at 5% level for testing the hypothesis the following table issued.

Table 2. ANOVA table

Source of variation	d.f.	S.S.	M.S.S	F (cal.) Value	F (tab) Value on 5% LOS
Due to replication	r-1	R.S.S.	M.R.S.S.	<u>M.R.S.S.</u> E.M.S.S.	F(t-1) (r-1) (t-1) 5%
Due to treatment	t-1	Tr.S.S.	M.T.S.S.	<u>M.T.S.S.</u> E.M.S.S.	
Due to error	(r-1) (t-1)	E.S.S.	E.M.S.S.		
Total	Rt-1	T.S.S.			

C.D. = S.Em. \pm x t-test (error d. f.) at 5%

$$S. Em. \pm = \left[\sqrt{\frac{2 E. M.ss}{r}} \right]$$

Where,

r = Replication

t = Treatment

T = Total

d.f. = Degree of freedom

MSS = Mean sum of squares

TrSS = Treatment sum of squares

TSS = Total sum of squares

RSS = Replication sum of squares

ESS = Error sum of squares

MRSS = Mean replication sum of squares

MTSS = Mean treatment sum of squares

F cal. = Calculated value of F

F tab. = Tabulated value of F

C.D. = S.E (D) x 't' error d. f. at 5 % level of significance

$$S. Ed. \pm = \sqrt{\frac{2 E. M.ss}{r}}$$

The significance and non-significance of the treatment effect was judged with the help of 'F' variance ratio test. Calculated 'F' value was compared with the table value 'F' at 5 % level of significance. If calculated value exceeds the table value the effect was considered to be significant. The significant difference between the means was tested critical differences at 5 % level of significance. For testing the hypothesis, the ANOVA table was used.

Results and Discussions

The data presented in table 7. and depicted in fig. 1 clearly shows clearly shows the response of total seed yield of green gram recorded as influenced by different levels NPK, fermented organic manure and mycorrhizal inoculation. The seed yield of green gram was found to be increased significantly with the increase in levels of N P K and Rhizobium seed inoculation. The maximum seed yield was recorded as 14.43 (q ha⁻¹) T₉ (NPK@RDF + M₂@ 10Kg ha⁻¹ 50% + F₂@ 50 l ha⁻¹ 50%) which was higher than any other treatment combination and the minimum seed yield was recorded as 7.88 (q ha⁻¹) in T₁ (control).

The combined application of NPK, fermented organic manure and mycorrhizal inoculation played a very important role. Application of organic manure increased supply of major assimilated as well as micro nutrient to plants, Mycorrhiza also perform better when soil is well applied with nutrients. Similar results were also reported by **Jat *et al.*, (2012)**, **Koushal and Singh (2011)**.

Table 3. Effect of different levels of N P K, fermented organic manure and mycorrhizal inoculation on number of grain pod⁻¹, number of pod plant⁻¹ and seed yield (q ha⁻¹) of Green gram var. PDM-139.

S. No.	Treatment combination	Seed yield (q ha ⁻¹)
T ₁	Absolute control	8.76
T ₂	NPK@RDF + M ₀ + F ₁ @ 25 l ha ⁻¹ 50%	9.82
T ₃	NPK@RDF + M ₀ + F ₂ @ 50 l ha ⁻¹ 100%	10.50
T ₄	NPK@RDF + M ₁ @ 5Kg ha ⁻¹ 50% + F ₀	11.70

T ₅	NPK@RDF + M ₁ @ 5Kg ha ⁻¹ 50% + F ₁ @ 25 l ha ⁻¹ 50%	12.88
T ₆	NPK@RDF + M ₁ @ 5Kg ha ⁻¹ 50% + F ₂ @ 50 l ha ⁻¹ 100%	13.55
T ₇	NPK@RDF + M ₂ @ 10Kg ha ⁻¹ 50% + F ₀	14.52
T ₈	NPK@RDF + M ₂ @ 10Kg ha ⁻¹ 50% + F ₁ @ 25 l ha ⁻¹ 50%	15.51
T ₉	NPK@RDF + M ₂ @ 10Kg ha ⁻¹ 50% + F ₂ @ 50 l ha ⁻¹ 100%	16.42
F-test		S
S.Em. (±)		0.15
C.D.@5%		0.32

Table 4. Cost of cultivation of green gram (₹ ha^{-1})

S. No.	Particular	Units	Rates (₹ ha^{-1})	Total cost
(A) Land Preparation				
1.	Ploughing with (Mould Board plough)	4 hr	@500 hr^{-1}	2000
2.	Ploughing with (Harrow)	4 hr	@500 hr^{-1}	2000
3.	Levelling of field (Leveller)	3 hr	@500 hr^{-1}	1500
4.	Preparation of layout	4 hr	@250 hr^{-1}	1000
(B) Sowing and Irrigation Charge				
1.	Green gram seed	20Kg/ha	@110 Kg^{-1}	2200
2.	Sowing charge	10 labours	@300 labour $^{-1}$	3000
3.	Irrigation charge (3 irrigation)	8hr/irrigation \times 3	@200 hr^{-1}	4800
4.	Irrigation labour charge	7 labours	@200/day/labour	4200
5.	Cost of urea for 25 Kg nitrogen	25Kg ha^{-1}	Rs. 9 kg^{-1}	225
6.	Cost of SSP for 50 Kg phosphorus	50Kg ha^{-1}	Rs. 12 kg^{-1}	600
7.	Cost of DOP for 25 Kg potassium	25Kg ha^{-1}	Rs. 16 kg^{-1}	400
8.	Cost of Fermented organic manure for 50 lit. ha^{-1}	50 lit ha^{-1}	Rs. 400 lit $^{-1}$	20000
9.	Cost of Mycorrhiza 10 kg ha^{-1}	10Kg ha^{-1}	Rs. 100 kg^{-1}	1000
10.	Fertilizer Application	8 labour	@300 Labour $^{-1}$	2400
(C) Weeding and harvesting				
1.	2 Weeding were done	25 labour	@250/day/labour	12500
2.	Harvesting	25 labour	@ 250/labour	6250
3.	Threshing by beating	10 labour	@200/labour	2000
4.	Field rent	3 months	@4500/month	13500
5.	Supervision	3 month	@3000/month	9000
Total				₹ 86,575

Table 5. Cost of Cultivation of Green gram as per the treatment

Treatment	Particular	Units	Rates (₹ ha ⁻¹)	Cost/Unit (₹ ha ⁻¹)	Cost of Cultivation	Total cost of Cultivation
T1	Absolute control	Absolute Control	-	-	86,575	86,575.00
T2	RDF + FOM @ 50% + Mycorrhiza @0%	@ 54.35 kg UREA @ 312.50 kg SSP @ 41.67 kg MOP @ 25 lit ha ⁻¹ FOM	9:12:16 kg ⁻¹ Fermented organic manure @ 400 lit ⁻¹	489.15 3750.00 666.72 10000.00	86575.00+ 4905.87+ 10000.00	101480.87
T3	RDF + FOM @ 100% + Mycorrhiza @0%	@ 54.35 kg UREA @ 312.50 kg SSP @ 41.67 kg MOP @ 50 lit ha ⁻¹ FOM	9:12:16 kg ⁻¹ Fermented organic manure @ 400 lit ⁻¹	489.15 3750.00 666.72 20000.00	86575.00+ 4905.87+ 20000.00	111480.87
T4	RDF + FOM @ 0% + Mycorrhiza @50%	@ 54.35 kg UREA @ 312.50 kg SSP @ 41.67 kg MOP @ 0 lit ha ⁻¹ FOM @ 5 kg ha ⁻¹ Mycorrhiza	9:12:16 kg ⁻¹ Fermented organic manure @ 400 lit ⁻¹ Mycorrhiza @ 100 kg ⁻¹	489.15 3750.00 666.72 0.00 500.00	86575.00+ 4905.87+ 500.00	91980.87
T5	RDF + FOM @ 50% + Mycorrhiza @50%	@ 54.35 kg UREA @ 312.50 kg SSP @ 41.67 kg MOP @ 25 lit ha ⁻¹ FOM @ 5 kg ha ⁻¹ Mycorrhiza	9:12:16 kg ⁻¹ Fermented organic manure @ 400 lit ⁻¹ Mycorrhiza @ 100 kg	489.15 3750.00 666.72 10000.00 500.00	86575.00+ 4905.87+ 500.00+ 10000.00	101980.87
T6	RDF + FOM @ 100% + Mycorrhiza @50%	@ 54.35 kg UREA @ 312.50 kg SSP @ 41.67 kg MOP @ 50 lit ha ⁻¹ FOM @ 5 kg ha ⁻¹ Mycorrhiza	9:12:16 kg ⁻¹ Fermented organic manure @ 400 lit ⁻¹ Mycorrhiza @ 100 kg	489.15 3750.00 666.72 20000.00 500.00	86575.00+ 4905.87+ 500.00+ 20000.00	111980.87
T7	RDF + FOM @ 0% + Mycorrhiza @100%	@ 54.35 kg UREA @ 312.50 kg SSP @ 41.67 kg MOP @ 0 lit ha ⁻¹ FOM @ 10 kg ha ⁻¹ Mycorrhiza .	9:12:16 kg ⁻¹ Fermented organic manure @ 400 lit ⁻¹ Mycorrhiza @ 100 kg ¹	489.15 3750.00 666.72 00.00 1000.00	86575.00+ 4905.87+ 1000.00	92480.87

T8	RDF + FOM @ 50% + Mycorrhiza @100%	@ 54.35 kg UREA @ 312.50 kg SSP @ 41.67 kg MOP @ 25 lit ha ⁻¹ FOM @ 10 kg ha ⁻¹ Mycorrhiza .	9:12:16 kg ⁻¹ Fermented organic manure @ 400 lit ⁻¹ Mycorrhiza @ 100 kg ¹	489.15 3750.00 666.72 10000.00 1000.00	86575.00+ 4905.87+ 1000.00+ 10000.00	102480.8 7
T9	RDF + FOM @ 100% + Mycorrhiza @100%	@ 54.35 kg UREA @ 312.50 kg SSP @ 41.67 kg MOP @ 50 lit ha ⁻¹ FOM @ 10 kg ha ⁻¹ Mycorrhiza .	9:12:16 kg ⁻¹ Fermented organic manure @ 400 lit ⁻¹ Mycorrhiza @ 100 kg ¹	489.15 3750.00 666.72 20000.00 1000.00	86575.00+ 4905.87+ 1000.00+ 20000.00	112480.8 7

Table 6. Effect of Different Cost benefit ratio (C:B) of different treatment combination with Green gram.

Treatment	Yield (q ha ⁻¹)	Selling price (₹ q ⁻¹)	Gross return (₹ ha ⁻¹)	Total cost of cultivation (₹ ha ⁻¹)	Net Profit (₹ ha ⁻¹)	Cost Benefit ratio (C:B)
T ₁	8.76	11000	96360	86,575.00	9785.00	1.1130
T ₂	9.82	11000	108020	101480.87	6539.13	1.0644
T ₃	10.50	11000	115500	111480.87	4019.13	1.0360
T ₄	11.70	11000	128700	91980.87	36719.13	1.3992
T ₅	12.88	11000	141680	101980.87	39699.13	1.3892
T ₆	13.55	11000	149050	111980.87	37069.13	1.3310
T ₇	14.52	11000	159720	92480.87	67239.13	1.7270
T ₈	15.51	11000	170610	102480.87	68129.13	1.6647
T ₉	16.42	11000	180620	112480.87	68139.13	1.6057

Conclusion

It is revealed from trial that the various level of NPK fermented organic manure and mycorrhizal inoculation used from in the experiment, the treatment combination T₉ - (NPK@RDF + M₂@ 10Kg ha⁻¹ 50% + F₂@ 50 l ha⁻¹ 100%) was found to be the best treatment gave highest gross return ₹ 180620.00 benefit of ₹ 67139.13 with highest cost benefit ratio 1:1.605 for green gram it could be recommended for profitable production of Green gram (*Vigna radiata* L.) var. PDM-139 and treatment T₉ is best in terms of soil physical and chemical properties. Further it increased soil fertility and nutrient uptake by the plant. Increasing in the level of NPK and Fermented organic manure there was a significant increase in yield attributes. The maximum yield was found in T₉ (NPK@RDF + M₂@ 10Kg ha⁻¹ 50% + F₂@ 50 l ha⁻¹ 100%) and the minimum yield was recoded in T₁ (control). However, as for the economy of different treatments is concerned, the treatment T₉ (NPK@RDF + M₂@ 10Kg ha⁻¹ 50% + F₂@ 50 l ha⁻¹ 100%) provide highest net profit of ₹ 68139.13 ha⁻¹ with highest cost benefit ratio is 1:1.605.

References

- Arumugam, T., Rajsekharan, S. and Nagarajan, S. M. (2011)** Response of arbuscular mycorrhizal fungi and Rhizobium inoculation on growth chlorophyll content of *Vigna unguiculata* L. walp var Pusa 151. *Journal of Applied Sciences and Environmental management* 14.
- Datta, S.K. and Tambat, H. (2022)** The role of N-Carb, a fermented organic manure in reclamation of soil having low organic carbon, *International Journal of Scientific Research Publication*, Volume 12,3.
- Imran, R.A., Shahib, Y.J. and Khyuon, A. (2020)** use of three types of fermented and non-fermented organic fertilizers in the growth and production of lettuce (*Lactuca sativa* L.). *International Journal of Stat. Sci.*, 16(1); 1681-1688
- Jat, P. C., Rathore, S. S. and Sharma, R. K. (2014)** Effect of integrated nitrogen management and intercropping systems on yield attributes and yield of maize. *Indian Journal of Hill Farming*, 27 (1): 91-99.

- Kumar, R., & Kushwaha, S. (2018).** Management of pest and diseases in green gram (*Vigna radiata*). *Journal of Pulses Research*, 41(2), 129-133.
- Nair, R., & Schreinemachers, P. (2020).** Sustainable production and consumption of pulses: Case of mungbean. *Field Crops Research*, 254, 107847.
- Natasha L. Hoover, Ji Yeow Law, Leigh Ann M. Long, Ramesh S. Kanwar and Michelle L. Soupir, (2019)** Long-term impact of poultry manure on crop yield, soil and water quality, and crop revenue. *Journal of Environmental Management*, 252: 109582
- Olsen, S.R., Watanale, F. S., (1954)** A method to determine a phosphorus adsorption measured by colorimetric method *soil sci.soc. Am.J.21,144-149*
- Owla. R.L. Chavan, B.N. and Singh, U. (2007)** Effect of plant densities and phosphorus levels on growth and yield of green gram (*Vigna radiata*) *Annals of Agricultural Research* 28(1): 95-97
- Rathore, G., Kaushal, R., Sharma, V., Sharma, G., Chaudhary, S. and Dhaliwal, S.S. (2023)** Evaluation of the Usefulness of Fermented Liquid Organic Formulations and Manures for Improving the Soil Fertility and Productivity *Agriculture* 2023, 13, 417.
- Ray, J.G., and Valsalakumar, N. (2010).** Arbuscular Mycorrhizal fungi and Piriformospora indica individually and in combination with rhizobium on green gram. *J. Plant Nutrition*, 33(2): 285-298.
- Sharma, K. L., Srinivas, K., Mandal, U. K., Vittal, K. P. R., Grace, K. J. and Maruthi, S.G.R.(2014)** Integrated Nutrient Management HT Strategies for Sorghum and Green gram in Semi-Arid Tropical Alfisol. *Indian J. Dryland Agric. Res. & Dev.* 2017 19(1), 13-23.
- Shraddha, A., Shukla, Y.R., Thakur, K. and Kumari, J. (2023)** Impact of Fermented Organic formulations combined with inorganic fertilizers on broccoli (*Brassica oleraceae* var. italica) cv. Palam Samridhhi. *Science Direct Heliyon*,9(2); 2031.
- Singh, R.P., Bisen. J.S., Yadav, P.K., Singh, S.N., Singh, R. K. and Singh, J. (2008)** Integrated use of sulphur and molybdenum on growth, yield and quality of black gram. *Legume Research* 31:214-217
- Smartt, J. (2013).** *Tropical Pulses* (2nd ed.). Cambridge University Press.
- USDA National Nutrient Database (2023).** Nutritional composition of mung beans.