

# Antiviral Potential of Some Medicinal Plants against Newcastle Disease Virus

---

## ABSTRACT

**Aims:** Newcastle disease virus (NDV) is a highly contagious paramyxovirus that infects poultry and can lead to high mortality rates. It can cause severe economic losses to farmers. As there is no effective treatment for NDV infection, this research focused on evaluating the possibility of using medicinal plant extracts against this virus.

**Place and Duration of Study:** The experiment was conducted at the Department of Physiology, Biochemistry, and Pharmacology and the Department of Microbiology and Veterinary Public Health, Chittagong Veterinary and Animal Sciences University (CVASU), Bangladesh, during a period from January to April 2017.

**Methodology:** In this study, aqueous extracts were prepared from neem bark, neem leaf, garlic, and ginger following standard procedures. Those were collected from the medicinal garden of CVASU. Then, the extracts were categorized into different groups to determine the possible effects of extract concentration and heat treatment on the antiviral potential of plant extracts. The plant extract was mixed with viable NDV (8 HA; field strain) and kept at 27.3 °C for 30 minutes. To determine virus inactivation, a 0.2-ml mixture was inoculated into nine-day-old embryonated chicken eggs and incubated. After 48 hours, the allantoic fluid was harvested, and a hemagglutination (HA) assay was performed to determine the virus HA titer.

**Results:** During this study, all plant extracts were found effective against the Newcastle disease virus. The antiviral activities of these extracts can vary due to the concentration and autoclaving of the extract.

**Conclusion:** To sum up, these plants can be a promising source for developing antiviral drugs against Newcastle disease.

*Keywords:* Newcastle disease, Anti-viral medicine, Medicinal plants, Aqueous extracts, Hemagglutination test.

## 1. INTRODUCTION

Newcastle disease virus (NDV) is a negative-sense single-stranded RNA virus under the family Paramyxoviridae (Mayo, 2002). It causes a severe disease in poultry named Newcastle disease or Ranikhet. This disease remains a serious economic challenge to all segments of the poultry industry because of its contagious and mortality (0–100%) records (Chollom et al., 2012; Sulaiman et al., 2013), although mortality varies depending on the pathotype of the virus. Medicinal plants are a rich source of natural compounds such as tannins, polyphenols, proanthocyanidins, sulfonamides, anti-adhesives, etc. that exhibit antiviral (Mukhtar et al., 2008; Gupta et al., 2014) and anti-inflammatory (Gupta et al., 2014) activities. Many reports showed that many indigenous communities used their herbal preparations for veterinary use (Madadgar et al., 2013). Like many other medicinal plants, Neem (*Azadirachta indica*) has been used in Ayurvedic medicine for more than 2000 years,

and now it is being used in modern medicine, cosmetics, and pharmaceuticals as the global scenario is changing towards the use of nontoxic plant products. Its medicinal values come from the fruits, seeds, leaves, roots, and bark (Biswas et al., 2002). Various preparations of neem obtained from its different parts have been found to exert anti-bacterial, anti-viral, anti-malarial, anti-oxidant, anti-fungal, anti-mutagenic, anti-carcinogenic, contraceptive, and antiulcer activity (Subapryya and Nagini, 2005). Garlic has been an interesting plant for centuries as a medicinal panacea. A broad range of pathogenic organisms, including bacteria, fungi, protozoa, and viruses, be sensitive to fresh, crushed garlic (Mehrbood et al., 2009). *Zingiber officinale* (family Zingiberaceae), commonly known as ginger, is commonly used as an effective medicine against coughs and colds. It was also found effective against Newcastle disease virus during an *in vitro* experiment (Kikuzaki et al., 1991; Sharma and Gupta, 1998). However, since there is a lack of sufficient information about the antiviral potential (and factors affecting this potential) of neem leaves, neem bark, garlic, and ginger against Newcastle disease virus, this study focused on finding out the effects of concentration and heat treatment (during autoclaving) on the anti-viral activity of these plant extracts. These plants had been used as traditional medicines by the natives for many years due to their antibacterial, antifungal, anti-allergic, anti-viral, and other important medicinal properties.

## **2. MATERIAL AND METHODS**

### **2.1 Sample collection**

The experiment was conducted at the Department of Physiology, Biochemistry, and Pharmacology and the Department of Microbiology and Veterinary Public Health, Chittagong Veterinary and Animal Sciences University (CVASU), Bangladesh, during a period from January to April 2017.

### **2.2 Virus sample**

A previously isolated and stored (-80 °C) field strain of the Newcastle disease virus was collected from a repository of the Department of Microbiology and Veterinary Public Health, CVASU. After thawing, the virus sample was treated with an antibiotic (Gentamicin) to prevent bacterial contamination. Then the virus samples were subjected to a hemagglutination (HA) test to confirm the viability and concentration of the virus.

### **2.3 Plant samples**

The bark and leaf of Neem (*Azadirachta indica*), Garlic (*Allium sativum*) and Ginger (*Zingiber officinale*) from the medicinal plant garden of Shahedul Alam Qudary Teaching Veterinary Hospital, CVASU, Bangladesh.

### **2.4 Plant extract preparation**

Collected neem leaves, garlic, and ginger were thoroughly cleaned with water to remove dirt and unwanted materials. Then 300-gram (g) samples were ground using an electrical grinder. The grinded samples (300 g) were mixed with distilled water at the ratio of 1:5 (sample: water) for neem leaf, garlic, and ginger (Iwalokun et al., 2004; Kwawukume et al., 2013; Gupta and Chaphalkar, 2015). In the case of neem bark, the collected barks were cut into small pieces and dried at room temperature (27°C) for several weeks. Then dried barks were pulverized using an electric grinder. Finally, 300 g of powder was mixed with distilled water in a 1:10 ratio (Mahmood et al., 2017). Sample mixtures were stirred for 1 hour by an electric stirrer (1000 rpm) and left overnight. All the mixtures were filtrated through Whatmanno. 1 filter paper separately. Then 300 ml of filtrates from different samples were taken into a round bottom flask of a rotator vacuum evaporator and condensed by the evaporation of solvent from the filtrate in a water bath at 56.7°C for 3-5 hours. After the

evaporation of solvent from the filtrate, the condensed extracts were preserved in a tightly corked-labeled bottle and stored at 4 °C.

### **2.5 Categorization of plant extracts**

Each plant extract was broadly categorized into two groups: Group-I: extract without heat treatment, and Group-II: extract treated with heat at 121°C for 15 min. Then every group was further divided into three sub-groups by mixing PBS with concentrated plant extract, such as sub-group I: 100% extract (concentrated extract); sub-group II: 75% extract (three parts extract and one part PBS); and sub-group III: 50% extract (one part extract and one part PBS).

### **2.6 Toxicity test of samples**

All types of plant extract were checked for any possible toxicity to chicken embryos by an embryonated egg inoculation assay. 0.2 ml of aqueous plant extract was inoculated in nine-day-old embryonated chicken eggs (5 eggs per concentration) collected from the Regional Poultry Farm (RPF), Chittagong. Then all eggs were incubated for 48 hours to check for embryo mortality.

### **2.7 Preparation of 8 HA unit virus**

A virus sample (0.2 ml) was propagated in embryonated chicken eggs. After propagation, allantoic fluid, collected after 48 hours of incubation of eggs, was subjected to an HA test to determine 1 HA unit. Based on the Hemagglutination (HA) test result, an 8 HA unit virus concentration was prepared by mixing 15 ml of sterile PBS with 1 ml of allantoic fluid.

### **2.8 Embryonated eggs inoculation**

Inoculums were prepared by properly mixing the plant extracts from all sub-groups with 8 HA units of virus at a 1:1 (500 µl extract with 500 µl virus) ratio in an eppendorf tube and were incubated at room temperature for 30 minutes. Five nine-day-old embryonated chicken eggs per concentration were inoculated with 0.2 ml of inoculum following standard procedure. The inoculated eggs were incubated for 48 hours. The eggs were candled after 24 hours of inoculation to check embryo mortality. After 48 hours of incubation, eggs were transferred to a chilling temperature and kept for 24 hours. About 10% control for both viruses and extracts were maintained during the whole procedure. After 24 hours of chilling, allantoic fluids were harvested from each egg using a sterile syringe. The fluid was taken into the falcon tube. Then the HA test was carried out using 1% chicken RBC, collected from a specific pathogen-free flock, to determine the virus titer in allantoic fluid.

### **2.9 Statistical analysis**

All data were entered into a spread sheet of MS Excel-2010. The data were sorted, cleaned and coded using the excel program before exporting for the analysis of Geometric mean titer (GMT). Finally, the GMT of HA were used to determine reduced HA titer (%) and the effect of two key factors-concentration and heat treatment, on antiviral potential of plant extract.

### 3. RESULTS

#### 3.1 Toxicity of Extract

In the toxicity testing assay, no plant extract caused mortality in chicken embryos within 48 hours of incubation (Table 1).

**Table 1: Determination of toxicity of extracts in embryonated eggs**

Plant extracts	Concentration (%)	Total no of eggs inoculated	Death observed in embryonated eggs			
			24 hours	48 hours	24 hours	48 hours
			without heat-treated plant extracts used		heat treated Plant extracts used	
Neem Leaf	100	20	A (0/5)	A(0/5)	A (0/5)	A(0/5)
	50	20	A (0/5)	A (0/5)	A (0/5)	A (0/5)
	33	20	A (0/5)	A (0/5)	A (0/5)	A (0/5)
Neem Bark	100	20	A (0/5)	A (0/5)	A (0/5)	A (0/5)
	50	20	A (0/5)	A (0/5)	A (0/5)	A (0/5)
	33	20	A (0/5)	A (0/5)	A (0/5)	A (0/5)
Garlic	100	20	A (0/5)	A (0/5)	A (0/5)	A (0/5)
	50	20	A (0/5)	A (0/5)	A (0/5)	A (0/5)
	33	20	A (0/5)	A (0/5)	A (0/5)	A (0/5)
Ginger	100	20	A (0/5)	A (0/5)	A (0/5)	A (0/5)
	50	20	A (0/5)	A (0/5)	A (0/5)	A (0/5)
	33	20	A (0/5)	A (0/5)	A (0/5)	A (0/5)

A= Indicates all alive eggs after 48 hr inoculation of extract

#### 3.2 Overall antiviral potential of extracts

The anti-viral effect (reduced HA) of plant extract was found to be variable when compared with the virus control group (Table-2). For the NDV field strain, the geometric mean of HA titer (GMT) of the virus control group was 128 HA units. 50% 33% of aqueous extracts of neem bark from group-I (without autoclave) exhibited 0 HA titer (GMT), which indicates complete (100%) inactivation of the virus in the embryonated chicken egg assay. Group II (using an autoclave) had GMT values of 16, 64, and 84, which correspond to decreases in viral titer of 87.5%, 50%, and 34.3% at 100%, 50%, and 33% concentrations, respectively. On the other hand, ginger extract was found to be the least effective medicinal plant against the ND virus. It showed high GMT values (101.59, 111.43, and 111.43 for group-I, and 97, 111.43, and 128 for group-II) that reflect the weak anti-viral potential (20.6%, 12.9%, and 12.9% GMT reduction by sub-groups of groups-I; 24.21%, 12.9%, and 0% GMT reduction by sub-groups of group-II) of ginger against NDV. All neem leaf extract sub-groups of group I showed similar anti-viral activity (GMT 64 and 50% GMT reduction) despite being different in concentration, although sub-groups of group II showed gradual decreased anti-viral activity due to dilution. The GMT values were 64, 84, and 111.43, reflecting 50%, 34.3%, and 12.9% virus titer reduction, respectively, for three sub-groups of neem leaf extract. Garlic extracts are good enough to reduce ND virus titers if they are not autoclaved. During this study, non-autoclaved extracts reduced 58% to 71.2% of the virus titers, whereas autoclaved extracts reduced 0% to 43.3% only.

**Table 2: HA activity of Plant extract against NDV in embryonated eggs**

Concentration	Plants Extract	No of eggs (n)	HA titer of virus treated with extracts Geometric mean-titer (GMT)		Control GMT	Reduced HA (%) =(*C-*E)/*C	
			Without heat treated	Heat treated		Without heat treated	Heat treated
100% (1:1)	Neem leaf	10	64	64	128	50	50
	Neem Bark	10	0	16		100	87.5
	Garlic	10	36.75	84		71.2	34.3
	Ginger	10	101.59	97		20.6	24.21
50% (1:2)	Neem leaf	10	64	84		50	34.3
	Neem Bark	10	0	64		100	50
	Garlic	10	45.25	97		64.6	24.2
	Ginger	10	111.43	111.43		12.9	12.9
33% (1:3)	Neem leaf	10	64	111.43		50	12.9
	Neem Bark	10	0	84		100	34.3
	Garlic	10	53.81	128		58	0
	Ginger	10	111.43	128		12.9	0


\*C= Haemagglutination titer of virus control group,

\*E = Haemagglutination titer of extracts treated embryonated group eggs

However, the antiviral activity of garlic also varied due to dilution (GMT 36.5, 45.25 and 53.81 in group-I; 84, 97 and 128 in group-II).


### 3.3 Effect of concentration and heat treatment on the antiviral potential of plant extract

There was a favorable correlation found between plant extract concentration and antiviral activity.




**Figure 1:** Regression lines reflecting effects of concentration of plant extracts on their anti-viral potential

The antiviral potential of all plant extracts decreases if the concentration of extract decreases. The regression analysis shows that concentration adversely affects the antiviral activity of neem bark ( $R^2=0.998$ ; slope=0.39), Neem leaf ( $R^2=0.872$ ; slope=0.25), Garlic ( $R^2=0.813$ ; slope=0.31) and Ginger ( $R^2=0.972$ ; slope=0.23) extracts (Figure 1). On the other hand, heat treatment has a negative effect on antiviral potential of plant extract (Figure 2). Due to autoclaving, the antiviral properties of plants decrease significantly ( $p=0.02$ ).



**Figure 2:** Regression line and box plot reflecting the effect of heat treatment (during autoclaving) on the antiviral activity of studied medicinal plant extracts

However, the effect of concentration on the virus titer reduction ability of plant extract is not modified by heat treatment (Figure 3).



**Figure 3:** Combined effect of concentration and heat treatment on anti-viral activity of studied medicinal plants

#### 4. DISCUSSION

Medicinal plants are the ultimate source for treating living organisms in humans and animals suffering from infectious and non-infectious diseases (Arunkumar and Muthuselvan, 2009). In this study, four aqueous extracts of three plant species neem (*Azadirachta indica*), garlic (*Allium sativum*), and ginger (*Zingiber officinale*) were prepared and tested against NDV,

which causes economic losses to the poultry industry across the world. In this study, the antiviral impact of plant extract is described in terms of HA titers, especially the geometric mean titer (GMT), the lower the GMT titer value, the stronger the antiviral effect of plant extract which was also described in another study of Sulaiman et al., 2011; Ong et al., 2014; Mahmood et al., 2017. In this research, all the studied plants were found to be non-toxic to the chicken embryo, and their antiviral activities against NDV can vary due to differences in species, concentration, and heat treatment. These findings provide solid proof of previous research findings (Keqiang and van Bruggen, 2001; Mahmood et al., 2017). Neem is comparatively more effective against NDV in comparison to other plant extracts; this may be due to the difference in their phytochemical properties (Song et al., 1997; Parida et al., 2002). Without autoclaving, the use of neem extracts (both leaf and bark) shows almost similar effectiveness against NDV, irrespective of extract concentration. But after autoclaving, the antiviral activities of neem extract decrease with the decrease in extract concentration (Sulaiman et al., 2011; Ong et al., 2014; Mahmood et al., 2017). This phenomenon indicates the negative effect of heat treatment on the antiviral potential of biochemical compounds (such as nimbidin, sodium nimbidate, epicatechin, catechin, etc.) that trigger antiviral activity (Biswas et al., 2002). Despite being affected by heat and concentration, the neem plant can be an incredible source of anti-viral therapeutics against NDV, as it also plays an important role in strengthening the immune system and inactivating viruses effectively (Sadekar et al., 1998; Awolu et al., 2013). Garlic is a rich source of allicin and ajoene (Ankri and Mirel an, 1999). Previously, garlic was found effective against infectious bronchitis virus during propagation in embryonated chicken eggs (Shojai et al., 2016). Besides, it can significantly increase the antibody titer against the ND and avian influenza viruses in poultry (Eid and Iraqi, 2014). This study found that garlic can also inactivate NDV, however, the extract's concentration and heat treatment have a major impact on its antiviral activity. Ginger, a source of different phenolic derivatives such as gingerol, paradole, bisabolene, zingerone, etc., has been used as an anti-viral agent in humans and animals for many years. Gupta and Chalkar (2015), described the anti-inflammatory and anti-viral potential, against NDV, of garlic. During this research, garlic was found to be able to inactivate NDV, but to a narrow extent. However, interestingly, the anti-viral activity of undiluted extract may be increased due to heat treatment. Due to the presence of gingerol, paradole, bisabolene, zingerone, zingiberol, and 6-shogaol, ginger (*Zingiber officinale*) can act as an anti-inflammatory and anti-viral agent. In previous research, ginger extracts were found effective against NDV (Mishra et al., 2012; Gupta and Chaphalkar, 2015), but during this study, ginger showed a lower antiviral effect against NDV, which may be due to the species differences of ginger and/or strain differences of virus used in these studies. Furthermore, the antiviral activity of ginger is conversely related to the dilution and autoclaving of the extract. However, as this study focused only on revealing the possibility of using local medicinal plants to treat ND in poultry, further detail is needed to identify the active ingredients of these plants and their possible effects on live birds.

## 5. CONCLUSION

The aqueous extracts of all studied plants are effective against Newcastle disease virus of poultry; although neems-bark extract is likely the most effective one among all four types of extracts. Heat treatment can affect the antiviral potential of medicinal plant extract which is worsen in case of garlic extract. Besides, concentration of extract also has a positive relationship with antiviral activities of studied plants. Although present study has not studied details, especially active compound, of studied plants, yet, it unveils the possibility of using common local medicinal plants to treat Newcastle disease in poultry.

## ETHICAL APPROVAL

Ethical Approval (CVASU/Dir(R&E) EC/2022/435(1)/7) was taken from Ethical Approval committee of Director of research of Extension, Chittagong Veterinary & Animal Sciences University.

## REFERENCES

1. Ankri, S., Mirelman, D. (1999). Antimicrobial properties of allicin from garlic. *Microbes and infection*1, 125-129.
2. Arunkumar, S., Muthuselvam, M. (2009). Analysis of phytochemical constituents and antimicrobial activities of Aloe vera L. against clinical pathogens. *World Journal of Agricultural Sciences*5, 572-576.
3. Awolu, O., Obafaye, R., Ayodele, B. (2013). Optimization of solvent extraction of oil from neem (*Azadirachta indica*) and its characterizations. *reactions*13, 68.
4. Biswas, K., Chattopadhyay, I., Banerjee, R.K. and Bandyopadhyay, U. (2002). Biological activities and medicinal properties of neem (*Azadirachta indica*). *Current science*, pp.1336-1345.
5. Chollom, S., Olawuyi, A., Danjuma, L., Nanbol, L., Makinde, I., Hashimu, G., Alesa, M., Esilonu, J., Ogundej, E., Kwatjel, J. (2012). Antiviral potential of aqueous extracts of some parts of *Momordica balsamina* plant against newcastle disease virus. *Journal of Advanced Pharmacy Education & Research* 2, 82-92.
6. Eid, K., Iraqi, M. (2014). Effect of garlic powder on growth performance and immune response for Newcastle and avian influenza virus diseases in broiler of chickens. *Animal Biotechnology (Poultry and Fish)*7, 7.
7. Gupta, A., Chaphalkar, S.R. (2015). Immunopharmacological activity of *Zingiber officinale* on human peripheral blood mononuclear cells. *Asian J Med Pharmac Res*5, 13-17.
8. Gupta, A., Khamkar, P., Chaphalkar, S. (2014). Applications and uses of active ingredients from medicinal plants. *Ind J Nov Drug Deli*6, 106-111.
9. Iwalokun, B., Ogunledun, A., Ogbolu, D., Bamiro, S., Jimi-Omojola, J. (2004). In vitro antimicrobial properties of aqueous garlic extract against multidrug-resistant bacteria and *Candida* species from Nigeria. *Journal of medicinal food*, 327-333.
10. Keqiang, C., van Bruggen, A.H. (2001). Inhibitory efficacy of several plant extracts and plant products on *Phytophthora infestans*. *Hebei Nongye Daxue Xuebao (China)*.
11. Kikuzaki, H., Usuguchi, J., Nakatani, N. (1991). Constituents of Zingiberaceae. I. diarylheptanoids from the rhizomes of ginger (*Zingiber officinale* Roscoe). *Chemical and Pharmaceutical Bulletin*39, 120-122.
12. Madadgar, O., Karimi, V., Nazaktabar, A., Kazemimanesh, M., Ghafari, M., Azimi Dezfouli, S., Hojjati, P. (2013). A study of Newcastle disease virus obtained from exotic caged birds in Tehran between 2009 and 2010. *Avian Pathology*42, 27-31.
13. Mahmood, M.S., Amir, H.W., Abbas, R.Z., Aslam, B., Rafique, A. (2017). Evaluation of Antiviral Activity of *Azadirachta indica* (Neem) Bark Extract against Newcastle Disease Virus. *Pakistan Veterinary Journal*, 25-28
14. Mahmoud, A. (2007). The antiviral effect of acacia *Nilotica* Extracts against the Mesogenic Strain of Newcastle Disease Virus (Doctoral dissertation, University of Khartoum).
15. Mayo, M. (2002). A summary of taxonomic changes recently approved by ICTV. *Archives of virology*147, 1655-1656.
16. Mehrbod, P., Amini, E., Tavasiti, K.M. (2009). Antiviral activity of Garlic Extract on influenza virus. *Iranian Journal of Virology* 3(1), pp.19-23.

17. Mishra, R.K., Kumar, A., Kumar, A. (2012). Pharmacological activity of *Zingiber officinale*. *International Journal of Pharmaceutical and Chemical Sciences*1, 1073-1078.
18. Mukhtar, M., Arshad, M., Ahmad, M., Pomerantz, R.J., Wigdahl, B., Parveen, Z. (2008). Antiviral potentials of medicinal plants. *Virus research*131, 111-120.
19. Ong, G., Faizul, F., Ramlan, M., Maizatul, Z., AH, M.H., Syamsiah, A., Chandrawathani, P., Leow, B., Zunaida, B. (2014). Antiviral effect of aqueous neem extract from branches of neem tree on Newcastle Disease virus.
20. Parida, M., Upadhyay, C., Pandya, G., Jana, A. (2002). Inhibitory potential of neem (*Azadirachta indica* Juss) leaves on dengue virus type-2 replication. *Journal of ethnopharmacology*79, 273-278.
21. Sadekar, R., Kolte, A., Barmase, B., Desai, V. (1998). Immunopotentiating effects of *Azadirachta indica* (Neem) dry leaves powder in broilers, naturally infected with IBD virus. *Indian journal of experimental biology*36, 1151-1153.
22. Sharma, G., Reddy, Y. (2002). Neem (*Azadirachta indica*) and its Medicinal value in livestock production. *Ind. J4*, 56-68.
23. Shojai, T.M., Langeroudi, A.G., Karimi, V., Barin, A., Sadri, N., 2016. The effect of *Allium sativum* (Garlic) extract on infectious bronchitis virus in specific pathogen free embryonic egg. *Avicenna journal of phytomedicine*6, 458.
24. Song, S., Song, J., Chang, M., Lee, J.S., Choi, Y. (1997). Identification of one of the major viruses infecting garlic plants, garlic virus X. *Molecules and cells*7, 705-709.
25. Subapriya, R., Nagini, S. (2005). Medicinal properties of neem leaves: a review. *Current Medicinal Chemistry-Anti-Cancer Agents*5, 149-156.
26. Sulaiman, L.K., Oladele, O.A., Shittu, I.A., Emikpe, B.O., Oladokun, A.T., Meseko, C.A.
27. (2011). In-ovo evaluation of the antiviral activity of methanolic root-bark extract of the African Baobab (*Adansonia digitata* Lin). *African Journal of Biotechnology*10, 4256-4258.
28. Sulaiman, L.K., Oladele, O.A., Shittu, I.A., Emikpe, B.O., Oladokun, A.T., Meseko, C.A. (2013). In-ovo evaluation of the antiviral activity of methanolic root-bark extract of the African Baobab (*Adansonia digitata* Lin). *African Journal of Biotechnology*10, 4256-4258.