

## Original Research Article

Effects of compost based on mangifera indica biomass combined with popularized mineral manure on soil chemical parameters and yield of maize (*zea mays* L.) in western Burkina Faso.

How about this topic

Comment [WU1]: State and explain how Mango biomass was formed in material and method

Comment [WU2]: Zea

Comment [WU3]: Effects of compost biomass using Mango (*Mangifera indica*) combined with popularized mineral manure on soil Chemical parameters on the yield of maize (*zea mays* L.) in western Burkina Faso

## ABSTRACT

**Aims :** To evaluate the effects of mango leaf compost and mineral manure on soil chemistry and maize yields.

**Study Design:** The present study was carried out in the context of the valorization of compost from *Mangifera indica* biomass and its combination with mineral manure, with a view to improving the chemical properties and productivity of maize in western Burkina Faso.

**Place and Duration of Study:** The study was conducted in Burkina Faso, for the Institute of Environment and Agricultural Research, Farako-Bâ station for twelve months.

**Methodology:** A Fisher block design was used, comprising four (4) replicates with five (5) treatments each. A single factor was studied, namely organo-mineral fertilization at five (5) levels of variation. T1: NPK (200 kg/ha) + Urea (150 kg/ha): GMF; T2: Compost (2.5 t/ha) + GMF; T3: Compost (5 t/ha) + GMF; T4: Compost (7.5 t/ha) + GMF and T5: Compost (10 t/ha) + GMF. Soil chemical and agronomic parameters of the maize were observed during the study. **Results:** The results showed that the different treatments did not totally positively influence soil chemical parameters and maize plant growth. However, the best results in terms of soil chemical parameters were obtained for available assimilable phosphorus and potassium before planting, with respective averages of (6.65 mg/kg and 72.14 mg/kg) and after planting by T2 (5.66 mg/kg) and T5 (58.13 mg/kg) respectively. CEC was high in treatment T3 with 3.86 Cmolc.kg<sup>-1</sup> and saturation rate (V) was high in treatment T5 with 3.86 Cmolc.kg<sup>-1</sup> And the saturation rate (V) was high in the T5 treatment at 59.76%. Plant growth was high, with the height at 45 days after planting by T5 (91.14 cm) The manures applied had a significant impact on maize yields. The highest biomass and grain yields came from the T5 treatment (10 t/ha compost + GMF), with 5.8 t/ha and 3.7 t/ha respectively.

**Conclusion:** This study suggests the use of 10 t/ha of compost + GMF to improve maize productivity in the western zone of Burkina Faso. It would be interesting to continue the study

in other agro-ecological zones of Burkina Faso in order to assess the impact of these fertilizers on the biological properties of the soil.

**Keywords:** Burkina Faso, *Mangifera indica* biomass, fertilizer, compost, soil, yield.

Comment [WU4]: Not important

## 1. INTRODUCTION

In Burkina Faso, the agricultural sector remains one of the levers of the Burkinabe economy. It is the basis of the food and nutrition security of the population and contributes to more than 60% of the income of agricultural households [1]. Maize is the most widely grown major food crop in sub-Saharan Africa [2]. It occupies more than 33 million hectares each year [2]. In Burkina Faso, even though maize is grown in all regions of the country, its production varies from one region to another [3]. In terms of production, maize is in 2nd place with an estimated production quantity of 143,085 tons and constitutes 20.6% of the land cultivated under cereals with a yield of 1920 kg/ha in 2015 [4]. Indeed, high-yielding varieties reaching 5-6 tons/ha have been developed by research. Indeed, high-yielding varieties reaching 5-6 tons/ha have been developed by research. Despite this high yield potential, maize cultivation faces low productivity, due to physical and socio-economic problems that hinder production. One of these major problems is undoubtedly soil degradation, which has a direct impact on production. The causes of this increased soil degradation include: the nature of the soil; climatic hazards determined by aggressive rainfall, severe winds and high temperatures. Faced with this major constraint, especially for a crop with high nutritional needs such as maize, would the addition of organic manure to mineral fertiliser improve maize productivity? Thus, the addition of these fertilizers has been popularized to producers with a view to improving soil productivity. However, it can be noted that for several reasons, this technology is insufficiently applied by producers. The adoption rate varies from one agricultural locality to another depending on the degree of supervision of producers, which is itself linked to the practice of cash crops, particularly cotton cultivation on farms [5]. The general objective of this study is to contribute to the maintenance and improvement of soil fertility as well as maize productivity with compost based on *Mangifera indica* biomass.

Comment [WU5]: In Burkina Faso, the agricultural sector remains one of the levers of the Burkinabe economy. It is the basis of the food and nutrition security of the population and contributes to more than 60% of the income of agricultural households.

Comment [WU6]: Morden researchers to increased productivity in the country.

Comment [WU7]: tivity

Comment [WU8]: The nature of the soil and climatic condition (aggressive rainfall, severe winds and high temperatures) which are the causes of soil degradation. Faced with this major constraint, maize crop with high nutritional needs would require additional organic manure mineral to fertilizer and improve, thus increase soil productivity.

## 2. MATERIAL ET METHODS

### 2.1. Presentation of the study site

Our study was conducted at the INERA Farako-Bâ agricultural research station, located on the Bobo-Banfora axis. According to [6] the geographical coordinates are 11°06' north

latitude and 4°20' west longitude. Its altitude above the sea is 405 meters. The Farako-Bâ station has vegetation that is characterized by a grassy to wooded savannah that is quite dense in places [7]. The soils of Farako-Bâ are mostly of the tropical ferruginous type [8]. The climate of the area is South Sudanian. In the last ten (10) years, 2018 was the rainiest year with 1303.8 mm in 70.

## 2.2. Material

The plant species used in the experiment is maize (*Zea mays* L.), Komsaya variety. It is a hybrid variety selected by INERA with a sowing-maturity cycle of 97 days and a potential yield of between 8 and 9.5 tonnes per hectare. The fertilizing material consisted of two types of manure (organic and mineral). These fertilizers used were one compost based on *Mangifera indica* biomass and the other with NPK-SB mineral fertilizer (14-18-18-6-1) and Urea (46%N).

Comment [WU9]: Full meaning

## 2.3. Methods

### Experimental design

The factor studied was fertilization, taken at 5 levels of variation. The levels of change in the factor constitute the treatments (Table I). The experimental set-up is a Fisher block comprising five (05) treatments with four (04) repetitions. Each treatment represented one elementary plot (EP), a total of 20. The dimensions of the entire system are 44 m x 19 m, i.e. a total area of 836 m<sup>2</sup> and those of the elementary plots is 8 m x 4 m, i.e. 32 m<sup>2</sup> each. The elementary plots were spaced 1 m apart, as were the blocks.

**Table I :** Treatments

Treatments	Significations	Code	Qtité/PE
Treatment 1	200 kg/ha NPK + 150 kg/ha Urea (MF : Minéral Fertiliser)	T1 :	640 g + 480 g
Treatment 2	2,5 t/ha of compost + MF	T2 :	8 kg +640 g+480g
Treatment 3	5 t/ha of compost + MF	T3 :	16 kg+640g+480 g
Treatment 4	7,5 t/ha of compost + MF	T4 :	24kg +640g+480g
Treatment 5	10 t/ha of compost + MF	T5 :	32 kg+640g+480g

## Data collection

### Plant height and corn yields

The height of the plants was taken on the 15th, 30th and 45th days after sowing. 12 plants in each elementary plot were randomly selected on the central lines and monitored on the 15th, 30th and 45th days after sowing (DAS), for the measurement of plant height using a graduated ruler. The number of grains per ear, the weight of 1000 grains, the biomass yield and the grain yield were determined by placing yield squares over 4 m<sup>2</sup> in each elementary plot. Biomass and grain yields were extrapolated to the hectare.

Comment [WU10]: Make interval of days

Comment [WU11]: Plant height was randomly measure with a meter ruler from the base to the tip, each 15days after planting for three consecutive times from twelve (12) elementary plots. The days were 15th, 30th and 45th after sowing.

Comment [WU12]: Superscrip

### Determination of soil chemical parameters

#### Soil sampling

Soil samples were taken before and after cultivation of the plots at a depth of 0-20 cm. Prior to crop establishment, a soil sample was taken from the entire trial plot following the three-point diagonal and after cultivation, soil samples were taken from each elementary plot. A total of 21 composite samples were collected for laboratory analysis.

Comment [WU13]: After and before ?

Comment [WU14]: It should be the same either 10 : 10 or 12 : 12 i.e before and after

Comment [WU15]: Did you just take the soil sample to the laboratory with sieving

Comment [WU16]: Make the work simple and direct

Comment [WU17]: The soil samples were collected after and before from ..... at a depth of 0-20cm using of soil auger, from ..... points with the farm or land and bulked. These were air dried, crushed and sieved through 2mm mesh, the sieved soil was then used to determine the physical and chemical properties of the soil. Part of the soil sample was use for pot experiments. A total of ... composite sample were collected.

#### Chemical soil analysis

The analyses of soil samples were carried out at the Soil-Water-Plant Laboratory of INERA Farako-Bâ. pH was measured from a soil suspension in water by the glass electrode pH meter electrometric method [9]. Organic Carbon (CO) was determined through the method from [10]. Total nitrogen (N<sub>tot</sub>): The Kjeldahl method was used to determine the total nitrogen. The C/N ratio was determined from the results of the total carbon and total nitrogen analyses. Assimilable phosphorus (Pass): it was determined according to the Bray I method [11]. Available potassium (K): it is first extracted with 30 ml of oxalic acid (H<sub>2</sub>C<sub>2</sub>O<sub>4</sub>) and concentrated hydrochloric acid (HCl) solution. The solution is completed to 1 litre in a volumetric flask. The assay is carried out by a flame photometer (JENWAY, PFP7 Flame Photometer). Sum of exchangeable bases (S): this is the amount of useful ions (Ca<sup>++</sup>, Mg<sup>++</sup>, K<sup>+</sup>, Na<sup>+</sup>) in the soil. The values of the exchangeable bases were obtained through displacement by a thiourea silver solution Ag(H<sub>2</sub>CSNH)<sub>2</sub> at 0.01 M and determined by

Comment [WU18]: surperscrip

spectrophotometry. Cation exchange capacity (CEC or T): It was made from the solution for extracting exchangeable bases. Saturation rate (V): it is the ratio of the sum of the exchangeable bases to the cation exchange capacity and corresponds to the percentage of the electronegative sites of the cation exchange capacity occupied by the exchangeable bases.

### **Statistical analysis of the data**

The data collected were entered with the Microsoft Excel 2016 spreadsheet and were subjected to an analysis of variance (ANOVA) using the GenStat edition 12.1, version 2009 software. The comparison and separation of the means was made possible thanks to the Student Newman-Keuls test (SNK) at the 5% probability threshold.

## **3. RESULTS AND DISCUSSION**

### **3.1. RESULTS**

#### **Effects of fertilisers on soil chemistry.**

#### **Effects of treatments on pH, organic C, nitrogen, C/N ratios, assimilable Phosphorus and available Potassium before and after culture**

The chemical characteristics of the soils collected before and after cultivation are presented according to the treatments in (Table II). In general, statistical analysis of the results revealed no significant variation between pre- and post-crop soil treatments for water pH, total organic Carbon, total nitrogen and C/N ratios. On the other hand, there are significant variations in the assimilable Phosphorus and Potassium available. The pH of the soils taken before cultivation and after cultivation show that the soils of the site are acidic with a pH between (5.04-5.30). Organic carbon and nitrogen levels in soils taken before and after cultivation remain low overall, regardless of the treatment. The C/N ratios of soils before and after cultivation are between 9.75 and 10.52. The available Phosphorus and Potassium contents of soils sampled before cultivation are 6.65 mg/kg and 72.14 mg/kg higher respectively than those of soils sampled after cultivation. In addition, the assimilable Phosphorus content of the T2 treatment (2.5 t/ha Compost + FV) increased by 7% compared to that of the T1 control (200 kg/ha NPK + 150 kg/ha Urea). On the other hand, in soils from the T4 (7.5 t/ha Compost + FV) and T5 (10 t/ha Compost + FV) treatments, the assimilable Phosphorus contents decreased by 20% compared to the control (T1). As for the available potassium, a 5% increase was achieved in the soils of the T5 and T3 treatments compared to the control soil (T1). The lowest available Potassium content was noted with the control treatment (T1). For the other parameters (pH,

**Comment [WU19]:** why not use real number (2)

total organic Carbon, total nitrogen and C/N ratio), compost treatments do not differ statistically from MF.

Table II. Soil chemical characteristics (pH, C, N, C/N, P, K) as a function of treatments.

Traitements	pH	C (%)	N-total (%)	C/N	Phosphorus (mg/kg)	Potassium (mg/kg)
Before cultivation	5.04±0.17	0.32±0.02	0.03±0.02	9.75±5.23	6.65a±0.44	72.14a±4.79
T1	5.25±0.13	0.39±0.06	0.04±0.01	9.92±0.24	4.95b±0.52	52.19b±3.87
T2	5.30±0.12	0.47±0.08	0.05±0.01	9.86±0.30	5.66ab±0.99	54.19b±7.62
T3	5.20±0.22	0.46±0.07	0.04±0.01	10.52±0.36	4.06b±0.91	57.99b±9.39
T4	5.26±0.13	0.42±0.05	0.04±0.01	10.47±0.35	3.28c±0.49	54.19b±7.42
T5	5.30±0.23	0.40±0.05	0.04±0.00	10.39±0.59	3.28c±0.64	58.13b±6.34
<i>Pr &gt;F</i>	<b>0.137</b>	<b>0.526</b>	<b>0.441</b>	<b>0.132</b>	<b>0.027</b>	<b>0.046</b>
<i>Signification</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>NS</i>	<i>S</i>	<i>S</i>

*Legend:* NS: non-significant, s: significant. The values assigned by the same letter(s) in the same column are not statistically different at the 5% significance level according to the Student-Newman-Keuls test.

### Effects of treatments on exchangeable bases, sum of exchangeable bases, cation exchange capacity, and saturation rate before and after soil sampling.

The results of the exchangeable bases, the sum of the exchangeable bases, the cation exchange capacity and the saturation rate are presented in (Table III). The analysis of variance showed no significant differences between pre- and post-cropping soils in terms of exchangeable bases and sum of exchangeable bases. On the other hand, there are significant variations between the soils of the plots before and after cultivation, with regard to the CEC and the saturation rate (V). Thus, the CEC and saturation rate contents of the plots before cultivation are lower than those of the soils from the plots after cultivation. The CEC rates of soils from the T3 (5 t/ha Compost + MF) and T2 (2.5 t/ha Compost + MF) treatments increased by 18% and 4% respectively compared to those from the T1 control soil (200 kg/ha NPK + 150 kg/ha Urea). The saturation rate from the soil of the T5 treatment (10 t/ha Compost + MF) increased by 16% and that from the soil of the T4 treatment (7.5 t/ha Compost + MF) increased by 6%, unlike the soil from the T2 treatment where a decrease of 2% was observed compared to that of the control soil T1.

Table III. The exchangeable bases, the sum of the exchangeable bases, the cation exchange capacity and the saturation rate according to the treatments.

Treatments	Ca <sup>2+</sup> (cmol <sub>c</sub> kg <sup>-1</sup> )	Mg <sup>2+</sup> (cmol <sub>c</sub> kg <sup>-1</sup> )	Na <sup>+</sup> (cmol <sub>c</sub> kg <sup>-1</sup> )	S (cmol <sub>c</sub> kg <sup>-1</sup> )	CEC (cmol <sub>c</sub> kg <sup>-1</sup> )	V (%)
Before cultivation	0.78±0.13	0.30±0.10	0.05±0.02	1.25±0.16	2.11b±0.46	45.50b±7.75
T1	0.67±0.11	0.33±0.07	0.02±0.01	1.14±0.15	2.69b±0.84	42.95b±6.80
T2	0.67±0.30	0.29±0.04	0.06±0.02	1.15±0.25	2.89b±0.47	40.92b±8.12
T3	0.77±0.21	0.39±0.07	0.05±0.01	1.34±0.23	3.86a±0.15	43.06b±9.07
T4	0.67±0.35	0.39±0.09	0.05±0.02	1.22±0.11	2.51b±0.20	48.12b±7.28
T5	0.73±0.52	0.41±0.08	0.05±0.01	1.33±0.14	2.43b±0.47	59.76a±8.12
<i>Pr &gt;F</i>	<b>0.989</b>	<b>0.901</b>	<b>0.478</b>	<b>0.989</b>	<b>0.047</b>	<b>0.048</b>
<b>Signification</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>S</b>	<b>S</b>

**Legend:** NS: non-significant, s: significant. The values assigned by the same letter(s) in the same column are not statistically different at the 5% significance level according to the Student-Newman-Keuls test. T1: 200 kg/ha of NPK + 150 kg/ha of Urea; T2: 2.5 t/ha of Compost + MF; T3: 5 t/ha of Compost + MF; T4: 7.5 t/ha of Compost + MF; T5: 10 t/ha of Compost + MF.

### Effect of treatments on the height of maize plants at the 15th, 30th and 45th DAS.

The evolution of the average height of maize plants as a function of the treatments at the 15th, 30th and 45th DAS is visible through the (Figure 1). Statistical analysis of variance revealed no significant variation between treatments in general at the 15th and 30th DAS. In contrast to the 45th DAS, there was a significant variation. Thus, the highest average heights were found in the T5 and T2 treatments with values of 91.14 cm and 83.95 cm compared to the control treatment which recorded the lowest average height with a value of 75.48 cm.




Figure 1: Average height of maize plants as a function of treatments at the 15th, 30th and 45th DAS.

### 3.2. Effect of treatments on the number of kernels per ear and on the weight of 1000 kernels

The number of grains per ear and the weight of 1000 grains were determined (Table IV). Statistical analysis of variance revealed a significant variation between the different treatments in general for the number of grains per ear as well as for the weight of 1000 grains. The best results in terms of number of grains per ear were obtained in the T5 treatment (10 t/ha of Compost + FV) with 438 grains/ear compared to the T1 control (200 kg/ha of NPK + 150 kg/ha of Urea) with 328.8 grains/ear. The same observation was observed with the weight of 1000 grains. The highest weight of 1000 grains comes from the T5 treatment with 248 g. On the other hand, the lowest weight of 1000 grains comes from the T1 control treatment with 221.8 g.

UNDER PEER REVIEW

Table IV. Average number of grains per ear and the average weight of 1000 grains depending on the treatments.

Treatments	Number of grains/ear	1000 grains weight (g)
T1	328.8b±127.04	221.8b±9.77
T2	436.5a±116.22	235.2ab±16.98
T3	389.5b±74.25	222.3b±20.02
T4	433.8a±114.58	232.3ab±22.74
T5	438.0a±98.74	248.0a±20.94
<b>Probability</b>	<b>0.042</b>	<b>0.049</b>
<b>Signification</b>	<b>S</b>	<b>S</b>

Legend: Means followed by the same letter in the same column are not significantly different at  $P = 0.05$ . S: Meaning.

### 3.3. Effect of treatments on biomass and grain yields

Average biomass and grain yield values were assessed (Table V). The analysis of variance showed a significant variation between treatments for both biomass and grain yield. The highest biomass and grain yields come from the T5 treatment with 5812 kg/ha and 3706 kg/ha respectively. The lowest biomass yield comes from the T1 control treatment with an average of 3031 kg/ha. As well as the lowest grain yield (2804 kg/ha).

Table V: Average biomass and grain yield as a function of treatments.

Treatments	Biomass yield (kg/ha)	Grain yield(kg/ha)
T1	3031c±108.53	2804b±160.05
T2	4906ab±195.49	2926b±197.98
T3	3875b±105.19	2838b±57.108
T4	4969ab±194.35	2950b±127,92
T5	5812a±153.59	3706a±267,32
<b>Probability</b>	<b>0.039</b>	<b>0.033</b>
<b>Signification</b>	<b>S</b>	<b>S</b>

Legend: Means followed by the same letter in the same column are not significantly different at  $P = 0.05$ . S: Meaning.

## 3.2. DISCUSSION

Effects of fertilisers on soil chemistry.

Effects of treatments on pH, organic C, nitrogen, C/N ratios, assimilable P and available K before and after culture.

No significant changes were reported in water pH, organic carbon, total nitrogen and C/N ratios. The analysis of the soils sampled before and after cultivation showed that the pH of our site is acidic ( $5.04 \leq \text{pH} \leq 5.30$ ). This acidity is due to the nature of the soil but also to the accumulation of certain chemicals (pesticides, inorganic fertilizers) used over several years in the soil of the site. The work of [12] showed that the use of inorganic fertilizers and pesticides causes a decrease in soil pH. The low level of organic carbon and nitrogen could be explained by the decrease in the accumulation of organic matter but also by tillage which leads to a loss of nitrogen through mineralization, leaching and water erosion [13]. The C/N ratio of the soil before and after sampling is low and is between 9.75 and 10.52, thus reflecting a strong mineralization of the organic matter in the soil according to BUNASOLS standards. For [14], a C/N ratio between 8 and 12 is an optimal ratio and a sign of good microbial activity and a good balance between humification and mineralization. The high levels of P and K available in the soil before cultivation compared to those in the soils after cultivation and the 20% decrease in the available phosphorus in the T4 and T5 treatments, would be due to the fact that the maize plants were able to take up a necessary quantity of these elements in the soil after cultivation and in these treatments. [15] also observed low levels of assimilable and total phosphorus in the soil under rhizospheric influence due to the uptake of bioavailable forms by the roots. This may also be linked to the fixation of phosphorus in the soil by other elements and to the low mineralisation due to a deficiency of organic matter and/or the unfavourable pH of the environment. Then, the 7% increase in assimilable P in T2 treatment and the 5% increase in K available in T5 and T3 treatments; would be explained by the combined action of organo-mineral fertilization in these treatments. Also, this could be linked to the role of organic amendments in improving the physicochemical properties of the soil through an increase in its exchange capacity [16]. The lowest available K content noted by the control treatment (T1) would be justified by the addition of exclusively mineral fertiliser.

**Effects of treatments on exchangeable bases, sum of exchangeable bases, cation exchange capacity, and saturation rate before and after culture.**

The results obtained after analysis show that the exchangeable bases and the sum of the exchangeable bases before and after cultivation are low overall. Also, these results show that the different fertilizers applied were not remarkable on these different parameters. This could be due to the non-restitution of organic waste to compensate for losses due to mineralization, resulting in a rapid decrease in soil organic matter and a decrease in exchangeable base contents, cation exchange capacity, acidification and an increase in exchangeable aluminum

content [17]. The high levels of CEC in the T3 (18%) and T2 (4%) treatments could be justified by the effect of compost associated or not with mineral fertilizers on the chemical parameters of organic matter, calcium and magnesium, similar to the previous results found by several authors [18] with different organic fertilizers. Indeed, with compost, the level of organic carbon, biological activity, soil moisture as well as nitrogen, phosphorus, potassium, magnesium and calcium in the soil increases [19]. The saturation rate increased by 16% in the treatment that received 10 t/ha of Compost + FMV and by 6% in the treatment that received 7.5 t/ha of Compost + FMV and this increase in these treatments confirms the ability of the compost enriched with mango leaf to restore the fertility of the soils of the study through its nutrient richness. This would come from the activity of fungi trapped in this compost which is an activator of the microbial activity of the soil. These results are in agreement with those of the authors [20], who showed the ability of compost to restore the properties of acidic soils in the Democratic Republic of Congo (DRC). The low saturation rate recorded by the treatment that received 2.5 t/ha of Compost + FMV is linked to the low amount of calcium and magnesium in this treatment.

#### **Effect of treatments on maize plant height.**

This height performance of maize plants from the T3 (5 t/ha of compost + FMV) and T5 (10 t/ha of compost + FMV) treatments could be justified by the availability of mineral elements and the improvement of the physicochemical properties of the soil due to the combination of organic and mineral fertilisation. These results are consistent with those of [21] who showed that organic matter retains nutrients on the surface while mineral fertilizer alone accelerates their vertical migration. The poor height performance of maize plants from the T1 treatment (200 kg/ha NPK + 150 kg/ha of Urea) is justified by the exclusive contribution of mineral fertiliser to the soil. This situation could result in a lack of available resources, in particular mineral elements, which are the main factors limiting agricultural production in the dry tropics [22].

#### **Effects of treatments on corn yield components**

The analysis of the results showed that the different fertilizers applied may have had a positive impact on the maize yield components. Indeed, these different components of maize yield have generally increased with the doses of 10 t/ha of compost + FMV. This result could be due to the effect of the dose of organic manure combined with mineral fertiliser, which was able to release a necessary amount of nutrient thus promoting good maize productivity. This is what is maintained ([23] and [24]) when they stipulate that nutrients (N P K) are very accessible to plants, thus ensuring a good diet for the plant and allowing the

best yields to be obtained in most cases. They attribute this to a high demand for mineral elements (especially N and P). Our results are similar to those obtained by [25] and [26] who state that decomposed litter could be a nutrient source for maize plants. In addition, this increase in yield could be linked to the presence of less lignified material in this compost, which reflects the availability of free organic matter for corn plants [27]. These results corroborate those obtained by [25] who found that potassium (K) stimulates the constitution of the nutrient reserve; Nitrogen is involved in the main yield determination processes, and phosphorus could accelerate seed planting and seed maturation. Our results are also in line with those of [28] who found that the best rice yields are obtained by organo-mineral fertilizations. However, the lowest yield components were reported by the treatment which received exclusively popularized mineral manure (200 kg/ha of NPK + 150 kg/ha of Urea). This would be justified by the low level of soil fertility in this treatment.

#### 4. CONCLUSION

The objective of this study was to evaluate the effects of mango biomass compost combined with popularized mineral fertilizer on soil chemical parameters and maize yields in western Burkina Faso. The aim is to contribute to the improvement of soil chemical properties and maize productivity. It appears that the fertilization applied did not have much effect on the chemical parameters of the soil and on the growth of the maize plants. Nevertheless, there was a significant improvement with the assimilable P, the available K, the CEC, the saturation rate and the height of the plants at the 45th JAS. Thus, the best results in terms of soil chemical parameters and plant growth were achieved with the T3 (5 t/ha compost + FMV) and T5 (10 t/ha compost + FMV) treatments. As for the yield components, our results showed that the different fertilizers applied were beneficial. Indeed, these different components have generally increased with the dose of 10 t/ha of compost + FMV (T5). However, the lowest yield components were reported by the treatment that received 200 kg/ha NPK + 150 kg/ha Urea (T1). We can therefore say that the combination of organo-mineral fertilisers makes it possible to increase the productivity of maize and to partly improve soil fertility. It would be interesting to continue the study in other agro-ecological zones of Burkina Faso in order to assess the impact of these fertilizers on the biological properties of the soil.

## REFERENCES

- [1] - SCADD, 2013. Mid-term performance report 2013. Ouagadougou, Burkina Faso. 56p.
- [2] - FAOSTAT, 2014. <http://Faostat.fao.org> consulted on June 6, 2023.
- [3] - DGESS, 2016. Final results of the agricultural campaign and the food and nutritional situation 2015/2016, 39p.
- [4] - MARHASA., 2015. General Report of the Campaign (RGC) minerals and organics: consequences on the yields of a continuous corn crop, 57.
- [5] - Bacyé B., Somda S. and Boro A., 2001. Diagnostic study of soil fertility in cropping systems in the western cotton zone of Burkina Faso. Research report, SADAOC Network. 70p.
- [6] - Pallo F.J.P. and Sawadogo N., 2010. Correlation test between the characteristics of the climate of vegetation and soils in Burkina Faso. *Int. 1 Biol. Chem.Sci* 4 (5). ISSN 1991-8631. 18391850.
- [7] - Guinko S., 1984. Vegetation of Upper Volta, State doctoral thesis, University of Bordeaux 2, France 318p.
- [8] - Bado B. V., 2002. Role of legumes on the fertility of tropical ferruginous soils in the Guinean and Sudanese zones of Burkina Faso. Ph.D. thesis, Department of Soils and Environment, Laval University, 148 p.
- [9] - AFNOR, 1999. Determination of pH. (French Standardization Association) NF ISO103 90, AFNOR Soil quality, Paris, pp. 339-348.
- [10] - Walkley A. Black R. N, 1934. An examination of the Degtjareff method for determining soil organic matter and to proposed modification of the chromic acid titration method. *Soil Science*, 37:29-38.
- [11] - Bray R.H, Kurtz L.T., 1945. Determination of total, organic, and available forms of phosphorus in soils. *Soil Science* 59, 39-45.
- [12] - Sradnick A., Murugan R., Oltmanns M., Raupp J., Joergensen RG. 2013: Changes in functional diversity of the soil microbial community in a heterogeneous sandy soil after long-term fertilization with cattle manure and mineral fertilizer. *Applied Soil Ecology*, 63, 23-28p.

- [13] - Koulibaly B., 2011. Characterization of soil acidification and fertility management of cotton agrosystems in Burkina. Doctoral thesis, University of Ouagadougou, Applied Sciences, Agro-pedology, Burkina Faso, 183 p.
- [14] - Giroux M. Audesse P. 2004. Comparison of two methods for determining organic carbon, total nitrogen and C/N ratios of various organic amendments and farm fertilizers. *Agrosol*, vol.15, no2, 107-110.
- [15] - Boudiaf-Nait kaci M., Hedde M., Mouas-Bourbia S., Siad D. and Derridj A., 2012. Bioavailability of phosphorus in the rhizosphere of the pomegranate (*Punica granatum L.*) Under arid climate: case of the orchards of Messaad Wilaya of Djelfa, UMMTO Algeria. 11th Soil Study Days, France. 7 p.
- [16] - Koulibaly, B., Dakuo, D., Ouattara, A., Traoré, O., Lompo, F., Zombré, P.N. and Kouamé, A.Y. 2015. Effects of the association of compost and mineral manure on the productivity of a cotton and corn-based cropping system in Burkina Faso. *Tropicultura*, 33, 125–134
- [17] - Mills A. J., Fey M. V., 2003. Declining soil quality in South Africa: effects of land use on soil organic matter and surface crusting. *South African Journal of science*. 99 pp 429-436.
- [18] – Sawadogo J., Coulibaly P. J. A., Bambara F. J., Savadog A. C., Compaoré E. Legma J. B. 2020. Effects of organic fertilizers on soil physiochemical parameters and on onion (*Allium cepa L.*) productivity in the Central West region of Burkina Faso. *Afrique SCIENCE* 17 (6): 44-57.
- [19] - Karuku G., Onwonga R., Kathumo V. 2018. Effects of tillage practices of cropping systems and organic inputs on soil nutrient content in Machakos Comet. *Journal of Agriculture and Sustainability*, 12 (1) ISSN 2201-4357. [20] - Mpundu Mubemba M., Useni Y., Nyembo L., Colinet G., 2014. *Biotechnology, Agronomy, Society and Environment*, 18 (3) 367 – 375.
- [21] - Grogan N, Diomandé M, Beugré G. A. M, Ouattara Y, Akaffou D.S, 2018. Study of the quality of symbiosis (*Anabaena azollae*, *Azolla caroliniana*), compost and NPK on vegetative growth and tomato yield in Daloa (Côte d’Ivoire). *Journal of Applied Biosciences* 129: 13004 -13014. ISSN 1997-5902
- [22] - Nolte C., Tiki-Manga T., Badjel Badjel S., Gockowski J. and Hauser S. 2005. Groundnut, maize and cassava yield in mixed food cropfields after *Calliandra* tree fallow in southern Cameroon. *Experimental Agriculture* 41 (1): 21-37.

- [23] - Hema M L., 2016. Effect of previous crops and manures on phosphorus availability and productivity of strict rainfed rice in the southern Sudanian zone of Burkina Faso. Thesis of Agricultural Engineer of the Institute of Environmental Sciences and Rural Development of the Polytechnic University Center of Dédougou. 53 p.
- [24] - Traoré, A., 2016. Effect of rotations and fertilization on the yield of strict rainfed rice in the western zone of Burkina Faso. Unique Doctorate Thesis in Soil Science. EDoctoral school in Natural Sciences and Agronomy, Polytechnic University of Bobo-Dioulasso. 132 P.
- [25] - Topan S., 2015. Influence of Jatropha hedges on the productivity of sorghum (*Sorghum bicolor*) in the South Sudanese zone of Burkina Faso. Thesis of Agricultural Engineer, Multipurpose Agricultural Center of Matourkou, 42p.
- [26] - Ouédraogo R C. 2016. Comparative effect of three types of composts on the yield of corn (*Zea mays*) in the Farako-Bâ research station. Report of the Higher Technician Certificate (BTS) Agriculture of the Agricultural Center (CAP) of Matourkou. 43 P.
- [27] - Larbi M., 2006. Influence of the quality of composts and their extracts on the protection of plants against fungal diseases, doctoral thesis, University of Neuchâte. Botanical Institute, 161 p.
- [28] - Bougma B. A., 2013. Effect of previous crops and manures on soil fertility and rainfed rice yields. Master II end-of-cycle dissertation, specializing in integrated soil fertility management, Polytechnic University of Bobo Dioulasso, 47p.