

Effect of Integrated sources of nutrients on Growth, Yield and Economics of Cowpea (*Vigna unguiculata*L.) var Ankur Gomati

ABSTRACT

A field experiment was conducted in randomized block design with 7 treatments and 3 replications to study the effect of fertilizers, organic manures, biochar and crop residues on growth, yield and economics of cowpea (*Vigna unguiculata* L.) in Inceptisols of Prayagraj, Uttar Pradesh, India in 2022. A research variety Ankur Gomati was taken for this research trial. Recommended doses of fertilizers (N, P₂O₅ and K₂O @ 25, 75 and 60 kg ha⁻¹) was applied with inorganic fertilizers (Zn and B), organic manures (FYM, vermicompost, neem cake and poultry manure), biochar and crop residues (rice straw and wheat straw). Growth parameters such as the maximum plant height 46.96 cm, 70.21 cm and 92.22 cm, maximum number of branch plant⁻¹ 5.82, 7.56 and 9.79, highest number of leaf plant⁻¹ 28.32, 30.56 and 35.22, highest length of leaf as 5.37, 7.31 and 9.18 were recorded in T₇ [RDF + Poultry Manure @2 t ha⁻¹ + Biochar @2.5 t ha⁻¹ + Vermicompost @2 t ha⁻¹ + Neem Cake @500 kg ha⁻¹]at 30 DAS, 55 DAS and 70 DAS respectively. And yield parameters such as high pod length as 21.72 and 20.01, highest number of pods plant⁻¹ 35.52 and 35.75, highest number of seed pod⁻¹ 10.01 and 10.21 and highest test weight 27.12 and 26.23 were recorded in T₇ [RDF + Poultry Manure @2 t ha⁻¹ + Biochar @2.5 t ha⁻¹ + Vermicompost @2 t ha⁻¹ + Neem Cake @500 kg ha⁻¹]at 55 DAS and 70 DAS respectively. Highest yield 160.12 were obtained in treatment T₇ [RDF + Poultry Manure @2 t ha⁻¹ + Biochar @2.5 t ha⁻¹ + Vermicompost @2 t ha⁻¹ + Neem Cake @500 kg ha⁻¹]. Maximum gross return ₹ 2,72,204 was recorded in treatment T₇ [RDF + Poultry Manure @2 t ha⁻¹ + Biochar @2.5 t ha⁻¹ + Vermicompost @2 t ha⁻¹ + Neem Cake @500 kg ha⁻¹]. Maximum C:B ratio 1:4.31 was recorded in T₂ [RDF + Boron @2 kg ha⁻¹ + Zinc @7 kg ha⁻¹]. But maximum net return ₹ 1,98,765.6 was recorded in treatment T₂ [RDF @100% + Boron @2 kg ha⁻¹ + Zinc @7 kg ha⁻¹]. The yield and growth parameters also significantly increased with the application of the combination of organic and inorganic fertilizers.

Keywords: Fertilizes, organic manures, biochar, crop residues, cowpea, growth, yield and economics.

INTRODUCTION

Cowpea [*Vigna unguiculata* (L.) Walp] is one of the important kharif pulse crops grown for vegetable, grain, forage and green manuring belonging to family Leguminaceae (Ahmed *et al.*, 2012). In India cowpea grown states are Uttar Pradesh, Punjab, Haryana, Rajasthan and Madhya Pradesh and cowpea cultivated in arid and semi-arid. Uttar Pradesh is the leading state among them (Meena *et al.*, 2014). Green tender pods of cowpea are used as vegetable, the vegetable cowpea pods contain 24.8% protein, 63.6% carbohydrate, 1.9% fat, 6.3% fiber, 7.4 ppm thiamine, 4.2 ppm riboflavin and 28.1 ppm niacin (Ahlawat and Shivkumar 2005).

Cowpea is highly responsive to fertilizers. The dose of fertilizer depends on the initial soil fertility status and moisture availability conditions. Although it gives good response to application of recommended dose of fertilizer (Ahlawat and Shivkumar 2005). Nutrient management with recommended dose of NPK and micronutrients B and Zn can augment the economic yield of cowpea both in green pod and seed production (Debnath *et al.*, 2018). Use of organic manure alone or in combination with chemical fertilizers, helps in improving physio-chemical properties of the soil, improves the efficient utilization of applied fertilizers resulted in higher seed yield and quality (Salem, 2012). Organic manures viz., farm yard manure (FYM), vermicompost (VC), poultry manure (PM) and oilcakes help in the improvement of soil structure, aeration and water holding capacity of soil (Joshi *et al.*, 2016). According to Behera *et al.*, (2007), it improves plant

growth by improving the physical and chemical characteristics of the soil, i.e., cation exchange capacity (CEC), bulk density, water holding capacity and permeability as well as biological properties, all contributing to an increased crop productivity. Crop residues are a potential source of organic matter in soils and Soil organic matter indirectly effects on the plant (Kumari *et al.*, 2019).

Previous researches on cowpea reveal growth and yield attributes enhancements through the use of inorganic nutrients sources (Debnath *et al.*, 2018), organic nutrient sources (Joshi *et al.*, 2016; Patel *et al.*, 2018), crop residues (Ndisoet *et al.*, 2018), and even with biochar application (Phares *et al.*, 2020). However, there is no systematic study that involves the integrated use of these nutrient and carbon sources on the performance of cowpea. In this light, the present study aimed to evaluate the effect of integrated sources (inorganic and organic) of nutrients and carbon (biochar) on i) growth and yield, and ii) economics of cowpea crop grown in an Inceptisol of Indo-Gangetic plain.

METHOD AND MATERIALS

The field experiment was conducted at research farm located at Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, Uttar Pradesh (25°24' N latitude, 81° 51' E longitude) during *Kharif* season of 2022. This area experiences sub-tropical to semi- arid climate with extremely hot summer and cold winter. The maximum temperature reaches up to 46– 48°C and seldom falls as low as 4–5°C. The average rainfall in this area is ~1100 mm most of which is received during July-September.

The experimental area was laid out in randomized block design with seven treatments and three replications. The treatments comprised of T₁- RDF @ 25:75:60 N:P:K, T₂- RDF + B @ 2 kg ha⁻¹ + Zn @ 7 kg ha⁻¹, T₃- RDF + Farm Yard Manure @ 5 t ha⁻¹, T₄- RDF + Vermicompost @ 2 t ha⁻¹ + Rice Straw @ 6 t ha⁻¹, T₅- RDF + Neem Cake @ 500 kg ha⁻¹ + Wheat Straw @ 6 t ha⁻¹, T₆- RDF + Biochar @ 2.5 t ha⁻¹ and T₇- RDF + Poultry Manure @ 2 t ha⁻¹ + Vermicompost @ 2 t ha⁻¹ + Biochar @ 2.5 t ha⁻¹ + Neem Cake @ 2 t ha⁻¹. The organic manures were incorporated into soil before three weeks of sowing and the inorganic fertilizers were applied at the time of showing. The sources of N, P, K, Zn and B were urea, SSP, MOP, zinc sulphate and zorax respectively. Half dose of urea was applied at the time of showing and the other half were applied after 21 days of showing as split dose. First picking was held on 55 DAS and the final harvesting was held on 70 DAS. All cultural practices were adopted uniformly in each plot as and when required. In general, no incidence of disease and pest was recorded during both the season. The data recorded during the course of the investigation will be subjected to statistical analysis by 3X 3 RBD, as per the method "Analysis of Variance (ANOVA) technique" as given by Fischer (1960). Experiment will be laid out in RBD and the treatment will be replicated three times. The significant and non-significant effect was judged with the help of "F" (variance ratio) table. The significant difference between the means was tested against the critical difference of 5% level.

Growth parameters like plant height (cm), number of branches plant⁻¹, no of leaf plant⁻¹, leaf length (cm) and pod length taken at 30 DAS, 55 DAS and 70 DAS and yield parameters like no of pod plant⁻¹, no of seed pod⁻¹, seed weight and pod yield (kg ha⁻¹) taken at 55 DAS and 70 DAS. Economics like net profit, gross return, cost of cultivation and cost benefit ratio (C:B) / return per rupee invested also calculated.

RESULT AND DISCUSSIONS

Effect on growth parameters of Cowpea

The data presented in Table 1 shows that growth parameters significantly increased by using the combination of inorganic fertilizers, organic manures, biochar and crop residues. Maximum plant height 46.96 cm, 70.21 cm and 92.22 cm, maximum number of branch plant⁻¹ 5.82, 7.56 and 9.79, highest number of leaf plant⁻¹ 28.32, 30.56 and 35.22, highest length of leaf as 5.37, 7.31 and 9.18 were recorded in T₇ [RDF + Poultry Manure @ 2 t ha⁻¹ + Biochar @ 2.5 t ha⁻¹ + Vermicompost @ 2 t ha⁻¹ + Neem Cake @ 500 kg ha⁻¹] at 30 DAS, 55 DAS and 70 DAS respectively followed by T₄ [RDF + Vermicompost @ 2 t ha⁻¹ + Rice Straw @ 6 t ha⁻¹] respectively and minimum result were found in T₁ [RDF]. This may be

because of adequate supply of nutrients by nutrient sources like inorganic fertilizers, organic manures, biochar and crop residues which helps in vigorous vegetative growth of plants (Khandagleetal., 2019).

Table 1. Effect of integrated sources of nutrients on growth parameters of Cowpea

Treatments	plant height (cm)			No of branch plant ⁻¹			No of leaf plant ⁻¹			Length of leaf (cm)		
	30 DAS	55 DAS	70 DAS	30 DAS	55 DAS	70 DAS	30 DAS	55 DAS	70 DAS	30 DAS	55 DAS	70 DAS
T ₁	39.93	58.31	78.12	3.26	5.07	7.21	20.13	24.21	26.33	4.05	6.02	7.66
T ₂	40.36	60.15	80.24	3.55	5.27	7.54	21.54	25.82	27.01	4.61	6.25	7.89
T ₃	43.29	64.23	85.16	4.85	6.02	8.36	25.55	27.16	28.67	5.01	7.01	8.51
T ₄	45.11	68.52	89.56	5.45	7.12	9.23	26.36	28.99	30.31	5.21	7.28	9.01
T ₅	44.82	65.86	86.77	4.98	6.40	8.76	26.01	27.91	29.12	5.04	7.15	8.71
T ₆	41.45	62.11	82.71	3.87	5.51	7.82	25.05	26.12	27.92	4.73	6.11	8.15
T ₇	46.96	70.21	92.22	5.82	7.56	9.79	28.32	30.56	32.22	5.37	7.31	9.18
F-test	S	S	S	S	S	S	S	S	S	S	S	S
S.Em. (±)	0.72	1.18	1.32	0.04	0.01	0.11	0.31	0.31	0.38	0.07	0.08	0.13
C.D.	2.12	3.47	3.89	0.14	0.31	0.34	0.93	0.91	1.12	0.23	0.26	0.40

Effect on yield parameters of Cowpea

The data presented in Table 2 shows that yield parameters significantly increased by using the combination of inorganic fertilizers, organic manures, biochar and crop residues. Highest average pod yield 161.12 q ha⁻¹ and yield parameters like high pod length 21.72 and 20.01, highest number of pods plant⁻¹ 35.52 and 35.75, highest number of seed pod⁻¹ 10.01 and 10.21 and highest test weight 27.12 and 26.23 and were recorded in T₇ [RDF + Poultry Manure @ 2 t ha⁻¹ + Biochar @ 2.5 t ha⁻¹ + Vermicompost @ 2 t ha⁻¹ + Neem Cake @ 500 kg ha⁻¹] at 55 DAS and 70 DAS respectively followed by T₄ [RDF + Vermicompost @ 2 t ha⁻¹ + Rice Straw @ 6 t ha⁻¹] respectively and were found to be significant. Minimum yield parameters observed in T₁ [RDF]. Increased in yield parameters may be due to adequate availability of nutrients during reproductive stage of crop results in the increased yield parameters (El-Sharawy *et al.*, 2018 and Joshi *et al.*, 2016).

Table 2. Effect of integrated sources of nutrients on yield parameters of Cowpea

Treatments	Pod length (cm)		No of pods plant ⁻¹		No of seed pod ⁻¹		Seed weight (g)		Pod Yield (qha ⁻¹)
	55 DAS	70 DAS	55 DAS	70 DAS	55 DAS	70 DAS	55 DAS	70 DAS	
T ₁	14.12	15.01	19.21	20.12	6.99	7.45	20.55	19.79	19.79
T ₂	15.63	14.92	21.98	22.11	7.45	8.14	21.21	20.13	20.13
T ₃	17.16	15.55	25.24	26.01	8.21	8.95	23.85	23.21	23.21
T ₄	19.36	19.12	28.74	28.89	9.74	9.86	26.01	25.13	25.13
T ₅	17.75	16.21	26.07	26.52	8.82	9.05	24.31	23.43	23.43
T ₆	16.55	15.16	23.54	23.03	8.12	8.56	22.45	21.54	21.54
T ₇	21.72	20.01	35.52	35.75	10.01	10.21	27.12	26.23	26.23
F-test	S	S	S	S	S	S	S	S	S
S.Em. (±)	0.25	0.26	0.50	0.48	0.14	0.12	0.44	0.39	0.39
C.D.	0.75	0.77	1.47	1.43	0.42	0.35	1.29	1.16	1.16

Effect on economics

The data presented in Table 3 shows that maximum gross return (₹ 2,41,680) was recorded in treatment T₇ [RDF + Poultry Manure @ 2 t ha⁻¹ + Biochar @ 2.5 t ha⁻¹ + Vermicompost @ 2 t ha⁻¹ + Neem Cake @ 500 kg ha⁻¹] followed by T₄ [RDF + Vermicompost @ 2 t ha⁻¹ + Rice Straw @ 6 t ha⁻¹] respectively and minimum gross return (₹ 1,51,020) recorded in treatment T₁ [RDF]. Maximum B:C ratio (2.80) was recorded in T₄ [RDF + Vermicompost @ 2 t ha⁻¹ + Rice Straw @ 6 t ha⁻¹] treatment followed by T₃ [RDF + Farm Yard Manure @ 5 t ha⁻¹] respectively and minimum B:C ratio (1.36) recorded in T₆ [RDF + Biochar @ 2.5 t ha⁻¹] treatment. But maximum net return (₹ 1,47,624.1) was recorded in treatment T₄ [RDF + Vermicompost @ 2 t ha⁻¹ + Rice Straw @ 6 t ha⁻¹] followed by T₅ [RDF + Neem Cake @ 500 kg ha⁻¹ + Wheat Straw @ 6 t ha⁻¹] respectively and the minimum net return (₹ 49,584.1) was recorded in T₆ [RDF + Biochar @ 2.5 t ha⁻¹].

Table 3. Economic of Cowpea of different treatment combination

Treatm ent	Yield(qha ⁻¹)	Yield(₹ha ⁻¹)	Grossr eturn (₹ha ⁻¹)	Totalcostofc ultivation (₹ha ⁻¹)	Netprofit (₹ha ⁻¹)	(C:B)
T ₁	100.68	1500	151020	56890.9	124129.1	1:2.65
T ₂	107.21	1500	160815	59991.4	100823.6	1:2.68
T ₃	128.51	1500	192765	69390.9	103874.1	1:2.77
T ₄	153.01	1500	229515	81890.9	147624.1	1:2.80
T ₅	141.56	1500	212340	78390.9	133949.1	1:2.70
T ₆	122.65	1500	183975	134390.9	49584.1	1:1.36
T ₇	161.12	1500	241680	175390.9	66289.1	1:1.37

CONCLUSION

Application of inorganic fertilizers, organic manures, biochar and crop residues gave the best economics like gross return, net return and cost benefit ratio (C:B) observed in T₂ [I @100% + B @100% + Z @100%] and gave best significant results of growth and yield parameters recorded in T₇ - [RDF + Poultry Manure @ 2 t ha⁻¹ + Biochar @ 2.5 t ha⁻¹ + Vermicompost @ 2 t ha⁻¹ + Neem Cake @ 500 kg ha⁻¹] and found at par than any other treatments. So, this is concluded that combination of inorganic, organic and carbon source increase growth and yield parameters of cowpea significantly.

REFERENCES

- Ahlawat, I. P. S. and Shivakumar, B. G. (2005) Kharif Pulses. In Textbook of Field Crops Production (R. Prasad, Ed.) ICAR, New Delhi, India.
- Ahmed, M. N., Abdelrhim, A. J., Salaheldeen, E. A. and Feisal, M. I. (2012) Determination of Suitable Variety and Plant Spacing of Cowpea (*Vigna unguiculata* L. Walp) in the Sandy Soil.
- Debnath, P., Pattanaik, S. K., Sah, D., Chandra, G. and Pandey, A. K. (2018) Effect of Boron and Zinc Fertilization on Growth and Yield of Cowpea (*Vigna unguiculata* L. Walp.) in Inceptisols of Arunachal Pradesh. *Journal of the Indian Society of Soil Science*. 66(2):229-234.
- El-Sharawy Huda, H., Mahmoud, M. I. and El-Tantawy, E. M. (2018) Organic farming of cowpea, international journal of soil science. 7(1):2314-6079.
- Joshi, Deepa, Gediya, K. M., Patel, J. S., Birari, M. M. and Gupta, Shivangini (2016) Effect of organic manures on growth and yield of summer cowpea [*Vigna unguiculata* (L.) Walp] under middle Gujarat conditions. *Agric. Sci. Digest*. 36(2): 134-137
- Kumari, R., David, A. A., Thomas, T., Bose, D. K. and Dawson, J. (2022) Response of Inorganic fertilizers, Crop residues, Organic manures and Biofertilizer on Soil health and Yield of Cowpea (*Vigna unguiculata* L.). *International Journal of Chemical Studies*. 6(3):2117-2119.
- Meena, J. S., Verma, H. P. and Pincholi, P. (2014) Effect of fertility levels and biofertilizer on Yield and Quality and economic of Cowpea [*Vigna unguiculata* (L.)]. *Agriculture for Sustainable Development*. 2(2): 162-164
- Ndiso, J. B., Cheminingwa, G. N., Olubayo, F. M. and Saha, H. M. (2018) Effect of cowpea crop residue management on soil moisture content, canopy temperature, growth and yield of maize - cowpea intercrops. *International Journal of Agriculture, Environment and Bio Research*. 3(5):231-250.
- Patel, Himani B., Shah, K. A., Barvaliya, M. M. and Patel, S. A. (2018) Response of Greengram (*Vigna radiata* L.) to Different Level of Phosphorus and Organic Liquid Fertilizer. *International Journal of Current Microbiology and Applied Sciences*. 6(10):3443-3451
- Pharesa, Christian Adler, Atiaha, Kofi, Frimponga, Kwame Agyei, Danquah, Andrews, Asareb, Aaron T., Aggor-Woananu, Samira (2020) Application of biochar and inorganic phosphorus fertilizer influenced rhizosphere soil characteristics, nodule formation and phytoconstituents of cowpea grown on tropical soil. *Heliyon*. 6: 1-9
- Yeboah, E., Asamoah, G., Ofori, P., Amoah, B., Agyeman, K. O. A. (2020) Method of biochar application affects growth, yield and nutrient uptake of cowpea. *DE GRUYTER*. 5: 352-360