

Original Research Article

Enhancement of sesamum yield with split application of nitrogen under irrigated conditions

ABSTRACT

Aim: to determine the optimum dose and best time of nitrogen application in sesamum in YSR Kadapa district of Andhra Pradesh.

Study design: Split plot design

Place and duration of study : Department of Agronomy, Agricultural Research Station, Utukur, Kadapa, Andhra Pradesh, India for three years *i.e.*, 2020-21 to 2022-23.

Methodology: . The treatments included four levels of nitrogen *viz.*, 40, 60, 80 and 100 kg ha⁻¹ and four split doses as 50:50 basal and 30 DAS, 25:75 basal and 30 DAS, 25:37.5:37.5 basal, 20 and 40 DAS and 0:50:50 basal, 20 and 40 DAS which were laid out in split plot design with levels of nitrogen in main plots and time of nitrogen application in sub plots in three replications. The variety YLM-66 was sown at a spacing of 30 cm × 10 cm. Plant height, number of branches per plant, number of capsules plant⁻¹, number of seeds capsule⁻¹ and seed yield were recorded.

Results: Maximum plant height, number of branches per plant, number of capsules plant⁻¹, number of seeds capsule⁻¹ and seed yield were recorded with application of 100 kg N ha⁻¹ which was on par with 80 kg N ha⁻¹. Test weight of sesamum was not significantly influenced by nitrogen management and regarding the time of application and number of splits, application of 100 kg N ha⁻¹ in three splits in the ratio of 25-37.5-37.5 at basal, 20 and 40 days after sowing has recorded maximum seed yield in sesamum under irrigated conditions which was on par with 80 kg N ha⁻¹ in three splits in the ratio of 25-37.5-37.5 at basal, 20 and 40 days after sowing.

Conclusion: Application of 80 kg N ha⁻¹ may be considered as an optimum dose of nitrogen in contrast to the present recommended dose of 40 kg N ha⁻¹ which has a yield advantage of 11.5 % over 100 % RDF.

Key words : Nitrogen, Sesamum, Split application, Time of application

INTRODUCTION

Sesamum (*Sesamum indicum* L.) called as queen of oil seeds is one of the most ancient oilseed crops in the tropics. It is an important source of high-quality oil and protein. The oil has excellent stability due to the presence of natural antioxidants such as sesamol and sesamin. The proteins in sesamum are rich in essential sulphur containing amino acids such as methionine and tryptophan. In India average production of sesamum was 741.56 Mt with area of 1594.41 Mha and productivity of 465 kg ha⁻¹ compared to the world average (535 kg ha⁻¹). Inappropriate use of fertilizers is one of the major production constraints in sesamum (Amin and Alireza, 2015, El-Sherif, 2016).

Sesamum is grown in an area of 3279 ha in YSR Kadapa district of Andhra Pradesh during *rabi* season with average productivity of 406 kg ha⁻¹. Among the different agronomic management components that influences the productivity of sesamum, nutrient management plays a key role out of which widely applied nitrogen management is little bit difficult due to its losses by ammonia volatilization during nitrification, denitrification, downward leaching and runoff with poor efficiency of 30–40 %. Nitrogen is considered as an essential element of bio-molecules such as amino acids, proteins, nucleic acids, phyto hormones and a number of enzymes and coenzymes. Nitrogen strongly stimulates growth, expansion of the crop canopy and interception of solar radiation. Nitrogen is the most dynamic and important nutrient required for survival of all living things (Rosolem *et al.*, 2017). Significant positive correlation was observed between nitrogen application rate and yield in sesamum. Patel *et al.* (2014) reported that varied responses of sesamum to nitrogen from 60 kg ha⁻¹ to 180 kg ha⁻¹ were observed at different places. Kushawala *et al.* (2022) reported the response up to 75 kg N ha⁻¹, Umar *et al.* (2012) studied the effects of nitrogen fertilizer rates (0, 40 and 80 kg ha⁻¹) on sesamum yield and concluded that 80 kg N ha⁻¹ as optimum for higher seed yield (0.79 t ha⁻¹) and oil content (45.9%). Damodar *et al.* (2015) observed increased sesamum yield up to 90 kg N ha⁻¹ and Vaghani *et al.* (2010) up to 100 kg ha⁻¹. Zeb and Jan (2021) reported significant increase in sesamum yield and yield attributes with increase in nitrogen levels up to 120 kg N ha⁻¹. Accordingly, methods have been recommended for improving the efficiency of fertilizer nitrogen *viz.*, selection of plants with high nitrogen efficiency, adjusting the timing of fertilizer application and developing more efficient fertilizers. Out of this, time of fertilizer application with synchronization of crop nutrient demand is largely important as the maximum potential rate of N uptake is determined by plant growth stage and lower N uptake occurs during the seedling stage and pre-harvest periods, while the vegetative growth and flower initiation demand higher N supply. Gebremariam (2015) reported significant influence of timing of nitrogen application on yield of sesamum. Hence the experiment was initiated to determine the optimum dose and best time of nitrogen application in sesamum in YSR Kadapa district of Andhra Pradesh.

METHODOLOGY

The experiment was conducted in the farm of Agricultural Research Station, Utukur, Kadapa, Andhra Pradesh, India which is situated at 14.45°N latitude and 78.81°E longitude during *rabi* season (December II FN to January I FN) of three years 2020-21, 2021-22 and 2022-23 to study the performance of sesamum under different levels and time of nitrogen application. The soil of the experimental site was red clayey in texture, low in available nitrogen (160 kg ha⁻¹), medium in available phosphorus (30 kg ha⁻¹) and rich in available potash (320 kg ha⁻¹). The treatments included four levels of nitrogen *viz.*, 40, 60, 80 and 100 kg ha⁻¹ and four split application of nitrogen as 50:50 basal and 30 DAS, 25:75 basal and 30 DAS, 25:37.5:37.5 basal, 20 and 40 DAS and 0:50:50 basal, 20 and 40 DAS which were laid out in split plot design with levels of nitrogen in main plots and time of application in sub plots with three replications. The clean seed of popularly grown YLM-66 was sown with a spacing of 30 cm × 10 cm in plot size of 5.0 m × 4.0 m. Recommended dose of phosphorus and potassium (25–25 kg ha⁻¹) were applied through single super phosphate and muriate of potash respectively and nitrogen in the form of urea was applied as per the treatments. At present, the recommended dose of nitrogen for sesamum during *rabi* season was 40 kg ha⁻¹. Two hand weeding were carried out at 20 and 40 DAS. All other agronomic and protective practices were followed as per the recommendations prescribed by ANGRAU. The observations on growth and yield attributes were taken from randomly selected five plants and yield from net plot area and finally subjected to statistical analysis.

RESULTS AND DISCUSSION

Effect of nitrogen

Successive increase in nitrogen from 40 to 100 kg ha⁻¹ influenced the growth parameters, yield attributes and yield of sesamum under irrigated conditions during *rabi* (Table 1). Significantly maximum plant height, number of branches plant⁻¹, number of capsules plant⁻¹ and seeds capsule⁻¹ were recorded with application of 100 kg N ha⁻¹ which was on par with 80 kg N ha⁻¹. Test weight in sesamum was not significantly influenced by the quantity of nitrogen. Significantly maximum seed yield (1355 kg ha⁻¹) was recorded with application of 100 kg N ha⁻¹ which was on par with 80 kg N ha⁻¹ (1299 kg ha⁻¹) (Table 2). Ahmad *et al.* (2018) also reported that an optimum dose of 100 kg N ha⁻¹ in sesamum resulted in increase of seed yield. Increase in nitrogen supply within limits are associated with increase in leaf area, carboxylases and chlorophyll content which in turn determines the photosynthetic activities in leaves and ultimately dry matter production and allocation to various organs of plant improving source sink relation (Maqsood *et al.*, 2016, Babajide and Oyeleke, 2014) which ultimately determines the yield.

Effect of time of application of nitrogen

Application of nitrogen in 25-37.5-37.5 ratio at basal, 20 and 40 DAS recorded significantly maximum plant height, higher number of capsules plant⁻¹ and number of seeds capsule⁻¹, number of branches plant⁻¹, capsule length however, test weight of sesamum was not significantly influenced by the time of nitrogen application (Table 1). Significantly higher seed yield of 1358 kg ha⁻¹ was recorded with application of nitrogen in three splits of 25-37.5-37.5 ratio at basal, 20 and 40 DAS (Table 2). Interaction of nitrogen dose and time of application was found non-significant.

Economics

Application of 100 kg N ha⁻¹ in three splits as 25-37.5-37.5 at basal, 20 and 40 DAS has recorded maximum net returns of Rs. 1,42,762/- per hectare and B:C ratio of 6.77 where as, 80 kg N ha⁻¹ in three splits as 25-37.5-37.5 at basal, 20 and 40 DAS has recorded net returns of Rs. 1,42,300/- per hectare and maximum B:C ratio of 6.81 (Table 3.). With the above economic output it could be concluded that application of 80 kg N ha⁻¹ in three splits as 25-37.5-37.5 at basal, 20 and 40 DAS in sesamum might result in higher B:C ratio.

Conclusion

The number of capsules plant⁻¹, number of seeds capsule⁻¹, seed yield, net returns were higher with application of 100 kg N ha⁻¹ which was on par with 80 kg N ha⁻¹ indicating this as an optimum dose of nitrogen fertilizer and regarding the time of application and number of splits, application of 80 kg N ha⁻¹ in three splits in the ratio of 25-37.5-37.5 at basal, 20 and 40 days after sowing may be recommended for obtaining maximum seed yield in sesamum under irrigated conditions during *rabi*.

References

- Ahmad, J., Ahmad, F., Iqbal, S., Shah, SMA, Ali M., Abbas, M.W., Nawaz, H., Mehmood, Z., Ali B., & Ali, S. (2018). Growth and oil yield of sesamum as influenced by sulphur and nitrogen. *Journal of Agricultural Research*, 3(7), 187.
- Amin, K., & Alireza, S. (2015). The effect of different levels and split application of nitrogen on yield and yield components of sesamum plant in Hamidiyeh weather conditions. *Indian Journal of Fundamental and Applied Life Sciences*, 5(2), 34–40.
- Babajide, P.A., & Oyeleke, O.A. (2014). Evaluation of sesamum (*Sesamum indicum*) for optimum nitrogen requirement under usual farmers' practice of basal organic manuring in the Savanna ecoregion of Nigeria. *Journal of Natural Sciences Research*, 4(17), 122–132.
- Damodar, R.R., Bhale, V.M., & Deshmukh, K.M. (2015). Yield, growth and quality of summer sesamum (*Sesamum indicum* L.) as influenced by irrigation and nitrogen levels. *International Journal of Agricultural Sciences*, 11(2), 301–306.
- El-Sherif, A. (2016). Sesamum (*Sesamum indicum* L.) yield and yield components influenced by nitrogen and foliar micronutrient applications in the Fayoum region, Egypt. *Egypt Journal of Agronomy*, 38(3), 355–367.
- Gebremariam, G. (2015). Growth yield and yield component of sesamum (*Sesamum indicum* L.) as affected by timing of nitrogen application. *Journal of Biology, Agriculture and Healthcare*, 5(5), 165–169.
- Kushawala, S., Singh, A.K., Singh, A., Singh, B.K., & Minz, S.D. (2022). Effect of different levels of sulphur and nitrogen on growth, yield and quality of sesamum (*Sesamum Indicum* L.) under Bael (*Aegle marmelos* L.) based agri-horti system. *Ecology Environment and Conservation*, 28(3), 1404–1410.
- Maqsood, M.A., Awan, U.K., Aziz, T., Arshad, H., Ashraf, N., & Ali, M. (2016). Nitrogen management in calcareous soils: Problems and solutions. *Pakistan Journal of Agricultural Sciences*, 53(1), 79–95.
- Patel, H.K., Patel, R.M., Desai, C.K., Patel, H.B. (2014). Response of summer sesamum (*Sesamum indicum* L.) to different spacings and levels of nitrogen under north Gujarat condition. *International Journal of Agricultural Sciences*, 10(1), 336–343.
- Rosolem, C.A., Ritz, K., Canterella, H., Galdos, M.V., Hawkesford, M., Whalley, R., & Mooney, S.J. (2017). Enhanced plant rooting and crop system management for improved nitrogen use efficiency. *Advances in Agronomy*, 146, 205–239.
- Umar, U.A., Mahmud, M., Abubakar, I.U., Babaji, B.U., & Idris, U.D. (2012). Effect of nitrogen fertilizer level and intra row spacing on growth and yield of sesamum (*Sesamum indicum* L.) varieties. *International Journal of Agronomy and Plant Production*, 3(4), 139-144.
- Vaghani, J.J., Polara, K.B., Chovatia, P.K., Thumar, B.V., & Parmar, K.B. (2010). Effect of nitrogen, potassium and sulphur on yield, quality and yield attributes of *kharif* sesamum (*Sesamum indicum* L.). *Asian Journal of Soil Science*, 5(2), 318–321.
- Zeb, A., & Jan, A. (2021). Response of sesamum (*Sesamum indicum* L.) to sowing methods, nitrogen and sulphur levels. *Sarhad Journal of Agriculture*, 37(1), 278–289.

Table 1. Growth and yield attributes of sesamum as influenced by nitrogen management over three years (pooled)

Treatments	Plant height (cm)	No. of branches/plant	No. of capsules/plant	Capsule length (cm)	No. of seeds/capsule	Test weight (g)
Nitrogen (kg/ha)						
40	92.1	3.2	46.5	2.09	56.1	0.31
60	98.2	3.6	53.6	2.17	60.7	0.32
80	103.8	3.9	58.9	2.41	65.2	0.32
100	112.6	4.2	68.2	2.67	67.4	0.32
S.Em_±	2.69	1.68	3.53	1.41	4.27	0.008
CD (P= 0.05)	6.29	3.93	8.26	3.29	9.99	NS
CV (%)	13.6	9.2	14.3	9.6	15.9	2.6
Time of application						
50-50 (basal, 30 DAS)	97.7	3.5	62.6	2.42	65.3	0.32
25-75 (basal, 30 DAS)	92.3	3.4	55.4	2.26	53.2	0.32
25-37.5-37.5 (basal, 20, 40 DAS)	109.5	3.9	66.5	2.59	69.7	0.33
0-50-50 (basal, 20, 40 DAS)	89.6	3.1	46.0	2.14	45.1	0.31
S.Em_±	2.22	1.94	3.64	1.92	5.23	0.006
CD (P= 0.05)	5.19	4.53	8.51	4.49	12.23	NS
CV (%)	12.6	8.8	14.9	8.3	16.2	1.9
Interaction of Nitrogen dose at Time of application						
S.Em_±	1.89	1.23	2.13	1.06	3.76	0.002
CD (P=0.05)	NS	NS	NS	NS	NS	NS
Interaction of Time of application at Nitrogen dose						
S.Em_±	1.20	1.37	2.27	1.26	4.27	0.004
CD (P=0.05)	NS	NS	NS	NS	NS	NS

Table 2. Seed yield (kg/ha) of sesamum as influenced by nitrogen management over three years

Treatments	2020-21	2021-22	2022-23	Pooled mean
Nitrogen (kg/ha)				
40	1188	1158	1108	1151
60	1248	1288	1206	1247
80	1310	1300	1288	1299
100	1372	1392	1300	1355
S.Em _±	26.8	20.6	62.0	42.6
CD (P= 0.05)	93.6	57.3	175.0	119.7
CV (%)	15.7	12.5	14.9	14.8
Time of application				
50-50 (basal, 30 DAS)	1295	1287	1232	1271
25-75 (basal, 30 DAS)	1223	1207	1101	1177
25-37.5-37.5 (basal, 20, 40 DAS)	1399	1359	1317	1358
0-50-50 (basal, 20 40 DAS)	1205	1078	1059	1114
S.Em _±	26.1	22.7	67.0	37.8
CD (P= 0.05)	76.8	61.5	190.0	106.2
CV (%)	12.6	14.9	13.2	13.1
Interaction of Nitrogen dose at Time of application				
S.Em _±	21.4	24.9	36.2	37.1
CD (P=0.05)	NS	NS	NS	NS
Interaction of Time of application at Nitrogen dose				
S.Em _±	24.9	20.9	25.6	30.6
CD (P=0.05)	NS	NS	NS	NS

Table 3. Economics of sesamum cultivation as influenced by nitrogen management over three years

Treatments	Cost of 'N' fertilizer (Rs./ha)	Cost of 'P' & 'K' fertilizers (Rs./ha)	Total cost of cultivation with fertilizers (Rs./ha)	Seed yield (kg/ha)	Gross returns (Rs./ha)	Net returns (Rs./ha)	B:C ratio
T ₁ : 40 kg N in 50-50	522	3456	23978	1205	144600	120622	6.03
T ₂ : 40 kg N in 25-75	522	3456	23978	1049	125880	101902	5.25
T ₃ : 40 kg N in 25-37.5-37.5	522	3456	23978	1250	150000	126022	6.26
T ₄ : 40 kg N in 0-50-50	522	3456	23978	1100	132000	108022	5.51
T ₁ : 60 kg N in 50-50	780	3456	24236	1196	143520	119284	5.92
T ₂ : 60 kg N in 25-75	780	3456	24236	1186	142320	118084	5.87
T ₃ : 60 kg N in 25-37.5-37.5	780	3456	24236	1401	168120	143884	6.94
T ₄ : 60 kg N in 0-50-50	780	3456	24236	1203	144360	120124	5.96
T ₁ : 80 kg N in 50-50	1044	3456	24500	1326	159120	134620	6.49
T ₂ : 80 kg N in 25-75	1044	3456	24500	1224	146880	122380	6.00
T ₃ : 80 kg N in 25-37.5-37.5	1044	3456	24500	1390	166800	142300	6.81
T ₄ : 80 kg N in 0-50-50	1044	3456	24500	1255	150600	126100	6.15
T ₁ : 100 kg N in 50-50	1302	3456	24758	1375	165000	140242	6.66
T ₂ : 100 kg N in 25-75	1302	3456	24758	1235	148200	123442	5.99
T ₃ : 100 kg N in 25-37.5-37.5	1302	3456	24758	1396	167520	142762	6.77
T ₄ : 100 kg N in 0-50-50	1302	3456	24758	1395	167400	142642	6.76