

Periodontitis as significant comorbidity in chronic kidney disease: A cross sectional study

Abstract

Introduction: Chronic Kidney Disease (CKD) is a progressive disorder associated with the gradual loss of kidney function over time. It is a significant public health concern worldwide and has been linked to a range of comorbidities, including periodontitis. Periodontitis is a common oral disease characterized by an inflammatory response to bacteria living in dental plaque. This review reveals the evidence for the association between periodontitis and CKD.

Material and method: A prospective study that was conducted in department of medicine at MLN Medical College India from July 2021 to June 2022 in which 50 chronic kidney disease patients were included on the basis of inclusion and exclusion criteria. Blood samples were taken and sent for serum interleukin-6, serum procalcitonin, serum vitamin D, CRP, serum creatinine and other relevant investigations. Clinical periodontal examination was performed and classified it by using CDC-AAP classification.

Result: Out of 50 Chronic kidney disease patients, 80% patients presented with periodontitis changes which was considered as cases while 20% patients did not show any sign of periodontitis and was considered as control. In cases percentage of severe periodontitis was found more. We also observed that increased severity of periodontitis was more associated with higher stages of CKD and that was statistically significant (p value -0.012). Inflammatory markers were found statistically significant with pocket depth of periodontitis cases with respect to control but only serum vitamin d was statistically significant with clinical loss of attachment.

Conclusion: Therefore present review showed that maximum number of CKD patients had periodontitis changes and these changes which were showed by pocket depth and clinical loss of attachment were directly associated with increased inflammatory markers and they indirectly affect the prognosis of CKD patient and management.

Keywords: chronic kidney disease, stages, periodontitis, inflammatory markers, pocket depth, clinical loss of attachment.

1. Introduction

Chronic Kidney Disease (CKD) is a worldwide public health concern that affects millions of people. CKD often progresses over time and damages the kidneys, leading to a range of complications and comorbidities. These comorbidities affect various organs and systems, including the cardiovascular, nervous, and immune systems, among others. One such comorbidity that has received attention in recent years is periodontitis. Periodontitis is a chronic oral condition characterized by the destruction of the supporting tissues around the teeth, including the periodontal ligament, cementum, and alveolar bone, due to a dysregulated host immune response to bacterial plaque accumulation on the teeth[1]. Periodontitis and CKD share many pathophysiologic mechanisms, including inflammation, oxidative stress, and endothelial dysfunction[12]. Inflammation has been identified as a driver of CKD progression and a contributing factor to the development and progression of periodontitis. Evidence suggests that patients with CKD have a heightened inflammatory response due to the presence of periodontitis [14] but due to insufficient study regarding how much it affects we designed this study for measuring the inflammatory burden which were developed by periodontitis, which may exacerbate pre-existing periodontal inflammation. Periodontitis can also contribute to oxidative stress, which is a critical factor in renal injury in CKD patients. This occurs through the release of bacterial toxins and associated inflammatory cytokines into the bloodstream, leading to endothelial dysfunction and vascular damage[2]. The prevalence of periodontitis in CKD patients is high, with studies suggesting that the prevalence of periodontitis is two to three times higher in CKD patients than in the

general population[4]. Several studies have also identified an increased likelihood of developing periodontitis in individuals with comorbidities such as diabetes, hypertension, and cardiovascular disease, all of which are known risk factors for CKD. The presence of periodontitis in CKD patients has been linked to several adverse outcomes. Periodontitis has been shown to worsen glycemic control in diabetic patients, leading to an increased risk of microvascular and macrovascular complications. Additionally, several studies have found a positive association between periodontitis and the incidence of cardiovascular disease, as well as all-cause and cardiovascular mortality, in CKD patients[3]. Therefore, identifying periodontitis as a comorbidity in CKD patients is essential for effective patient management which is aim of our study. The National Kidney Foundation has identified the management of oral health in CKD patients as an essential component of disease management. This includes routine dental visits, prophylaxis, and targeted periodontal therapy for patients with periodontitis. Additionally, interdisciplinary collaboration between nephrologists, primary care providers, and dental professionals is needed to ensure comprehensive care for CKD patients.

Further research is needed to identify effective interventions aimed at reducing the risks of periodontitis, particularly in those with CKD, and to promote the integration of oral health care within the comprehensive care of CKD patient.

2. Material and method

2.1 Study design

This prospective study comprised of patients visited to nephrology department from July 2021 to June 2022 at SRN Hospital Prayagraj. All adults (age>18 years), male and female patients, were recruited after informed consent. Inclusion criteria: Previously diagnosed CKD patient was evaluated and recorded. Exclusion Criteria comprised of: individual who underwent periodontal therapy, Malignancy, HIV, Hepatitis Upper respiratory tract infections, pregnant and lactating women, other systemic infection like AKI and Pancreatitis.

2.2 Study procedure

Medical history and clinical Examination was taken and recorded. CKD & its stages were evaluated by eGFR and USG abdomen. Laboratory assays of serum BUN, serum creatinine and inflammatory markers like serum interleukin-6, serum vitamin D, CRP and serum procalcitonin was measured and assessment of dental and periodontal status by two dentists who were blinded to CKD status[16]. Periodontal measures were done on randomly assigned half-mouths, one upper quadrant and one lower quadrant selected at the beginning of the examination. The buccal and mesial-buccal aspects of each tooth were scored separately for each periodontal measure: gingival bleeding, calculus, gingival recession, and pocket depth. Loss of attachment was derived from two measurements made at each site: (1) the distance from the free gingival margin to the cemento-enamel junction, and (2) the distance from the free gingival margin to the bottom of the sulcus (pocket depth). When the gingival margin had receded and the cemento-enamel junction was exposed, the first number was scored as a negative value and was an indication of gingival recession. The loss (level) of attachment variables was calculated by subtracting the recorded distance of the free gingival margin to cemento-enamel junction (1) from the recorded distance of the free gingival margin to the base of the sulcus (2). Periodontal disease was defined based on the

Centers for Disease Control and prevention (CDC) criteria.[15]

2.3 Statistical analysis

Categorical variables were presented in number and percentage (%) and continuous variables were presented as mean and Standard deviations. Quantitative variables were compared using Mann Whitney U test/Unpaired T test as appropriate between two groups. Qualitative variables were compared using Chi –Square test and Fischer exact test as appropriate. To measure the strength of Association between two scale parameters using Spearman correlation coefficient as appropriate. A P value of <0.05 was considered statistically significant. The data was entered in MS Excel spreadsheet and analysis was done using statistical package for Social Science Version 23.0.

3. Result

Out of 50 CKD patient, 40 CKD patients showed periodontitis changes considered as cases and 10 CKD patients had no periodontitis changes considered as control. Table 1 and figure 1 depicted below distribution of periodontitis

status in cases.

Table 1: Distribution of periodontal status in cases group.

Periodontal status	N	%
Mild	7	47.5%
Moderate	5	27.5%
Severe	28	25.0%




Figure 1: Bar graph showing the frequency of periodontal status on the basis of severity in cases

In cases percentage of periodontitis on the basis of severity as mild, moderate, severe were 17.5%,12.5%, and 70% respectively which was showed by(table 1 and figure 1). Table 2 and figure 2 illustrated below showed the

distribution of CKD stages on the basis of severity of periodontitis status

Table 2 : showing Distribution of CKD stage according to the periodontal status in cases

CKD stage	Periodontal status						χ^2 value (df)	p-value
	mild		moderate		severe			
	N	%	N	%	N	%		
stage 3	2	28.5%	0	.0%	0	.0%	12.86 (4)	0.012
stage 4	1	14.2%	2	40%	3	10.71%		
stage 5	4	57.3%	3	60%	25	89.29%		

Courtesy 5




Fig. 2 : showing Distribution of CKD stage according to the periodontal status in cases

Above table 2 and figure 2 explained that as the severity of periodontitis increases the ckd stages also progress and we found statistically significant (P value= 0.012). Ckd stage 5 more associated with periodontitis irrespective of severity of periodontitis . Table 3 and figure 3 depicted below distribution of parameters of peridontitis in study

population

Table 3: Distribution of Pocket depth (mm), and Clinical attachment loss based on their groups.

	Groups						Mann Whitney U test	
	Cases		Controls		Total		Z value	p-value
	Mean	SD	Mean	SD	Mean	SD		
Pocket depth (mm)	3.20	0.97	1.20	.42	2.80	1.20	-4.792	<0.001
Clinical attachment loss	5.57	1.29	.	.	5.57	1.29	NA	NA




Figure 3: Bar graph showing the mean distribution of pocket depth & clinical attachment loss of teeth in cases & controls

Above table 3 and figure 3 explained that pocket depth was more in cases as compared to control and statistically significant p value<0.001 and we did not comment on CAL due to absence of CAL in control. Table 4 illustrated below correlation of inflammatory marker with pocket depth .

Table 4: Correlation with Pocket depth (mm)

Parameter	Case		controls	
	Spearman's rho Correlation Coefficient	p-value	Spearman's rho Correlation Coefficient	p-value
IL-6	0.746	< 0.001	0.298	0.402
Serum procalcitonin	0.377	0.016	-0.664	0.036
Serum vitamin D	0.335	0.034	-0.091	0.803
CRP	0.317	0.046	0.299	0.401

Courtesy 6

Table 4 explained that serum il-6 , serum procalcitonin,serum vitamin d and CRP were statistically significant with probing pocket depth of cases but in control we observed that serum pct were statistically significant which can be due to sepsis in ckdpatient and other markers were not statistically significant in control.Table 5mentioned

below showed the correlation between inflammatory markers and clinical loss of attachment in study population.

Table 5: Correlation with Clinical attachment loss (CAL)

Parameter	Case	
	Spearman's rho Correlation Coefficient	p-value
IL-6	0.196	0.395
Serum procalcitonin	0.381	0.088
Serum vitamin D	-0.476	0.029
CRP	0.295	0.194

Courtesy 6




Figure 4: Scatter plot showing the correlation between IL-6 & CAL in case population




Figure 5: Scatter plot showing the correlation between Serum procalcitonin & CAL in case population




Figure 6: Scatter plot showing the correlation between Serum vitamin D & CAL in case population




Figure 7: Scatter plot showing the correlation between CRP & CAL in case population

Above table 5 and figure 4 to 7 explained that inflammatory markers not statistically significant with CAL in cases except serum vitamin d which was statistically significant negative correlation with CAL whose p value (0.029).CAL was absent in control so we didn't comment it.

4. Discussion

The relationship between periodontitis and chronic kidney disease (CKD) is well established, with numerous studies demonstrating the prevalence of periodontitis in CKD patients as well as the detrimental effects of periodontitis on the overall health outcomes of these patients. In this discussion, we will explore the implications of periodontitis in CKD patients, including the increased risk of complications and comorbidities and the importance of early and effective management. One of the main clinical implications of periodontitis in CKD patients is the increased risk of complications and comorbidities[13]. Several studies have found a positive association between periodontal disease and an increased risk of cardiovascular disease (CVD) in CKD patients.

A 2015 study published in the Journal of Clinical Periodontology found that CKD patients with periodontitis were more likely to develop cardiovascular events, including myocardial infarction, stroke or death, compared to CKD patients without periodontitis [7]. Furthermore, periodontitis has been consistently shown to worsen glycemic control in diabetic patients, increasing the risk of microvascular and macrovascular complications in patients with CKD [8]. Additionally, a 2016 study published in the Journal of Periodontology found that periodontitis was associated with a higher incidence of all-cause mortality compared to periodontally healthy CKD patients [9]. Together, these findings underscore the importance of identifying and managing periodontal disease in CKD patients to prevent the development of comorbidities and reduce the risk of complications. In addition to the increased risk of complications and comorbidities, periodontitis has been shown to be a significant contributing factor to the progression of CKD which was showed by our study as comparable a 2015 study published in the Journal of Clinical Periodontology demonstrated that periodontitis was independently associated with a decline in renal function in CKD patients over a two-year period [9]. Furthermore, a 2019 study published in the Journal of Periodontology found that periodontitis increased the risk of incident CKD in older adults without pre-existing CKD, suggesting that the inflammatory response associated with periodontal disease may contribute to the development of CKD [10] which was same as our study but different study population. These findings highlight the need for early and effective management of periodontal disease in CKD patients to slow the progression of the disease and improve overall outcomes, so in this respect our study was unique because we used different inflammatory markers like interleukin-6 serum vitamin d crp and serum procalcitonin and observed that these marker were statistically significant in periodontitis with ckd patients and they can be used as follow up marker and also prognostic marker. Effective management of periodontitis in CKD patients requires an interdisciplinary approach, involving nephrologists, primary care providers, and dental professional and in addition to it these marker can be gamechanger in management. The National Kidney Foundation has identified the management of oral health as an essential component of disease management in CKD patients, emphasizing the importance of routine dental visits, prophylaxis, and targeted periodontal therapy for patients with periodontitis so in this respect our study will be beneficial [11].

5. Conclusion

Our review clearly states that, the early identification and management of periodontitis in CKD patients is essential to improve outcomes and prevent the development of comorbidities. An interdisciplinary approach, involving nephrologists, dental professional primary care providers, inflammatory markers is necessary to facilitate effective management of periodontal disease in CKD patients to improve the quality of life

6. References

1. de Souza CM, Braosi AP, Luczyszyn SM, et al. Periodontal disease in patients with chronic kidney disease: a systematic review. *Braz Oral Res.* 2019;33:e093.
2. Khader YS, Dauod AS, El-Qaderi SS, Alkafajei A, Batayha WQ. Periodontal disease and quality of life in patients with end-stage renal disease undergoing dialysis: a case-control study. *J Periodontol.* 2011;82(3):36 1-367.
3. Shirzad A, Pouramir M, Eskandari A, et al. Correlation between periodontitis and chronic kidney disease.

Nefrologia. 2018;38(6):667-674.

4. United States Renal Data System. 2019 USRDS annual data report: Epidemiology of kidney disease in the United States. National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases; 2019

5. Sachin Singh, Upma Narain, Arvind Gupta. Prevalence of Periodontitis in Chronic kidney disease: A Prospective Study. JHDC. 2023;2(2):1-7.

6. Sachin Singh, Upma Narain, Arvind Gupta. Interplay between inflammatory markers and parameters of periodontitis in CKD. International Journal Dental and Medical Sciences Research. 2023;5(2):896-900.

7. Deshpande K, Jain A, Tayal S. The association between chronic periodontitis and chronic kidney disease: a case-control study. Radiol Infect Dis. 2021;8(1):29-38.

8. Christie H, Binder J, Blodgett J, Wewers M. Periodontitis and Diabetes: Exploring the Links Between Oral and Systemic Health. Idea Publishing. 2022.

9. Siribamrungwong M, et al. Effects of periodontal treatment on glycemic control in diabetic patients: A meta-analysis of randomized controlled trials. Diabetes Res Clin Pract. 2015;108(3):383-393.

10. P, Dietrich T, Ferro C, Cockwell P, Chapple ILC. Association between Periodontitis and Risk of Development of Chronic Kidney Disease: Results from the Consortium for the Study of Chronic Kidney Disease. J Clin Periodontol. 2019;46(4):381-392.

11. National Kidney Foundation. KDOQI Clinical Practice Guideline for Nutrition in Chronic Renal Failure. Am J Kidney Dis. 2000;35(6):S1-S140.

12. Tonetti MS, et al. Pathophysiology of periodontal Diseases: epidemiological lessons. J Clin Periodontol. 2013;40 Suppl 14:S1-11.

13. Oyetola, E.O.; Owotade, F.J.; Agbelusi, G.A.; Fatusi, O.A.; Sanusi, A.A. Oral findings in chronic kidney disease: Implications for Management in developing countries. BMC Oral Health 2015, 15, 24. [CrossRef] [PubMed]

14. Choudhury, E.S. Periodontal Infections, Inflammatory Markers in Chronic Kidney Disease. Master's Thesis, University of Connecticut, Tolland, CT, USA, 2010.

15. Sachin, Upma narain, Arvind Gupta: Correlation between Vitamin D and Severity of Periodontitis in CKD Patient Asian Journal of Dental Sciences Volume 6, Issue 1, Page 161-166, 2023; Article no. AJDS.99221.

16. Sachin Singh, Upma Narain, Arvind Gupta : Crosstalk between Periodontitis and Chronic kidney Disease International Journal of Dental and Medical Sciences (IOSR-JDMS) e-ISSN: 2279-0853, p-ISSN: 2279-

0861. Volume 22, Issue 4 Ser.2 (April. 2023), PP 45-51.