

Original Research Article

Standardization of Grafting Techniques: Unlocking the Potential of Pomegranate (*Punica granatum* Linn.)

ABSTRACT

An Experiment was conducted at Department of Agriculture, School of Agriculture and Veterinary Science, Shridhar University, Pilani, Rajasthan, India during 2023-2024. Softwood grafting and (T₁ on June 15; T₂ on June 30; T₃ on July 15; T₄ on July 30; T₅ on August 15; T₆ on August 30), tongue grafting (T₇ on June 15; T₈ on June 30; and T₉ on July 15, T₁₀ on 30th July, T₁₁ on August 15, T₁₂ on August 30) these were the 12 grafting techniques that used as treatments in the study. For this study, one-year-old Ganesh seedlings were utilized as the rootstock. Experiment T₃, which entailed grafting softwood on July 15th, had the highest outcomes in terms of early bud break (15.16 days), improved sprouting percentage (57.32%) at 150 DAG, and grafting success (74%), at 60 DAG. When softwood and whip & tongue grafting methods were compared, softwood grafting produced better results in terms of bud break, sprouting percentage, and grafting success.

Keywords: Pomegranate, grafting methods, graft success, grafting time, softwood grafting, tongue grafting

INTRODUCTION

The pomegranate, *Punica granatum* (L.), commonly referred to as the Chinese apple, is a plant belonging to the Punicaceae family with chromosomes no. $2n=18$. One of the most revered and ancient beloved fruits. A common term for it is "friendship fruit." Customers all across the world seek for this well-known commercial fruit because of its exceptional combination, **super** dessert quality, and sweet-acid flavor. Due to its robust nature, pomegranates may thrive in a broad range of climates, including tropical, subtropical, desert, and temperate ones. The world's largest producer of pomegranates is India. There are over 760 varieties of pomegranate. Each pomegranate has about 840 seeds. One of the richest and oldest fruits is the pomegranate. It is grown commercially in the state of Maharashtra, Gujrat, Karnataka, Andhra Pradesh, Tamil Nadu, Madhya Pradesh and Rajasthan **and** the 1.79% of India's entire production is exported. It has enormous medicinal, nutritional value and one of the richest sources of anti-oxidant.

Currently, 275,500 hectares are planted with this fruit crop, which yields 3 million metric tons (MT) of fruit year (NHB statistics 2021-22). With over 75% of the country's pomegranate output and area, Maharashtra leads the field, followed by Andhra Pradesh and Karnataka. The most common propagation strategy for pomegranates is air layering. Examining the entire matter, a task was posed to evaluate pomegranate vegetative proliferation using grafting (softwood and whip/tongue grafting).

MATERIALS AND METHODS

The current study was conducted in 2023–24 at the Department of Agriculture (experimental field), School of Agriculture and Veterinary Science, Shridhar University, Pilani, Rajasthan, India. The study was conducted in the IV agro-climatic zones of Rajasthan (Transitional Plain of Inland Drainage), which are located at an altitude of 533 meters above mean sea level and at 28° 27 North latitude and 75° 63 East longitude. In the current study, 12 treatment combinations totaling 2 grafting methods tongue grafting and wedge grafting were included at every fortnightly interval throughout the months of June, July, and August. The treatments were set up in a totally randomized design with three replications. About 15 cm long scion of 'Bhagawa' was used for grafting. A one-year-old seedling of Ganesh were used. Rootstocks of pencil thickness (0.6-0.8 cm diameter) having uniform growth were selected. The rootstock was headed back when the scion was chosen, leaving a 15–17 cm long stem above the polythene bag. The beheaded rootstock was split open around 2.5-5.0 cm deep through the center from the cut end of the rootstock with a grafting knife. A wedge-shaped cut, slanting from both the sides (5 cm long), was made on the proximal end. The scion was inserted into the stock and pressed it so that cambium tissue of the stock and scion both are together come into contact. Then, the stock and scion combination were tied with the help of a 150-gauge polythene strip, 1.5 cm in width and 30-40 cm in length. Immediately after grafting, the graft was covered by a 2.5×16 cm long white polythene cap that was tied with thread at the lower end. After grafting, the percentage of sprouting was noted at 30days, 60days, 90days, 120days, and 150 days. The percentage of success was measured at 60 days after grafting, and the length of the scion was measured at 30days, 60days, 90days, 120days, and 150 days. The Panse and Sukhatme (1978) protocol were followed in doing the statistical analysis of the data.

RESULT & DISCUSSION

Out of all the treatments, T₃ (softwood grafting) had the highest sprouting percentage (57.32%) at 150 DAG, whereas T₉ (whip and tongue grafting) had the lowest sprouting percentage (11.4%) at 150 Days after grafting (Table 1). In terms of grafting success percentage, the softwood grafting (T₃) on July 15th had the highest grafting success percentage (74%) 60 days after grafting (Table 2). On July 15, whip and tongue grafting in T₉ had the lowest grafting success rate (11.49%) (Table 2). The maximum shoot height (in cm) in T₃ at 150 DAG was measured at 56.00 cm. However, T₅ (softwood grafting on August 15) reported the largest shoot length (24.64 cm) at 60 DAG. In treatments including whip and tongue grafting, the minimum shoot length was noted (Table 3.) On July 30, at 150 DAG, the highest scion diameter (5.82 mm) was measured in softwood grafting (T₄), and on August 30, in whip and tongue grafting (T₁₂), the lowest scion diameter (5.9mm) was measured (Table 4).

Notably, T₃ (softwood grafting) showed the highest proportion of pomegranate graft sprouting (57.32%), whereas tongue and whip grafting showed lower percentages of sprouting (11.4%) at 150 DAG (Dubey et al., 2022; Das et al., 2024). The results show that the percentage of pomegranate grafts that sprouted was significantly influenced by the procedure and season. The extreme humidity and temperatures that were present at the time might account for this. Graft uptake was significantly impacted by the high humidity and warmth (Hartman and Kester, 1986). The results indicate that the percentage of pomegranate grafts that sprouted was significantly influenced by the procedure and season (Raj et al., 2024). The extreme humidity and temperatures that were present at the time might account for this. The high humidity and warmth had a major effect on graft absorption (Hartman and Kester, 1986; Patil et al., 2010; Sonawane et al., 2012) have confirmed the results of this study. It is plausible that incorrect interlocking of the scion and stocks causes less intimate contact between the two, preventing callus formation and reducing the sprouting percentage in tongue and whip grafting. A parallel result was defined by Hussain et al. (2016). The percentage of graft success (%) was highly impacted by the grafting method.

In contrast, T₉ had the lowest grafting success (%) of any species when whip and tongue grafting were used. The two primary factors that have a greater influence on the formation of graft unions are the optimal temperature and relative humidity, and they are to blame. In July,

higher humidity helps to keep the scion from desiccating, which keeps the cells turgid and facilitates the creation of a callus between the scion and stock quickly. Similar to what Ghosh et al. (2010) observed, Nitish et al. (2019) saw increased graft success rates in July (72%) for softwood grafting in sapota. The highest grafting success (63.33%) in softwood grafted plants in sapota was reported by (Patil et al., 2010; Patil, 2019) also observed that the month of July (80.00%) had a higher grafting success percentage for softwood grafting. At 150 DAG, T₃ had a significant increase in shoot length (cm) of 56.00, whereas T₁₁ had the lowest shoot length (44.15). Table 3 makes it clear that, in contrast to whip/tongue grafted plants, softwood grafted plants produced the longest shoots. It might be the result of increased warmth and humidity within the shade net, which raises cell metabolism and causes the scion's development to extend. Nitish et al. (2019) in sapota found the highest growth of scion in softwood grafting method performed in July month (Dubey et al., 2021; Sunil et al., 2023). There is a correlation between those research' findings and ours. Softwood grafting unquestionably provided the highest scion diameter (5.82 cm), as compared to whip and tongue grafting (5.9 cm) (Table 4). It occurred because softwood grafted plants have more leaves and early development, which leads to a larger accumulation of carbohydrates. Additionally, the humidity and temperature in July were favorable for the grafts to flourish. The results of this study are in line with those of Patel et al. (2010), who examined the highest scion diameter of softwood grafted plants in July while doing research in Mandarin.

Table 1: Sprouting percentage at various days of grafting

Treatments	Sprouting percentage (%) at DAG				
	30	60	90	120	150
T ₁	24.00 (29.00)	32.32 (34.16)	28.16 (31.58)	28.15 (31.58)	24.00 (28.48)
T ₂	32.32 (34.16)	32.32 (34.16)	32.32 (34.16)	32.32 (34.16)	32.32 (34.16)
T ₃	53.16 (46.40)	74.00 (59.53)	61.49 (51.23)	57.32 (48.81)	57.32 (48.82)

T ₄	11.49 (19.69)	24.00 (29.00)	24.00 (29.00)	24.00 (29.00)	24.00 (29.00)
T ₅	36.49 (36.75)	36.49 (36.75)	36.49 (36.75)	36.49 (36.75)	36.49 (36.75)
T ₆	49.00 (44.00)	53.16 (46.40)	49.00 (44.00)	44.82 (41.57)	44.82 (41.58)
T ₇	0.00	0.00	0.00	0.00	0.00
T ₈	0.00	0.00	0.00	0.00	0.00
T ₁₀	0.00	0.00	0.00	0.00	0.00
T ₁₁	40.67 (39.16)	53.16 (46.40)	53.16 (46.40)	53.16 (46.40)	53.16 (46.46)
T ₁₂	40.66 (39.16)	36.49 (36.75)	32.32 (34.16)	32.32 (34.16)	32.32 (34.16)
S. E_±	1.42	2.11	1.83	1.79	2.12
C.D 5%	4.17	6.21	5.38	5.26	6.24

Table 2: Graft success percentage at 60 days after grafting

Treatments	Percentage of graft success
T ₁	32.32 (34.16)
T ₂	32.32 (34.16)
T ₃	74.00 (59.53)
T ₄	24.00 (29.00)
T ₅	36.49 (36.75)
T ₆	53.16 (46.40)
T ₇	0.00 (0.00)

T ₈	0.00 (0.00)
T ₉	11.49 (19.69)
T ₁₀	0.00 (0.00)
T ₁₁	53.16 (46.40)
T ₁₂	36.49 (36.75)
S. E\pm	2.11
C.D 5%	5.21

Table 3: Impact of grafting on length of shoot at various days of grafting

Treatment	Shoot length (cm) at DAG				
	30	60	90	120	150
T ₁	15.61	20.00	32.42	38.33	47.14
T ₂	14.35	18.80	29.72	38.65	48.14
T ₃	16.57	20.49	40.19	44.19	56.00
T ₄	16.29	20.24	28.84	35.17	51.52
T ₅	11.77	24.65	27.08	33.15	44.21
T ₆	14.75	21.41	27.58	43.61	49.75
T ₇	0.00	0.00	0.00	0.00	0.00
T ₈	0.00	0.00	0.00	0.00	0.00

T ₉	11.72	17.52	26.00	30.79	42.06
T ₁₀	0.00	0.00	0.00	0.00	0.00
T ₁₁	8.92	19.81	28.86	35.23	44.15
T ₁₂	14.72	19.74	24.65	34.22	47.84
S. E_±	0.69	1.00	1.27	2.15	2.24
C.D 5%	2.06	2.92	3.75	6.33	6.59

Table 4: Impact of grafting on scion diameter at various days after grafting (DAG)

Treatment	Scion diameter (mm) at DAG				
	30	60	90	120	150
T ₁	4.62	4.82	5.06	5.32	5.44
T ₂	4.71	5.08	5.32	5.45	5.76
T ₃	5.17	5.21	5.28	5.46	5.55
T ₄	5.73	5.37	5.44	5.81	5.82
T ₅	4.65	4.92	5.28	5.43	5.65
T ₆	4.68	4.77	4.97	5.15	5.22
T ₇	0.00	0.00	0.00	0.00	0.00

T ₈	0.00	0.00	0.00	0.00	0.00
T ₉	5.07	5.18	5.26	5.37	5.46
T ₁₀	0.00	0.00	0.00	0.00	0.00
T ₁₁	4.53	4.73	4.79	5.01	5.41
T ₁₂	4.57	4.47	5.05	5.02	5.9
S. E\pm	0.12	0.18	0.06	0.9	0.10
C.D 5%	0.37	0.54	0.18	0.29	0.31

DAG - Days after grafting

CONCLUSION

In summary according to the current research, grafting might be used to propagate pomegranates. Based on all the parameters examined, the softwood grafting that was done on July 15th produced the best results under shade net conditions. As a result, softwood grafting may be employed commercially by employing rootstocks resistant to different serious diseases, such as pomegranate wilt.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version,

model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1. The authors declare that they have not used any AI technology for the writing, editing and for the analysis of data for this manuscript.
2. The authors declare that they have not used any AI technology for the writing, editing and for the analysis of data for this manuscript.
3. The authors declare that they have not used any AI technology for the writing, editing and for the analysis of data for this manuscript.

REFERENCES

- Chandra R, Jadhav VT, Sharma J, Marathe RA. Effect of grafting methods and time on scion sprouting, graft success and subsequent growth of grafted pomegranate (*Punica granatum* L.) 'Bhagawa'. *Acta Hort.* 2011; 890:83-86.
- Das A, Kadawla K, Nath H, Chakraborty S, Ali H, Singh S, Dubey VK. Drone-Based Intelligent Spraying of Pesticides: Current Challenges and Its Future Prospects. In *Applications of Computer Vision and Drone Technology in Agriculture 4.0*. Springer Nature Singapore; 2024;199-223.
- Dubey VK, Kalleshwaraswamy CM, Joshi S, Shivanna, BK. 2022. Diversity and Diagnostics of Sternorrhynchan Insect Pests Infesting Arecanut. *Indian J Entomol.* 2022;84(3):509-515.
- Dubey VK, Kalleshwaraswamy CM, Shivanna BK. Seasonal incidence of major sternorrhynchan insect pests infesting arecanut in South India. *Indian J Agric. Res.* 2021;47(5):436-440.
- Ghosh SN, Bera B, Roy S, Banik BC. Effect of cultivars and season on grafting success in sapota under Paschim Midnapur conditions of West Bengal. *J Hort. Sci.* 2010; 5(2):138-139.
- Gehlot, T., Mishra, S. R., Yadav, A. K., & Chaudhary, P. (2024). Studies on Crop Weather Calendar of Brinjal Crop in Eastern Uttar Pradesh, India. *Journal of Experimental Agriculture International*, 46(6), 816-823.
- Hartmann HS, Kester DE. *Plant propagation*. Prentice Hall of India, Pvt. Ltd. New Delhi. India, 1986, pp. 399-447.
- Hussain S, Mir MM, Bhat R, Wani SA, Shameem R. Standardization of propagation techniques in chestnut (*Castanea sativa* Mill.). *Eco. Env. And Cons.* 2016; 22(2):911-915.

- Jamoh, O. (2021). Study of standardization of grafting techniques in pomegranate (*Punica granatum L.*) in Northern Dry zone of Karnataka.
- Kar PL, Singh RP, Chadha TR. Studies on the standardization of top working technique in wild pomegranate trees. *Hort. J.* 1989; (1):68-70.
- Karimi HR, Homayoun F. study of pomegranate (*Punica granatum L.*) propagation using Bench Grafting. *J of fruit and ornamental plant res.* 2011; 19(2):67-72.
- Karimi HR, Nowrozy M. Effect of rootstock and scion on graft success and vegetative parameters of pomegranate. *Scientia Horti.* 2017; 214:280-287.
- Karimi R. Stenting (cutting & grafting): A new technique for propagating pomegranate (*Punica granatum L.*). *J, Fruit Ornam, Plant Res.* 2011; 19:73-79.
- Nitish HT, Murthy V, Goudappanavar PB, Deeksha RN. Standardization of softwood grafting techniques in sapota (*Manilkara achras L.*) on invigorated khirni rootstock under polyhouse and shade net conditions. *International J Chemical studies.* 2019; 7(2):2079-2081.
- Padeshetti, B., Kulapati, K. H., Gollagi, S. G., Pattepur, S. and Jamoh, O. (2020). Effect of grafting methods on physiology, time on Bud break, scion sprouting and graft success of pomegranate (*Punica granatum L.*) Cv. Kesar. *Journal of Pharmacognosy and Phytochemistry*, 9(2), 297-299.
- Patil SN, Suryawanshi AB, Phad GN. Effect of season of grafting on percent graft-take and growth of scion shoot of sapota on khirni rootstock. *International J Pl. Sci.* 2010; 5(1):6-9.
- Raj M, Lal K, Satdev, Kumari P, Kumari S, Dubey VK, Kumar S. Potential nutrient cycling and management in agroforestry. In *Agroforestry to Combat Global Challenges: Current Prospects and Future Challenges.* Springer Nature Singapore; 2024;71-92.
- Sonawane GR, Khandekar RG, Korake GN, Haldankar PM, Mali PC. Effect of season on softwood grafting in carambola (*Averrhoa carambola L.*). *The Asian J Hort.* 2012; 7(2):412-415.
- Sunil V, Majeed W, Chowdhury S, Riaz A, Shakoori FR, Tahir M, Dubey VK. Insect Population Dynamics and Climate Change. In *Climate Change and Insect Biodiversity.* CRC Press; 2023;121-146.
- Vazifeshenas M, Khayyat M, Jamalian S. Effects of different scion rootstocks combinations on vigour, tree size, and yield and fruit quality on three Iranian cultivars of pomegranate. *Acta Hort.* 2009; 463:143-152.