

Use of Fern Leaf Extract (*Diplazium Esculentum*) in Raw Fish Feed on Growth and Feed Conversion Ratio in Mangrove Crab (*Scylla serrata*)

ABSTRACT

Mangrove crab or known as *Scylla* sp. is one of the biota whose habitat is in coastal waters, especially in mangrove forest areas. Currently, the need for mangrove crabs still largely relies on capture in nature which is fluctuating. It is necessary to carry out mangrove crab cultivation in a controlled manner that can support effective, efficient and economically profitable mangrove crab cultivation efforts. In addition, cultivation can increase mangrove crab production to meet domestic and foreign market demands. An alternative that can be chosen for mangrove crab cultivation is an apartment system that is arranged vertically in the form of a storage rack. The water medium used in this apartment system cultivation uses a recirculation system. One of the efforts to accelerate growth in crabs is by using steroid hormones. Ecdisteroid is the main steroid hormone that has the main function as a molting hormone, besides that it also regulates physiological functions, such as growth, metamorphosis, and reproduction. Fern leaves are one type of fern plant that contains the hormone ecdisteroid, the hormone is useful for accelerating molting of mangrove crabs. Therefore, this study was conducted to determine the effect of using fern leaf extract (*Diplazium esculentum*) in low fish feed with different doses on growth and feed conversion ratio in mud crabs. This research method uses a field experimental method with a complete randomized design (CRD) consisting of 5 treatments and 3 replicates with different dosing percentages, namely: PA : Percentage of 0 ml/Kg, PB: Percentage of 100 ml/Kg, PC: Percentage of 125 ml/Kg, PD: Percentage of 150 ml/Kg, PE: Percentage of 175 ml/Kg. The results of the research that has been done that the use of fern leaf extract in mangrove crab (*Scylla serrata*) feed has a significant effect on the growth of absolute weight, Absolute Length, SGR (Specific Growth Rate) and FCR (Feed Conversion Ratio) where the most influential treatment is treatment E, namely the highest dose of 175 ml / kg. This shows that treatment E is the best dose treatment for the growth and feed conversion ratio of mangrove crabs (*Scylla serrata*).

Kata kunci: mangrove crab (*Scylla serrata*), Leaf extract (*Diplazium esculentum*), apartmentSystem.

1. INTRODUCTION

Mangrove crab or known as *Scylla* sp. is one of the biota whose habitat is in coastal waters, especially in mangrove forest areas. Currently, the demand for mangrove crabs mostly relies on capture in nature, which fluctuates. The current world production of *Scylla* sp. is mostly derived from mangrove crab catches in the world in 2011 reaching 44,670 tons but began to decline to 38,055 tons in 2013. The reduction in catches also affected the number of Indonesian mangrove crab exports which decreased from 34,172 tons in 2013 to 28,080 tons in 2014 [1].

It is necessary to carry out mangrove crab cultivation in a controlled manner that can support an effective, efficient and economically profitable mangrove crab farming business[2]. In addition, cultivation can increase mangrove crab production to meet domestic and foreign market demands. An alternative that can be chosen for mangrove crab cultivation is an apartment system that is vertically arranged in the form of a storage rack. The water medium used in this apartment system cultivation uses a recirculation system[3]

One of the efforts to accelerate growth in crabs is by using steroid hormones. Ecdysteroid is the main steroid hormone that has the main function as a molting hormone, besides that it also regulates physiological functions, such as growth, metamorphosis, and reproduction [2]. Fern leaves are one type of fern plant that contains the hormone ecdysteroid, which is useful for accelerating molting of mangrove crabs [4]. Therefore, this study was conducted to determine the effect of using fern leaf extract (*Diplazium esculentum*) in raw fish feed with different doses on growth and feed conversion ratio in mud crabs.

2. RESEARCH METHODS

This research was conducted in June-July 2024 at Empol Preparation Village, Sekotong Tengah, Sekotong District, West Lombok Regency. This research method uses a field experimental method with a complete randomized design (CRD) consisting of 5 treatments and 3 replications with different dosing percentages, namely:

1. PA : Persentase 0 ml/Kg
2. PB : Persentase 100 ml/Kg
3. PC : Persentase 125 ml/Kg
4. PD : Persentase 150 ml/Kg
5. PE : Persentase 175 ml/Kg

The five treatments were repeated 3 times, resulting in 15 experimental units. Each experimental unit was then invited into a crab apartment unit. The construction of the research layout design can be seen in Figure 1.




Fig 1. Experimental designs

2.1 Research Procedure

The containers used in the study were apartments assembled using jirgens arranged vertically in the shape of a storage rack. Before stocking, it should be cleaned using soap and rinsed with fresh water. The apartment will be filled with seawater that has been diluted with fresh water so that it becomes brackish water with a salinity of 22-25 ppt. The maintenance container is filled with water to a height of 3-5 cm. Subsequently, the rearing container is immersed for 1 day.

The test animals are mangrove crabs (*Scylla serrata*) which are selected before stocking. Furthermore, the crabs that have been stocked will be acclimatized for 2-3 days. The crabs used had an average initial weight of 30-40 g/head, which was obtained from the catch of the community in the village of Empol Preparation, Central Sekotong, Sekotong Sub-district, West Lombok Regency. The fern leaves used are old fern leaves and are not eaten (not consumed). Then washed and air dried, then dried in the hot sun 3-4 days and then ground until smooth. Extraction was carried out at the Basic Chemistry Laboratory, Faculty of Mathematics and Natural Sciences, Mataram University.

The feed used is kurisi (*Nemipterus*) mixed with fern leaf extract. Before mixing, the fish was first cut into smaller sizes. Then given the addition of fern leaf extract with as much as 0 ml / kg, 100 ml / kg, 125 ml / l, 150ml / kg, 175ml / kg according to the treatment dose. Feed that has been mixed with fern leaf extract is stirred until homogeneous. Feeding is given once a day at 17:00 WITA. Measurement of absolute weight and length of mangrove crabs is done every 1 week. To measure the weight using analytical scales. Meanwhile, to measure the absolute length of the crab, the crab used a vernier caliper. Then water quality observations were made, namely measuring physical parameters such as temperature, chemical parameters such as pH, salinity and dissolved oxygen (DO) which were carried out once a week except for DO parameters carried out at the beginning and end of maintenance.

2.2 Research parameters

Parameters measured in this study include absolute weight growth, absolute length measurement, survival rate (SR), daily growth rate (SGR), feed utilization efficiency (EPP), feed conversion ratio (FCR), blood profile and data analysis.

Absolute body weight growth is measured by the formula $W_m = W_t - W_o$. W_m = Absolute weight growth (gr); W_t = Final average weight (gr); W_o = Initial average weight (gr). [5]

Percentage of survival with the formula $SR = N_t / N_o \times 100\%$; SR = Survival rate (%); N_t = Number of mud crabs at the end of rearing (fish); N_o = Number of mud crabs at the beginning of rearing (fish) [6]

Specific growth rate formula $SGR = \ln W_t - \ln W_o / T \times 100\%$; SGR = Daily growth rate (%); W_o = Weight of test animals at the beginning of the study (g); W_t = Weight of test animals at the end of the study (g); T = Study time (days)[7]

Feed efficiency is calculated based on the formula $EP = (W_t - W_o) / F \times 100\%$; EP = Feed efficiency; W_t = Weight of test animals at the end of the study (g) W_o = Weight of test animals at the beginning of the study (g) F = Feed given (g) [7]

Food Conversion Ratio (FCR) refers to the formula $FCR = F / (W_t - W_o)$; FCR = Feed Conversion Ratio; F = Amount of feed consumed (g); W_t = Weight of test animals at the end of the study (g); W_o = Weight of test animals at the beginning of the study (g)[8]

The hemocyte assay can be calculated using the formula Hemocyte Cell Type Percentage = Number of Each Hemocyte Cell / Total Hemocytes x 100% [9]

2.2 Data Analysis

The data obtained from this study are absolute weight growth, absolute length measurement, daily growth rate (SGR), feed utilization efficiency (EPP), feed conversion ratio (FCR) and blood profile. Will be analyzed using Analysis of Variance (ANOVA) with 95% confidence level. Furthermore, it is continued with the Ducau test and homogeneity test if significantly different results are obtained. While water quality data is presented descriptively.

3. RESULTS AND DISCUSSION

3.1 Absolute Weight Growth

The research was conducted for 43 days in Empol Preparation Village, Central Sekotong, Sekotong District, West Lombok Regency. Feeding the crab with fern leaf extract at different doses showed that the average absolute weight of the crab obtained ranged from 5.37%-11.62% as can be seen in Figure 2.




Figure 2. Absolute Weight Chart

Based on the results of observations on the growth parameters of absolute weight, the highest was produced from treatment E ($11.62 \pm 0.78c$) with a dose of 175mg / kg, while the lowest absolute weight growth was in treatment A control ($5.37 \pm 0.77a$). This shows that PE energy obtained in consumed feed can be utilized properly. Fern leaves contain ecdysteroids which are not only useful for accelerating molting but also stimulating growth. Fern leaves, one type of fern plant, contain the hormone ecdysteroid which is useful for accelerating molting [4]. Ecdysteroid is a steroid hormone that has a main function as a molting hormone, but also regulates physiological functions, such as growth, metamorphosis, and reproduction. Judging from the results of the study, the dose of fern leaf extract was 175ml/kg[2].




Figure 3. Absolute Length

3.2 Absolute Length

Based on the research conducted, the highest absolute length gain was PE ($4.2 \pm 0.72b$) and the lowest was PA and PB ($0.0 \pm 0a$). It is suspected that the absolute length growth of mangrove crabs increased in PE because some mangrove crabs in PE experienced molting and the dose of fern leaf extract was greater than other treatments. Absolute length increase is seen if the molting process has occurred[10]. Absolute length increase does not occur in the absence of molting, this molting phenomenon is periodic, because for growth crustaceans must expand their body volume by molting. The growth process must be assisted by the absorption of large amounts of water[11].




Figure 4. SR (*Survival Rate*)

3.3 SR (*Survival Rate*)

Based on the results of the study, it shows that feeding the type of fish with and without giving different doses of fern leaf extract has no significant effect ($P > 0.05$) on the survival rate of mangrove crabs. The high survival rate is thought to be due to the controlled environment, adequate feeding and the absence of competition between other crabs. A well-controlled maintenance environment, sufficient amount of feed, no competition for space and feed can support high survival of mangrove crabs during the maintenance period [12].




Fig 5. SGR (*Specific Growth Rate*)

3.4 SGR (*Specific Growth Rate*)

Based on the results of the research conducted that the highest growth rate of PE ($0.27 \pm 0.02c$) and the lowest PA ($0.12 \pm 0a$) giving a dose of fern leaf extract of 175ml / kg to PE into the raw fish feed can accelerate growth compared to other treatments given a lower dose of extract and without the use of fern leaf extract. The specific growth rate is closely related to the body weight gain derived from the feed consumed. The greater the specific growth rate, the better the feed is utilized for growth [13]. Protein plays an important role in feed for the growth of cultivated animals. Raw fish feed has high completeness and nutritional value, so that it can meet the nutritional needs of the mangrove crab body [5]. Fresh raw fish has a high nutritional content, which has a crude protein content of 64.33%, carbohydrates 1.14%, fat 7.40%, and Ca 4.15% [14].

In addition, fern leaves contain ecdysteroids that can accelerate growth in mud crabs. Ecdysteroid is the main steroid hormone in arthropods which has the main function as

a molting hormone, besides that it also regulates physiological functions, such as growth, metamorphosis, and reproduction and good feed utilization can maintain the body condition of mud crabs so that the feed given is used for growth[15]. the growth rate of cultural organisms depends on the species, feed and environment[16]




Figure 6. EPP (Feed Utilization Efficiency)

3.5 EPP (Feed Utilization Efficiency)

Based on the observations made, the highest feed utilization efficiency (EPP) value of mud crab (*Scylla serrata*) in PE was 38.66% and the lowest in PA was 24.06%. This shows that PE utilizes feed well for higher growth of mangrove crabs compared to other treatments given lower extracts and without the use of fern leaf extract. The high value of feed utilization efficiency in PE indicates that the highest dose of 175ml/kg has an effect on growth supported by the use of fresh fish feed which has good nutrition and is not easily destroyed in water compared to artificial feed so that the efficiency of feed utilization is quite high. The quality of feed can be seen from the EPP value which states that the smaller the EPP value, the lower the quality of feed consumed. The efficiency of food use by cultivated cultivars shows the percentage of food that can be utilized by the body. The amount and quality of food greatly affects growth and lack of essential fatty acids can be shown by symptoms of weight loss, small feed efficiency, increased mortality[17]. High feed consumption will provide high growth for the body[18].




Figure 7. Feed Conversion Ratio

3.6 FCR (Feed Conversion Ratio)

Based on the observations made, the highest feed conversion ratio (FCR) value of mud crab (*Scylla serrata*) in PA was 88.80% and the lowest in PE was 20.84%. This shows that the difference in specific growth rate (SGR) in each treatment is supported by the difference in feed conversion ratio (FCR) results. Growth is closely related to feed conversion ratio (FCR) [13]. The difference in the amount of feed consumed by mangrove crabs to the weight produced in each treatment, influences the difference in the average value of feed conversion ratio (FCR). The higher the feed conversion ratio, the less efficient the feed utilization for growth [16].

The level of feed consumption can also affect growth, where the quality and quantity of feed consumed will determine the metabolic process in the cultivator's body, the amount of energy stored in the body. In addition, the amount of energy used for body activities also determines the ability of the cultivator's body to utilize the nutrients obtained from the feed consumed [10].




Figure 8: Total haemocyte count (THC) of mud crab (*Scylla serrata*)

Table 1 Differential haemocyte count (DHC) of mud crabs (*Scylla serrata*)

Perlakuan	DHC					
	Awal		Akhir			
	Hialin %	Granulosit%	Semi Granulosit	Hialin	Granulosit	Semi Granulosit
A	35	35	30	38	33	29
B	39	35	27	38	35	27
C	37	31	31	38	33	30
D	47	31	22	46	31	23
E	38	35	27	39	35	27

3.7 Mangrove Crab Blood Profile

Total haemocyte count (THC) is a picture of blood in crustaceans to show the anti-body response in the body, especially in fighting diseases that attack the body or environmental conditions. Total haemocyte count analysis results obtained at the beginning of maintenance ranged from 2.13×10^6 - 2.80×10^6 and at the end of maintenance ranged from 2.70×10^6 - 3.07×10^6 . Shows that the results obtained are still within the normal range of THC of mangrove crabs. The normal hemocyte count in Crustaceans under normal circumstances is 1.85×10^6 cells/ml. The abundance of the number and type of hemocyte cells in crustaceans can vary due to various things including sex, water temperature, molting cycles, starvation conditions. In addition, the increased THC is the mangrove crab's effort to increase the body's immunity to the surrounding environment or foreign objects that enter the body [19]. The observation results obtained the value of differential haemocyte count obtained at the beginning of maintenance, namely hyaline ranging from 35-47%, granulocytes ranging from 31-39%, and semi granulocytes 20-30% and at the end of maintenance, namely hyaline ranging from 35-49%, granulocytes ranging from 29-43%, and semi granulocytes 20-31%. These 3 types of hemocyte cells, namely hyaline, semigranulocytes and granulocytes have different shapes and sizes. Hyaline cells have a smaller size, spherical without granules or few granules that have a phagocytic role while oval-shaped semigranulocytes play a role in encapsulation, phagocytosis and cytotoxic response, Granulocytes are round, many granules play a role in cytotoxic activity. [9].

Table 2. Water quality parameters of mud crab (*Scylla serrata*)

Parameters	Obtained Range	Ideal Range	Reference
Salinitas (ppt)	22-25	15-25	Winestri <i>et al.</i> , [20]
Suhu ($^{\circ}$ C)	27-29	29-30	Adila <i>et al.</i> , [8]
pH	7,0-7,7	7-8	Asyhariyati <i>et al.</i> , [7]
DO (ppm)	5,2-6,3	3-15	Suryani <i>et al.</i> , [21]

3.8 Water Quality Parameters

Water quality is one of the important factors supporting the success of mangrove crab cultivation and also an important factor affecting the survival rate of mangrove crabs apart from the feed factor. The salinity range obtained during the study ranged from 22-25 ppt. where this range is still within normal limits for mangrove crab cultivation. The appropriate salinity for mangrove crab rearing is 15-25 ppt [20]. Mangrove crabs can live in a salinity range of less than 15 ppt to greater than 30 ppt [22].

The results of temperature measurements during the study ranged from 27-29 $^{\circ}$ C, which is within the optimal range for mangrove crab cultivation. The temperature tolerance limit for mangrove crab life is 29-30 $^{\circ}$ C [8]. Measurement of acidity (pH) during the study ranged from 7.0 to 7.7, the range is still within the optimal range. Good acidity (pH) levels for crab farming range from 7-8 [7].

Dissolved oxygen (DO) measurements during maintenance were found to range from 5.2-6.3 ppm. The range of dissolved oxygen suitable for mangrove crab life ranges from 3-15 ppm [21].

4. CONCLUSION

The conclusion of the research that has been done is that the use of fern leaf extract in mangrove crab (*Scylla serrata*) feed has a significant effect on the growth of absolute weight, Absolute Length, SGR (Specific Growth Rate) and FCR (Feed Conversion Ratio) where the most influential treatment is treatment E, which is the highest dose of 175 ml/kg. This shows

that treatment E is the best dose treatment for the growth and feed conversion ratio of mangrove crabs (*Scylla serrata*).

CONSENT

All authors declare that 'written informed consent' has been obtained from the patient (or other consenting party) for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editorial office/Chief Editor/Editorial Board of this journal.

ETHICAL APPROVAL

The principles of laboratory animal care" (SNI 9057-3:2023) have been followed, as well as applicable national laws. All experiments were reviewed and approved by the appropriate ethics committee.

REFERENCES

- [1] H. Kiswanto, "Karakteristik Lingkungan Tambak Tumpang Sari di Subang, Jawa Barat Sebagai Acuan Pengembangan Budidaya Kepiting Bakau *Scylla serrata*," no. 1, pp. 1–27, 2015.
- [2] R. Ario, A. Djunaedi, I. Pratikto, P. Subardjo, and F. Farida, "Perbedaan Metode Mutilasi Terhadap Lama Waktu Molting *Scylla serrata*," *Bul. Oseanografi Mar.*, vol. 8, no. 2, p. 103, 2019, doi:10.14710/buloma.v8i2.24886.
- [3] A. Kurniawan, M. Haikal, N. Rahmadina, and S. Berliani, "Model Budidaya Kepiting Soka Skala Rumah Tangga Sistem Apartemen Sebagai Sarana Edukasi Masyarakat Pulau Bangka," *Literasi J. Pengabd. Masy. dan Inov.*, vol. 2, no. 1, pp. 8–14, 2022, doi: 10.58466/literasi.v2i1.155.
- [4] A. Romadhon, E. Prasetyono, and A. M. Farhaby, "Laju Pertumbuhan Dan Kecepatan Molting Kepiting Bakau (*Scylla serrata*) Dengan Pemberian Ekstrak Daun Pakis Hutan (*Diplazium caudatum*)," *J. Trop. Mar. Sci.*, vol. 5, no. 1, pp. 9–18, 2022, doi:10.33019/jour.trop.mar.sci.v5i1.2312.
- [5] I. Samidjan, D. Rachmawati, S. Dosen, P. S. Budidaya, and P. Fpik, "Rekayasa Budidaya Kepiting Bakau Melalui Pemotongan Kaki Jalan Dalam Upaya Peningkatan Produksi Kepiting Soka (Soft shell)," pp. 103–121, 2015.
- [6] L. S. S. Sagala, M. Idris, and M. N. Ibrahim, "Perbandingan pertumbuhan kepiting bakau (*Scylla serrata*) jantan dan betina pada metode kurungan dasar," *J. Mina Laut Indones.*, vol. 03, no. 12, pp. 46–54, 2013.
- [7] A. I. Asyhariyati, I. Samidjan, and D. Rachmawati, "Pemberian Kombinasi Pakan Keong Macan dan Ikan Rucah terhadap Pertumbuhan dan Kelulushidupan Kepiting Bakau (*Scylla paramamosain*)," *J. Aquac. Manag. Technol.*, vol. 2, no. 4, pp. 131–138, 2013, [Online]. Available: <https://ejournal3.undip.ac.id/index.php/jamt/article/view/4814>.
- [8] A. Adila, S. Septifitri, and M. Ali, "Penggemukan Kepiting Bakau (*Scylla Serrata*) Dengan Pakan Yang Berbeda," *J. Ilmu-ilmu Perikan. dan Budid. Perair.*, vol. 15, no. 2, pp. 86–94, 2020, doi: 10.31851/jipbp.v15i2.5086.
- [9] A. H. W. Sari and R. Ekawaty, "Profil Hemosit dan Aktifitas Fagositosis Kepiting Bakau (*Scylla sp.*) yang Terserang Ektoparasit di Ekosistem Mangrove Kuta Selatan, Bali," *J. Mar. Aquat. Sci.*, vol. 2, no. 1, p. 34, 2016, doi: 10.24843/jmas.2016.v2.i01.34-39.
- [10] L. Ode, M. Harisud, E. Bidayani, and F. Syarif, "Performa Pertumbuhan dan Kelangsungan Hidup Kepiting Bakau (*Scylla sp .*) dengan Pemberian Kombinasi Pakan Keong Mas dan Ikan Rucah Growth Performance and Survival of Mud Crab (*Scylla sp .*) Feeding With Combination of Golden Snail and Trash Fishes," *J. Trop. Mar. Sci.*, vol. 2, no. 2, pp. 43–50, 2019.
- [11] K. D. P. A. P. Nova, M. Agustini, S. Sumaryam, and S. O. Madyowati, "Pengaruh Jenis Substrat yang Berbeda Terhadap Pertumbuhan Berat Mutlak dan Panjang Mutlak Kepiting Bakau (*Scyllas serrata*) dalam Bak Pemeliharaan," *Juv. Ilm. Kelaut. dan Perikan.*, vol. 4, no. 3, pp. 246–253, 2023, doi: 10.21107/juvenil.v4i3.20508.
- [12] M. Agus, H. Pranggono, and H. Murtadlo, "Pengaruh Pemberian Pakan Keong Mas terhadap Pertumbuhan dan Kelangsungan Hidup Kepiting Bakau Sistem Single Room," *Pena J. Ilmu-ilmu Pengetah. dan Teknol.*, vol. 21, no. 1, pp. 68–74, 2015, [Online]. Available: <https://jurnal.unikal.ac.id/index.php/pena/article/view/54>.

- [13] B. P. Aditya, Sunaryo, and A. Djunaedi, "Pemberian Pelet dengan Ukuran Berbeda terhadap Pertumbuhan Kepiting Bakau (*Scylla serrata* Forsskål, 1775)," *J. Mar. Res.*, vol. 1, no. 1, pp. 146–152, 2012, [Online]. Available: <http://ejournal-s1.undip.ac.id/index.php/jmr>.
- [14] A. Hanif and S. Herlina, "Persentase Pemberian Pakan Ikan Rucah yang Berbeda Terhadap Pertumbuhan Kepiting Bakau (*Scylla* spp)," *J. Ilmu Hewani Trop.*, vol. 10, no. 1, pp. 1–5, 2021, [Online]. Available: <https://unkripjournal.com/index.php/JIHT/article/view/183>.
- [15] S. Aslamyah and Y. Fujaya, "Stimulasi Molting dan Pertumbuhan Kepiting Bakau (*Scylla* sp.) Pendahuluan," *Ilmu Kelaut. Indones. J. Mar. Sci.*, vol. 15, no. September, pp. 170–178, 2010, [Online]. Available: <https://ejournal.undip.ac.id/index.php/ijms/article/view/1679>.
- [16] M. I. Abadi, S. S. Agustina, and L. D. Khartiono, "Pemberian jenis pakan terhadap pertumbuhan dan sintasan kepiting bakau (*Scylla serrata*)," *ZAB J. Zo. Akuatik Banggai*, vol. 1, no. 2, pp. 1–8, 2020.
- [17] Y. Wahyuningsih, Pinandoyo, and L. L. Widowati, "Pengaruh Berbagai Jenis Pakan Segar Terhadap Laju Pertumbuhan dan Kelulushidupan Kepiting Bakau (*Scylla serrata*) Cangkan Lunak dengan Metode Popeye," *J. Aquac. Manag. Technol.*, vol. 4, no. 2, pp. 109–116, 2015, [Online]. Available: <https://ejournal3.undip.ac.id/index.php/jamt/article/view/8549>.
- [18] A. T. Ambarwati, D. Rachmawati, and I. Samidjan, "Pengaruh Penambahan Vitamin C Dengan Dosis yang Berbeda pada Pakan Buatan Terhadap Pertumbuhan dan Kelulushidupan Kepiting Bakau (*Scylla* sp)," *J. Aquac. Manag. Technol.*, vol. 3, no. 4, pp. 26–33, 2014.
- [19] T. Pratiknyo, "Profil Respon Fisiologis dan Kinerja Produksi Kepiting Bakau *Scylla serrata* pada Wadah Terang–Gelap dalam Sistem Resirkulasi," 2018, [Online]. Available: <https://agris.fao.org/agris-search/search.do?recordID=ID2021104543>.
- [20] J. Winestri, D. Rachmawati, and I. Samidjan, "Pengaruh Penambahan Vitamin E Pada Pakan Buatan Terhadap Pertumbuhan dan Kelulushidupan Kepiting Bakau (*Scylla paramamosain*)," *J. Aquac. Manag. Technol.*, vol. 3, no. 4, pp. 40–48, 2014, [Online]. Available: <https://ejournal3.undip.ac.id/index.php/jamt/article/view/6639/6407>.
- [21] N. D. P. I. Suryani, P. G. S. Julyantoro, and A. P. W. K. Dewi, "Panjang Karapas dan Laju Pertumbuhan Spesifik Kepiting Bakau (*Scylla serrata*) yang diberi Jenis Pakan Berbeda di Area Ekowisata Kampung Kepiting, Bali," *J. Mar. Aquat. Sci.*, vol. 4, no. 1, p. 38, 2017, doi: 10.24843/jmas.2018.v4.i01.38-46.
- [22] B. Katiandagho, "Analisis fluktuasi parameter kualitas air terhadap aktifitas molting kepiting bakau (*Scylla* sp)," *Agrikan J. Agribisnis Perikan.*, vol. 7, no. 2, pp. 21–25, 2014, doi: 10.29239/j.agrikan.7.2.21-25.