

GONADAL HISTO-MORPHOLOGIES AND FERTILITY POTENTIAL IN MALE STZ-INDUCED DIABETIC RATS TREATED WITH CURCUMA LONGA

ABSTRACT

The study was aimed at investigating the effects of ethanolic extract of *Curcuma longa* (Turmeric) on fertility profile and gonadal histo-morphology of Streptozotocin induced hyperglycemic male Wistar rats.

Semen parameters (count, motility and viability) and testicular histology of streptozotocin (STZ)-induced hyperglycemic male Wistar rats were investigated following the administration of *Curcuma longa* rhizome extract. Twenty-four rats weighing averagely 110-180g were randomly assigned into four groups of six rats each. Group A animals served as the normal control and they were given rat pellets and distilled water. Group B served as the diabetic control and the animals were administered 65mg/kg body weight of STZ intraperitoneally. Group C served as the STZ + Metformin treated group and the animals were administered 250mg/kg body weight of metformin. Group D served as the STZ + *Curcuma longa* treated group given 500mg/kg body weight of *Curcuma longa*. The administration of metformin and *C. longa* lasted for twenty eight days. Histo-morphology of the testis and semen analysis (sperm count, sperm motility, sperm viability) were carried out. The diabetic group showed seminiferous tubules with reduced germinal epithelium and distorted interstitial connective tissue with a significant ($p < 0.05$) reduction in sperm motility, count, viability compared to the normal group that showed normal histological features and normal sperm parameters (motility, count and viability). However, the administration of *C. longa* showed significant ($p < 0.05$) increase in levels of sperm parameters (motility, count and viability) and the progressive restoration of histological integrity within the testes.

In Conclusion, ethanolic extract of *C. longa* has the potential to reverse the damage caused by hyperglycaemia in male reproductive function.

Key words: *Curcuma longa*, Semen analysis, testis, diabetes mellitus

INTRODUCTION

Diabetes mellitus is a metabolic disorder with a characteristic high level of blood sugar (hyperglycemia) over a prolonged period of time¹ due to two major mechanisms which are either inability of the pancreas to produce insulin or the inability of the cells of the body to take up the insulin produced by the pancreas. Insulin is produced by the beta cells of the pancreas and its function is to regulate the uptake of glucose from blood into the different cells and tissues.²

Three major symptoms of Diabetes Mellitus include frequent urination (Polyuria), increased hunger and appetite (Polyphagia) and increased thirst (Polydipsia).¹ DM is also associated with complications which include diabetic ketoacidosis, Cardiovascular disease, foot ulcers, damage to nerves and eyes, encephalopathy, Cardiomyopathy^{1,3,4} and reproductive dysfunction.⁵

The diabetic complications on different cells, tissues and organs have been linked to the reactive oxygen species (ROS) that is generated due to hyperglycemia.⁶

Prolonged and poorly controlled diabetes has been reported to cause sexual dysfunction in males and females. Sexual dysfunction can also be an early sign of diabetes.⁷ In males, diabetes has effects on the endocrine control of spermatogenesis, causes erectile dysfunction and impaired ejaculation.⁸ It also negatively impacts sperm parameters such as volume, count, motility and morphology.⁹ Due to the above mentioned complications, it became imperative to find affordable solutions by scientists around the world.

Studies have shown that to achieve a good metabolic control of diabetes and energy balance, a combination of lifestyle, nutrition, exercise and pharmaceutical treatment are important.¹⁰ And to some, they are the major factors in the management of diabetes.¹¹

Pharmaceutically, Metformin is considered to be the first choice agent for treatment of diabetes.¹² Another class of drugs of choice is Sulfonylureas¹³ These drugs however are seen to

have undesirable side effects. On account of these side effects, there is advocacy for the use of medicinal plants¹⁴, because they have been found to have little or no side effects and are easily available and affordable. Some of these plants and herbs possess antioxidant properties. They contain Carotenoids, Flavonoids, alkaloids, glycosides and some are said to have anti diabetic effects¹⁵ One of such plants is *Curcuma longa* commonly called Turmeric.

Curcuma longa, the turmeric plant is used as a spice commonly but has been recognized by the scientific community. In Asia, it has been used as a natural therapeutic medicine since ancient times.¹⁶ Its main constituent is curcuminoid¹⁷ which gives it a wide range of pharmacological properties including antioxidant, anti-protozoan, anti-venom, anti-inflammatory¹⁷, antibacterial, antidiabetic, ~~and~~ antiviral, and anticancer activities ¹⁵. It would therefore be interesting to evaluate its antidiabetic effects and its effects on the fertility profile of streptozotocin induced hyperglycemic male Wistar rats.

It has been estimated that 8-12% of couples worldwide during their reproductive life suffer some form of infertility, an estimated 50-80 million couples.¹⁸ Previously, infertility was thought to be a problem of the females but studies have now shown that 20-30% of infertility is linked to the man¹⁹ commonly due to poor semen quality or quantity²⁰

Diabetes has also shown an increase in prevalence over the years with an estimated 425 million persons worldwide and nearly 50% of that number is not diagnosed²¹ Studies have shown that there is a decline in fertility in males with diabetes⁵ leaving a high percentage of them with reproductive dysfunctions including a reduced libido and impotence^{Reference}.

The use of medicinal plants as an alternative medicine has in the last millennium, been accepted all over the world²² In the United States of America for instance, about 38% of the population uses herbal medicine²³ ~~and~~²³ and in Turkey, 48.8% of the population are said to use

Formatted: Not Superscript/ Subscript

Formatted: Not Superscript/ Subscript

Formatted: Superscript

herbal medicines²⁴ to treat chronic diseases and illnesses. In Africa, the World ~~health-Health~~ organization-Organization (WHO) states that at least 80% of the population relies on medicinal plants²⁵ basically due to low cost, availability and apparently low toxicity²⁶

Diabetes Mellitus is said to cause reproductive dysfunctions in both males and females.⁵ Karimi et al.²⁷ showed that diabetic men may observe a reduction in sperm count, motility and morphology due to hyperglycaemic induced ~~Oxidative-oxidative~~ stress which causes damage to nuclear DNA of the sperm cells.^{27,28} In 2015, La Vignera et al.⁸ found out that type 1 diabetic patients showed a lower level of spermatozoa with increased motility, altered mitochondrial function and post ejaculatory dysfunction of the epididymis.

Streptozotocin (STZ) is an alkylating agent that produces pancreatic islet β -cell destruction and is widely used experimentally to produce a model of type 1 diabetes mellitus (T1DM).²⁹ It is a cytotoxic glucose analogue which has been used as a chemotherapeutic agent in the treatment of metastasizing pancreatic islet cell tumours and other malignancies³⁰ and its effect can be seen within seventy-two hours after administration depending on doses administered.³¹ STZ has been one of the chemical agents used for the induction of diabetes in experimental animals.

Streptozotocin functions as DNA (Deoxyribonucleic acid) synthesis inhibitor in bacterial and mammalian cells.³² The selective pancreatic beta cell toxicity and diabetic condition, resulting from STZ induction, is related to the glucose moiety in its chemical structure which enables STZ to enter the beta cell via the low affinity glucose -2- transporter in the plasma membrane.³³

Semen analysis serves as a pivotal and indispensable procedure in the assessment of male fertility status, carrying substantial importance within the field.³⁴ When conducted

meticulously and in an in-depth manner, it possesses the ability to elucidate the underlying causes contributing to male infertility, thereby offering valuable insights and diagnostic clarity.³⁵ The comprehensive evaluation of semen encompasses the analysis of diverse parameters, including the measurement of ejaculate volume, quantification of sperm count, assessment of sperm motility and movement patterns, evaluation of sperm morphology and structural integrity, as well as the investigation of the composition and makeup of seminal secretions.³⁵

Methods

Plant ~~Collection-collection~~ and extract preparation

Rhizomes of *Curcuma longa* (Turmeric) were bought at watt market in Calabar, Cross River State. It was identified and authenticated by a taxonomist from the department of botany, University of Calabar, Calabar and a voucher number (Bot/Herb/UCC/201) was given.

The fresh turmeric rhizomes were cleaned, chopped into tiny pieces and air dried for 7days after which they were grounded into powdered form. A measured amount of 950g of powdered rhizomes were extracted using 2 liters of 95% ethanol for 24 hours. The extract was first double filtered with ~~Chess-chess~~ cloth, then with filtered paper (Whatman No.1 filter paper). The filtrate (extract) was concentrated under reduced pressure at 45°C in rotary evaporator to 10% volume and then to complete dryness using vacuum water bath yielding 58.9g (6.2%) of crude extract. The crude extract (paste) obtained was stored in a refrigerator until it was required.

Experimental Animals

~~Twenty four~~Twenty-four (24) adult male Wistar rats with average weight of 160g were used for this research. The rats were kept in clean cages and divided into four groups designated A, B, C and D with six rats in each group. The rats were allowed to acclimatize for two weeks in

Formatted: Font: Italic, Complex Script Font: Italic

animal house, University of Calabar and allowed unrestricted access to commercially available chow (livestock feed) and water.

Experimental Design

Table 1 shows the experimental design of the research. ~~Twenty four~~ Twenty-four animals were divided randomly into four groups containing six rats each, including. ~~The groups as shown in the table includes the N~~ normal control, the ~~Diabetic-diabetic~~ control, the ~~Metformin-metformin~~ treated group, and the ~~Extract-extract~~ (*C. longa*) treated group.

Table 1: Experimental Design

S/N	NO	GROUPS	TREATMENT
1.	6	A	Normal control Distilled <u>(distilled water)</u>
2.	6	B	Diabetic control <u>(65mg/kg -BW of STZ)</u>
3.	6	C	STZ + 250mg/kg -bw <u>BW of metformin</u>
4.	6	D	STZ + 500mg/kg -bw <u>BW of <i>Curcuma longa</i></u>

Formatted: Font: Italic, Complex Script Font: Italic

Induction of Hyperglycaemia

Streptozotocin (STZ) was administered intra-peritoneally after fasting for twelve hours in order to induce hyperglycaemia. STZ was reconstituted in 0.5M Sodium citrate and administered at a dose of 65mg/kg ~~-bw~~ BW.³⁶

Confirmation of ~~Diabetes-~~ diabetes Mellitus mellitus

Diabetes was confirmed three days after administration of STZ using Accu-Check glucometer with blood samples obtained from tails of the Wistar rats. The blood glucose levels (mg/dl) was checked (before and after) induction and every seven (~~7~~) days during administration of the ethanolic extract of *Curcuma longa* to ascertain hyperglycaemic state. It was observed that the blood glucose of all the animals in the diabetic groups were above 170 mg/dl as compared to that of the normal control which had values less than 91 mg/dl.

Administration of ~~Extrae~~extract

Curcuma longa extract administration commenced three days after induction of hyperglycaemia by oral gastric intubation and lasted for 28 days.

Formatted: Font: Italic, Complex Script Font: Italic

Termination of ~~Experiment~~experiment

At the end of treatment period, the experimental animals were weighed and sacrificed through ~~Chloroform-~~chloroform inhalation. The anterior abdominal wall was incised. The testis was excised, weighed and preserved in 10% formal saline for tissue processing using Haematoxylin haematoxylin and eosin staining method. Semen was extracted from the caudal epididymis for analysis using WHO ~~Recommended-~~recommended method.

Ethical clearance

According to the international guideline for handling of laboratory animals and the relevant University ethical clearance committee on laboratory animal use, all animal experiments were carried out in accordance with the approved institutional and WHO guidelines for the care and

use of animals. Ethical clearance was obtained from the Faculty Animal Research Ethics Committee (FAREC-FBMS), Faculty of Basic Medical Science, University of Calabar, Calabar with the number: 220ANA2423

Determination of weight

All the animals were weighed before and after induction of hyperglycaemia and every three days during the course of administration of the extract.

Analysis of sperm parameters

The epididymis was dissected out and placed in a physiological saline in a ratio of 1:10 weight (g) by volume (ml), and was then macerated using surgical blade to release the sperm cells. The suspension was then filtered with 80um stainless mesh after pipetting according to the method of [Strugała et al.](#)³⁷ The following sperm parameters were estimated as follows;

Commented [T1]: ????

- i) **Sperm motility (%)**: Two drops of sperm suspension were put on a clean and labelled microscope slide and covered with a coverslip. This was mounted on light microscope and the number of motile cells divided by the total number of sperm cells counted and was expressed in percentage.
- ii) **Sperm viability (%)**: This was done using the eosin-nigrosin staining technique. The sperm suspension was mixed with equal volume of the stain and smeared on glass slides. Live sperm excluded the stain and appeared lightly coloured, while dead sperm took up the stain and appeared pink in colour. The counts of live sperm were divided by the total number of sperm cells and multiply by 100.

iii) **Sperm count (x10⁶/ml)**: Improved ~~nueber-Nueber~~ haemocytometer was used for the sperm count. A capillary tube was used to pipette the sperm suspension into the counting chamber of the ~~nueber-Nueber~~ haemocytometer. The ~~nueber-Nueber~~ haemocytometer was placed on a light microscope and the counting was estimated by multiplying the number of cells counted by the dilution factor and by the haemocytometer volume.

Gonadal histo-morphology was assessed using Haematoxylin ~~&-and~~ Eosin.

Statistical Analysis

Data obtained from the experiment was ~~analysed~~analyzed using one-way Analysis of Variance and Duncan post hoc test using a Statistical Package for Social Science, SPSS version 26.0 for Windows. The results were presented as mean \pm standard error of mean and considered statistically significant at $p < 0.05$.

RESULTS

Assessment of ~~Blood-blood~~ Glucoseglucose

Table 2 shows the effect of STZ on fasting blood sugar levels. ~~At-On~~ day zero, blood glucose level in all experimental groups was considered normal ranging from about 70 mg/dl to 107 mg/dl. Elevated blood glucose concentration was seen in all ~~the~~ groups that were induced for diabetes (B, C and D) following STZ administration ~~of 65mg/kg.bw of streptozotocin~~. Blood glucose levels in these groups were significantly higher when compared with the normal control ($p < 0.05$).

Along the course of the experiment through to its termination (Day 28), all diabetic animals still had significantly higher values (± 19.53) of blood sugar ~~level increased significantly higher~~

than (± 19.53) compared with the normal Control group ($p < 0.05$). Animals in Groups groups C and D animals placed on standard anti-diabetic drug (Metformin) and extracts of *curcuma longa* showed a significant ($p < 0.05$) decrease (± 2.98) in blood glucose level compared to both diabetic control group and normal control group ($p < 0.05$) (Tables 2a and 2b).

Table 2a: Daily blood glucose concentration (mg/dl) of the different experimental groups (mg/dl).

Group	<u>blood glucose concentration (mg/dl)Day</u>				
	<u>Day 1</u>	<u>Day 4</u>	<u>Day 7</u>	<u>Day 10</u>	<u>Day 13</u>
Normal Control	82.20 ± 5.45	87.60 ± 4.39	90.60 ± 4.39	86.80 ± 3.84	82.20 ± 4.02
Diabetic Control	274.80 $\pm 8.73^*$	260.00 $\pm 6.52^*$	260.40 $\pm 7.78^*$	219.40 $\pm 12.32^*$	235.00 ± 10.03
Metformin Treated	310.60 $\pm 24.43^*$	264.60 $\pm 10.41^*$	215.20 $\pm 9.72^*$	199.60 $\pm 5.31^*$	177.80 $\pm 7.66^{*,a}$
Extract group	337.40 $\pm 19.68^{*,a}$	285.80 $\pm 17.62^*$	246.80 $\pm 11.91^*$	213.40 $\pm 4.23^*$	174.80 $\pm 8.82^{*,a}$

Values are expressed in Mean \pm SEM. N = 4.

* = Values are significantly decreased when compared to ~~Normal-normal Control-control~~ at $p < 0.05$.

a = Values are significantly increased when compared to ~~Diabetic-diabetic Control-control~~ at $p < 0.05$.

Table 2b: Continuation of daily blood glucose concentration (mg/dl) of the different experimental groups(mg/dl).

Group	<u>blood glucose concentration (mg/dl)Day</u>					
	16	19	22	25	28	Change
Normal Control	89.40 ±4.53	86.40 ±4.84	89.60 ±2.48	95.40 ±3.57	94.80 ±5.47	12.60 ±8.23
Diabetic Control	222.60 ±18.24*	264.67 ±5.34*	249.33 ±19.53*	268.33 ±5.68*	279.67 ±6.49	-24.40 ±26.61
Metformin	142.00 ±5.49* ^a	123.60 ±6.95* ^a	107.80 ±6.90* ^a	101.20 ±4.35 ^a	83.00 ±4.35 ^a	-227.60 ±24.32* ^a
Extract group	129.20 ±6.61* ^a	115.00 ±2.98* ^a	134.20 ±18.33* ^a	114.40 ±9.53 ^a	100.80 ±5.67 ^a	-236.60 ±25.14* ^a

Values are expressed as mean ±SEM, n = 5.

* = significantly different from control at p<0.05

a = significantly different from diabetic control at p<0.05

Figs. 1 & 2 show the blood glucose levels of all groups throughout the experimental period. Elevated blood glucose concentration was seen in all diabetic groups (B, C and D) following administration of 65mg/kg ~~bw~~BW of streptozotocin. Blood glucose level in these groups was remarkably higher ($p < 0.05$) when compared with the normal control. At termination of the experiment (Day 28), the diabetic control animals still had ~~values of blood sugar level~~ significantly ($p < 0.05$) higher ~~values of blood sugar level~~ ($p < 0.05$) when compared with normal control. The blood glucose of the Metformin treated group and extract treated group however reduced significantly and had values which had no statistical significance when compared with normal control group.




Figure 1: Daily blood glucose level of the different experimental groups.

Values are expressed as mean +SEM, n = 5.

* = p<0.05 vs control

z = p<0.05 vs other groups

The blood glucose level of the diabetic control and metformin groups declined daily




Figure 2: Initial and final blood glucose concentrations of the different experimental groups.

Values are expressed as mean +SEM, n = 5.

Assessment of ~~Body-body Weightweight~~

There was a significant decrease in body weight of the experimental rats induced with diabetes (± 6.57) as compared to the normal control ($p < 0.05$). On administration of the standard drug (metformin), and extract of *curcuma longa*, results showed a significant increase (± 4.81) ~~increase~~ of body weight compared to diabetic control group ($p < 0.05$) and progressive change as observed from day 7 of treatment.

At termination of the experiment (Day 28), the diabetic control animals had significantly ($p < 0.05$) ~~values of lighter~~ body weight ~~significantly lower ($p < 0.05$) when compared with those~~ in normal control which also had no difference significantly from the metformin treated group and extract treated group (Tables 3a and 3b) (Figs. 3 & ~~and~~ 4).

Commented [T2]: Results must be presented only in figures or Tables only

Table 3a: Daily body weights of the different experimental groups

Group	Day				
	1	4	7	10	13
Normal Control	158.80 ± 3.98	160.20 ± 3.37	162.60 ± 3.47	164.20 ± 4.28	166.80 ± 4.83
Diabetic Control	183.00 $\pm 6.57^*$	169.60 ± 5.35	163.40 ± 4.48	158.40 ± 4.96	154.20 ± 4.33
Metformin	163.00 $\pm 4.81^{*a}$	156.60 ± 4.51	158.80 ± 5.62	161.60 ± 5.54	160.00 ± 5.61

Extract group	165.40	166.40	161.40	164.00	167.20
	$\pm 5.34^*$	± 5.57	± 5.38	± 5.37	± 5.00

UNDER PEER REVIEW

Table 3b: Continuation of daily body weights of the different experimental groups

Group	Day					Change
	16	19	22	25	28	
Normal Control	169.60	172.20	174.40	176.80	178.80	20.00
	±4.58	±4.69	±4.50	±4.99	±4.14	±1.14
Diabetic Control	152.20	148.67	146.33	142.67	142.67	-37.00
	±4.08	±5.77*	±5.66*	±6.35*	±6.09*	±5.75*
Metformin	159.20	161.40	160.80	162.20	163.80	0.80
	±4.79	±5.37	±6.27	±6.40	±6.01	±1.80* ^a
Extract group	165.40	164.80	163.20	162.40	164.20	-1.20
	±6.28	±5.34	±5.82	±3.57	±4.07* ^a	±2.08* ^a

Values are expressed as mean ±SEM, n = 5.

* = significantly different from control at p<0.05

a = significantly different from diabetic control at p<0.05




Figure 3.: Daily body weights of the different experimental groups.

Values are expressed as mean +SEM, n = 5.

* = significantly different from control at $p < 0.05$

The body weight of the diabetic control declined daily




Figure 4: Initial and final body weights of the different experimental groups.

Values are expressed as mean +SEM, n = 5.

* = significantly different from control at $p < 0.05$

a = significantly different from diabetic control at $p < 0.05$

SEMEN ANALYSIS

Sperm Count

Results showed a significant ($p < 0.05$) decrease (~~37.70 ± 7.5 million/ml and 47.10 ± 1.7 million/ml~~) of sperm count ($\times 10^6$) in the diabetic control and metformin treated groups (~~37.70 ± 7.5 and 47.10 ± 1.7 /ml~~, respectively compared to the normal control group ~~that had~~ (69.6 ± 1.5 million/ml) sperm count. However, ~~65.40 ± 1.4 million/ml of~~ sperm count ($\times 10^6$) was ~~65.40 ± 1.4 /ml recorded~~ in Group D animals (placed on 500mg/kg.bw of *C. longa*). This increase in sperm concentration was statistically significant ($p < 0.05$) when compared with the diabetic control ($p < 0.05$) (Table 4, Fig 5).


Formatted: Space Before: 12 pt

Formatted: Highlight

Table 4: Sperm count of experimental groups expressed in million/ml.

GROUPS	Treatment	Sperm Count <u>count</u> (million/ml)
A	Normal Control	69.6 ± 1.5
B	Diabetic Control	37.70 ± 7.5
C	STZ + 250mg/kg b.w <u>BW</u> metformin	47.10 ± 1.7
D	STZ + 500mg/kg b.w <u>BW</u> <i>curcuma longa</i>	65.40 ± 1.4

Values are expressed in Mean ± SEM. N = 4.



NC - Normal control
DC - Diabetic control
MET - Metformin treated group
T - Tumeric treated group

Figure 5: Sperm count in the different experimental groups.

Values are expressed in Mean \pm SEM. N = 4.

* = Values are significantly decreased compared to Normal Control ($p < 0.05$).

a = Values are significantly increased compared to Diabetic Control at $p < 0.05$.

Sperm Motility

Table 5 and Fig 6 show the result of sperm motility for the experimental animals. Animals in the diabetic control group and the group placed on metformin recorded significant decrease in sperm motility values when compared with the normal control group ($p < 0.05$). However, Group D animals (placed on 500mg/kg.bw of *C. longa*), showed an increase in sperm motility which was statistically significant when compared with the diabetic control and the metformin treated group.


UNDER PEER REVIEW

Table 5: Sperm motility of different experimental groups

GROUPS	TREATMENT	SPERM MOTILITY
A	Normal Control	90.0 ± 0.0
B	Diabetic Control	50.0 ± 0.0
C	STZ + 250mg/kg b.w metformin	57.50 ± 2.50
D	STZ + 500mg/kg b.w <i>curcuma longa</i>	87.50 ± 2.50

Values are expressed in Mean ± SEM. N = 4.

UNDER PEER REVIEW



NC - Normal Control
 DC - Diabetic Control
 MET - Metformin treated group
 T - Turmeric Treated group

Figure 6: Sperm motility in the different experimental groups.

Values are expressed in Mean \pm SEM. N = 4.

* = Values are significantly decreased compared to Normal Control (p<0.05).

a = Values are significantly increased compared to Diabetic Control (p<0.05).

Sperm Viability

Experimental animals in the diabetic control group and the group placed on metformin recorded low sperm viability values which was significantly decreased when compared with the normal control group ($p < 0.05$). However, Group D animals (placed on 500mg/kg.bw of *C. longa*), showed an increase in sperm viability of which was statistically significant when compared with the diabetic control (Table 6 & Fig 7).


UNDER PEER REVIEW

Table 6: Sperm Viability of different experimental groups

GROUPS	TREATMENT	Sperm Viability
A	Normal Control	96.5 ± 0.5
B	Diabetic Control	58.5 ± 3.5
C	STZ + 250mg/kg b.w metformin	68.0 ± 3.0
D	STZ + 500mg/kg b.w <i>curcuma longa</i>	94.50 ± 0.5

Values are expressed in Mean ± SEM. N = 4.

UNDER PEER REVIEW



NC- Normal control
 DC- Diabetic control
 MET- Metformin Treated group
 T- Tumeric treated group

Figure 7: sperm Viability in the different experimental groups.

Values are expressed in Mean \pm SEM. N = 4.

* = Values are significantly decreased when compared to Normal Control at $p < 0.05$.

a = Values are significantly increased when compared to Diabetic Control at $p < 0.05$.

HISTOLOGICAL OBSERVATIONS

Histological study on sections of testes in the normal control (X400) showed prominent seminiferous tubules with regular and consistent cellular arrangements on the germinal epithelium. The lumens of seminiferous tubules were filled with tails (flagella) of newly formed spermatozoa. Basement membrane was intact and the intervening interstitium which contained blood vessels and clusters of Leydig cells showed regular pattern (Plate 1).

Section of testes from the diabetic control group (Group B) showed significant alteration in histological patterns in the testes when compared with the normal control. Irregular shaped seminiferous tubules with shrunken lumen and decreased tubular diameter were observed (Plate 2).

Section of testes in diabetic animals treated with Metformin (Group C) showed normal seminiferous tubules although its germinal epithelium was distorted with differentiating cells arranged in a coarse pattern. The interstitial connective tissue was distorted (plate 3).

For the group placed on 500mg/kg.bw of *C. longa*, Seminiferous tubules were prominent and intact with basement membrane although some alterations were still observed. there is a progressive restoration observed.




PLATE 1: TESTIS X400 NORMAL CONTROL X400

Section of testes showing normal seminiferous tubules with germinal cell layers (**doubleheads arrow**) on the basal lamina to spermatozoa filled lumen (**dotted circular lines**). The interstitial connective (**#**) tissue was well preserved. H&E stain, x400 Magnification.




PLATE 2: H&E section of testis of Diabetic control (X400)

Section of testes showing seminiferous tubules with reduced germinal epithelium (doublehead arrow), distorted interstitial connective tissue (#).H&E stain, X400 magnification




PLATE 3: H& E section of TESTIS DC+METFORMIN X400

Section of testes showing normal seminiferous tubules tubules with germinal cell layers (**doubleheads arrow**) starting from spermatogonia (**arrowhead**) on the basal lamina and spermatocytes filled lumen (**dotted circular lines**). The interstitial connective (**#**) tissues was distorted. H&E stain, X400 Magnification




PLATE 4: H& E section of TESTIS X400 *C.longa* ONLY (500mg/kg.bw)

Section showed different stages of seminiferous tubules and the interstices. The seminiferous with germinal cell layers (**doubleheads arrows**) showed lumen partly & completely filled (**dotted circular line**) with mature spermatozoa. Although alterations are still noticed, there is a progressive restoration observed. H&E, X400 magnification

DISCUSSION, SUMMARY AND CONCLUSION,

This discussion of the work, conclusion, contribution to knowledge and recommendation for further studies are well presented in this chapter

Discussion

Complex interplay of various pathological processes, prominently characterized by oxidative stress damage, inflammation, and apoptosis are involved in diabetic testicular damage.³⁸ Curcumin, a bioactive compound derived from *Curcuma longa*, has emerged as a potential therapeutic agent with protective properties against diabetic testicular injury. However, the precise underlying mechanisms through which curcumin exerts its beneficial effects in this context remain a subject of ongoing investigation. The present study examined the effect of *C. longa* on the histology of the testis and semen parameters of streptozotocin induced hyperglycemic male rats.

The sustained increase in the blood glucose levels of diabetic animals observed at the end of the study was corroborated by studies by³⁹ and⁴⁰ which revealed hyperglycemic response in STZ treated animals due to impaired glucose oxidation, which causes NAD⁺ depletion, ultimately culminating in the inhibition of insulin biosynthesis and secretion. The reduction in blood glucose levels in the metformin and extract-treated groups, aligns with the study by⁴¹, who reported a significant reduction in fasting blood glucose levels in animals administered curcumin. Studies by^{42,17} have corroborated the anti-diabetic efficacy of the extract, attributing it to its antioxidant and anti-inflammatory properties. Findings reported by⁴³ also revealed the potential of the methanol fraction of *C. longa*, to significantly reduce in blood glucose levels in experimental animals.

The final bodyweight of the experimental animals in the extract-treated group showed a significant increase when compared to both the diabetic and control groups. The overall changes in bodyweight statistically indicated that both the metformin and extract-treated groups experienced significant increases in their respective bodyweights when compared to both the control and diabetic groups. These findings are consistent with a study conducted by ⁴⁴ and ⁴⁵ which reported a significant decrease in bodyweight in streptozotocin-injected rats. Furthermore, the observed decrease in bodyweight of the diabetic rats aligns with the findings of ^{46, 47} who observed a similar effect on diabetic animals induced with streptozotocin. The underlying mechanisms contributing to the reduction in bodyweight within the diabetic group can be attributed to the intricate interplay of factors such as the degradation of structural proteins and muscle wasting, as reported by ⁴⁸. Conversely, the ameliorative effects of *Curcuma longa* on bodyweight is supported in the work of ⁴³ where the oral administration of the plant extract demonstrated remarkable improvement in the bodyweight of experimental animals.

Semen analysis revealed a significant reduction in the total count of sperm cells in the diabetic control group, compared to the normal control group. This observation underscores the detrimental impact of diabetes mellitus, regardless of whether it is type 1 or type 2 diabetes, on male fertility. The findings of ⁴⁹ demonstrated the adverse effects of experimentally induced diabetes on sperm parameters. Also a study conducted by ⁵⁰ showed significantly reduced sperm count in diabetic animals. While the scientific basis underlying the pathophysiological mechanisms linking diabetes to semen parameters, particularly sperm count, remains diverse in the literature, prolonged hyperglycemia, triggers the body's oxidative stress response and leads to endothelial injury in blood vessels, including those within the testis and epididymis ⁵¹. Moreover, elevated blood glucose levels can disrupt the regulatory function of the

hypothalamic-pituitary-gonadal axis, resulting in alterations in the number and morphology of testicular interstitial cells, degeneration of Sertoli cells, reduced synthesis and secretion of testosterone, impaired sperm development and maturation, and ultimately, compromised reproductive functions. However, the administration of the extract in the treated group led to a notable improvement in the total sperm count, which was comparable to that observed in the normal control group. These findings are in line with the research conducted by ⁵² who demonstrated that replacing a ketogenic diet with curcumin supplementation improved semen quality. Additionally, ⁵³ reported an increase in total sperm count among infertile men participating in a randomized clinical trial after receiving Curcumin. Together, these studies provide further support for the positive effects of curcumin and its potential in ameliorating the adverse impacts of diabetes on semen parameters, including sperm count.

The experimental findings revealed a significant decline in the percentage concentration of motile sperm cells in the diabetic group compared to the control group, however both the extract-treated and metformin-treated groups showed an increase which was not statistically significant compared to the normal control animals. These observed outcomes are consistent with the outcomes of prior investigations conducted by ^{54, 55} who reported that the addition of curcumin in frozen-thawed Angora goat semen improved both progressive motility and acrosome integrity, while also enhancing the progressive motility and functional integrity of sperm plasma membrane in frozen bull semen. Similarly, ⁵⁵ reported a higher percentage of progressive motile sperm cells associated with the supplementation of curcumin.

Sperm viability values which decreased significantly in the DC and metformin groups compared with the normal control group. In the extract treated group there was an increase in sperm viability which was statistically significant when compared with the diabetic control.

This finding aligns with reports by ⁵⁶ who observed that the percentage of sperm viability in STZ-induced diabetic rats improved mildly in *Curcuma longa* treated group.

Remarkable histological alterations were explicitly observed in the tissue sections of the experimental rats belonging to the diabetic group. These histopathological manifestations were characterized by irregularly shaped seminiferous tubules exhibiting a shrunken lumen and decreased tubular diameter, which were indicative of severe edema and pronounced hyperemia within the veins. These findings align harmoniously with the seminal research conducted by ⁵⁷ who expounded upon the frequent occurrence of abnormal histology within the testes of diabetic animals, with discernible alterations observed in the cytoarchitecture of the seminiferous epithelium as well as disruptions in the occlusive distribution pattern. Conversely, the tissue sections derived from the extract-treated group exhibited remarkable therapeutic efficacy, showcasing prominent seminiferous tubules characterized by prominent basement membranes. Studies by ^{58, 59} documented the positive influence of *Curcuma longa* on the cytoarchitecture of the testes. This remarkable therapeutic outcome can be attributed to the myriad of phytochemical compounds present within the extract, including saponins, flavones, tannins, and terpenes. The collective action of these bioactive compounds, either individually or synergistically, confers an enhanced functionality to the testes. Notably, the extract's insulin-like action, as well as its proficiency in inducing DNA repair systems owing to its potent antioxidant activities, effectively curtails or prevents the generation of deleterious free radicals, thus exerting a protective effect on the testis and semen parameters.

Summary

The study examined the potential hypoglycemic effect of *Curcuma longa* (*C. longa*), known as turmeric, on the histology of the testis and reproductive parameters in male Wistar rats with streptozotocin-induced hyperglycemia. The study was aimed at investigating the effects of ethanolic extract of *Curcuma longa* (Turmeric) on fertility profile of Streptozotocin induced hyperglycemic male Wistar rats.

Twenty four male Wistar rats were divided into four groups of six rats each. The groups included: normal control group given normal chow and distilled water only, diabetic group administered with 65mg/kg.bw of STZ, metformin treated group which was given 250/kg.bw of metformin and the extract treated group to which 500mg/kg.bw of *curcuma longa* was administered. The study lasted for about forty five days of two weeks acclimatization of the animals, three days for confirmation of DM after administration of STZ and twenty eight days for administration of metformin and *C. longa* to the experimental groups. The research investigated various parameters related to testicular histology and reproductive function, such as sperm count, sperm motility, sperm viability. The findings from the study showed alterations in the cytoarchitecture of the testis with a significant decrease in sperm motility, count and viability in the diabetic group compared to the Normal control group. However, administration of *C. longa* as a therapeutic intervention exhibited promising outcomes, as evidenced by the increase in levels of sperm parameters (motility, count and viability) and the progressive restoration of histological integrity within the testes. This underscores the potential of *C. longa* to effectively counteract the detrimental effects induced by hyperglycemia.

Conclusion

The findings of the study suggests that ethanolic extracts of *Curcuma longa* has the potential to improve fertility in diabetic male animals as well as protect against the detrimental effects of hyperglycemia on the male reproductive systems.

REFERENCES

1. World Health Organization "About diabetes" . Archived from the original on 31 March 2014. Retrieved 4 April 2014.
2. Shoback DG, Gardner D, eds.. "Chapter 17". Greenspan's basic & clinical endocrinology (9thed.). New York: McGraw-Hill Medical. 2011; ISBN 978-0-07-162243-1.
3. Saedi, E., Gheini, M. R., Faiz, F. & Arami, M. A. "Diabetes Mellitus and cognitive impairments". *Word Journal of Diabetes*. 2016; 7 (17): 412-422.
4. Leszek J., Trypka E., Tarasov V. V., Ashraf G. M., Aliev G. Type 3 diabetes mellitus: A novel implication of Alzheimers disease. *Current Topics in Medicinal Chemistry*, 2017; 17(12), 1331–5.
5. Codner E, Merino PM, Tena-Sempere M. Female reproduction and type 1 diabetes: from mechanisms to clinical findings. *Hum Reprod*. 2012;18(5):568-85. Doi: 10.1093
6. Jaganjac, M., Tirosh, O., Cohen, G., Sasson, S. & Zarkovic, N. "Reactive aldehydes second messengers of free radicals in diabetes mellitus," *Free Radical Research*, 2013; 47(1):39–48.
7. Miočić J, Car N, Metelko Ž. Sexual dysfunction in women with diabetes mellitus. *Diabetol Croat*. 2008 ;37:35–42
8. La Vignera S, Condorelli RA, Di Mauro M, *et al*. Reproductive Function in male patients with type 1 diabetes mellitus. *Andrology*. 2015; 3(6): 1082-7.
9. Alves M.G, Martins A.D, Rato L, Moreira P.L , Socorro S, Oliveira P.F, Molecular mechanisms beyond glucose transport in diabetes-related male infertility. *BiochimBiophysActa* 2013b ;1832: 626-635.

10. Toumpanakis A, Turnbull T, Alba-Barba I “Effectiveness of plant-based diets in promoting well-being in the management of type 2 diabetes: a systematic review”. *British Medical Journal*. 2018; 6 (1) Open Diabetes.
11. Cappuccio. F.P., cooper.D.,D’Elia. L.,Strazzullo.P.& Miller.M.A Sleep duration predicts cardiovascular outcomes: a systematic review and meta-analysis of prospective studies. *European Heart Journal*, .2011; 32:1484-1492.
12. Inzucchi, S.E., Bergenstal. R.M., Buse.J.B.,Diamant, M., Ferrannini, E., Nauck, M., Peters, A. L., Tsapas A., Wender, R. & Matthews, D. R. Management of hyperglycemia in type 2 Diabetes: a patient-centered approach: position statement of the American Diabetes Association (ADA) and the European Association for the study of Diabetes (EASD). *Diabetes care*. 2012; 35:1364-1379.
13. Eldor, R. &Raz, I. Diabetes therapy--focus on Asia: second-line therapy debate: insulin/secretagogues. *Diabetes Metabolism Research Review*. 2012; 28 Suppl 2:85–89.
14. Kooti, W., Moradi, M., Ali-Akbari, S., Sharafi-Ahvazi, N., Asadi-Samani, M., &Ashtary-Larky, D. Therapeutic and pharmacological potential of *Foeniculumvulgare* Mill: a review. *Journal of HerbMed Pharmacology*, 2015; 4(1), 1-9.
15. Afrisham. R.,Aberomand. M.,Ghaffari. M.A.,Siahpoosh. A. &jamalan. M.. Inhibitory Effect of *Heracleumpersicum* and *Ziziphusjujuba* on Activity of Alpha-Amylase. 2015; *Journal of Botany*, 1-8.
16. Matias, J. N., Achete, G., Campanari, G., Guiguer, É. L., Araújo, A. C., Buglio, D. S. *et al.*, A Systematic Review of the Antidepressant Effects of Curcumin: Beyond Monoamines Theory. *Aust New Z J Psychiatry*, 2021; 4867421998795.
17. Kunnumakkara, A. B., Bordoloi, D., Padmavathi, G., Monisha, J., Roy, N. K., Prasad, S., & Aggarwal, B. B. Curcumin, the golden nutraceutical: multitargeting for multiple chronic diseases. *British Journal of Pharmacology*, 2017; 174(11), 1325-1348.
18. Mascarenhas MN, Flaxman SR, Boerma T, Vanderpoel S, Stevens GA. National, regional, and global trends in infertility prevalence since 1990: a systematic analysis of 277 health surveys. *PLoS Med*. 2012; 9:e1001356
19. Chowdhury, S. H., Cozma, A. I., Chowdhury, J. H. Infertility. *Essentials for the Canadian Medical Licensing Exam: Review and Prep for MCCQE Part I*. 2nd edition. 2017; Wolters Kluwer. Hong Kong.
20. Cooper TG, Noonan E, von Eckardstein S, Auger J, Baker HW, Behre HM, Haugen TB, Kruger T, Wang C, Mbizvo MT, Vogelsong KM. “World Health Organization reference

- values for human semen characteristics” (PDF). Human Reproduction Update. 2010; (3): 231–45. Doi:10.1093/humupd/dmp048. PMID 19934213.
21. International Diabetes Federation. Diabetes atlas. 8th ed. Brussels: Internatinal Diabetes Federation; 2017.
 22. Wu, C. H., Wang, C. C. & Kennedy, J. Changes in herb and dietary supplement use in the U.S. adult population: A comparison of the 2002 and 2007 National health interview surveys. *Clinical Therapy*, 2011; 33:1749–58.
 23. Soner, B. C., Şahin, A. S. &Şahin, T. K. A survey of Turkish hospital patients’ use of herbal medicine. *European Journal of Integrated Medicine*, 2013; 547–52.
 24. Ekor M. The growing use of herbal medicines: issues relating to adverse reactions and challenges in monitoring safety . *Frontiers in Pharmacology*. 2014; 4 : 177.
 25. Parmer, V. Herbal Medicine: Its Toxic Effect and Drug Interactions. The Indian Anaesthetics’ Forum, 2005; pp. 1-9.
 26. Mallidis C, Agbaje I, McClure N, Kliesch S. Einfluss des Diabetes Mellitus auf die Fertilität des Mannes. *Der Urologe A*. 2011; 50(1): 33-7.
 27. Karimi J, Goodarzi M, Tavilani H, Khodadadi I, Amiri I. Increased Receptor for advanced glycation end products in spermatozoa of Diabetic men and its association with sperm nuclear DNA Fragmentation. *Andrologia*. 2012; 44(s1): 280-6.
 28. Furman B. L. Streptozotocin-Induced Diabetic Models in Mice and Rats. *Current protocols in pharmacology*, 2015; 70, 5.47.1–5.47.20.
 29. Lenzen, S. Alloxan Monohydrate and Streptozotocin Diabetes. *Diabetologia*. 2007; 51:216-226.
 30. Junod A, Lambert AE, Orci L, Pictet R, Gonet AE, Renold AE. Studies of the diabetogenic action of streptozotocin. *Proc Soc Exp Biol Med*. 1967; 126(1):201-5. Doi: 10.3181/00379727-126-32401. PMID: 4864021.
 31. Bolzán AD, Bianchi MS. Genotoxicity of streptozotocin. *Mutat Res*. 2002; 512(2-3):121-34. Doi: 10.1016/s1383-5742(02)00044-3.
 32. Elsner M, Guldbakke B, Tiedge M, Munday R, Lenzen S: Relative importance of transport and alkylation for pancreatic beta-cell toxicity of streptozotocin. *Diabetologia*. 2007; 43: 1528–1533.
 33. Barratt C. L. Semen analysis is the cornerstone of investigation for male infertility. *The Practitioner*, 2007; 251(1690), 8–17.

34. Leslie S.W, Soon-Souton, T.L, Khan, M. AB. Male Infertility . StatPearls.2023
35. Nna, V. U., Bakar, A. B. A., Ahmad, A., & Mohamed, M. Diabetes-induced testicular oxidative stress, inflammation, and caspase-dependent apoptosis: the protective role of metformin. *Archives of Physiology and Biochemistry*, 2020; 126(5), 377–388. <https://doi.org/10.1080/13813455.2018.1543329>.
36. Tangvarasittichai S. Oxidative stress, insulin resistance, dyslipidemia and type 2 diabetes mellitus. *World Journal of Diabetes*, 2015; 6(3), 456–480. <https://doi.org/10.4239/wjd.v6.i3.456>.
37. Strugała, P., Dzydzan, O., Brodyak, I., Kucharska, A. Z., Kuroпка, P., Liuta, M., Kaleta-Kuratewicz, K., *et al.* Antidiabetic and Antioxidative Potential of the Blue Congo Variety of Purple Potato Extract in Streptozotocin-Induced Diabetic Rats. *Molecules*, 2019; 24(17), 3126.
38. Marton, L. T., Pescinini-E-Salzedas, L. M., Camargo, M. E. C., Barbalho, S. M., Haber, J. F. D. S., Sinatora, R. V., Detregiachi, C. R. P., Girio, R. J. S., Buchaim, D. V., & Cincotto Dos Santos Bueno, P. The Effects of Curcumin on Diabetes Mellitus: A Systematic Review. *Frontiers in endocrinology*, 2021; 12, 669448. <https://doi.org/10.3389/fendo.2021.669448>.
39. Pulido-Moran, M., Moreno-Fernandez, J., Ramirez-Tortosa, C., & Ramirez-Tortosa, M. Curcumin and Health. *Molecules (Basel, Switzerland)*, 2016; 21(3), 264. <https://doi.org/10.3390/molecules21030264>
40. Mohammed, A., Wudil, A. M., Alhassan, A. J., Imam, A. A., Muhammad, I. U., Idi, A. Hypoglycemic Activity of Curcuma longa Linn Root Extracts on Alloxan Induced Diabetic Rats. *The Saudi Journal of Life Sciences*, 2017; 2(2), 43-49.
41. Akpaso, M. I., Atangwho, I. J., Akpantah, A., Fischer, V. A., Igiri, A. O., & Ebong, P. E. Effect of Combined Leaf Extracts of Vernonia amygdalina (Bitter Leaf) and Gongronema latifolium (Utazi) on the Pancreatic β -Cells of Streptozotocin-Induced Diabetic Rats. *Journal of Advances in Medicine and Medical Research*, 2011; 1(1), 24–34. <https://doi.org/10.9734/BJMMR/2011/215>
42. Sharma, S., Kulkarni, S. K., & Chopra, K. Curcumin, the active principle of turmeric (Curcuma longa), ameliorates diabetic nephropathy in rats. *Clinical and Experimental Pharmacology & Physiology*, 2006; 33(10), 940–945. <https://doi.org/10.1111/j.1440-1681.2006.04468>.
43. Akpaso, M., Igiri, A., Odey, P. A Comparative Study on the Effect of Combined Methanolic Leaf Extracts of Vernonia amygdalina and Gongronema latifolium and Metformin on

- the Pancreatic beta cells of Streptozocin Induced Diabetic Wistar Rats. *Asian Journal of Pharmacy, Nursing and Medical Sciences*, 2017; 5,(2)42-48
44. Shenoy, A. G., & Ramesh, K. G. Improvement of insulin sensitivity by perindopril in spontaneously hypertensive and streptozotocin Induced diabetic rats. *Indian Journal Pharmacology*, 2002; 34, 156- 164.
 45. Szudelski, T. S. The mechanism of alloxan and streptozotocin action in β cells of the rat pancreas. *Journal of Physiology Research*, 2001; 50, 536- 546.
 46. Ding, G. L., Liu, Y., Liu, M. E., Pan, J. X., Guo, M. X., Sheng, J. Z., & Huang, H. F. The effects of diabetes on male fertility and epigenetic regulation during spermatogenesis. *Asian Journal of Andrology*, 2015; 17(6), 948–953. <https://doi.org/10.4103/1008-682X.150844>
 47. Zhong, O., Ji, L., Wang, J., Lei, X., & Huang, H. Association of diabetes and obesity with sperm parameters and testosterone levels: a meta-analysis. *Diabetology & Metabolic Syndrome*, 2021; 13(1), 109. <https://doi.org/10.1186/s13098-021-00728-2>
 48. He, Z., Yin, G., Li, Q. Q., Zeng, Q., & Duan, J. Diabetes Mellitus Causes Male Reproductive Dysfunction: A Review of the Evidence and Mechanisms. *In vivo* (Athens, Greece), 2021; 35(5), 2503–2511. <https://doi.org/10.21873/invivo.12531>
 49. Tsao, C. W., Ke, P. S., Yang, H. Y., Chang, T. C., Liu, C. Y. Curcumin Remedies Testicular Function and Spermatogenesis in Male Mice with Low-Carbohydrate-Diet-Induced Metabolic Dysfunction. *International Journal of Molecular Sciences*, 2022; 23, 10009
 50. Alizadeh, F., Javadi, M., Karami, A. A., Gholaminejad, F., Kavianpour, M., & Haghghian, H. K. Curcumin nanomicelle improves semen parameters, oxidative stress, inflammatory biomarkers, and reproductive hormones in infertile men: A randomized clinical trial. *Phytotherapy Research*, 2017; 32(3), 514–521.
 51. Bucak, M. N., Tuncer, P. B., Sarıözkan, S., Başpınar, N., Taşpınar, M., Cayan, K., Bilgili, A., Akalın, P. P., Büyükleblebici, S., Aydos, S., Ilgaz, S., Sunguroğlu, A., & Oztuna, D.. Effects of antioxidants on post-thawed bovine sperm and oxidative stress parameters: antioxidants protect DNA integrity against cryodamage. *Cryobiology*. 2010; 6(7), 248-253. <https://doi.org/10.1016/j.cryobio.2010.09.001>
 52. Chanapiwat, P., & Kaeoket, K. The effect of Curcuma longa extracted (curcumin) on the quality of cryopreserved boar semen. *Animal Science Journal*, 2015; n/a–n/a. doi:10.1111/asj.12395
 53. Olarenwaju . J., Olatunji, S. Y., Owolabi, J. O., Oluwatosin, A. T., Amaechi, W. C., Desalu, A. B. O. Invivo Evidences of Curcuma longa on oxidative stress in STZ-induced

diabetes on sperm parameters in male Wistar rats. *Anatomy Journal of Africa*, 2017; 6(3): 1045-1051

54. Ricci, G., Catizone, A., Esposito, R., Pisanti, F. A., Vietri, M. T., & Galdieri, M. Diabetic rat testes: morphological and functional alterations. *Andrologia*, 2009; 41(6), 361–368. <https://doi.org/10.1111/j.1439-0272.2009.00937.x>

55. Zha, W., Bai, Y., Xu, L., Liu, Y., Yang, Z., Gao, H., & Li, J. Curcumin Attenuates Testicular Injury in Rats with Streptozotocin-Induced Diabetes. *BioMed Research international*, 2018; 7468019. <https://doi.org/10.1155/2018/7468019>

56. Belhan, S., Yıldırım, S., Huyut, Z., Özdek, U., Oto, G., & Algül, S. Effects of curcumin on sperm quality, lipid profile, antioxidant activity and histopathological changes in streptozotocin-induced diabetes in rats. *Andrologia*, 2020; 52(6), e13584. <https://doi.org/10.1111/and.13584>

UNDER PEER REVIEW