

Review Article

Colonic Volvulus: An Update on Management- Narrative review article

Abstract

Colonic volvulus is a rare cause of large bowel obstruction, and it is commonly seen in older patients. The clinical presentation is abdominal pain and abdominal distension which is followed by absolute constipation. The diagnosis is usually obtained by performing abdominal Xray or computerized tomography. Sigmoid volvulus is the most common cause followed by cecal volvulus and transverse colon volvulus. The management of colonic volvulus is divided into endoscopic detorsion and surgical therapy which involves surgical resection. We have conducted this review article to look at the various causes of colonic volvulus and the management of this condition.

Keywords- “colonic volvulus”, “sigmoid volvulus”, “cecal volvulus”, “ileo-sigmoid knotting”, “endoscopic detorsion”, “sigmoid resection” and “right hemicolectomy”.

Introduction

Colonic volvulus is a condition that is characterized by segmental twisting of a segment of the colon on its narrow and fixed mesentery, leading to a closed loop obstruction caused by a redundant loop of colon. It accounts for 10%-15% of cases of colonic obstruction and the most common site is the sigmoid colon followed by the cecum, transverse colon and splenic flexure. The most common clinical presentation is abdominal distension, constipation and abdominal pain. Sigmoid volvulus is commonly seen in elderly patients with chronic constipation in western countries whereas it is seen in younger patients in Central Asia, South Asia, the middle east and Africa where there is a high fiber diet. Cecal volvulus is seen in younger patients and predominantly female, as is patients with transverse colon and splenic flexure volvulus(1,2).

The diagnosis of colonic volvulus is by performing abdominal radiographs which can demonstrate a loop of dilated intestine like a bent inner tube or omega loop in sigmoid volvulus. If the diagnosis is not confirmed, then a computerized tomography is performed. In cecal volvulus plain abdominal radiographs are usually not diagnostic and the diagnosis is confirmed by computerized tomography. Computerized tomography can detect the degree of colonic distension and direct signs of intestinal ischemia in the twisted loop of intestine. The treatment of colonic volvulus depends on the location with patients with sigmoid volvulus initially being

treated with endoscopic decompression followed by an elective definitive treatment like sigmoid resection and anastomosis. For patients with complicated sigmoid volvulus a Hartmann's procedure may be performed. Cecal volvulus is treated with a right hemicolectomy or a cecostomy and endoscopic detorsion are rarely performed(3–7).

The American Society of Colon and Rectal Surgeons Clinical practical guidelines for the management of colonic volvulus has recommended that for patients with sigmoid volvulus a sigmoidoscopy and detorsion should be performed for stable patients and for unstable patients a laparotomy and sigmoid resection should be performed with either an anastomosis or stoma being performed. Treatments like detorsion followed by sigmoidopexy or mesosigmoidoplasty are not recommended. For patients with cecal volvulus endoscopic detorsion is not recommended with surgical resection like a right hemicolectomy or ileocecal resection being the preferred treatment(8).

The management of colonic volvulus depends on the site of volvulus, and it can be divided into endoscopic therapy and operative treatment. For patients with sigmoid volvulus the treatment can be divided into endoscopic detorsion followed by definitive therapy in the form of sigmoid resection while for patients with cecal volvulus surgical resection is the primary form of therapy. We have conducted this review article to investigate the various treatment options in the management of sigmoid volvulus, ileo-sigmoid knotting, cecal volvulus and transverse and splenic flexure volvulus. We conducted a literature review using PUBMED, Cochrane database of clinical reviews and Google scholar looking for clinical trial, observational studies, cohort studies systemic reviews, and meta-analysis from 1980 to 2024. We used the following keywords, “Sigmoid volvulus”, “Cecal volvulus”, “Colonic volvulus”, “endoscopic detorsion”, “sigmoid resection”, “right hemicolectomy” and “ileo- sigmoid knotting”. All articles were in English language only. Further articles were obtained by manual cross referencing of the literature. Case reports and studies with less than 10 patients and editorials were excluded. Adult male and female patients were included in this study. Pregnant patients and pediatric patients were excluded.

Discussion

Sigmoid volvulus

Sigmoid volvulus accounts for 2% to 5% of acute colonic obstruction in western countries and 20% to 50% of patients in Africa, central Europe and central Asia. The etiology is multifactorial, and it is due to the redundancy of the sigmoid colon with a mesentery that is wider than long and

narrowing of the base of the mesentery. Other factors include advancing age and constipation in elderly patients, and intake of high fiber diet in younger patients especially in countries in Africa(9).Among the factors that can lead to sigmoid volvulus include increased bowel motility, excessive body motion and overeating following a period of starvation(10–12).

The clinical presentation of sigmoid volvulus is abdominal pain, abdominal distension and constipation, with or without signs of peritonitis. An abdominal Xray may reveal a massively dilated colon, but in cases where the diagnosis is in doubt a computerized tomography or Barium enema may be performed(13,14).The World Society Of Emergency Surgeons(WSES) in their consensus guidelines on the management of sigmoid volvulus has recommended that uncomplicated sigmoid volvulus patients should undergo an emergency sigmoidoscopy and detorsion, followed by the insertion of a flatus tube. Once the patient's clinical condition has improved, a definitive sigmoid resection should be performed to prevent recurrence. For unstable patients with suspected sigmoid ischemia emergency surgical resections should be performed with a Hartmann's procedure being the most common operation that is performed(15).

Endoscopic treatment with a sigmoidoscopy is both diagnostic to establish the diagnosis of sigmoid volvulus and therapeutic in that detorsion of the volvulus can be done. Once endoscopic detorsion is done a flatus tube should be inserted to prevent early recurrence(16–18).Endoscopic detorsion is associated with minimal morbidity and mortality with a success rate of 80% to 90% but it has a high recurrence rate of up to 40%. Hence a definitive surgical therapy should be offered to the patient after successful endoscopic detorsion to ensure better survival and quality of life(19–22).

The surgical management for sigmoid volvulus can be divided into sigmoid resection followed by anastomosis, which is performed for patients with uncomplicated sigmoid volvulus. For complicated sigmoid volvulus the Hartmann's procedure is the most common operation that is performed. If there is extensive ischemia of the sigmoid colon than a subtotal colectomy may be performed(23–27).The patients who underwent sigmoid resection for complicated sigmoid volvulus were associated with a higher morbidity and mortality than those who underwent elective sigmoid resection(28).A retrospective study by Ifversen et al comparing the mortality and recurrence of patients who underwent endoscopic therapy versus surgical therapy concluded that patients who underwent surgical resection had the best survival rate and lowest recurrence rate(29).

Sigmoid colectomy with primary anastomosis was compared with Hartmann's procedure in the management of sigmoid volvulus, in several retrospective studies and there was no difference with regards to the post operative complications and mortality. The Hartmann's procedure was associated with a shorter stay in the hospital(30,31).A systemic review and meta-analysis was conducted by Awedew et al comparing primary resection with anastomosis versus the Hartmann's procedure in complicated sigmoid volvulus.11 studies with 724 patients were included in the study and the mortality from primary anastomosis was 15% and Hartmann's

procedure was 19%. This study concluded that there was no difference with regards to morbidity and mortality from the choice of resection and primary anastomosis or the Hartmann's procedure in the management of complicated sigmoid volvulus(32).

Percutaneous endoscopic colostomy is another procedure that can be performed in patients who have undergone endoscopic decompression but are not fit to undergo surgery and a systemic review by Jackson et al did highlight the advantage of this procedure in unfit patients, but further studies will be needed to evaluate its efficacy(33). Tube sigmoidostomy is another alternative to sigmoidopexy in the management of sigmoid volvulus in patients who are not fit for surgery, but further studies are required to assess its efficacy(34,35).

Table 1 : This table shows the mortality between sigmoid resection and anastomosis against the Hartmann's procedure in the surgical management of sigmoid volvulus

Study	Study type	Year	N=numbers	Mortality from resection & anastomosis (%)	Mortality from the Hartmann's procedure (%)
Coban et al	Retrospective study	2008	47	8%	8%
Shahmoradi et al	Retrospective study	2020	102	0%	1.8%
Awedew et al	Systemic review & meta-analysis	2023	724	15%	19%

Ileo-sigmoid knotting

This condition is characterized by the wrapping of the ileum and sigmoid colon around one another and its underlying mesentery. It is a rare but severe form of intestinal obstruction and is seen in middle aged patients in central, south Asia and Africa. It is seen in a hypermobile bowel with an elongated mesentery and narrow base. The consumption of a high bulk diet with an empty small bowel are predisposing factors to this condition(36,37). Ileo- sigmoid knotting can

be classified into class 1 where the ileum revolves around the sigmoid colon, class 2 where the sigmoid colon revolves around the ileum, class 3 the ileocecal junction revolves around the sigmoid colon and the undetermined type where the cause of obstruction cannot be determined. The diagnosis of ileo-sigmoid knotting is usually confirmed by performing a computerized tomography (CT) or magnetic resonance imaging (MRI)(38,39).

The management of ileo-sigmoid knotting is by fluid resuscitation and failure to pass the sigmoidoscope beyond the sigmoid colon should alert one to the possibility of this condition. Surgery is the primary form of treatment of this condition, with untwisting of the knot and assessing the viability of the bowel. Bowel resection and anastomosis can be performed if the bowel is viable, and this can involve small bowel resection and a sigmoid colectomy with anastomosis. If there is bowel perforation with contamination of the peritoneal cavity, then a resection with a stoma may be performed with a Hartmann's procedure being the preferred operation(40–43).non-resection surgical procedures like sigmoidoplasty, sigmoidopexy and cecopexy are rarely performed due to their high recurrence rate. The mortality rate from uncomplicated ileo-sigmoid knotting is 8% -10% but for complicated cases the mortality approaches 60% to 80%(44–47).

Cecal Volvulus

Cecal Volvulus is the second most cause of colonic volvulus accounting for 10% to 40% of cases. It can be divided into 2 subgroups, the first is a loop axial ileocolic volvulus and the second is a cecal bascule. Predisposing factors for cecal bascule include constipation, previous abdominal surgery, prolonged immobility and high fiber intake(48).The clinical presentation is right sided abdominal pain and abdominal distension which may progress to generalized abdominal pain. The diagnosis can be confirmed by performing a computerized tomography which may demonstrate the “coffee bean”, “bird beak” and “whirl sign”(49).The treatment of cecal volvulus is by surgical means and can be divided into resection and non-resection surgery depending on the viability of the cecum. Right hemicolectomy and ileocecal resection are the most common resection surgeries with cecopexy and cecostomy being the common non - resection procedures. Colonoscopy decompression is not usually performed due to the high failure rate and difficulty(50,51).

Cecal bascule is a rare form of cecal volvulus that accounts for 5%-10% of cases, and it is characterized by distended cecum fold anteriorly on the ascending colon without any torsion. Adhesion, abdominal surgeries and mobile cecum have been implicated in the etiology of cecal bascule and the clinical presentation is right sided abdominal pain and abdominal distension. The diagnosis is confirmed by computerized tomography. The treatment of cecal bascule is by performing a right hemicolectomy or cecopexy(52–54).

Transverse colon and splenic flexure colon volvulus

These are rare causes of volvulus and are seen in less than 5% of cases of colonic volvulus. The clinical presentation is abdominal pain and abdominal distension which may be followed by generalized abdominal pain. The diagnosis is confirmed by computerized tomography and the treatment is surgical resection as detorsion is usually not performed. The decision to perform an anastomosis or stoma formation will depend on the viability of the colon segment that is involved(55–59).

Conclusion

Colonic volvulus is one of the causes of large bowel obstruction that is seen in older patients and due to the delay in diagnosis, this condition often presents as an acute emergency. Sigmoid volvulus is the most common presentation of colonic volvulus with endoscopic detorsion being the first line therapy and surgical resection being done to prevent recurrence. The choice of performing an anastomosis or a stoma will depend on the clinical condition of the patient. Cecal volvulus is the second most common cause of colonic volvulus, and it is always treated with surgical resection with the right hemicolectomy being the most common operation. The treatment of colonic volvulus is important due to the high mortality if there is a delay in diagnosis and treatment.

References

1. Rakinic J. Colonic Volvulus. In: The ASCRS Textbook of Colon and Rectal Surgery [Internet]. New York, NY: Springer New York; 2011. p. 395–406. Available from: http://link.springer.com/10.1007/978-1-4419-1584-9_23
2. Margolin DA, Whitlow CB. The Pathogenesis and Etiology of Colonic Volvulus. *Semin Colon Rectal Surg.* 2007 Mar;18(1):79–86. <https://doi.org/10.1053/j.scrs.2006.12.013>
3. Kapadia MR. Volvulus of the Small Bowel and Colon. *Clin Colon Rectal Surg.* 2017 Feb;30(1):40-45. doi: 10.1055/s-0036-1593428. PMID: 28144211; PMCID: PMC5179272.
4. Jones IT, Fazio VW. Colonic volvulus. Etiology and management. *Dig Dis.* 1989;7(4):203-9. doi: 10.1159/000171220. PMID: 2663257.
5. Bauman ZM, Evans CH. Volvulus. *Surg Clin North Am.* 2018 Oct;98(5):973-993. doi: 10.1016/j.suc.2018.06.005. Epub 2018 Jul 31. PMID: 30243456.
6. Valsdottir E, Marks JH. Volvulus: small bowel and colon. *Clin Colon Rectal Surg.* 2008 May;21(2):91-3. doi: 10.1055/s-2008-1075856. PMID: 20011403; PMCID: PMC2780196.

7. Perrot L, Fohlen A, Alves A, Lubrano J. Management of the colonic volvulus in 2016. *J Visc Surg.* 2016 Jun;153(3):183-92. doi: 10.1016/j.jviscsurg.2016.03.006. Epub 2016 Apr 28. PMID: 27132752.
8. Alavi K, Poylin V, Davids JS, Patel SV, Felder S, Valente MA, Paquette IM, Feingold DL; Prepared on behalf of the Clinical Practice Guidelines Committee of the American Society of Colon and Rectal Surgeons. The American Society of Colon and Rectal Surgeons Clinical Practice Guidelines for the Management of Colonic Volvulus and Acute Colonic Pseudo-Obstruction. *Dis Colon Rectum.* 2021 Sep 1;64(9):1046-1057. doi: 10.1097/DCR.0000000000002159. PMID: 34016826.
9. Atamanalp SS. Sigmoid volvulus. *Eurasian J Med.* 2010 Dec;42(3):142-7. doi: 10.5152/eajm.2010.39. PMID: 25610145; PMCID: PMC4261258.
10. Dişçi E, Atamanalp SS. Factors precipitating volvulus formation in sigmoid volvulus. *Ulus Travma Acil Cerrahi Derg.* 2022 Mar;28(3):281-284. doi: 10.14744/tjtes.2020.03762. PMID: 35485550; PMCID: PMC10493538.
11. Raveenthiran V, Madiba TE, Atamanalp SS, De U. Volvulus of the sigmoid colon. *Colorectal Dis.* 2010 Jul;12(7 Online):e1-17. doi: 10.1111/j.1463-1318.2010.02262.x. Epub 2010 Mar 10. PMID: 20236153.
12. Bhandari TR, Shahi S, Poudel R. Colonic Volvulus: An Experience at Tertiary Care Hospital in Nepal. *Cureus.* 2019 Jul 18;11(7):e5165. doi: 10.7759/cureus.5165. PMID: 31528515; PMCID: PMC6743661.
13. Gingold D, Murrell Z. Management of colonic volvulus. *Clin Colon Rectal Surg.* 2012 Dec;25(4):236-44. doi: 10.1055/s-0032-1329535. PMID: 24294126; PMCID: PMC3577612.
14. Humbert C, Grillet F, Malakhia A, Meuriot F, Lakkis Z, Piton G, Vuitton L, Loffroy R, Calame P, Delabrousse E. Stratification of sigmoid volvulus early recurrence risk using a combination of CT features. *Diagn Interv Imaging.* 2022 Feb;103(2):79-85. doi: 10.1016/j.diii.2022.01.005. Epub 2022 Jan 24. PMID: 35086786.
15. Tian BWCA, Vigutto G, Tan E, van Goor H, Bendinelli C, Abu-Zidan F, Ivatury R, Sakakushev B, Di Carlo I, Sganga G, Maier RV, Coimbra R, Leppäniemi A, Litvin A, Damaskos D, Broek RT, Biffl W, Di Saverio S, De Simone B, Ceresoli M, Picetti E, Galante J, Tebala GD, Beka SG, Bonavina L, Cui Y, Khan J, Cicuttin E, Amico F, Kenji I, Hecker A, Ansaloni L, Sartelli M, Moore EE, Kluger Y, Testini M, Weber D, Agnoletti V, Angelis ND, Coccolini F, Sall I, Catena F. WSES consensus guidelines on sigmoid volvulus management. *World J Emerg Surg.* 2023 May

- 15;18(1):34. doi: 10.1186/s13017-023-00502-x. PMID: 37189134; PMCID: PMC10186802.
16. Atamanalp SS, Atamanalp RS. The role of sigmoidoscopy in the diagnosis and treatment of sigmoid volvulus. *Pak J Med Sci.* 2016 Jan-Feb;32(1):244-8. doi: 10.12669/pjms.321.8410. PMID: 27022384; PMCID: PMC4795878.
 17. Atamanalp SS, Dişçi E, Peksöz R, Korkut E, Aksungur N, Altundaş N, Kara S. Recurrence-Preventive Role of Flatus Tubes Following Endoscopic Decompression in Sigmoid Volvulus. *Turk J Gastroenterol.* 2023 Apr;34(4):371-377. doi: 10.5152/tjg.2023.22201. PMID: 36635912; PMCID: PMC10210850.
 18. Negm S, Farag A, Shafiq A, Moursi A, Abdelghani AA. Endoscopic management of acute sigmoid volvulus in high risk surgical elderly patients: a randomized controlled trial. *Langenbecks Arch Surg.* 2023 Aug 28;408(1):338. doi: 10.1007/s00423-023-03071-4. PMID: 37635200; PMCID: PMC10460710.
 19. Lou Z, Yu ED, Zhang W, Meng RG, Hao LQ, Fu CG. Appropriate treatment of acute sigmoid volvulus in the emergency setting. *World J Gastroenterol.* 2013 Aug 14;19(30):4979-83. doi: 10.3748/wjg.v19.i30.4979. PMID: 23946604; PMCID: PMC3740429.
 20. Slack Z, Shams M, Ahmad R, Ali R, Antunes D, Dey A, Patel M, Shabana A, Bond-Smith G, Tebala GD. Prognostic factors in the decision-making process for sigmoid volvulus: results of a single-centre retrospective cohort study. *BMC Surg.* 2022 Mar 14;22(1):95. doi: 10.1186/s12893-022-01549-4. PMID: 35287640; PMCID: PMC8919604.
 21. Moro-Valdezate D, Martín-Arévalo J, Pla-Martí V, García-Botello S, Izquierdo-Moreno A, Pérez-Santiago L, Pedrós-Giménez JM, Villagrasa R, Peña A, Espí-Macías A. Sigmoid volvulus: outcomes of treatment and predictors of morbidity and mortality. *Langenbecks Arch Surg.* 2022 May;407(3):1161-1171. doi: 10.1007/s00423-022-02428-5. Epub 2022 Jan 14. PMID: 35028738; PMCID: PMC9151547.
 22. Abdelrahim A, Zeidan S, Qulaghassi M, Ali O, Boshnaq M. Dilemma of sigmoid volvulus management. *Ann R Coll Surg Engl.* 2022 Feb;104(2):95-99. doi: 10.1308/rcsann.2021.0123. Epub 2021 Dec 3. PMID: 34860119; PMCID: PMC10335211.
 23. Safioleas M, Chatziconstantinou C, Felekouras E, Stamatakos M, Papaconstantinou I, Smirnis A, Safioleas P, Kostakis A. Clinical considerations and therapeutic strategy for sigmoid volvulus in the elderly: a study of 33 cases.

World J Gastroenterol. 2007 Feb 14;13(6):921-4. doi: 10.3748/wjg.v13.i6.921. PMID: 17352024; PMCID: PMC4065930.

24. Lee K, Oh HK, Cho JR, Kim M, Kim DW, Kang SB, Kim HJ, Park HC, Shin R, Heo SC, Ryoo SB, Park KJ; Seoul Colorectal Research Group (SECOG). Surgical Management of Sigmoid Volvulus: A Multicenter Observational Study. *Ann Coloproctol.* 2020 Dec;36(6):403-408. doi: 10.3393/ac.2020.03.23. Epub 2020 Dec 31. PMID: 33486909; PMCID: PMC7837394.
25. Cirocchi R, Farinella E, La Mura F, Morelli U, Trastulli S, Milani D, Di Patrizi MS, Rossetti B, Spizzirri A, Galanou I, Kopanakis K, Mecarelli V, Sciannone F. The sigmoid volvulus: surgical timing and mortality for different clinical types. *World J Emerg Surg.* 2010 Jan 13;5:1. doi: 10.1186/1749-7922-5-1. PMID: 20148115; PMCID: PMC2820010.
26. Ağaoğlu N, Yücel Y, Türkyılmaz S. Surgical treatment of the sigmoid volvulus. *Acta Chir Belg.* 2005 Aug;105(4):365-8. doi: 10.1080/00015458.2005.11679737. PMID: 16184717.
27. Hardy NP, McEntee PD, McCormick PH, Mehigan BJ, Larkin JO. Sigmoid volvulus: definitive surgery is safe and should be considered in all instances. *Ir J Med Sci.* 2022 Jun;191(3):1291-1295. doi: 10.1007/s11845-021-02713-0. Epub 2021 Jul 29. PMID: 34327621; PMCID: PMC9135785.
28. Traoré D, Sanogo ZZ, Bengaly B, Sissoko F, Coulibaly B, Togola B, Traoré I, Goïta D, Keïta S, Togo AP, Diallo G, Sangaré D, Ongoïba N, Koumaré AK. Acute sigmoid volvulus: results of surgical treatment in the teaching hospitals of Bamako. *J Visc Surg.* 2014 Apr;151(2):97-101. doi: 10.1016/j.jvisc Surg. 2014.01.010. Epub 2014 Mar 4. PMID: 24618365.
29. Ifversen AK, Kjaer DW. More patients should undergo surgery after sigmoid volvulus. *World J Gastroenterol.* 2014 Dec 28;20(48):18384-9. doi: 10.3748/wjg.v20.i48.18384. PMID: 25561806; PMCID: PMC4277976.
30. Kazem Shahmoradi M, Khoshdani Farahani P, Sharifian M. Evaluating outcomes of primary anastomosis versus Hartmann's procedure in sigmoid volvulus: A retrospective-cohort study. *Ann Med Surg (Lond).* 2021 Jan 19;62:160-163. doi: 10.1016/j.amsu.2021.01.019. PMID: 33520215; PMCID: PMC7820798.

31. Pattanaik SK. Emergency Management of Sigmoid Colon Volvulus in a Volvulus Belt Population and a Review of Literature. *Indian Journal of Surgery*. 2018 Dec 12;80(6):599–605. DOI:[10.1007/s12262-017-1699-7](https://doi.org/10.1007/s12262-017-1699-7)
32. Awedew AF, Asefa Z, Enkoye BD. Comparing Resection and Primary Anastomosis versus Hartmann's Stoma on the Mortality and Morbidity of Gangrenous Sigmoid Volvulus: Systematic Review and Meta-Analysis. *Ethiop J Health Sci*. 2023 Nov;33(6):1087-1096. doi: 10.4314/ejhs.v33i6.19. PMID: 38784481; PMCID: PMC11111268.
33. Jackson S, Hamed MO, Shabbir J. Management of sigmoid volvulus using percutaneous endoscopic colostomy. *Ann R Coll Surg Engl*. 2020 Nov;102(9):654-662. doi: 10.1308/rcsann.2020.0162. Epub 2020 Aug 11. PMID: 32777932; PMCID: PMC7591603.
34. Gupta SS, Singh O, Paramhans D, Mathur RK. Tube sigmoidostomy: a valuable alternative to sigmoidopexy for sigmoid volvulus. *J Visc Surg*. 2011 Apr;148(2):e129-33. doi: 10.1016/j.jviscsurg.2011.02.003. Epub 2011 Apr 14. PMID: 21497150.
35. Imakita T, Suzuki Y, Ohdaira H, Urashima M. Colonoscopy-assisted percutaneous sigmoidopexy: a novel, simple, safe, and efficient treatment for inoperable sigmoid volvulus (with videos). *GastrointestEndosc*. 2019 Sep;90(3):514-520. doi: 10.1016/j.gie.2019.04.246. Epub 2019 May 8. PMID: 31077700.
36. Atamanalp SS. Ileosigmoid knotting. *Eurasian J Med*. 2009 Aug;41(2):116-9. PMID: 25610081; PMCID: PMC4261469.
37. Mandal A, Chandel V, Baig S. Ileosigmoid knot. *Indian J Surg*. 2012 Apr;74(2):136-42. doi: 10.1007/s12262-011-0346-y. Epub 2011 Nov 17. PMID: 23542502; PMCID: PMC3309095.
38. Atamanalp SS, Peksöz R, Dişçi E. Sigmoid Volvulus and Ileosigmoid Knotting: An Update. *Eurasian J Med*. 2022 Dec;54(Suppl1):91-96. doi: 10.5152/eurasianjmed.2022.22310. PMID: 36655451; PMCID: PMC11163360.
39. Abebe K, Sherefa K, Teshome H, Abebe E. Ileosigmoid Knotting: Analysis of Patients Clinical Profiles and Determinants of Outcomes. *Surg Res Pract*. 2020 Jul 23;2020:3826138. doi: 10.1155/2020/3826138. PMID: 32775608; PMCID: PMC7396096.

40. Atamanalp, S. S., Peksöz, R., Dişçi, E., Atamanalp, R. S., & Tatar Atamanalp, C. (2024). Management of Ileosigmoid Knotting: A Literature Review. *European Journal of Therapeutics*, 30(4), 525–530. <https://doi.org/10.58600/eurjther2271>
41. Atamanalp SS, Korkut E, Karadeniz E, Aksungur N. Ileosigmoid Knotting: Changing Trends Over 50 Years. *Indian Journal of Surgery*. 2018 Oct 1;80(5):470–3.
42. Molla YD, Yasin MO, Kassa SA. Ileo-sigmoid knotting: A case series of 25 patients. Vol. 58, *International Journal of Surgery Open*. Elsevier Ltd; 2023. <http://dx.doi.org/10.1016/j.ijso.2023.100664>
43. Bayleyegn NS, Zelelew AN, Sisay AL. Evaluation of clinical profiles, surgical experience and outcomes of ileosigmoid knotting in low-resource setup: A retrospective cohort study at Jimma University Medical Center. *World J Surg*. 2024 Jun;48(6):1331-1347. doi: 10.1002/wjs.12155. Epub 2024 Mar 25. PMID: 38526512.
44. Mallick IH, Winslet MC. Ileosigmoid knotting. *Colorectal Dis*. 2004 Jul;6(4):220-5. doi: 10.1111/j.1463-1318.2004.00361.x. PMID: 15206962.
45. Atamanalp SS, Disci E, Peksoz R, Atamanalp RS, Atamanalp CT. Ileosigmoid knotting: A review of 923 cases. *Pak J Med Sci*. 2022 Mar-Apr;38(3Part-I):711-715. doi: 10.12669/pjms.38.3.5320. PMID: 35480527; PMCID: PMC9002437.
46. Selçuk Atamanalp S. Treatment for ileosigmoid knotting: a single-center experience of 74 patients. *Tech Coloproctol*. 2014 Mar;18(3):233-7. doi: 10.1007/s10151-013-1046-3. Epub 2013 Jul 10. PMID: 23839796.
47. Ooko PB, Saruni S, Oloo M, Topazian HM, White R. Ileo-sigmoid knotting: a review of 61 cases in Kenya. *Pan Afr Med J*. 2016 Apr 15;23:198. doi: 10.11604/pamj.2016.23.198.6255. PMID: 27347287; PMCID: PMC4907762.
48. Hasbahceci M, Basak F, Alimoglu O. Cecal volvulus. *Indian J Surg*. 2012 Dec;74(6):476-9. doi: 10.1007/s12262-012-0432-9. Epub 2012 Mar 14. PMID: 24293902; PMCID: PMC3537995.
49. Consorti ET, Liu TH. Diagnosis and treatment of caecal volvulus. *Postgrad Med J*. 2005 Dec;81(962):772-6. doi: 10.1136/pgmj.2005.035311. PMID: 16344301; PMCID: PMC1743408.
50. Madiba TE, Thomson SR. The management of sigmoid volvulus. *J R Coll Surg Edinb*. 2000 Apr;45(2):74-80. PMID: 10822915.
51. Mahendran V, Reddy B, Jaradat I. The Unfolding Situation of Caecal Volvulus: A Retrospective Analysis of 36 Cases From a Single Center. *Cureus*. 2022 Jan

10;14(1):e21071. doi: 10.7759/cureus.21071. PMID: 35028246; PMCID: PMC8744366.

52. Lung BE, Yelika SB, Murthy AS, Gachabayov M, Denoya P. Cecal bascule: a systematic review of the literature. *Tech Coloproctol.* 2018 Feb;22(2):75-80. doi: 10.1007/s10151-017-1725-6. Epub 2017 Nov 20. PMID: 29159782.
53. Park JS, Ng KS, Young CJ. Caecal bascule: a case series and literature review. *ANZ J Surg.* 2018 May;88(5):E386-E389. doi: 10.1111/ans.13898. Epub 2017 Mar 20. PMID: 28318090.
54. Majeski J. Operative therapy for cecal volvulus combining resection with colopexy. *Am J Surg.* 2005 Feb;189(2):211-3. doi: 10.1016/j.amjsurg.2004.11.004. PMID: 15720993.
55. Plevin RE, Campbell AR. Colon volvulus. In: *Emergency General Surgery: A Practical Approach.* Springer International Publishing; 2018. p. 333–8. DOI-10.1007/978-3-319-96286-3_28
56. Beck, D.E., Roberts, P.L., Rombeau, J.L., Stamos, M.J., Wexner, S.D. (2009). Colonic Volvulus. In: Wexner, S., Stamos, M., Rombeau, J., Roberts, P., Beck, D. (eds) *The ASCRS Manual of Colon and Rectal Surgery.* Springer, New York, NY. https://doi.org/10.1007/b12857_19
57. Truitt MS, Gutierrez T. Volvulus. In: *Common Problems in Acute Care Surgery* [Internet]. Cham: Springer International Publishing; 2017. p. 349–55. Available from: http://link.springer.com/10.1007/978-3-319-42792-8_33
58. Sharifi A, Tafti SMA, Keramati M, Kazemeini A, Behboudi B. Splenic flexure volvulus, a rare etiology of colonic obstruction: Case report. *Int J Surg Case Rep.* 2021 Sep;86:106128. doi: 10.1016/j.ijscr.2021.106128. Epub 2021 Jun 19. PMID: 34500250; PMCID: PMC8430379.
59. Niksch L, Lockwood M, van Rooyen PL, Niksch NA, Lorentz L. Transverse colon volvulus - a case report and literature review. *S Afr J Surg.* 2023 Nov;61(4):237-239. doi: 10.36303/SAJS.4069. Epub 2023 Oct 25. PMID: 38450699.