

Synthesis, Characterization and X-Ray Structure of Mononuclear Cu(II) Complex of The Schiff Base 2,2'-[1,1'-(Ethane-1,2-Diyl)bis(nitriloethylidyne)]diphenol

ABSTRACT

The reactions of the Schiff base 2,2'-[1,1'-(ethane-1,2-diyl)bis(nitriloethylidyne)]diphenol (H_2L), which was synthesised by condensation of 2-hydroxyacetophenone and ethylenediamine in the 2:1 ratio, with acetate copper (II) Salt, afforded a new mononuclear complex formulated as $[Cu(L)] \cdot (H_2O)_{0.5}$. The ligand and the complex have been characterized by elemental analysis, FTIR, UV-Vis and 1H and ^{13}C NMR spectroscopies. The characterization of the copper complex was completed by molar conductivity and room temperature magnetic measurements. The structure of the complex has been resolved by X-ray crystallography technic. The mononuclear copper (II) complex crystallises in the monoclinic space group $C2/c$ with the following unit cell parameters $a = 26.131(6) \text{ \AA}$, $b = 7.293(5) \text{ \AA}$, $c = 17.256(4) \text{ \AA}$, $\beta = 106.10(5)^\circ$, $V = 3160(2) \text{ \AA}^3$, $Z = 4$, $R_1 = 0.077$ and $wR_2 = 0.145$. In the structure of the complex the ligand acts in tetradentate fashion and the coordination environment of the copper atom is best described as distorted square planar geometry.

Keywords: 2-hydroxyacetophenone, Schiff base, Copper, Mononuclear, FTIR, XRD

1. INTRODUCTION

"Schiff bases are organic ligands widely studied in coordination chemistry. They are derived from the condensation of amine derivatives and ketone or aldehyde derivatives and constitute a wide range of organic ligands" (Haj *et al.*, 2020; Mighani, 2020; Segura *et al.*, 2016)." These ligands, which can have biological or physical properties, are often used in the preparation of transition metal complexes that can enhance the properties of the free ligands" (Al Zoubi *et al.*, 2018; El-Sonbati *et al.*, 2019; Gul *et al.*, 2024; Hosny *et al.*, 2024; Venkatesh *et al.*, 2024). Schiff bases can generate mononuclear (Middya *et al.*, 2025; Paul *et al.*, 2024), dinuclear (Uyada *et al.*, 2024; Huang *et al.*, 2024) or polynuclear complexes (Chakraborty *et al.*, 2024; Mahapatra *et al.*, 2024; Nowicka *et al.*, 2024). "The interest of chemists in these compounds has increased considerably in recent years because of the exceptional properties observed for these molecules. Some coordination compounds derived from Schiff bases have been tested for their biological activities and have been shown to be antibacterial" (Anacona *et al.*, 2021; Kargar *et al.*, 2022), antifungal (Amirthaganesan *et al.*, 2022; Frei *et al.*, 2021), anticancer (Alorini *et al.*, 2022; Jiang *et al.*,

2024), or antituberculosis (Bufarwa *et al.*, 2024; Malav & Ray, 2024). In these compounds, the presence of several metal ions can induce cooperation between the metal centers and improve physical properties such as magnetism (Chen *et al.*, 2024), fluorescence (Dong *et al.*, 2024) or optics (Turan *et al.*, 2024) and catalytic properties (Parada *et al.*, 2024; Rezaie *et al.*, 2024). "Schiff bases and their coordination compounds have also been used in the field of biomimetic chemistry to understand the behavior of some metalloenzymes" (Mandal *et al.*, 2020; Mureseanu *et al.*, 2024; Shahraki, 2022). "In particular, hydroxyacetophenone is a keto-precursor that enables the synthesis of Schiff bases used in the preparation of biomimetic complexes" (Banerjee & Paine, 2020; Chandra, 1984; Mir *et al.*, 2018; Nasr-Esfahani *et al.*, 2009). "In our previous studies, we reported the preparation and use of 2-hydroxyacetophenone-derived Schiff bases to prepare transition metal complexes with novel structures and interesting properties such as antioxidation" (Fall *et al.*, 2020; Faye *et al.*, 2021; Kébé *et al.*, 2021; Mamour *et al.*, 2018). This study aims to isolate and characterize a copper(II) Schiff base complex obtained by the condensation reaction between 2-hydroxyacetophenone and ethylenediamine.

MATERIALS AND METHODS

EXPERIMENTAL ANALYSIS

"All chemicals and solvents were of analytical reagent grade and were used directly without further purification. Elemental analyses of C, H and N were recorded on a VxRio EL Instrument. Infrared spectra were obtained on a FTIR Spectrum Two of Perkin Elmer spectrometer in the 4000-400 cm^{-1} region. The UV-Visible spectra were recorded on a Perkin Elmer Lambda UV-Vis spectrophotometer. The ^1H NMR spectra were recorded at 300 MHz and $^{13}\text{C}\{^1\text{H}\}$ NMR spectra at 75 MHz on a Bruker AC-300 instrument. The molar conductance of 1×10^{-3} M in DMF solutions of the metal complex was measured at 25 °C using a WTW LF-330 conductivity meter with a WTW conductivity cell. Room temperature magnetic susceptibility of the powdered sample was measured using a Johnson Matthey scientific magnetic susceptibility balance (Calibrant: $\text{Hg}[\text{Co}(\text{SCN})_4]$)" (Kumar, 2017).

1.1. Synthesis of 2,2'-[1,1'-(ethane-1,2-diyl)bis(nitriloethylidene)]diphenol (H_2L)

The Schiff base H_2L was synthesized by mixing 2-hydroxyacetophenone (6.02 mL, 0.050 mmol) and ethylenediamine (1.70 mL, 0.025 mmol) in 50 mL of absolute ethanol. The reaction mixture was heated under reflux for four hours. After cooling the mixture at room temperature, yellowish powder appears. The compound was filtered and recrystallized from ethanol (Kumar, 2017). Yield, 83 %, M. P. 199 °C.

1.2. Synthesis of the copper(II) complex

In a 100 mL round bottomed flask containing 20 mL of methanol, H_2L (0.2961 g; 1 mmol) was introduced with stirring. Then 10 mL of methanolic solution containing $\text{Cu}(\text{OAc})_2 \cdot 4(\text{H}_2\text{O})$ (0.2 g; 1 mmol) was added and the mixture was heated to reflux for two hours. After cooling, the brown solution was filtered, and the

filtrate was left to evaporate slowly for two weeks. Brown single crystals suitable for XRD are collected by filtration (Kumar, 2017). Yield, 67%.

1.3. Crystal structure determination

“Details of the X-rays crystal structure solution and refinement are given in Table 1. Diffraction data were collected using an ENRAF NONIUS Kappa CCD diffractometer with graphite monochromatized MoK α radiation ($\lambda = 0.71073 \text{ \AA}$). All data were corrected for Lorentz and polarization effects. No absorption correction was applied. Complex scattering factors were taken from the program package SHELXTL (Sheldrick, 2015a). The structures were solved by direct methods which revealed the position of all non-hydrogen atoms. All the structures were refined on F^2 by a full-matrix least-squares procedure using anisotropic displacement parameters for all non-hydrogen atoms” (Sheldrick, 2015b). “The hydrogen atoms of water molecules and NH groups were located in the Fourier difference maps and refined. Others H atoms (CH and CH₃ groups) were geometrically optimized and refined as riding models by AFIX instructions. Molecular graphics were generated using ORTEP-3” (Farrugia, 2012).

Table 1. Crystallographic data and refinement parameter for the complex.

Chemical formula	[2(C ₁₈ H ₁₈ CuN ₂ O ₂)]·H ₂ O
M_r	733.80
Crystal system, space group	Monoclinic, $C2/c$
Temperature (K)	293
a (Å)	26.131 (6)
b (Å)	7.293 (5)
c (Å)	17.256 (4)
β (°)	106.10 (5)
V (Å ³)	3160 (2)
Z	4
Radiation type	Mo $K\alpha$
μ (mm ⁻¹)	1.40
Crystal size (mm)	0.19 × 0.08 × 0.05
No. of measured reflections	16310
No. of independent reflections	4504


No. of observed [$I > 2\sigma(I)$] reflections	2135
R_{int}	0.077
$R[F^2 > 2\sigma(F^2)]$	0.077
$wR(F^2)$	0.145
GOF	1.11
No. of reflections	4504
No. of parameters	215
$\Delta\rho_{\text{max}}, \Delta\rho_{\text{min}}$ ($e \text{ \AA}^{-3}$)	1.03, -0.72

2. RESULTS AND DISCUSSION

2.1. General study

The symmetrical Schiff base ligand 2,2'-[1,1'-(ethane-1,2-diyl)bis(nitriloethylidyne)]diphenol (H_2L) was prepared by refluxing acetophenone and ethylene diamine in 2:1 ratio as shown in Scheme 1. **The spectroscopic data are summarized in Table 2.** The ^1H NMR spectrum of the ligand exhibits a broad signal pointed at 12.93 ppm attributed to the phenolic protons. The signals of the aromatic protons appear in the range 6.80-7.55 ppm as a complex multiplet. The signals due to the methylene and the methyle protons appear, respectively, at 3.99 ppm and 2.52 ppm. The ^{13}C NMR spectrum reveal two signals at 172.39 and 163.63 ppm attributed to the carbon atoms of the azomethine group and the $C_{\text{ipso-OH}}$ of the aromatic ring. The signals due to the other aromatic carbon atoms appear in the range 117.49-132.79 ppm. The signals due to the methyle and the methylene carbon atoms are pointed, respectively, at 14.80 and 50.12 ppm. The FTIR of the free ligand shows a broad band at 3423 cm^{-1} attributed to the $\nu_{\text{O-H}}$ stretching of the phenolic group. The $\nu_{\text{C=N}}$ stretching band is pointed at 1612 cm^{-1} . The $\nu_{\text{C-O}}$ stretching appears at 1285 cm^{-1} . The bands which appear in the range $1550\text{-}1450 \text{ cm}^{-1}$ are attributed to the C=C of the aromatic ring. The electronic spectrum of the free ligand shows two bands pointed at 243 nm and 324 nm which are attributed to the $\pi \rightarrow \pi^*$ and $n \rightarrow \pi^*$ of the phenyl ring and -C=N- bond (Kendel *et al.*, 2020), respectively. The copper complex was synthesized by mixing H_2L ligand in 15 mL of methanol with copper acetate in a molar ratio 1:1. Elemental analyses gives result in agreement with the chemical formula obtained from X-ray diffraction study. The FTIR spectrum of the complex reveals the absence of the band due to $\nu_{\text{O-H}}$ **which appeared** in the spectrum of the free ligand at 3423 cm^{-1} . This observation is indicative of the deprotonation of the phenolic groups. The band due to $\nu_{\text{C=N}}$ stretching shifts to low frequencies and appears at 1588 cm^{-1} . This fact is indicative of the involvement of the nitrogen atom of the azomethine moiety in the coordination. The band due to $\nu_{\text{C-O}}$, which appears at 1285 cm^{-1} in the spectrum of the free ligand is pointed at 1220 cm^{-1} , **suggesting** the coordination of the phenolate oxygen to the copper(II). The

electronic spectrum of the complex shows four bands. The bands at 230 and 279 nm are attributed to the $\pi \rightarrow \pi^*$ and $n \rightarrow \pi^*$ of the aromatic ring and the $-C=N-$ bond. The band at 372 nm is due to the MLCT (Saha *et al.*, 2014). The electronic spectrum displays also a $d \rightarrow d$ band at 554 nm which is attributed to ${}^2B_{1g} \rightarrow {}^2A_{1g}$ transition (Dey *et al.*, 2021). The magnetic moment value of $1.76 \mu_B$ of the copper(II) complex is in good agreement with paramagnetic copper(II) complexes (Srivastava *et al.*, 2021) having one unpaired electron with d^9 system. The conductimetry measurement ($10 \text{ S}\cdot\text{cm}^2\cdot\text{mol}^{-1}$) showed that the copper (II) complex is neutral electrolyte type (Geary, 1971). On standing for two weeks the conductance remains quite constant, indicating good stability of the complex in DMF solution.



Scheme 1. Chemical diagrams for the ligand and the complex.

Table 2. Spectroscopic data for the ligand and the complex.

H ₂ L		[Cu(L)]·(H ₂ O) _{0.5}
FTIR (ν (cm^{-1}))		
$\nu_{\text{O-H}}$	3423	3458
$\nu_{\text{C=N}}$	1612	1588
$\nu_{\text{C=C}}$	1550-1450	1455-1550
$\delta_{\text{C=O}}$	1285	1220
UV-Visible (λ (nm))		
$\pi \rightarrow \pi^*$	243	230
$n \rightarrow \pi^*$	324	279
MLCT	-	372
${}^2B_{1g} \rightarrow {}^2A_{1g}$	-	554
${}^1\text{H NMR}$ (δ (ppm))		Λ ($\text{S}\cdot\text{cm}^2\cdot\text{mol}^{-1}$)
δ_{CH_3}	2.52	Fresh solution 10
δ_{CH_2}	3.99	Old solution 12
δ_{HAr}	6.80-7.55	Magnetic Moment

δ_{OH}	12.93	$\mu_{\text{eff}} (\mu_{\text{B}})$	1.76
^{13}C NMR (δ (ppm))			-
δ_{CH_3}	14.80		-
δ_{CH_2}	50.12		-
δ_{CAr}	117.49-132.79		-
$\delta_{\text{CAr-OH}}$	163.63		-
$\delta_{\text{C=N}}$	172.39		-
Elemental analyses			
% C : Calc. (Found)	72.95 (72.91)	58.92 (58.88)	
% H : Calc. (Found)	6.80 (6.77)	5.22 (5.17)	
% N : Calc. (Found)	9.45 (9.40)	7.64 (7.59)	

2.2. Description of the structure of the complex $[\text{Cu}(\text{C}_{18}\text{H}_{18}\text{O}_2\text{N}_2)] \cdot (\text{H}_2\text{O})_{0.5}$

The molecular structure of the copper(II) complex presented in this work was solved by X-ray diffraction and is illustrated Fig. 1. The main bond lengths and angles are shown in Table 3. The asymmetric unit is composed of one copper(II) ion, one dinegative ligand molecule and a half uncoordinated water molecule. The ligand is coordinated through two nitrogen atoms of azomethine moieties and two phenolate oxygen atoms, yielding a tetracoordinated copper(II) ion which is situated in an N_2O_2 environment. For elucidating the geometry around the metal ion in tetracoordinated complexes, the trigonality parameter τ_4 ($\tau_4 = [360^\circ - (\alpha + \beta)]/141$ (α and β are the largest angles around the metal center)) is used : $\tau_4 = 1$ define a perfect tetrahedral geometry and $\tau_4 = 0$ describe a perfect square planar geometry) (Singh *et al.*, 2017). The τ_4 value of 0.068 is indicative of a distorted square planar geometry around the copper(II) for the title complex. The *cisoid* angles which are in the range $[86.83 (14)^\circ - 92.99 (15)^\circ]$ deviate from the ideal value of 90° . The *transoid* angles values of $174.91(16)^\circ$ [N1-Cu1-O2] and $175.55 (17)^\circ$ [N2-Cu1-O1] deviate from the ideal value of 180° . The sum of the angles subtended by the atoms defining the square planar geometry is 360.39° . The coordinating atoms N1/N2/O1/O2, from the ligand are, approximately, in the same plane (planar rms deviation of 0.0785 \AA) and the copper atom is located 0.004 \AA out of this plane. Upon coordination of the ligand to the copper(II), one five-membered ring and two six-membered rings were formed. The atoms defining the five membered-ring Cu1/N1/C9/C10/N2 are coplanar with rms deviation of 0.0835 \AA . The atoms defining the six membered-ring Cu1/O1/C1/C6/C7/N1 and Cu1/O2/C18/C13/C11/N2 are coplanar with rms deviation of 0.0676 \AA and 0.0462 \AA , respectively. The mean plane of the five-membered ring forms dihedral angle of $5.852(2)^\circ$ and $4.591(2)^\circ$ with the means planes of the two six-membered-rings, respectively. The means planes of the two six-membered-rings form a dihedral angle of $4.472(1)^\circ$. The Cu-N bond lengths [Cu1-N1 = $1.931 (4) \text{ \AA}$ and Cu1-N2 = $1.934 (4) \text{ \AA}$] are longer than the Cu-O [Cu1-O1 = $1.878 (3) \text{ \AA}$ and Cu1-O2 = $1.876 (3) \text{ \AA}$] distances. These values are close proximity to the values reported for the complex Chlorido(2-{1-[(2-morpholinoethyl)-imino]ethyl}phenolato- $\kappa^3\text{N,N,O}$)-copper(II) (Ikmal Hisham *et al.*, 2011). The bond length of $1.299(5) \text{ \AA}$ for C1-O1 and $1.303(5)$ for C18-O2 are compatible with data reported for C-O_{phenolate} distances (Zhang *et*

al., 2009; Zhao *et al.*, 2008). Additionally, C7-N1 [1.286 (5) Å] and C11-N2 [1.292 (6) Å] have double bond character and the values of 1.471(5) Å and 1.461(5) Å for C9-N1 and C10-N2 are indicative of single bond character (Seck *et al.*, 2020). No hydrogen-bonds are found in the structure. The packing diagram of the structure is illustrated in Fig. 2.




Fig. 1. Crystal structure of the mononuclear copper (II) complex.




Fig. 2. The packing of the complex in the crystal structure.

Table 3. Selected geometric parameters (Å, °).

Cu1—O2	1.876 (3)	O1—C1	1.299 (5)
Cu1—O1	1.878 (3)	N2—C11	1.292 (6)
Cu1—N1	1.931 (4)	N2—C10	1.461 (5)
Cu1—N2	1.934 (4)	N1—C7	1.286 (5)
O2—C18	1.303 (5)	N1—C9	1.471 (5)
O2—Cu1—O1	86.83 (14)	O2—Cu1—N2	92.99 (15)
O2—Cu1—N1	174.91 (16)	O1—Cu1—N2	175.55 (17)
O1—Cu1—N1	92.54 (15)	N1—Cu1—N2	88.03 (15)

3. CONCLUSION

The prepared organic ligand H_2L coordinates to copper(II) ion yielding a mononuclear complex. The compounds are characterized by elemental analysis, FTIR, UV-Vis and NMR spectroscopies, room temperature magnetic moment measurement and molar conductivity. The structure of the mononuclear copper (II) complex is established by single-crystal X-ray diffraction. The complex is non-ionic electrolyte in DMF solution. The organic molecule H_2L acts in its **dinegative** form as tetradentate ligand through two nitrogen azomethine atoms and two phenolate oxygen atoms. Considering the magnetic moment, infrared spectra and the electronic spectrum data for Cu(II) complex, square planar geometry is proposed for the Cu(II) complex as observed in X-ray diffraction study.

4. SUPPORTING INFORMATION

CCDC-1844618 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via Available: <https://www.ccdc.cam.ac.uk/structures/>, or by, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

REFERENCES

Al Zoubi, W., Al-Hamdani, A. A. S., Ahmed, S. D., & Ko, Y. G. (2018). Synthesis, characterization, and biological activity of Schiff bases metal complexes. *Journal of Physical Organic Chemistry*, 31(2), e3752.

Available:<https://doi.org/10.1002/poc.3752>

Alorini, T. A., Al-Hakimi, A. N., Saeed, S. E.-S., Alhamzi, E. H. L., & Albadri, A. E. A. E. (2022). Synthesis, characterization, and anticancer activity of some metal complexes with a new Schiff base ligand. *Arabian Journal of Chemistry*, 15(2), 103559.

Available:<https://doi.org/10.1016/j.arabjc.2021.103559>

Amirthaganesan, K., Vadivel, T., Dhamodaran, M., & Chandraboss, V. L. (2022). In vitro antifungal studies of Ruthenium (III) complex derived from chitosan Schiff bases. *Materials Today: Proceedings*, 60, 1716–1720.

Available:<https://doi.org/10.1016/j.matpr.2021.12.265>

Anacona, J. R., Santaella, J., Al-shemary, R. K. R., Amenta, J., Otero, A., Ramos, C., & Celis, F. (2021). Ceftriaxone-based Schiff base transition metal(II) complexes. Synthesis, characterization, bacterial toxicity, and DFT calculations. Enhanced antibacterial activity of a novel Zn(II) complex against *S. aureus* and *E. coli*. *Journal of Inorganic Biochemistry*, 223, 111519.

Available:<https://doi.org/10.1016/j.jinorgbio.2021.111519>

Banerjee, S., & Paine, T. K. (2020). Bioinspired iron(II)- β -diketonate and iron(II)- α -hydroxy ketone complexes of a carbanionic N_3C ligand: Oxidation of metal center vs C-C bond cleavage of co-ligand with dioxygen. *Inorganica Chimica Acta*, 501, 119200.

Available:<https://doi.org/10.1016/j.ica.2019.119200>

Bufarwa, S. M., El-Sefait, R. M., Thbayh, D. K., Belaidi, M., Al-Shemary, R. K., Abdusamea, R. M., El-Ajaily, M. M., Fiser, B., Bader, H. A., Saleh, A. A., & Bufarwa, M. M. (2024). Antituberculosis, antimicrobial, antioxidant, cytotoxicity and anti-inflammatory activity of Schiff base derived from 2,3-diaminophenazine moiety and its metal(II) complexes: Structural elucidation, computational aspects, and biological evaluation. *Reviews in Inorganic Chemistry*.

Available:<https://doi.org/doi:10.1515/revic-2024-0007>

Chakraborty, P., Chatterjee, A., Benmansour, S., Gómez-García, C. J., & Dey, S. K. (2024). Synthesis, characterization and magnetic properties of a hexanuclear copper (II) complex with hydroxido, alkoxido and phenoxido bridges. *Inorganica Chimica Acta*, 122442.

Available:<https://doi.org/10.1016/j.ica.2024.122442>

Chandra, S. (1984). Copper (II) complexes with some semicarbazones; possible biomimetic models. *Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry*, 14(7), 957–967.

Available:<https://doi.org/10.1080/00945718408058277>

Chen, J.-T., Yan, H., Wang, T., & Sun, W.-B. (2024). Heteronuclear complexes $[MDyM]$ ($M=Cu$; Zn ; Ni) constructed by Schiff base ligands with different amine backbone exhibiting significant single-molecule magnets. *Journal of Molecular Structure*, 1311, 138473.

Available:<https://doi.org/10.1016/j.molstruc.2024.138473>

Dey, L., Rabi, S., Hazari, S. K. S., Roy, T. G., Buchholz, A., & Plass, W. (2021). Copper(II) and nickel(II) complexes of an N-pendent bis-(cyanoethyl) derivative of an isomeric hexamethyl tetraazamarocyclic ligand: Synthesis, characterization, electrolytic behavior and antimicrobial studies. *Inorganica Chimica Acta*, 517, 120172.

Available:<https://doi.org/10.1016/j.ica.2020.120172>

Dong, Q.-W., Yang, L.-Z., Yue, T.-C., Wang, D.-Z., & Wang, L.-L. (2024). New complexes based on reduced Schiff base carboxylic acid ligand: Electric properties, fluorescence properties and detection of Fe³⁺. *Journal of Molecular Structure*, 1295, 136747.

Available:<https://doi.org/10.1016/j.molstruc.2023.136747>

El-Sonbati, A. Z., Mahmoud, W. H., Mohamed, G. G., Diab, M. A., Morgan, Sh. M., & Abbas, S. Y. (2019). Synthesis, characterization of Schiff base metal complexes and their biological investigation. *Applied Organometallic Chemistry*, 33(9), e5048.

Available:<https://doi.org/10.1002/aoc.5048>

Fall, N., Faye, F. D., Gaye, A. A., Diouf, O., & Gaye, M. (2020). Synthesis of mono and bis-substituted asymmetrical compounds, 1-(2'-hydroxy-3'-methoxybenzylidene)carbonohydrazide and 1-(2'-hydroxy-3'-methoxybenzylidene)-5-(1'pyridylmethylene)carbonohydrazide: Structural characterization and antioxidant activity study. *Journal of Applied Chemistry (IOSR-JAC)*, 13(12), 22–30.

Available:<https://doi:10.9790/5736-1312022230>

Farrugia, L. J. (2012). WinGX and ORTEP for Windows: An update. *Journal of Applied Crystallography*, 45(4), 849–854.

Available:<https://doi.org/10.1107/S0021889812029111>

Faye, F., Guèye, A., Camara, P. S., Gaye, A. A., Tamboura, F. B., Gruber, N., & Gaye, M. (2021). Synthesis of Schiff bases compounds from oxamic hydrazide: spectroscopic characterization, X-ray diffraction structure and antioxidant activity study. *American Journal of Applied Chemistry*, 9(1), 6–12.

Available:<https://doi.org/10.11648/j.ajac.20210901.12>

Frei, A., King, A. P., Lowe, G. J., Cain, A. K., Short, F. L., Dinh, H., Elliott, A. G., Zuegg, J., Wilson, J. J., & Blaskovich, M. A. T. (2021). Nontoxic cobalt(III) Schiff base complexes with broad-spectrum antifungal activity. *Chemistry – A European Journal*, 27(6), 2021–2029.

Available:<https://doi.org/10.1002/chem.202003545>

Geary, W. J. (1971). The use of conductivity measurements in organic solvents for the characterisation of coordination compounds. *Coordination Chemistry Reviews*, 7(1), 81–122.

Available:[https://doi.org/10.1016/S0010-8545\(00\)80009-0](https://doi.org/10.1016/S0010-8545(00)80009-0)

Gul, S., Jan, F., Alam, A., Shakoor, A., Khan, A., AlAsmari, A. F., Alasmari, F., Khan, M., & Bo, L. (2024). Synthesis, molecular docking and DFT analysis of novel bis-Schiff base derivatives with thiobarbituric acid for α -glucosidase inhibition assessment. *Scientific Reports*, 14(1), 3419.

Available:<https://doi.org/10.1038/s41598-024-54021-z>

Haj, N. Q., Mohammed, M. O., & Mohammad, L. E. (2020). Synthesis and Biological Evaluation of Three New Chitosan Schiff Base Derivatives. *ACS Omega*, 5(23), 13948–13954.

Available:<https://doi.org/10.1021/acsomega.0c01342>

Hosny, S., Abdelfatah, A., & Gaber, G. A. (2024). Synthesis, characterization, synergistic inhibition, and biological evaluation of novel Schiff base on 304 stainless steel in acid solution. *Scientific Reports*, 14(1), 470.

Available:<https://doi.org/10.1038/s41598-023-51044-w>

Huang, M., Jiao, L., Yang, J., Ning, D., Xu, J., Wu, Q., & Weng, Z. (2024). Regulating effect of the halogen atoms on the cofacial dinuclear Schiff base complexes: Synthesis, spectroscopy, electrochemistry, and DFT calculations. *Applied Organometallic Chemistry*, 38(7), e7536.

Available:<https://doi.org/10.1002/aoc.7536>

Ikmal Hisham, N. A., Suleiman Gwaram, N., Khaledi, H., & Mohd Ali, H. (2011). Chlorido(2-{1-[(2-morpholinoethyl)imino]ethyl}phenolato- κ^3 N,N',O)copper(II). *Acta Crystallographica Section E*, 67(1), m57.

Available:<https://doi.org/10.1107/S1600536810051160>

Jiang, M., Su, X., Zhong, X., Lan, Y., Yang, F., Qin, Y., & Jiang, C. (2024). Recent development of Schiff-base metal complexes as therapeutic agents for lung cancer. *Journal of Molecular Structure*, 1318, 139403.

Available:<https://doi.org/10.1016/j.molstruc.2024.139403>

Kargar, H., Ashfaq, M., Fallah-Mehrijardi, M., Behjatmanesh-Ardakani, R., Munawar, K. S., & Tahir, M. N. (2022). Unsymmetrical Ni(II) Schiff base complex: Synthesis, spectral characterization, crystal structure analysis, Hirshfeld surface investigation, theoretical studies, and antibacterial activity. *Journal of Molecular Structure*, 1265, 133381.

Available:<https://doi.org/10.1016/j.molstruc.2022.133381>

Kébé, M., Thiam, I. E., Sow, M. M., Diouf, O., Barry, A. H., Sall, A. S., Retailleau, P., & Gaye, M. (2021). Hexanuclear copper(II) complex of 2-hydroxy-*N,N*-bis[1-(2-hydroxyphenyl)ethylidene]propane-1,3-diamine incorporating an open-cubane core. *Acta Crystallographica Section E*, 77(7), 708–713.

Available:<https://doi.org/10.1107/S2056989021005570>

Kendel, A., Miljanić, S., Kontrec, D., Soldin, Ž., & Galić, N. (2020). Copper(II) complexes of aroylhydrazones: Preparation and structural characterization. *Journal of Molecular Structure*, 1207, 127783.

Available:<https://doi.org/10.1016/j.molstruc.2020.127783>

Kumar, C. (2017). Synthesis, characterization and biological evaluations of heterobinuclear complexes of Cu^{II} and Ni^{II} Schiff base of ethylenediamine and *o*-hydroxyacetophenone with alkali metal salts of *o*-nitrophenol. *Journal of Indian Chemical Society*, 94, 857-861.

Available:<https://doi.org/10.5281/zenodo.5595126>

Mahapatra, T. S., Roy, B. C., Dutta, B., Lengyel, J., Shatruck, M., & Ray, D. (2024). Structures and magnetic properties of a trinuclear angular [Ni₃] and a heptanuclear wheel-like [Ni₇] complexes with a Schiff base ligand. *Polyhedron*, 249, 116782.

Available:<https://doi.org/10.1016/j.poly.2023.116782>

Malav, R., & Ray, S. (2024). Recent developments on the synthesis of copper and cobalt-Schiff base complexes and their assessment as anti-tuberculosis drugs. *Chemical Papers*, 78(8), 4623–4646.

Available:<https://doi.org/10.1007/s11696-024-03425-2>

Mamour, S., Mayoro, D., Elhadj Ibrahima, T., Mohamed, G., Aliou Hamady, B., & Ellena, J. (2018). {1-[1-(2-Hydroxyphenyl)ethylidene]-2-(pyridin-2-yl-κN)hydrazine-κ²N, O}-[1-(2-oxidophenyl)ethylidene]-2-(pyridin-2-yl-κN)hydrazine-κ²N, O}nickelate(II) nitrate hemihydrate. *Acta Crystallographica Section E*, 74(5), 642–645.

Available:<https://doi.org/10.1107/S2056989018005261>

Mandal, A., Sarkar, A., Adhikary, A., Samanta, D., & Das, D. (2020). Structure and synthesis of copper-based Schiff base and reduced Schiff base complexes: A combined experimental and theoretical investigation of biomimetic catalytic activity. *Dalton Transaction*, 49(43), 15461–15472.

Available:<https://doi.org/10.1039/D0DT02784G>

Middya, P., Sarkar, S. D., & Chattopadhyay, S. (2025). Use of tridentate NNO donor salicylidine Schiff base ligands for the synthesis of mononuclear cobalt complexes having the general formula [Co(ABC)(DE)X]: Synthetic strategy, structure and application. *Journal of Molecular Structure*, 1322, 140242.

Available:<https://doi.org/10.1016/j.molstruc.2024.140242>

Mighani, H. (2020). Schiff Base polymers: Synthesis and characterization. *Journal of Polymer Research*, 27(6), 168.

Available:<https://doi.org/10.1007/s10965-020-02080-x>

Mir, J. M., Roy, S., Vishwakarma, P. K., & Maurya, R. C. (2018). *cis*-Dioxomolybdenum(VI) complex of *N*-*o*-hydroxyacetophenone-isonicotinic acid hydrazide as nosocomial anti-infectious agent: Experimental and theoretical study. *Journal of the Chinese Advanced Materials Society*, 6(3), 282–300.

Available:<https://doi.org/10.1080/22243682.2018.1466727>

Mureseanu, M., Bleotu, I., Spînu, C.-I., & Cioatera, N. (2024). Anchoring of Copper(II)–Schiff base complex in SBA-15 matrix as efficient oxidation and biomimetic catalyst. *International Journal of Molecular Sciences*, 25(2).

Available:<https://doi.org/10.3390/ijms25021094>

Nasr-Esfahani, M., Montazerzohori, M., & Akhlaghi, P. (2009). Efficient Biomimetic Oxidative Decarboxylation of Some Carboxylic Acids Catalyzed by a Manganese (III) Schiff Base Complex. *Bulletin Korean Chemical Society*, 30(7), 1583–1587.

Available:[https://doi.org/chrome-](https://doi.org/chrome-extension://efaidnbmninnibpcjpcglclefindmkaj/Available:https://web.archive.org/web/20170819211938id_/http://ocean.kisti.re.kr/downfile/volume/chemical/JCGMCS/2009/v30n7/JCGMCS_2009_v30n7_1583.pdf)

[extension://efaidnbmninnibpcjpcglclefindmkaj/Available:https://web.archive.org/web/20170819211938id_/http://ocean.kisti.re.kr/downfile/volume/chemical/JCGMCS/2009/v30n7/JCGMCS_2009_v30n7_1583.pdf](https://web.archive.org/web/20170819211938id_/http://ocean.kisti.re.kr/downfile/volume/chemical/JCGMCS/2009/v30n7/JCGMCS_2009_v30n7_1583.pdf)

Nowicka, D., Kubicki, M., Patroniak, V., Łuczak, T., & Gorczyński, A. (2024). Self-assembly of simple Schiff base ligand into unique saddle-type [4x4] tetranuclear architecture and its application as selective voltammetric dopamine sensor in aqueous conditions. *Electrochimica Acta*, 476, 143754.

Available:<https://doi.org/10.1016/j.electacta.2023.143754>

Parada, J., Atria, A. M., Avilés, C., Mendoza, J., Basualto, C., Parra-Melipan, S., Zúñiga, C., & Aguirre, P. (2024). Recyclable Ni(II), Cu(II), and Co(II) complexes containing Schiff base immobilized on nanosilica as catalysts in oxidation of hydrocarbons. *Inorganica Chimica Acta*, 572, 122303.

Available:<https://doi.org/10.1016/j.ica.2024.122303>

Paul, S., Sanjurani, T., Talukdar, J., Barman, P., Das, A., & Pandey, P. (2024). Unraveling the Potential of Mononuclear Zn(II) and Cu(II) Schiff Base Metal Complexes: Microwave-Aided Synthesis, Theoretical Analysis and Application in Biomimetic Catalysis, CT-DNA Interaction, and In Vitro Biological Assays. *Applied Organometallic Chemistry*, 38(12), e7749.

Available:<https://doi.org/10.1002/aoc.7749>

Rezaie, M.-A., Khojastehnezhad, A., & Shiri, A. (2024). Post-synthetic modification of Zr-based metal organic framework by schiff base zinc complex for catalytic applications in a click reaction. *Scientific Reports*, 14(1), 24644.

Available:<https://doi.org/10.1038/s41598-024-76199-y>

Saha, S., Sasmal, A., Choudhury, C. R., Gómez-García, C. J., Garribba, E., & Mitra, S. (2014). A new linear double phenoxide-bridged trinuclear Cu(II) Schiff base complex: Synthesis, crystallographic elucidation, magneto-structural correlation and DFT Study. *Polyhedron*, 69, 262–269.

Available:<https://doi.org/10.1016/j.poly.2013.12.006>

Seck, T. M., Faye, F. D., Gaye, A. A., Thiam, I. E., Diouf, O., Gaye, M., & Retailleau, P. (2020). Synthesis of mono and bis-substituted asymmetrical compounds, (1-(pyridin-2-yl)ethylidene)carbonohydrazide and 1-(2'-hydroxybenzylidene)-5-(1'-

pyridylethylidene)carbonohydrazone: Structural characterization and antioxidant activity study. *European Journal of Chemistry*, 11(4), Article 4.

Available:<https://doi.org/10.5155/eurjchem.11.4.285-290.2023>

Segura, J. L., Mancheño, M. J., & Zamora, F. (2016). Covalent organic frameworks based on Schiff-base chemistry: synthesis, properties and potential applications. *Chem. Soc. Rev.*, 45(20), 5635–5671.

Available:<https://doi.org/10.1039/C5CS00878F>

Shahraki, S. (2022). Schiff base compounds as artificial metalloenzymes. *Colloids and Surfaces B: Biointerfaces*, 218, 112727.

Available:<https://doi.org/10.1016/j.colsurfb.2022.112727>

Sheldrick, G. M. (2015a). Crystal structure refinement with SHELXL. *Acta Crystallographica Section C*, 71(1), 3–8.

Available:<https://doi.org/10.1107/S2053229614024218>

Sheldrick, G. M. (2015b). SHELXT – Integrated space-group and crystal-structure determination. *Acta Crystallographica Section A*, 71(1), 3–8.

Available:<https://doi.org/10.1107/S2053273314026370>

Singh, Y. P., Patel, R. N., Singh, Y., Choquesillo-Lazarte, D., & Butcher, R. J. (2017). Classical hydrogen bonding and stacking chelate rings in new copper(ii) complexes. *Dalton Trans.*, 46(9), 2803–2820.

Available:<https://doi.org/10.1039/C6DT04661D>

Srivastava, A. K., Yadav, P., Srivastava, K., & Prasad, J. (2021). Synthesis, characterization, biological and electrochemical investigation of copper (II) complexes containing 4-chloro-2-[2, 6-diisopropylphenylimino) methyl] phenol Schiff base ligand and aromatic diinines. *Chemical Data Collections*, 32, 100659.

Available:<https://doi.org/10.1016/j.cdc.2021.100659>

Turan, N., Seymen, H., Gündüz, B., Buldurun, K., & Çolak, N. (2024). Synthesis, characterization of Schiff base and its metal complexes and investigation of their electronic and photonic properties. *Optical Materials*, 148, 114802.

Available:<https://doi.org/10.1016/j.optmat.2023.114802>

Uyada, K. A. H, Mahesha, Pampa, K. J., Harohally, N. V., Krishnamurthy, C., Jathi, K., Ahmad, A., Alshammari, M. B. & Lokanath, N. K. (2024). Structural Investigation of Schiff Base Ligand and Dinuclear Copper Complex: Synthesis, Crystal Structure, Computational, and Latent Fingerprint Analysis. *ACS Omega*, 9(28), 30109–30119.

Available:<https://doi.org/10.1021/acsomega.3c07536>

Venkatesh, G., Vennila, P., Kaya, S., Ahmed, S. B., Sumathi, P., Siva, V., Rajendran, P., & Kamal, C. (2024). Synthesis and Spectroscopic Characterization of Schiff Base Metal Complexes, Biological Activity, and Molecular Docking Studies. *ACS Omega*, 9(7), 8123–8138.

Available:<https://doi.org/10.1021/acsomega.3c08526>

Zhang, J., Chen, X.-D., Zhang, H.-H., & Sun, B.-W. (2009). Bis(μ -2-[1-(2-Pyridylmethylimino)ethyl]phenolato)bis-[azidocopper(II)]. *Acta Crystallographica Section E*, 65(9), m1136.

Available:<https://doi.org/10.1107/S1600536809030475>

Zhao, G.-Q., Xue, L.-W., Hao, C.-J., Chen, L.-H., & Wu, H.-T. (2008). [Bis(3,5-dimethylpyrazol-1-yl)methane] $\{N$ -[1-(2-oxidophenyl)ethylidene]-DL-alaninato)copper(II) monohydrate. *Acta Crystallographica Section E*, 64(12), m1553.

Available:<https://doi.org/10.1107/S1600536808037264>