

Nutrient release pattern and soil enzyme activities in calcareous and non-calcareous soils as influenced by PROM

ABSTRACT

A study was conducted at Post Graduate Laboratory, Division of Soil Science, College of Agriculture, Pune to study the nutrient release pattern and soil enzyme activities in calcareous and non-calcareous soils as influenced by various levels of phosphorus using organic and inorganic sources. An incubation experiment was laid out in factorial completely randomized design with seven treatments replicated three times. The study demonstrated that applying 100% P_2O_5 through Phosphate Rich Organic Manure (PROM) significantly improved soil nutrient status, particularly in non-calcareous soils, with reduction in soil pH and calcium carbonate, improving organic carbon, and enhancing the availability of macronutrients (nitrogen, phosphorus, potassium). Enzyme activities, viz urease, dehydrogenase and alkaline phosphatase were higher in non-calcareous soils, correlating positively with nutrient availability, particularly with phosphorus and micronutrients such as iron and manganese. Phosphate Rich Organic Manure (PROM) application was more effective than chemical fertilizers in nutrient mobilization and improving soil enzyme activity over 120 days incubation. Strong correlations were observed between enzyme activity and micronutrient availability, especially in treatments receiving 100% P_2O_5 through PROM, indicating its potential to enhance both nutrient cycling and soil fertility.

Keywords: Calcareous soil, PROM, incubation, nutrient release, enzyme activities, interaction and correlation.

1. INTRODUCTION

Phosphorus is the second most essential macronutrient for plants, after nitrogen (Reddy *et al.*, 2004) and is crucial for root development, nodulation, nitrogen fixation, and various physiological functions such as energy transfer, cell division, and photosynthesis. However, in calcareous and alkaline soils, phosphorus availability is limited due to its fixation with calcium, iron, and aluminium, leading to poor plant uptake. Lack of phosphorus in calcareous and alkaline soils is a major global problem. In calcareous/alkaline soils, phosphorus availability to plant is restricted by its reduced mobility in soils and higher fixation (Shen *et al.*, 2011). Phosphorus solubilizing microorganisms (PSB) and soil enzymes like phosphatase play a crucial role in converting immobilized P into plant-available forms, especially in phosphorus-deficient soils. Non-traditional phosphorus fertilizers like Phosphate Rock (PR) and Phosphate Rich Organic Manure (PROM) offer sustainable alternatives to

conventional chemical fertilizers by maintaining phosphorus availability for longer periods and enhancing crop productivity. PROM, in particular, has been recognized as an eco-friendly solution, approved by the Government of India, to improve phosphorus availability in calcareous soils. Despite India's significant rock phosphate reserves, only a fraction is suitable for fertilizer production. Therefore, further research on optimizing PROM's production and its effects on phosphorus availability in different soil types is essential for improving phosphorus use efficiency and sustainable agriculture.

2. EXPERIMENTAL DETAILS

An incubation study was conducted in Post Graduate Laboratory, Division of Soil Science, College of Agriculture, Pune. The soil samples for the incubation study were collected from the PG Instructional Farm, College of Agriculture, Pune. The CaCO_3 less than 5 % and more than 10 % were selected as non-calcareous and calcareous soils for study.

list 1 :Treatments Details:

Tr.No.	Factor 'A' (Soil type)	Tr.No.	Factor 'B' (Nutrient management)
1	Highly calcareous soil (A_1)	1	Absolute control
2	Low calcareous soil (A_2)	2	Recommended dose of fertilizers through DAP (50:75:45 N: P_2O_5 : K_2O kg ha ⁻¹)
		3	Recommended dose of fertilizers through SSP (50:75:45 N: P_2O_5 : K_2O kg ha ⁻¹)
		4	25 % P_2O_5 through PROM
		5	50 % P_2O_5 through PROM
		6	75 % P_2O_5 through PROM
		7	100 % P_2O_5 through PROM

Note:

1. The recommended dose of soybean (50:75:45 N: P_2O_5 : K_2O kg ha⁻¹) was used for incubation study.
2. The uniform dose of N and K_2O was common to all the treatments except absolute control.

The recommended dose of fertilizer (50:75:45 kg ha⁻¹ of N, P_2O_5 and K_2O) of soybean was used for this incubation study except absolute control treatment. The recommended dose of phosphorus was applied through PROM (organic), DAP and SSP (inorganic) sources. The nitrogen and potassium were supplied through urea and muriate of potash, respectively. The PROM was prepared at Division of Soil Science, College of Agriculture, Pune and its proximate analysis was done before start of incubation.

Table: 1 Proximate analysis of PROM

Sr. No	Parameters	Unit	PROM
1	pH (1:10)	-	7.18
2	EC	(dS m ⁻¹)	1.74
3	Moisture	(%)	24.02
4	Organic Carbon	(%)	21.06
5	Total N	(%)	0.78
6	Total P	(%)	14.57
7	Total K	(%)	0.37
8	Total Fe	(mg kg ⁻¹)	12.70
9	Total Mn	(mg kg ⁻¹)	0.67
10	Total Zn	(mg kg ⁻¹)	2.69
11	Total Cu	(mg kg ⁻¹)	0.46
12	C:N ratio	-	24:1
13	C:P ratio	-	1.44:1

Plastic bowls of 2 kg capacity were filled with 1kg soil (2 mm) in one hundred sixty eight bowls. The calculated quantity of PROM, DAP, SSP, urea and MOP was added as per treatment. The moisture in the soil was maintained at field capacity. The set of experiment was incubated for 30, 60, 90 and 120 days in three replications using discarding method. The incubation study was conducted by using factorial completely randomized design. (Panse and Sukhatme,1985).

3. RESULTS AND DISCUSSION

3.1 Chemical properties of soil

Low soil pH was observed in non-calcareous soil and with 100% P₂O₅ applied through PROM. The lowest pH (7.60) was recorded at 120 days with non-calcareous soil and 100% P₂O₅ through PROM. Organic matter played an important role in reducing the pH of both soils as negative correlation between pH and organic matter is reported by Talashilkaret *al.* (2000). Soil EC in calcareous soil was higher, with slight improvement for RDF through DAP and SSP. At 120 days, the highest EC (0.51 dS m⁻¹) was observed in non-calcareous soil with RDF applied through DAP, showing significant interaction effects. Interaction effects with the combination of non-calcareous soil and RDF through DAP recorded higher EC (0.51 dSm⁻¹) at 120 days.

Soil organic carbon was higher in non-calcareous soil, increasing from 0.71% at 30 days to 0.80% at 120 days. The higher organic carbon content was observed due to application of 100%

P₂O₅ through PROM, with significant interaction at 30 days only. The application of rock phosphate with organic materials in gypsiferous soils increased the organic carbon and phosphorous status of soil. (Muhawish and Al-Kafaje, 2017). Non-calcareous soil recorded lower CaCO₃ than calcareous soil, decreasing from 4.12% at 30 days to 3.55% at 120 days with 100% P₂O₅ through PROM. Significant reductions were observed across all PROM treatments (100%, 75%, and 50%) over time.

Table:2 Influence of soil calcareousness and PROM on periodical changes in pH of soil

Days →	30			60			90			120		
Levels of P ₂ O ₅ ↓	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B
B ₁	8.21	7.79	8	8.17	7.73	7.95	8.17	7.73	7.95	8.15	7.69	7.92
B ₂	8.18	7.76	7.97	8.16	7.7	7.93	8.15	7.7	7.93	8.13	7.67	7.90
B ₃	8.19	7.77	7.98	8.17	7.69	7.93	8.17	7.69	7.93	8.14	7.64	7.89
B ₄	8.21	7.73	7.98	8.19	7.68	7.94	8.16	7.68	7.92	8.15	7.64	7.9
B ₅	8.19	7.72	7.96	8.16	7.69	7.93	8.14	7.69	7.92	8.13	7.62	7.88
B ₆	8.18	7.72	7.95	8.13	7.69	7.91	8.13	7.69	7.91	8.12	7.62	7.87
B ₇	8.17	7.72	7.95	8.12	7.68	7.9	8.12	7.68	7.9	8.1	7.6	7.85
Mean A	8.19	7.74	--	8.16	7.7	--	8.15	7.69	--	8.13	7.64	--
Factors	(A)	(B)	(AXB)	(A)	(B)	(AXB)	(A)	(B)	(AXB)	(A)	(B)	(AXB)
C.D. at (5%)	0.006	0.011	0.016	0.006	0.011	0.015	0.004	0.008	0.011	0.008	0.015	0.021
SE(m)±	0.002	0.004	0.006	0.002	0.004	0.005	0.001	0.003	0.004	0.003	0.005	0.007

Table:3 Influence of soil calcareousness and PROM on periodical changes in electrical conductivity of soil

Days →	30			60			90			120		
Levels of P ₂ O ₅ ↓	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B
B ₁	0.37	0.35	0.36	0.39	0.38	0.38	0.4	0.43	0.41	0.41	0.44	0.42
B ₂	0.44	0.37	0.41	0.46	0.42	0.44	0.49	0.46	0.48	0.51	0.47	0.49

B ₃	0.43	0.38	0.4	0.46	0.43	0.45	0.48	0.45	0.47	0.49	0.48	0.49
B ₄	0.39	0.36	0.37	0.4	0.39	0.39	0.41	0.41	0.41	0.43	0.4	0.42
B ₅	0.4	0.37	0.39	0.41	0.4	0.41	0.43	0.4	0.42	0.44	0.43	0.44
B ₆	0.41	0.37	0.39	0.43	0.41	0.42	0.44	0.43	0.43	0.46	0.44	0.45
B ₇	0.41	0.38	0.39	0.44	0.42	0.43	0.46	0.44	0.45	0.48	0.46	0.47
Mean A	0.41	0.37	--	0.43	0.41	--	0.44	0.43	--	0.46	0.45	--
Factors	(A)	(B)	(AXB)	(A)	(B)	(AXB)	(A)	(B)	(AXB)	(A)	(B)	(AXB)
C.D. at (5%)	0.009	0.017	NS	0.01	0.019	NS	0.01	0.019	0.028	0.011	0.02	0.029
SE(m)±	0.003	0.006	0.008	0.003	0.006	0.009	0.004	0.007	0.009	0.004	0.007	0.01

Table:4 Influence of soil calcareousness and PROM on periodical changes of organic carbon in soil

Days →	30			60			90			120		
Levels of P ₂ O ₅ ↓	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B
B ₁	0.55	0.66	0.61	0.58	0.68	0.63	0.63	0.7	0.67	0.64	0.73	0.69
B ₂	0.67	0.68	0.68	0.59	0.71	0.65	0.65	0.73	0.69	0.66	0.75	0.71
B ₃	0.65	0.7	0.67	0.62	0.73	0.67	0.66	0.75	0.71	0.68	0.77	0.73
B ₄	0.56	0.71	0.64	0.64	0.75	0.7	0.68	0.77	0.72	0.69	0.79	0.74
B ₅	0.59	0.73	0.66	0.66	0.77	0.71	0.7	0.78	0.74	0.71	0.82	0.77
B ₆	0.61	0.75	0.68	0.69	0.79	0.74	0.71	0.79	0.75	0.73	0.84	0.79
B ₇	0.63	0.77	0.7	0.7	0.82	0.76	0.73	0.83	0.78	0.75	0.88	0.82
Mean A	0.61	0.71	--	0.64	0.75	--	0.68	0.76	--	0.70	0.80	--
Factors	(A)	(B)	(AXB)	(A)	(B)	(AXB)	(A)	(B)	(AXB)	(A)	(B)	(AXB)
C.D. at (5%)	0.01	0.018	0.026	0.009	0.017	NS	0.009	0.016	NS	0.009	0.017	NS
SE(m)±	0.003	0.006	0.009	0.003	0.006	0.008	0.003	0.006	0.008	0.003	0.006	0.008

Table:5 Influence of soil calcareousness and PROM on periodical changes of CaCO₃ in soil

Days →	30			60			90			120		
Levels of P ₂ O ₅ ↓	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B
B ₁	12.45	4.21	8.33	12.37	4.17	8.27	12.3	4.08	8.19	12.24	4.01	8.13
B ₂	12.33	4.07	8.2	12.23	4.12	8.18	11.81	3.98	7.9	10.75	3.82	7.29
B ₃	12.36	4.08	8.22	12.08	4.14	8.11	11.75	4.01	7.88	10.58	3.98	7.28
B ₄	12.28	4.17	8.22	11.82	4.09	7.96	11.33	3.82	7.58	10.5	3.77	7.14
B ₅	12.15	4.14	8.15	11.33	4.08	7.71	11	3.77	7.39	10.08	3.68	6.88
B ₆	11.91	4.1	8.01	11.31	4.06	7.69	10.75	3.68	7.22	10.07	3.57	6.82
B ₇	11.42	4.09	7.75	11.25	4.04	7.64	10.58	3.57	7.08	9.8	3.55	6.68
Mean A	12.13	4.12	--	11.77	4.1	--	11.36	3.85	--	10.58	3.77	--
Factors	(A)	(B)	(AXB)	(A)	(B)	(AXB)	(A)	(B)	(AXB)	(A)	(B)	(AXB)
C.D. at (5%)	0.124	0.233	0.329	0.128	0.24	0.339	0.118	0.221	0.312	0.172	0.321	0.454
SE(m)±	0.043	0.08	0.113	0.044	0.082	0.116	0.04	0.076	0.107	0.059	0.11	0.156

3.2 Influence of PROM on soil macronutrients

The higher available nitrogen (250.51–260.16 kg ha⁻¹) was recorded in non-calcareous soil than calcareous soil (135.50–145.42 kg ha⁻¹) over the incubation period. Application of RDF through DAP recorded higher nitrogen levels, followed by SSP and 100% P₂O₅ through PROM, with non-significant interaction effects. Pandey *et al.* (2024) reported that ammonical nitrogen, the first available organic form of nitrogen for plants and microorganisms, was significantly higher over the incubation study compared to the control.

Table:6 Influence of soil calcareousness and PROM on release of nitrogen in soil

Days →	30			60			90			120		
Levels of P ₂ O ₅ ↓	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B
B ₁	116.6	237.5	177.1	119	242.4	180.7	120.4	247.3	183.8	123.6	248.4	186
B ₂	145.5	265.3	205.4	146.3	266.3	206.3	152.2	262.1	207.2	159	268.3	213.6
B ₃	142.6	259.1	200.8	143.3	260.3	201.8	147.8	268.3	208	152.6	271.2	211.9
B ₄	126.3	242.4	184.3	137.8	247.3	192.6	139.4	248.4	193.9	140.4	253.9	197.2
B ₅	137.8	247.3	192.6	139.4	248.6	194	140.4	253.9	197.2	142.8	255.2	199
B ₆	139.4	248.6	194	140.4	254.4	197.4	142.8	255.2	199	143.3	257.1	200.2
B ₇	140.3	253.3	196.8	142.8	256.5	199.6	150.2	261.6	205.9	156.3	267	211.7
Mean A	135.5	250.5	--	138.4	253.7	--	141.9	256.7	--	145.4	260.2	--
Factors	(A)	(B)	(AXB)	(A)	(B)	(AXB)	(A)	(B)	(AXB)	(A)	(B)	(AXB)
C.D. at (5%)	7.455	13.947	NS	7.409	13.862	NS	5.425	10.15	NS	4.546	8.504	NS
SE(m)±	2.56	4.79	6.774	2.545	4.76	6.732	1.863	3.486	4.929	1.561	2.921	4.13

The soil available phosphorus content (21.50–25.54 kg ha⁻¹) was observed higher in non-calcareous soil compared to calcareous soil (12.50–15.07 kg ha⁻¹) over the incubation period. The highest phosphorus (28.24 kg ha⁻¹ at 120 days) was observed with 100% P₂O₅ through PROM in non-calcareous soil, showing significant interaction effects. Kumar *et al.* (2015) reported that the increased availability of phosphorus in organically amended soils results from a large reduction in phosphorus sorption.

Table:7 Influence of soil calcareousness and PROM on release phosphorous in soil

Days →	30			60			90			120		
Levels of P ₂ O ₅ ↓	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B
B ₁	11.95	20.39	16.17	12.17	21.6	16.89	12.43	22.02	17.23	12.68	22.06	17.37
B ₂	12.67	21.59	17.13	13.02	22.45	17.74	15.71	25.14	20.43	16.11	26.25	21.18
B ₃	12.37	22.07	17.22	12.48	23.04	17.76	14.97	26.11	20.54	15.28	28.02	21.65

B ₄	12.22	20.68	16.45	12.25	21.8	17.02	12.97	22.06	17.52	14.03	22.65	18.34
B ₅	12.27	21.38	16.83	12.37	22.01	17.19	13.34	22.65	18	14.78	23.34	19.06
B ₆	12.99	22.18	17.59	14.28	23.05	18.67	15.76	26.18	20.97	16.31	28.21	22.26
B ₇	13	22.2	17.6	14.3	23.12	18.71	15.84	26.25	21.04	16.33	28.24	22.28
Mean A	12.5	21.5	--	12.98	22.44	--	14.43	24.35	--	15.07	25.54	--
Factors	(A)	(B)	(AXB)	(A)	(B)	(AXB)	(A)	(B)	(AXB)	(A)	(B)	(AXB)
C.D. at (5%)	0.011	0.021	0.03	0.068	0.127	0.18	0.049	0.092	0.131	0.018	0.034	0.048
SE(m)±	0.004	0.007	0.01	0.023	0.044	0.062	0.017	0.032	0.045	0.006	0.012	0.016

The higher available potassium content (515.21–525.67 kg ha⁻¹) was observed in non-calcareous soil than calcareous soil (487.05–493.90 kg ha⁻¹) over the incubation period. RDF through DAP recorded the highest potassium levels, with significant interaction effects, as non-calcareous soil with RDF through DAP reached 533.88 kg ha⁻¹ at 120 days. Potassium content is more readily available in manured soil than in unmanured soil was also observed by Brar *et al.* (2015).

Table:8 Influence of soil calcareousness and PROM on release of potassium in soil

Days →	30			60			90			120		
Levels of P ₂ O ₅ ↓	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B
B ₁	482.2	511.2	496.7	483.3	512.4	497.8	486.3	512.9	499.6	486.2	515.3	500.7
B ₂	490.4	519	504.7	493.9	522.3	508.1	499	526.7	512.9	502.4	530.3	516.4
B ₃	489.8	519	504.4	493.8	522.1	507.9	496.1	528.3	512.2	500.1	532.4	516.2
B ₄	484.4	511.9	498.1	486.3	515.9	501.1	487.4	518.5	503	488.8	521.3	505.1
B ₅	486.3	512.4	499.3	487.4	517.3	502.3	488.8	519.2	504	490.1	522.3	506.2
B ₆	487.4	515.9	501.7	488.8	518.5	503.7	491	521.3	506.2	491.3	524.3	507.8
B ₇	488.8	517.3	503	491.3	521.1	506.2	495.7	528.8	512.3	498.4	533.9	516.2
Mean A	487.1	515.2	--	489.3	518.5	--	492.1	522.3	--	493.9	525.7	--
Factors	(A)	(B)	(AXB)	(A)	(B)	(AXB)	(A)	(B)	(AXB)	(A)	(B)	(AXB)
C.D. at (5%)	0.24	0.449	0.635	0.176	0.329	0.465	1.127	2.109	2.983	0.932	1.744	2.466
SE(m)±	0.082	0.154	0.218	0.06	0.113	0.16	0.387	0.724	1.024	0.32	0.599	0.847

3.3 Urease activity

At 0 days, non-calcareous soil reported higher urease activity ($40.03 \mu\text{g NH}_4^+ \text{-N g}^{-1} \text{hr}^{-1}$) than calcareous soil ($33.07 \mu\text{g NH}_4^+ \text{-N g}^{-1} \text{hr}^{-1}$), with non-significant interaction. By 120 days, non-calcareous soil maintained higher activity ($48.85 \mu\text{g NH}_4^+ \text{-N g}^{-1} \text{hr}^{-1}$), and 100% P_2O_5 through PROM resulted in the highest activity ($53.01 \mu\text{g NH}_4^+ \text{-N g}^{-1} \text{hr}^{-1}$). Significantly higher activity ($58.74 \mu\text{g NH}_4^+ \text{-N g}^{-1} \text{hr}^{-1}$) at 120 days of incubation was reported for interaction A_2B_7 (non-calcareous soil + 100% P_2O_5 through PROM). R. Srinivasan *et al.* (2016) found that the addition of organic sources is good source of energy and carbon to heterotrophs causing an increase in enzymatic activity and a rise in population.

Table:9 Influence of soil calcareousness and PROM on urease activity of soil

Days →	Urease ($\mu\text{gNH}_4^+ \text{-N g}^{-1} \text{hr}^{-1}$)					
	0 DAI			120 DAI		
Treatment	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B
B ₁	33.06	40.04	36.55	33.01	41.20	37.11
B ₂	33.07	40.02	36.54	35.21	41.22	38.21
B ₃	33.09	40.02	36.55	36.16	41.15	38.66
B ₄	33.08	40.03	36.55	39.77	45.35	42.56
B ₅	33.06	40.01	36.54	46.97	56.32	51.65
B ₆	33.05	40.04	36.55	47.05	57.95	52.50
B ₇	33.09	40.02	36.56	47.28	58.74	53.01
Mean A	33.07	40.03	--	40.78	48.85	--
Factors	(A)	(B)	(A X B)	(A)	(B)	(A X B)
C.D. (5%)	0.051	NS	NS	0.341	0.638	0.902
SE(m) ±	0.017	0.033	0.046	0.117	0.219	0.31

3.4 Dehydrogenase activity

At 0 days, non-calcareous soil reported higher dehydrogenase activity ($16.05 \mu\text{g TPF g}^{-1} \text{24 hr}^{-1}$) than calcareous soil ($14.34 \mu\text{g TPF g}^{-1} \text{24 hr}^{-1}$), with non-significant interaction, while at 120 days, non-calcareous soil recorded higher activity ($18.76 \mu\text{g TPF g}^{-1} \text{24 hr}^{-1}$). Application of 100% P_2O_5 through PROM recorded higher dehydrogenase activity ($18.18 \mu\text{g TPF g}^{-1} \text{24 hr}^{-1}$). The interaction effect was significant at 120 days and combination of A_2B_7 (non-calcareous soil + 100% P_2O_5 through PROM) found higher activity ($20.11 \mu\text{g TPF g}^{-1} \text{24 hr}^{-1}$).

Table:10 Influence of soil calcareousness and PROM on dehydrogenase activity of soil

	Dehydrogenase Activity ($\mu\text{gTPFg}^{-1} 24\text{hr}^{-1}$)					
Days →	0 DAI			120 DAI		
Treatment	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B
B ₁	14.34	16.01	15.18	14.52	16.20	15.36
B ₂	14.36	16.00	15.18	16.06	17.70	16.88
B ₃	14.46	15.96	15.21	15.99	18.50	17.24
B ₄	14.34	15.99	15.16	15.84	19.12	17.48
B ₅	14.31	16.01	15.16	15.95	19.74	17.85
B ₆	14.24	16.19	15.22	16.19	19.92	18.06
B ₇	14.35	16.20	15.28	16.26	20.11	18.18
Mean A	14.34	16.05	--	15.83	18.76	--
Factors	(A)	(B)	(A X B)	(A)	(B)	(A X B)
C.D. (5%)	0.091	NS	NS	0.101	0.189	0.268
SE(m) ±	0.031	0.059	0.083	0.035	0.065	0.092

3.5 Alkaline phosphatase activity

At 0 days of incubation alkaline phosphatase activity ($9.66 \mu\text{g PNP g}^{-1} \text{hr}^{-1}$) of calcareous soil was higher than non-calcareous soil ($8.51 \mu\text{g PNP g}^{-1} \text{hr}^{-1}$), with non-significant interaction. At 120 days, non-calcareous soil showed higher activity ($16.66 \mu\text{g PNP g}^{-1} \text{hr}^{-1}$), with 100% P₂O₅ through PROM leading to the highest activity ($18.08 \mu\text{g PNP g}^{-1} \text{hr}^{-1}$). The significant interaction effect at 120 days for A₂B₇ (non-calcareous soil + 100% P₂O₅) recorded higher alkaline phosphatase activity ($18.44 \mu\text{g PNP g}^{-1} \text{hr}^{-1}$). Waldrip *et al.* (2012) reported that organic manure may enhance phosphatase activity by providing soil microbes with carbon, nitrogen, and phosphorus.

Table: 11 Influence of soil calcareousness and PROM on alkaline phosphatase activity of soil

	Alkaline Phosphatase Activity ($\mu\text{g PNP g}^{-1} \text{hr}^{-1}$)					
Days →	0 DAI			120 DAI		
Treatment	A ₁	A ₂	Mean B	A ₁	A ₂	Mean B
B ₁	9.63	8.48	9.05	12.11	14.07	13.09

B ₂	9.69	8.49	9.09	14.24	15.32	14.78
B ₃	9.69	8.51	9.10	14.66	15.58	15.12
B ₄	9.63	8.50	9.07	15.35	17.31	16.33
B ₅	9.67	8.51	9.09	17.03	17.57	17.30
B ₆	9.61	8.52	9.07	17.70	18.35	18.03
B ₇	9.67	8.56	9.12	17.72	18.44	18.08
Mean A	9.66	8.51	--	15.55	16.66	--
Factors	(A)	(B)	(A X B)	(A)	(B)	(A X B)
C.D. (5%)	0.072	NS	NS	0.124	0.232	0.328
SE(m) ±	0.025	0.046	0.065	0.043	0.08	0.113

3.6 Correlation among soil enzyme activities and nutrient release

At 0 days, soil urease activity recorded a significant positive correlation with nitrogen ($r = 0.880$) and a non-significant negative correlation with phosphorus ($r = -0.619$) and potassium ($r = 0.277$). At 120 days, urease activity had a positive significant correlation with nitrogen ($r = 0.645$) and non-significant positive correlations with phosphorus ($r = 0.404$) and potassium ($r = 0.034$).

For micronutrients, at 0 days, it was negatively correlated with iron, zinc, and copper, while at 120 days, it showed significant positive correlations with all micronutrients: iron ($r = 0.981$), manganese ($r = 0.945$), zinc ($r = 0.972$), and copper ($r = 0.959$). Kaptanoglu Berber *et al.* (2014) reported in the study that there was not a statistically significant relationship between organic matter and urease enzyme activity

Table: 12 Correlation between soil enzyme activities and nutrient release at 0 days of incubation

0 DAI	N	P	K	Fe	Mn	Zn	Cu	Urease	Dehydrogenase	Alkaline phosphatase
N	1									
P	0.400	1								
K	0.025	-0.381	1							
Fe	0.448	0.445	0.525	1						
Mn	0.066	-0.127	0.854**	0.697*	1					
Zn	0.407	-0.695*	0.263	0.766**	0.324	1				
Cu	-0.026	0.232	0.392	0.528	0.756*	0.419	1			
Urease	0.880**	-0.619	0.277	-0.227	0.020	-0.553	-0.500	1		
Dehydro-	-0.353	0.847**	0.498	-0.437	0.212	-0.636*	-0.177	0.673*	1	

genase										
Alkaline Phosphatase	-0.015	0.787**	0.844**	0.186	0.730*	0.048	0.433	0.343	0.633*	1

At 0 days, soil dehydrogenase activity showed a significant positive correlation with phosphorus ($r = 0.847$) and a non-significant negative correlation with nitrogen ($r = -0.353$). Potassium had a non-significant positive correlation ($r = 0.498$), while zinc showed a significant negative correlation ($r = -0.636$). At 120 days, phosphorus continued to show a significant positive correlation ($r = 0.636$), with nitrogen ($r = 0.516$) and potassium ($r = 0.423$) showing non-significant positive correlations. Non-significant negative correlations were found with iron ($r = -0.437$) and copper ($r = -0.177$), and a non-significant positive correlation with manganese ($r = 0.212$). Phosphorus is a key factor affecting dehydrogenase activity over time. It was observed that the increase in dehydrogenase activity and microbial biomass were proportional to the addition of the number and quantity of nutrients. (Manjaiah and Singh, 2001).

At 0 days, soil alkaline phosphatase activity showed a significant positive correlation with phosphorus ($r = 0.787$) and potassium ($r = 0.844$), but a non-significant correlation with nitrogen ($r = -0.015$). At 120 days, it remained positively correlated with phosphorus ($r = 0.736$), with non-significant correlations to nitrogen ($r = 0.300$) and potassium ($r = 0.217$). At 0 days, alkaline phosphatase also showed a significant positive correlation with manganese ($r = 0.730$) and non-significant correlations with iron ($r = 0.186$), zinc ($r = 0.048$), and copper ($r = 0.433$). At 120 days, it showed significant correlations with iron ($r = 0.942$) and manganese ($r = 0.929$), and non-significant correlations with zinc ($r = 0.239$) and copper ($r = 0.137$). Tarafdaret *et al.* (1987) reported a significant correlation between organic P depletion and phosphatase activity in wheat ($r = 0.99^{**}$) and clover (0.97^{**}) rhizosphere soils.

Table: 13 Correlation between soil enzyme activities and nutrient release at 120 days of incubation

<i>O</i> DAI	N	P	K	Fe	Mn	Zn	Cu	Urease	Dehydrogenase	Alkaline phosphatase
N	1									
P	0.212	1								
K	0.687 *	0.818**	1							
Fe	0.014	0.328	-0.051	1						
Mn	-0.043	0.213	-0.127	0.984**	1					
Zn	0.010	-0.248	-0.091	0.990**	0.998**	1				
Cu	-0.014	0.267	-0.090	0.995**	0.994**	0.298	1			
Urease	0.645*	0.404	0.034	0.981**	0.945**	0.959**	0.972**	1		
Dehydrogenase	0.516	0.636*	0.423	0.801**	0.299	0.811**	0.237	0.156	1	

Alkaline Phosphatase	0.300	0.736*	0.217	0.942**.	0.929**	0.239	0.137	0.239	0.254	1
----------------------	-------	--------	-------	----------	---------	-------	-------	-------	-------	---

4. CONCLUSION

The application of Phosphate Rich Organic Manure (PROM) reduced soil calcium carbonate levels in both calcareous and non-calcareous soils with significantly enhancing the availability of macronutrients (nitrogen, phosphorus, and potassium) compared to inorganic fertilizers. Over the 120-day incubation period, all three enzymes-urease, dehydrogenase, and alkaline phosphatase exhibited a gradual increase in activity, indicating active microbial processes and ongoing soil organic matter decomposition; notably, the increase in dehydrogenase activity suggests enhanced organic matter breakdown, which can improve soil structure and nutrient availability. Additionally, the correlation analysis highlighted the essential role of these enzymes in nutrient cycling, particularly the significant positive correlation between alkaline phosphatase and phosphorus at 120 days, underscoring its potential contribution to phosphorus mineralization and cycling within the soil ecosystem.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1.

2.

3.

REFERENCES

1. Brar, B.S., Singh, J., Singh, G. and Kaur, G. (2015). Effects of long-term application of inorganic and organic fertilizers on soil organic carbon and physical properties in maize–wheat rotation. *Agronomy*, 5(2),220-238.
2. Hapse, D.G. (1993). Organic farming in the light of reduction in use of chemical fertilizers. *Proceedings of 43rd Annual Deccan Sugar Technology Association, Pune, Part, 1*.
3. Kaptanoğlu Berber, A.S., Farasat, S. and Namli, A. (2014). Afforestation effects on soil biochemical properties. *Eurasian Journal of Forest Science*, 1,25-34.
4. Kumar, S., Srivastava, A. and Gupta, A. (2015). Effect of organic amendments on availability of different chemical fractions of phosphorus. *Agricultural Science Digest-A Research Journal*, 35(2),83-88.
5. Mandal, L.N. and Mitra, R.R. (1982). Transformation of iron and manganese in rice soils under different moisture regimes and organic matter applications. *Plant and Soil*, 69,45-56.
6. Manjaiah, K.M., Singh, D. (2001). Soil organic matter and biological properties after 26 years of maize-wheat-cowpea cropping as affected by manure and fertilization in a cambisol in semiarid region of India. *Agriculture Ecosystem and Environment*. 86,155-162.
7. Motaghian, H.R. and Hosseinpour, A.R. (2017). The effects of cow manure and vermicompost on availability and desorption characteristics of zinc in a loamy calcareous soil. *Communications in Soil Science and Plant Analysis*, 48(18),2126-2136.
8. Muhawish, N.M. and Al-Kafaje, R.K. (2017). Soil organic carbon and phosphorus status after combined application of phosphate rock and organic materials in a gypsiferous soil. *Iraqi Journal of Agricultural Sciences*, 48.

9. Pandey, R., Bargali, S.S., Bargali, K., Karki, H. and Chaturvedi, R.K. (2024). Dynamics of nitrogen mineralization and fine root decomposition in sub-tropical Shorearobusta Gaertner f. forests of Central Himalaya, India. *Science of The Total Environment*, 921,170896.
10. Panse VG, Sukhatme PV. (1967). *Statistical Methods for Agricultural Workers*. Indian Council of Agricultural Research, New Delhi, India.,272-279.
11. R. Srinivasan, K. Jeevan Rao, S. K. Reza, Shelton Padua, D. Dinesh and S. Dharumarajan. (2016). Influence of Inorganic Fertilizers and Organic Amendments on Plant Nutrients and Soil Enzyme Activities under Incubation. *International Journal of Bio-resource and Stress Management*, 7(4),924-932.
12. Reddy, A.R., Chaitanya, K.V. and Vivekanandan, M. (2004). Drought-induced responses of photosynthesis and antioxidant metabolism in higher plants. *Journal of plant physiology*, 161(11),1189-1202.
13. Safarzadeh, S., Kasmaei, S.L. and Ahmad, A.Z. (2018). Effect of organic substances on iron-release kinetics in a calcareous soil after basil harvesting. *Journal of the Serbian Chemical Society*, 83(7-8),941-952.
14. Shen, J., Yuan, L., Zhang, J., Li, H., Bai, Z., Chen, X., Zhang, W. and Zhang, F. (2011). Phosphorus dynamics: from soil to plant. *Plant physiology*, 156(3),997-1005.
15. Talashilkar, S.C., Sawant, D.S., Dhamapurkar, V.B. and Savant, N.K. (2000). Yield and nutrient uptake by rice as influenced by integrated use of calcium silicate slag and UB-DAP in acid lateritic soil. *Journal of the Indian Society of Soil Science*, 48(4),847-849.
16. Tarafdar, J.C. and Jungk, A. (1987). Phosphatase activity in the rhizosphere and its relation to the depletion of soil organic phosphorus. *Biology and Fertility of Soils*, 3,199-204.
17. Tella, M., Bravin, M.N., Thuriès, L., Cazevieuille, P., Chevassus-Rosset, C., Collin, B., Chaurand, P., Legros, S. and Doelsch, E. (2016). Increased zinc and copper availability in organic waste amended soil potentially involving distinct release mechanisms. *Environmental pollution*, 212,299-306.
18. Waldrip, H.M., He, Z. and Griffin, T.S. (2012). Effects of organic dairy manure on soil phosphatase activity, available soil phosphorus, and growth of sorghum-sudangrass. *Soil science*, 177(11),629-637.