

Fungicide Sensitivity of *Alternaria brassicicola*: An *In vitro* Study for Managing Alternaria Blight in Brassica juncea

Abstract

Alternaria brassicicola, a phytopathogenic fungus, incites Alternaria leaf blight in mustard, leading to substantial yield and quality losses. This study aimed to investigate the efficacy of seven chemical fungicides against *A. brassicicola* *in vitro*. Seven chemical fungicides were evaluated at concentrations of 50, 100, 250, and 500 ppm using the poisoned food technique and Completely Randomized Design (CRD). Tebuconazole + Trifloxystrobin and Tebuconazole exhibited 100% growth inhibition at 100-500 ppm. Mancozeb + Metalaxyl and Mancozeb demonstrated 100% growth inhibition at 250-500 ppm. Chlorothalanyl displayed 83.33% growth inhibition at 500 ppm, while Thiophanate methyl showed the least efficacy (48.00% growth inhibition) at 50 ppm. The findings suggest that Tebuconazole-based fungicides are highly effective against *A. brassicicola*. Lower concentrations (50-100 ppm) of certain fungicides also exhibited promising results.

Key Words: *Alternaria brassicicola*, Alternaria leaf blight, mustard, chemical fungicides, *in vitro* evaluation.

Introduction

Mustard (*Brassica juncea*) is a prominent Rabi oilseed crop in India, ranking second in importance to groundnut in terms of area and production. It is extensively cultivated as a pure crop and intercrop in marginal and sub-marginal soils across eastern, northern, and north-eastern states. The cool and moist climate during winter months facilitates luxuriant growth and productivity of mustard in these regions. Mustard seeds are known by various local names, including sarson, rai, raya, torai, and lahi. However, the crop is susceptible to numerous bacterial, fungal, and viral diseases. Among these, Alternaria blight, caused by *Alternaria brassicicola*, is a globally widespread and economically significant disease. Alternaria blight caused by *Alternaria brassicicola* has been reported to inflict heavy yield losses to the tune of 35-60% (Kolte et al., 1987) in mustard crop. Typical symptoms of Alternaria blight disease are the formation of spots on leaves, stems and siliquae. The concentric black spots produced were

usually grey coloured and their characters varied with host and environmental factors. Initially symptoms appeared on the lower leaves as black points, later which enlarged to develop into prominent, round, concentric spots of various sizes. As disease progressed, the lower leaves defoliated and disease appeared on middle and upper leaves. At the later stage of the plant growth, spots appear on siliquae and stem. The spots were round black and quite conspicuous and the siliquae turn completely black. On the stem, black, elongated spots became visible in the form of black streaks with or without necrotic gray centres. Considering economic importance of the mustard and destructive nature of *Alternaria* blight incited by *Alternaria brassicicola* mustard, present studies on in vitro evaluation of fungicides against the test pathogen were undertaken.

Materials and Methods

The experiment was carried out in Completely Randomized Design using poisoned food technique. Seven chemical fungicides i.e. Mancozeb, Mancozeb + Metalaxyl, Tebuconazole + Trifloxystrobin, Thiophanate methyl, Tebuconazole, Chlorothalanil and Carbendazim were evaluated at four different concentrations such as 50, 100, 250 and 500 ppm respectively. Each of the treatment was replicated three times.

Isolation, purification and maintenance of pure culture

Pathogen *Alternaria brassicicola* was isolated from infected leaf of mustard collected from field. Spores were teased from infected portion for microscopic examination to check the presence of pathogenic fungus. After confirming the presence of *Alternaria brassicicola*, leaves were cut into small pieces (1-1.5cm) with sterile blade. These pieces were disinfected with 0.5% sodium hypochlorite (NaOCl) solution for two minutes followed by three washings with distilled water and excessive moisture was removed using sterile blotting paper. The sterilized leaf pieces were placed on PDA medium using sterilized forceps and incubated at 27 ± 1 °C for 7 days. On the basis of morphological characters of conidia as described by Yu (2015) pathogen was identified as *Alternaria brassicicola*. Then the culture was purified by transferring small piece of agar containing spore to another Petri plate containing media and incubated at 27 ± 1 °C for 7 days. The pathogen was sub cultured three times to obtain pure culture and pure culture thus obtained was preserved in PDA slant at 4 °C.

In-vitro evaluation of fungicides

For the evaluation of chemical fungicides, calculated amount of stock solution was mixed in sterilized PDA to make final concentration of 50ppm, 100ppm, 250ppm and 500ppm. Twenty ml of amended PDA was poured in each 90mm sterilized petri plate and allowed to solidify. A check was also maintained where no fungicide was added in the medium. A circular disc of 7mm diameter from 9 days old culture of *Alternaria brassicicola* was cut with sterilized cork borer and inoculated in the centre of solidified amended as well as control media. Each treatment was replicated in three petri plates. Then the petriplates were incubated at 27 ± 1 °C for seven days.

Growth inhibition test

The observation on mycelial growth was recorded after 7 days of incubation in each treatment using Vernier calliper scale. The percent growth inhibition of mycelial growth over control was calculated by using the formula given by Vincent (1947).

$$\text{Per cent inhibition} = \frac{(C-T)}{C} \times 100$$

Where,

C = Diameter of the colony in control (mm)

T = Diameter of the colony in treatments (mm)

Results and Discussion

In Vitro Evaluation of Fungicides

The efficacy of different fungicides was shown in the table showing 7 days of incubation. The efficacy of different chemical fungicides against test fungus was evaluated in vitro using poisoned food technique. The data on inhibition percent is presented in table. An insight into data reveals that all the tested chemical fungicides showed significant effect against pathogen growth over control (78.00 mm). The extent of mycelial growth inhibition increased with increase in their concentration. The tables show the percent mycelial growth and percent mycelial growth inhibition at different concentrations at seventh days after incubation. The data

showed (Table 1) that increase in concentration of the fungicides caused increased inhibition of mycelial growth of the fungal pathogen (*Alternaria brassicicola*). Among those, T3(Tebuconazole 50% + Trifloxystrobin 25%), showed highest inhibition per cent (87.82, 100, 100 and 100) followed by T5Tebuconazole (86.53, 100,100 and 100%) , T2Mancozeb 64% + Metalaxyl 8% (75.00, 89.74, 100 and 100%), and T1 Mancozeb (71.79, 79.48, 85.89 and 100%) at 50, 100 , 250 and 500 ppm concentrations, respectively.

Whereas, minimum inhibition per cent was showed by T4Thiophanate methyl (38.46, 57.05, 64.14 and 66.02%) followed by T7 Carbendazim (41.02, 60.89, 63.46 and 69.23%) and T6Chlorothalanyl(53.84, 69.23, 78.20 and 83.33%) at different 50, 100 , 250 and 500 ppm concentrations. Our results are in accordance with the findings of Meena *et al.* (2020), who tested different fungicides, *viz.*, Mancozeb, Metalaxyl MZ, Copper oxy-chloride, Copper hydroxyl-chloride, Carbendazim, Azoxystrobin, Tebuconazol, Nativo and leaf extracts of *A. indica*, *P. pinnata* and *M. alliacea* was performed following the “poisoned food technique”. Among the tested fungicides Tebuconazole + Trifloxystrobin and Tebuconazol (0.1%) were found most effective to inhibit the mycelium growth up to 89 per cent. Hussain *et al.* (2018), observed that the Topsin-M and Topas Fungicides did not proved better as compare to Nativo and Cabriotop fungicides. Similar result was recorded in findings of Panwar *et al.*, (2013) who reported complete growth inhibition of *Alternaria* in tebuconazole followed by mancozeb and least inhibition in carbendazim. Similarly, Tu (2015) recorded complete inhibition of *A. brassicae* by tebuconazole, mancozeb at 250, 500 and 1000 ppm and by metalaxyl + mancozeb at 500 and 1000 ppm and least inhibition at carbendazim. Kumar and Kumar (2022), Biswas and Ghosh (2018) and Kantwaet *al.*,(2014) obtained significant growth inhibition effect of *Alternaria* sp. in mancozeb. Synthetic fungicides bring about the inhibition of pathogens either by destroying their cell membrane or its permeability or by inhibiting metabolic processes of the pathogen and hence are effective (Kakraliya *et al.*, 2018). Higher inhibition effect of tebuconazole is due to inhibition of ergo sterol biosynthesis, controlling the growth and reproduction of fungal pathogen (Muhamad *et al.*, 2010).

Table 1: *In vitro* evaluation of fungicides on mycelial growth inhibition of *Alternaria brassicicola*

Treatments	Fungicides	Mean colony diameter (mm)	Growth inhibition (%) at
------------	------------	---------------------------	--------------------------

		at different concentrations				different concentrations (ppm)			
		(ppm)							
		50	100	250	500	50	100	250	500
T1	Mancozeb	22.0	16.0	11.0	0	71.79	79.48	85.89	100.00
T2	Mancozeb + Metalaxyl	19.5	8.0	0	0	75.00	89.74	100.00	100.00
T3	Tebuconazole + Trifloxystrobin	9.5	0	0	0	87.82	100.00	100.00	100.00
T4	Thiophanate methyl	48.0	33.5	28.0	26.5	38.46	57.05	64.10	66.02
T5	Tebuconazole	10.5	0	0	0	86.53	100.00	100.00	100.00
T6	Chlorothalanil	36.0	24.0	17.0	13.0	53.84	69.23	78.20	83.33
T7	Carbendazim	46.0	30.5	28.5	24.0	41.02	60.89	63.46	69.23
T8	Control	78.00	78.00	78.00	78.00				

Conclusion

Alternaria leaf blight is a worldwide disease of economic importance in mustard crop. Different chemical fungicides are commercially available in market to control this disease. This study revealed significant inhibition effect of all the tested chemical fungicides over control. Tebuconazole + Trifloxystrobin and Tebuconazole proved to be the most effective chemical fungicide recording 100% growth inhibition at 100ppm, 250ppm and 500ppm concentrations whereas, Thiophanate methyl was least effective in reducing fungal growth. Indiscriminate application of chemical fungicides have resulted several health hazards, negative impacts in environment so, the use of effective chemical at possible lower concentrations could be safer way to minimize health hazards and environmental pollutions. The effectiveness of certain chemical fungicides, even at lower concentrations (50 ppm and 100 ppm), in controlling *Alternaria brassicicola*. Utilizing these fungicides can minimize hazardous effects and provide a viable solution for managing *Alternaria* leaf blight in mustard crops.

References

- Biswas MK, Ghosh T (2018). Evaluation of phyto-extracts, biological agents and chemicals against the development of *Alternaria brassicicola* in vitro and vivo. *European Journal of Medicinal Plants* 22(9):1-9.
- Hussain S, Ayub M, Rasheed M, Hassan S, Hussain K, Hussain S, Hussain SI, Hassan M, Hussain A, Hussain T, Imtiyaz M (2018). Efficacy of different fungicides against *Alternaria tusi* cause leaf blight of peas (*Alternaria tusi*) was first time discovered from Kharmang Olding and Skardu Baltistan. *Journal of Entomology and Zoology Studies* 6: 2255-2260
- Kakraliya SS, Choskit D, Pandit D, Abrol S (2018). Effect of bio-agents, neem leaf extract and fungicides against *Alternaria* leaf blight of wheat (*Triticum aestivum* L.).
- Kantwa SL, Tetarwal JP, Shekhawat KS (2014). In vitro effect of fungicides and phyto-extracts against *Alternaria alternata* causing leaf blight of groundnut. *IOSR Journal of Agriculture and Veterinary Science* 7(6):28-31.
- Kolte SJ, Awasthi RP, Vishwanath V (1987). Assessment of yield losses due to *Alternaria* blight in rapeseed and mustard. *Indian Phytopathology*, 40(2): 209-211.
- Kumar Sanjeev and Kumar Amarendra (2022). Evaluation of Efficacy of Fungicides against *Alternaria* Blight Disease of Linseed. *Environment and Ecology* 40 (2A) : 584—588.
- Meena RP, Saran PWL, Kalariya KA, Manivel P (2020). Efficacy of fungicides and plant extracts against *Alternaria alternata* causing leaf blight of chandrasur (*Lepidium sativum*). *Indian Journal of Agricultural Sciences* 90(2): 337-340
- Muhamad H, Zainol M, Sahid I. (2010). Determination of hexaconazole in field samples using gas chromatography –electron capture detection (GC-ECD).
- Panwar V, Yadav RS, Singh H, Shukla AC (2013). In vitro evaluation of fungicides and bio agents against *Alternaria alternata* – an incident of leaf spot of Aloe vera.
- Tu VD (2015). Studies on *Alternaria* leaf spot(*Alternaria brassicicola* (Schw.) Wiltshire) of cabbage (*Brassica oleraceavar. capitata*L.). Ph.D. Thesis (Unpublished), University of Agricultural Sciences, Bengaluru.
- Vincent JM (1947). Distortion of fungal sac hyphae in the presence of certain inhibitors. *Nature* 159: 850

UNDER PEER REVIEW