

Variability studies in soybean genotypes under different water regimes

Abstract:

Soybean (*Glycine max* L.) is economically one of the most important oilseed crop worldwide which have high protein and oil content. Its productivity is frequently affected due to drought occurrence in India. Variability for root shoot traits under early seedling stage under drought stress may be important selection criteria for development of drought tolerant genotypes. The present investigation was carried out at Net house of Experimental Farm, Mandasaur University, Mandasaur. Sixty genotypes, procured from ICAR-IISR, were used to know the extent of genetic variability under different water regimes [normal, 100 ml (100%), S₁, 50 ml (50%), S₂, 25 ml (25%) and S₃, 0 ml (0%)] for root-shoot traits and relative leaf water content under early seedling stage. Significant genetic variability was recorded for all the traits among genotypes studied under normal and water stress conditions respectively which showed the presence of ample variability in studied material. High Phenotypic Coefficient of Variation than their corresponding Genotypic Coefficient of Variation for the entire trait under different water regimes denoted influence of environment on these traits. The large genetic variation found in these genotypes may be used to develop varieties with better drought tolerance behavior and used as drought donor lines in drought breeding programs. We measured high heritability for selected characters under non water stress condition (75.05% to 100%) and water stress conditions S₁ (81.40% to 100%), S₂ (85.69% to 100%) and S₃ (92.60% to 99.26%) respectively. Additive gene action interacted to control characters under different water regimes because high broad sense heritability along with high genetic advance as percentage of mean were recorded for most of traits. It is indicating that simple phenotypic selection would be effective for these traits under water stress condition.

Keyword: Heritability, soybean, variability, water regimes, water stress.

Introduction

Soybean (*Glycine max* L. Merrill) is the world's most important oilseed crop (Narayanan S and Fallen B., 2019; Mishra and Patidar, 2023; Bairagi *et al.*, 2023). Soybean oil is secondly most widely consumed oil and rich source of protein in world (Sunaryo *et al.*, 2016). In International market soya oil trading is next only to palm oil (Bhuva *et al.* 2020). It contributes to 25 % of the global edible oil, about two-thirds of the world's protein concentrate for livestock feeding. The soybean is not only known for its high total protein (5-52%) content but the quality of soy protein which is higher than that of other grain legume plant proteins (25-46%) and like animal protein (Hughes *et al.* 2011; Vollmann 2016). Soybean is known as the "Golden Bean" because of its health and nutritional benefits such as low Glycaemic index, low saturated fat, and cholesterol-free and is widely used as oilseed (Kumawat *et al.*, 2023). It is primarily produced by the United States, Brazil, Argentina, China, and India (Thu *et al.*, 2014). Soybean is also a very important *kharif* season crop of India. Kharif season is rainy season which start after onset of monsoon in India and considered from June- July to October-November. The top three soybean growing states in India are Madhya Pradesh (52 Lakh hectares; 55.40 Lakh ton), Maharashtra (45 Lakh hectares; 50.17 Lakh ton) and Rajasthan (11.13 Lakh hectares; 10.53 Lakh ton). Madhya Pradesh state is leading state in both area and production point of view in all over India (118.32 Lakh hectares; 125.82 Lakh ton) and emerged as India's Soy State (SOPA 2023; Kumawat *et al.*, 2023; Mishra and Patidar, 2023). Madhya Pradesh has 45% share in soybean production in the country. Soybean meal is a valuable ingredient in formulated feeds for poultry and fish.

Drought stress causes a decrease of 50% of the total yield soybean production (Sunaryo *et al.*, 2016). Water deficit affects physiological and agronomic traits of soybean plants, thus negatively influencing plant growth and development, resulting in grain yield reduction (Stolf-Moreira *et al.*, 2010; Giordani *et al.*, 2019, Nair *et al.* 2023; Kumawat *et al.*, 2023). Scientific forecasts draw a future with dark scenarios of water restrictions around the world (Dai, 2013; Mishra and Patidar, 2023).

Information on the magnitude of variability and extent, to which desirable characters are heritable, is important for planning breeding programme and ascertaining the scope of its improvement (Sileshi, 2019). The success of phenotypic selection depends upon the range of genetic diversity available in the population, whereas estimate of heritability and genetic advance are useful in inferring the genetic factors. Root traits affect the amount of water and nutrient absorption, and are important parameters for maintaining crop yields indirectly under water

stress conditions (Fenta *et al.*, 2014; Mishra *et al.*, 2016; Mishra *et al.*, 2017; Vijay *et al.*, 2018; Aski *et al.*, 2022). Despite the importance of root traits in drought tolerance, few breeding programs take these traits into account when developing drought-tolerant soybean varieties. Information regarding the soybean genetic variability of the root traits is limited, and the exploitation of this variability can assist soybean breeding programs in the development of varieties with desired root traits for drought tolerance.

Keeping in view the above fact the aim of present study was to observe genetic variability and heritability among sixty genotypes of soybean in terms of root traits under seedling stage for the purpose of selection of parental genotypes for further crosses in order to develop progenies with drought tolerant types with increased early vigor of roots and stems.

Materials and methods

➤ Experimental Site:

The experiment was conducted at Net house facility situated at Campus-I, Faculty of Agriculture Sciences, Mandsaur University, Mandsaur (M.P.). Net house and laboratory experiments were conducted to achieve the objective of present experiment. Seedlings were raised in 480 polybags during September- October 2022.

➤ List of genotypes and experimental design:

Total sixty lines collected from ICAR- Indian Institute of Soybean Research, Indore were sown on dated 02/09/2023 in four different water regimes (nonstress-100 ml, 50 ml, 25 ml and 0 ml) using completely randomized block design with two replications for each set of experiment. The bags were filled with standard soil having mixture of 1% sand, 1% FYM and 1% clay loamy soil. Every polybag was measure by using electronic weighing balance having capacity of 20 Kg. Each bag was filled with 8 Kg soil. The soil was also treated with readymade 1/4th MS Media for obtaining healthy plants. The list of genotypes is presented in **Table 1**.

Table 1: Name of 60 genotypes of *Glycin max L.* used for study

S.No	Name of genotypes	S.No	Name of genotypes	S.No	Name of genotypes	S.No	Name of genotypes
1	GW-34	16	GW-234	31	NRC-37(CHECK)	46	RSC-1107(CHECK)
2	GW-371(K-21C)	17	GW-196	32	GW-178	47	GW-212

3	GW-63(K-21)	18	GW-382	33	GW-87	48	NRC-138
4	GW-237(K-25)	19	GW-134	34	GW-45	49	GW-214
5	GW-155	20	AMS-2014-1(CHECK)	35	GW-89	50	NRC-142
6	GW-159	21	AGS-218	36	JS-2069	51	NRC-127
7	GW-99	22	GW-108	37	GW-207	52	JS-9560
8	GW-164	23	GW-132	38	GW-188	53	GW-203
9	GW-312	24	PK-472(CHECK)	39	GW-185	54	JS-20-116
10	GW-143	25	GW-100	40	GW-52	55	JS-2034
11	GW-152(K-21-C)	26	GW-10	41	GW-286	56	GW-253
12	GW-15 (CHECK)	27	IC-073710	42	GW-223	57	GW-225
13	GW-51(K-21)	28	GW-17	43	GW-251	58	TGX-9336E
14	GW-21	29	GW-13	44	GW-291	59	SQL-110
15	GW-161	30	GW-28	45	GW-221	60	AGS-25

➤ **Water stress imposition:**

Regular irrigation was continued till two leaf stage of plants under both normal and water stress conditions (**Figure 5**). After two leaf stage of plants (after 7 days), drought was imposed by applying water in proportion of 100% means 100 ml in non-water stress (control), 50% means 50 ml (stress S₁), 25% means 25 ml (stress S₂) and 0% (stress S₃) conditions till 30 days after sowing. After seed sowing at 15-20 days, the extra plants were rouged out except one to two healthy seedlings so that there will not any competition for nutrient and for healthy growth of plant for observation. The whole plant from bags were picked up by making vertical cut on bags, washed with tap water and data were observed for root length (cm), shoot length (cm), root fresh weight (gm), shoot fresh weight (gm), root dry weight (gm), shoot dry weight (gm), root shoot ratio by length, root shoot ratio by weight and relative leaf water content (%) respectively.

➤ **Statistical Analysis**

The replicated values were subjected to statistical analysis of variance (ANOVA) as prescribed by Panse and Sukhatme (1978) for individual characters for each environment.

Components of variance, coefficient of variation, broad sense heritability, and genetic advance were evaluated separately for each trait both in water stress and non-stress conditions.

Phenotypic Coefficient of Variation (PCV) and Genotypic Coefficient of Variation (GCV) were calculated in range of low (if it is less than 10 percent), moderate (10-20 percent) and high (If more than 20 percent) as classified by Shivasubramanian and Menon (1973).

The range (low- < 30%, medium-30%-60%, high > 60%) of heritability and genetic advance (GA) as percentage of mean range (low- < 10%, medium-10%-20%, and high > 20%) was calculated as suggested by Johnson *et al.*, (1955). The data were analyzed by using OPSTAT software.

Results and discussion

The increasing population, abiotic factors are threatening the global food security (Saha and Choyal, 2022). The sudden change in environmental factors further reduces the water availability and causes drought stress in major agro-systems especially rain fed ecosystem worldwide. It affects severely the crop production and productivity including soybean which varies with the crop stage of the crop as well as with the local environmental conditions (Okunlola *et al.*, 2017). Water stress affects the seed germination, vegetative and reproductive growth and maturity stage of a crop especially seedling and germination stage by reducing seedling length and seedling biomass, seedling water content, biochemical and molecular attributes depending upon the frequency and duration of drought stress (Anjum *et al.*, 2017). The use of morpho-physiological traits in relation to drought tolerance has been suggested by many researchers (Painwadee *et al.* 2009) because the inheritance of these characters is simpler than pod yield. Genetic variability is essential to know response to selection pressure. It has also been reported that the magnitude of genetic variability present in base population of any crop species is important in crop improvement and must be exploited by plant breeder for yield improvement (Akram *et al.*, 2011; Mehra *et al.*, 2020). The information related to heritability of characters is important for plant breeders to formulate appropriate breeding strategies to achieve breeding objectives.

Results of genetic variability under non-water stress and water stress conditions have been presented in following heads:

➤ **Genetic variability under non-water stress (normal water) condition:**

Many researchers have reported variability in root traits among soybean cultivars. This variability was observed during early growth (Manavalan *et al.*, 2009; Fenta *et al.*, 2014; Thu *et al.*, 2014; Fried *et al.*, 2018; Falk *et al.*, 2020; Dayoub *et al.*, 2021; Syiem *et al.*, 2022), at flowering stage (Zhao *et al.*, 2004; Mwamlima *et al.*, 2019) or at maturity (Ao *et al.*, 2010).

➤ **Analysis of variance non stress**

The results from ANOVA (**Table-2**) in non-stress condition (100 ml water) indicated that high amount of variability was present among the genotypes for all the traits investigated because significant difference were observed among sixty genotypes. Similar results under non-water stress condition were reported by Meena *et al.* (2014) in chickpea for plant height and relative leaf water content, Prince *et al.* (2015), Falk *et al.* (2020), Yan *et al.* (2020), Mishra *et al.* (2021), Dayoub *et al.* (2021), Syiem *et al.* (2022) in soybean; Kumar *et al.* (2023) for root length, shoot length and shoot dry weight in wheat; for RWC, Bayoumi *et al.* (2008) in wheat and Kumar *et al.* (2021) in chickpea; Shankar *et al.*, (2019) for plant height, root length, root shoot ratio by length and RWC in groundnut, Priya *et al.* (2021) and Reddy *et al.* (2023) for root length in lentil, green gram and black gram respectively. Presence of sufficient amount of variability in the studied germplasm provides ample scope for selection superior and desired genotypes.

Table 2: Analysis of variance (mean sum of squares) for nine characters in soybean under non-stress (normal water) condition

Source of variation	DF	Root length (cm)	Shoot length (cm)	Root fresh weight (g)	Shoot fresh weight (g)	RWC %	Root dry weight (g)	Shoot dry weight (g)	Root shoot ratio by length	Root shoot ratio by weight
Replication	1	0.09	0.00	0.000	0.047	0.74	0.000	0.001	0.001	0.007
Treatment	59	16.28**	394.02**	0.017**	0.269**	835.44**	0.0002**	0.008**	0.034**	0.046**
Error	59	1.62	2.86	0.000	0.014	1.78	0.000	0.000	0.001	0.001
Total	119	8.88	196.77	0.009	0.141	415.10	0.000	0.004	0.017	0.023

** Significant at 1% (P= 0.01) level of significance

* Significant at 5% (P=0.05) level of significance

➤ **Genetic variability parameters under non water stress condition:**

The estimates of variability parameters for all the traits are shown in **Table 3 and Figure 1** for non-water stress condition. PCV values for all traits had generally greater, but closer values to their corresponding GCV values which indicated presence of lower environmental influence on the expression of these traits. Similar results were also reported by Kumar *et al.* (2023) in wheat.

The root length was ranged 5.05 cm to 18.20 cm with grand mean of 10.42 cm. The estimated values of PCV and GCV recorded were 24.48 and 21.21 per cent, respectively. High heritability (bs) 75.05 per cent coupled with medium genetic advance over percentage of mean 37.85 per cent were noticed for this trait. Similarly, high (More than 20%) PCV and GCV, (>60%) high heritability and high genetic advance (>30%) was recorded by Gobu *et al.* (2017) in eggplant, Kumar *et al.* (2023) in wheat; Shankar *et al.*, 2019 in groundnut and Hoque *et al.* (2021) in rice and Syiem *et al.* (2022) in soybean under normal water condition. **High heritability for this trait was also reported by Reddy *et al.* (2023) in green gram (89%) and black gram (89%).**

A wide significant range of variation (12.20 cm to 79.45 cm with an overall mean of 46.94 cm) was found among all the germplasm accessions for shoot length. The recorded PCV and GCV for this character were 24.59 and 24.32 per cent, respectively. Similar results of high PCV and GCV were reported by Bayoumi *et al.* (2008) in wheat. This character revealed high h^2 (97.85 per cent) along with high GA (49.57 per cent). Similar result of high heritability for this trait was also reported by Riaz *et al.* (2013) in cotton; Meena *et al.*, (2014) in chickpea. Similarly, high (10-30%) PCV and GCV, (>60%) high heritability and high genetic advance (>30%) was recorded by Gobu *et al.* (2017) in eggplant, Shankar *et al.*, 2019 in groundnut, Hoque *et al.* (2021) in rice and Syiem *et al.* (2022) in soybean under normal water condition.

The root fresh weight was ranged 0.042 gm to 0.389 gm with total mean of 0.199 gm. The phenotypic coefficient of variation (38.53) and genotypic coefficient of variation (36.95), heritability (91.93 per cent) and genetic advance over percentage of mean (72.98 per cent) respectively were found for this trait. Result of high GCV, PCV, heritability and high GA as

percentage of mean was also noticed by Syiem *et al.* (2022) in soybean for root fresh weight under non-water stress condition. Similar result of high heritability for this trait was also recorded by Riaz *et al.* (2013) in cotton.

A wide range of variation (0.041 to 1.592 gm with an overall mean of 0.807 gm) was found among all the germplasm accessions for shoot fresh weight. The PCV (39.00), GCV (36.15), heritability (h^2) (85.95%) and genetic advance over mean of (69.05%) were recorded respectively for shoot fresh weight. Similar findings for soybean for shoot fresh weight under seedling stage were also reported by Syiem *et al.* (2022) in soybean in normal water. Similar result of high heritability for this trait was also found by Riaz *et al.* (2013) in cotton.

The physiological character, relative leaf water content (RWC) varied from 18.765 to 93.150% with an overall mean of 62.851%. The PCV (26.61), GCV (26.52), heritability (h^2) (99.36%) and genetic advance over mean of percentage (54.46%) were observed respectively for this parameter. Similar results on heritability for this trait was also obtained by Bayoumi *et al.* (2008) in wheat, Garg *et al.* (2017) in rice and Kumar *et al.* (2021) in chickpea.

The root dry weight ranged from 0.009 to 0.079 with an overall mean of 0.025 gm. Recorded values of PCV and GCV were 39.41 and 39.41 per cent respectively. The values of high heritability (100.00%) together with high genetic advance as per cent mean (81.19) were recorded for this trait. Similar finding on heritability for this trait was also found by Riaz *et al.* (2013) in cotton; Lalitha *et al.* (2015) in chickpea.

The grand mean of shoot dry weight was 0.130 with a range of 0.0250 to 0.300. High phenotypic (40.10%) and genotypic coefficient of variability (39.72%) along with heritability (98.13%) and genetic advance as per cent mean (81.05) were recorded for this trait. Similar PCV and GCV (>30%), heritability (>60%) and genetic advance (>30%) was recorded by Kumar *et al.* (2023) in wheat and by Lalitha *et al.* (2015) in chickpea

Character, root shot ratio by length ranged from 0.101 to 1.023 with a grand mean value of 0.246. The PCV and GCV observed were 44.60 and 42.45 per cent, respectively. Heritability (bs) 90.59 per cent coupled with high genetic advance over percentage of mean 83.23 per cent were noticed for this trait. This is according to finding as reported by Gobu *et al.* (2017) in eggplant, Shankar *et al.* (2019) in groundnut and Hoque *et al.* (2021) in rice for this trait under normal water condition.

A perusable mean value of different genotypes of trait root shoot ratio by weight revealed that overall mean value of it was 0.234. High PCV (53.60) and high GCV (52.20) were recorded for this parameter. The high heritability (94.85%) coupled with high genetic advance (104.73%) were noticed for it.

UNDER PEER REVIEW

Table 3: Determination of mean, range and genetic variability parameters for nine traits in soybean under non-stress (normal water) condition

Genotypes	Mean	Min	Max	var (g)	var (p)	Heritability (%)	GA	GA% mean	GCV (%)	PCV (%)
Root length (cm)	10.424	5.050	18.200	4.89	6.51	75.05	3.95	37.85	21.21	24.48
Shoot length (cm)	46.943	12.200	79.450	130.39	133.24	97.85	23.27	49.57	24.32	24.59
Root fresh weight (g)	0.199	0.042	0.389	0.005	0.006	91.93	0.15	72.98	36.95	38.53
Shoot fresh weight (g)	0.807	0.041	1.592	0.085	0.099	85.95	0.56	69.05	36.15	39.00
Rwc %	62.851	18.765	93.150	277.89	279.67	99.36	34.23	54.46	26.52	26.61
Root dry weight (g)	0.025	0.009	0.079	0.0001	0.0001	100.00	0.020	81.19	39.41	39.41
Shoot dry weight (g)	0.130	0.025	0.300	0.0027	0.0027	98.13	0.105	81.05	39.72	40.10
Root shoot ratio by length	0.246	0.101	1.023	0.0109	0.0121	90.59	0.205	83.23	42.45	44.60
Root shoot ratio by weight	0.234	0.043	0.756	0.0150	0.0158	94.85	0.245	104.73	52.20	53.60

** Significant at 1% (P= 0.01) level of significance

* Significant at 5% (P=0.05) level of significance




Figure1.: Determination of mean, range and genetic variability parameters for nine traits in soybean under non-stress (normal water) condition

➤ **Genetic variability under drought conditions**

Genetic variability for drought tolerance has been reported by many researchers in different crops like (Thu *et al.*, 2014; Falk *et al.*, 2020; Yan *et al.*, 2020; Mishra *et al.*, 2021; Dayoub *et al.*, 2021; Bui *et al.*, 2022 in soybean), (Raina *et al.*, 2019; Amarapali 2022 in green gram), (Widuri *et al.*, 2018; Langat *et al.*, 2020 in common beans), (Upadhyay, 2005; Songsri *et al.* 2009 and Painwadee *et al.*, 2009 in pea), (Ali *et al.*, 2010 in chickpea) (Thakur *et al.*, 2013; Vaidya *et al.*, 2015; Shankar *et al.*, 2016; Barathi *et al.*, 2022 in groundnut), (Anita Kumari *et al.*, 2019; Aneja *et al.*, 2021 in mustard), (Praveen *et al.*, 2021 in sunflower) (Rajarajan *et al.* 2018 in sorghum), (Dutta and Borua, 2017; Hoque *et al.* 2020 in rice), (Li *et al.*, 2015 in maize), (Narayanan *et al.*, 2014; Fernandes *et al.*, 2020 in wheat) (Handi and Katageri, 2016 in cotton).

➤ **Analysis of variance**

Mean sum of squares of accessions in water stress condition S₁(50% water imposition), S₂ (25% water imposition) and S₃ (0% water imposition) indicated significant difference among all the genotypes for all the nine traits showed good deal of variability in the material used (**Table 4,6 and 8**). Similarly significant differences for trait RWC was reported by Bayoumi *et al.* (2008) in wheat, Kanvi *et al.* (2020) in green gram and Ajayi (2022) in cow pea; Shankar *et al.*, (2019) for plant height, root length, root shoot ratio by length, root fresh weigh, shoot fresh weight, root dry weight, shoot dry weight and RWC in groundnut; Langat *et al.* (2019) for plant height in common beans (*Phaseolus vulgaris* L.); Priya *et al.* (2021) for root length in lentil and Wattoo *et al.* (2018) for relative leaf water content, root length, shoot length, root fresh weight, shoot fresh weight, root dry weight and shoot dry weight in maize.

Table 4: Analysis of variance (mean sum of squares) for nine characters in soybean under S₁ (50% water) stress condition

Source of variation	DF	Root length (cm)	Shoot length (cm)	Root fresh weight (g)	Shoot fresh weight (g)	RWC %	Root dry weight (g)	Shoot dry weight (g)	Root shoot ratio by length	Root shoot ratio by weight
Replication	1	0.20	0.71	0.001	0.002	7.87	0.000	0.001	0.000	0.000
Treatment	59	10.40**	279.82**	0.046**	0.196**	719.33**	0.0001**	0.005**	0.015**	0.047**
Error	59	0.74	4.11	0.001	0.013	8.36	0.000	0.000	0.001	0.001
Total	119	5.52	140.78	0.023	0.104	360.85	0.000	0.003	0.008	0.024

** Significant at 1% (P= 0.01) level of significance

* Significant at 5% (P=0.05) level of significance

Table 5: Estimation of mean, range and genetic variability parameters for nine quantitative characters in Soybean under S₁ (50% water) stress condition

Genotypes	Mean	Min	Max	var (g)	var (p)	Heritability (%)	GA	GA% mean	GCV (%)	PCV (%)
Root length (cm)	9.587	4.550	16.100	3.22	3.96	81.40	3.34	34.80	18.72	20.75
Shoot length (cm)	47.468	15.050	74.000	91.90	96.01	95.72	19.32	40.70	20.20	20.64
Root fresh weight (g)	0.198	0.044	0.995	0.015	0.016	95.15	0.25	123.88	61.65	63.20
Shoot fresh weight (g)	0.769	0.072	1.727	0.061	0.074	82.81	0.46	60.32	32.18	35.36
Rwc %	63.320	20.555	93.265	236.99	245.35	96.59	31.17	49.22	24.31	24.74
Root dry weight (g)	0.021	0.006	0.044	0.0001	0.0001	76.92	0.014	64.72	35.82	40.84
Shoot dry weight (g)	0.123	0.041	0.298	0.0018	0.0018	100.00	0.087	70.55	34.25	34.25
Root shoot ratio by length	0.217	0.101	0.641	0.0049	0.0055	90.08	0.137	63.20	32.32	34.06
Root shoot ratio by weight	0.205	0.032	0.942	0.0153	0.0166	92.63	0.246	119.97	60.51	62.87




Figure 2: Estimation of mean, range and genetic variability parameters for nine quantitative characters in Soybean under S₁ (50% water) stress condition

Table 6: Analysis of variance (mean sum of squares) for nine characters in Soybean under S₂ (25% water) stress condition

Source of variation	DF	Root length (cm)	Shoot length (cm)	Root fresh weight (g)	Shoot fresh weight (g)	RWC %	Root dry weight (g)	Shoot dry weight (g)	Root shoot ratio by length	Root shoot ratio by weight
Replication	1	0.01	2.22	0.001	0.003	0.10	0.000	0.001	0.000	0.007
Treatment	59	13.03**	264.65**	0.030**	0.283**	657.89**	0.001**	0.008**	0.015**	0.118**
Error	59	0.69	3.61	0.001	0.014	1.86	0.000	0.000	0.000	0.001
Total	119	6.80	133.03	0.015	0.147	327.11	0.000	0.004	0.008	0.059

** Significant at 1% (P= 0.01) level of significance

* Significant at 5% (P=0.05) level of significance

Table 7: Estimation of mean, range and genetic variability parameters for nine quantitative characters in Soybean under stress S₂ (25% watering) condition

Genotypes	Mean	Min	Max	var (g)	var (p)	Heritabili	GA	GA%	GCV (%)	PCV (%)
-----------	------	-----	-----	---------	---------	------------	----	-----	---------	---------

						ty (%)		mean		
Root length (cm)	9.546	3.750	16.450	4.12	4.80	85.69	3.87	40.52	21.25	22.96
Shoot length (cm)	46.749	17.900	73.250	87.01	90.63	96.01	18.83	40.28	19.95	20.36
Root fresh weight (g)	0.191	0.011	0.885	0.010	0.010	94.36	0.20	103.04	51.49	53.01
Shoot fresh weight (g)	0.757	0.104	1.739	0.090	0.104	86.22	0.57	75.66	39.55	42.60
Rwc %	64.274	26.435	92.115	218.68	220.54	99.16	30.33	47.19	23.01	23.11
Root dry weight (g)	0.025	0.006	0.084	0.0002	0.0002	100.00	0.026	107.55	52.21	52.21
Shoot dry weight (g)	0.123	0.032	0.319	0.0027	0.0027	100.00	0.107	87.33	42.39	42.39
Root shoot ratio by length	0.218	0.095	0.471	0.0048	0.0052	91.21	0.136	62.29	31.66	33.15
Root shoot ratio by weight	0.259	0.030	1.635	0.0389	0.0398	97.74	0.402	154.97	76.09	76.96




Figure 3: Estimation of mean, range and genetic variability parameters for nine quantitative characters in Soybean under stress S₂ (25% watering) condition

Table 8: Analysis of variance (mean sum of squares) for nine characters in Soybean under S₃ (0% water) stress condition

Source of variation	DF	Root length (cm)	Shoot length (cm)	Root fresh weight (g)	Shoot fresh weight (g)	RWC %	Root dry weight (g)	Shoot dry weight (g)	Root shoot ratio by length	Root shoot ratio by weight
Replication	1	0.03	4.73	0.000	0.004	5.40	0.000	0.002	0.000	0.007

Treatment	59	27.15**	411.19**	0.026**	0.497**	756.20**	0.001**	0.013**	0.028**	0.065**
Error	59	0.70	5.91	0.000	0.012	1.87	0.000	0.000	0.000	0.001
Total	119	13.81	206.84	0.013	0.252	375.90	0.000	0.007	0.014	0.033

** Significant at 1% (P= 0.01) level of significance

* Significant at 5% (P=0.05) level of significance

UNDER PEER REVIEW

➤ Genetic variability parameters under water stress conditions

Genetic variability for drought tolerance has been reported by many researchers in different crops like in soybean (Dayob *et al.*, 2021), (Raina *et al.*, 2019; Amarapali 2022 in green gram), (Widuri *et al.*, 2018; Langat *et al.*, 2020 in common beans), (Upadhyay, 2005; Songsri *et al.* 2009 and Painwadee *et al.*, 2009 in pea), (Ali *et al.*, 2010 in chickpea) (Thakur *et al.*, 2013; Vaidya *et al.*, 2015; Shankar *et al.*, 2016; Barathi *et al.*, 2022 in groundnut), (Anita Kumari *et al.*, 2019; Aneja *et al.*, 2021 in mustard), (Praveen *et al.*, 2021 in sunflower) (Rajarajan *et al.* 2018 in sorghum), (Dutta and Borua, 2017; Hoque *et al.* 2020 in rice), (Li *et al.*, 2015 in maize), (Narayanan *et al.*, 2014; Fernandes *et al.*, 2020 in wheat) (Handi and Katageri, 2016 in cotton).

The estimates of variability parameters of all genotypes for root shoot characters under water stress have been presented in **Table 5, 7 and 9 and Figure 2,3 and 4.**

Among abiotic factors, the scarcity of water great impact on root distribution system and root length. A dry soil surface with high water stress often forces roots to increase their growth deep into the soil profile where water is more available (Adiku *et al.*, 1996, Amarapali, 2022). The deep and dense root system show better drought tolerance by extracting water from deeper soil layers (Parameshwarappa *et al.*, 2012). The root length showed significant good amount of genetic variation under stress conditions among the genotypes used for investigation. Under stress **S₁ (50% water)**, the range of mean was 4.55 cm to 16.10 cm with overall mean of 9.59 cm. **In stress condition S₂ (25% water)** the range of root length variability was observed from 3.75 cm to 16.45 cm with the grand mean of 9.55 cm. **Under stress condition S₃ (0% water)** highest root length was calculated by genotype NRC138 (20.50 cm) whereas lowest was recorded by GW312 (3.40 cm) with the overall mean of 0.41 cm. It indicates that germplasm lines are genetically variable. Similar increasing range of variation under water stress in comparison to normal condition was also noticed by Dayoub *et al.* (2021) in soybean; Raina *et al.* (2019) in mung bean. PCV and GCV recorded were (20.75% and 18.72%) under stress **S₁ (50% water)**; (22.96 and 21.25 per cent) under **stress condition S₂ (25% water) and** (22.96 and 21.25 per cent) **under stress condition S₂ (25% water) respectively.** Similar high variability was reported by Manickvelu *et al.* (2006), Dutta and Borua (2017), Sallleh *et al.* (2021) in rice; Rajkumar and Fakrudin (2018) in sorghum, Shankar *et al.*, (2019) in groundnut, Bayoumi *et al.* (2008) in wheat, Mishra and Sharma, (2015) in muskmelon, Wattoo *et al.* (2019) in maize. High heritability along with high genetic advance as percentage of mean was observed (81.40% and 34.80%) under stress **S₁ (50% water)**; (85.69% and 40.52%) **in stress condition S₂ (25% water) and** (92.60% and 56.52%) **in stress condition S₃ (0% water)** respectively. High heritability

coupled with high genetic advance represent that character root length is governed by additive gene action. It is **indicating that simple selection would be effective for this trait under stress condition**. Similar result of high heritability and high genetic advance for this trait was also observed by Irum *et al.*, (2011) and Riaz *et al.* (2013) in cotton, Dutta and Borua (2017) in rice, Gobu *et al.* (2017) in eggplant, Rajkumar and Fakrudin (2018) in sorghum, Shankar *et al.* (2019) in groundnut, Mishra and Sharma, (2015) in muskmelon, Gedam *et al.* (2021) in onion and **Kumar *et al.* (2023) in wheat under stress condition**. Bayoumi *et al.* (2008) in wheat, Wattoo *et al.* (2019) in maize and Pavitra *et al.* (2022) in rice for high heritability.

A wide variation was found among the germplasm accessions for shoot length. This character varied from 15.05 cm to 74.00 cm with an overall mean of 47.47 cm in **S₁** ; 17.90 cm to 73.25 cm with the overall mean of 46.74 cm in **S₂** and 14.10 to 90.50 cm with a grand mean of 49.36 cm in stress conditions in **S₃**. The PCV and GCV were 20.64 and 20.20 per cent; 20.36 and 19.95 per cent; 24.06 and 23.55 per cent respectively in stress conditions **S₁ (50% water)**, **S₂ (25% water)** and **S₃ (0% water)**. Similarly, high (More than 20%) PCV and GCV, was recorded by Mishra *et al.*, (2015) in muskmelon, Bayoumi *et al.* (2008), Ashfaq *et al.* (2022) and Kumar *et al.* (2023) in wheat. The estimates of h^2 (95.72%, 96.01% and 95.81 per cent) together with an expected GA over mean of (40.70%, 40.28% and 47.48 per cent) were recorded for this character in stress conditions **S₁ (50% water)**, **S₂ (25% water)** and **S₃ (0% water)** respectively. Similar data on heritability for this trait was also obtained by Meena *et al.* (2014) in chickpea, Mishra and Sharma, (2015) in muskmelon, Gobu *et al.* (2017) in eggplant, Shankar *et al.* (2019) in groundnut, Wattoo *et al.* (2019) in maize, Gedam *et al.* (2021) in onion; for high h^2 and high GA as percentage of mean by Riaz *et al.* (2013) in cotton and by Kumar *et al.* (2023) in wheat for this trait.

Root biomass and root length that aid in greater soil moisture extraction were identified as important root traits during terminal drought (Kashiwagi *et al.*, 2006; Varshney *et al.*, 2011). High PCV, GCV, heritability and genetic advance as percentage of mean was observed for root fresh weight under stress condition **S₁**, **S₂** and **S₃** respectively. In stress conditions **S₁ (50% water)** the range of genetic variability for the trait root fresh weight was found from 0.04 gm to 0.10 gm with the overall mean (0.20 gm), **in stress condition S₂ (25% water)** the grand mean of root fresh weight was 0.19 gm whereas highest root fresh weight was depicted by genotype GW312 (0.885 gm) and lowest by genotype GW212 (0.011 gm), **in stress condition S₃ (0% water)** the overall mean of it was 0.19 gm while highest root fresh weight was produced by genotype GW312 (0.79 gm) and lowest by GW212 (0.01 gm). The observed values of PCV, GCV, heritability and GA over percentage of mean were (63.20, 61.65 per cent, 95.15% and 123.28%); (53.01 and 51.49 per cent, 94.36% and 103.04%); (50.34, 49.06 per cent, 94.99% and 98.50%) under stress conditions **S₁**, **S₂** and **S₃** respectively. The

observed values of PCV, GCV, heritability and GA over percentage of mean were according to findings of by Riaz *et al.* (2013) in cotton and Rajkumar and Fakrudin (2018) in sorghum; by Wattoo *et al.* (2019) in maize and by Shankar *et al.* (2019) in groundnut.

A wide variation was observed among the germplasm accessions for shoot fresh weight. The character shoot fresh weight varied from (0.072 to 1.73 gm; 0.10 to 1.74 gm; 0.13 to 2.35 gm) with an overall mean of (0.77 gm; 0.76 gm; 0.85 gm) under **S₁**, **S₂** and **S₃** respectively. The PCV, GCV, h^2 together with an expected GA over mean under **S₁**, **S₂** and **S₃** were (35.36%, 32.18%, 82.81% and 60.08%); (42.60%, 39.55%, 86.22% and 75.66 %); (48.82%, 47. %, 86.22 % and 75.66 %) respectively. Similar result of high PCV and GCV and heritability for this trait was also recorded by Wattoo *et al.* (2019) in maize; for high heritability by Irum *et al.* (2011), Riaz *et al.* (2013), Handi and Katageri (2016) in cotton; for high heritability and high genetic advance by Shankar *et al.* (2019) in groundnut.

RWC is the appropriate measure of plant water status in terms of the physiological consequence of cellular water deficit. In case of absorption of water by the plant by roots from soil, water potential is useful in dealing with water transport in the soil-plant-atmosphere continuum but does not account for osmotic adjustment (OA). Osmotic adjustment (OA) mechanism is responsible for conserving cellular hydration under drought stress and RWC expresses the effect of osmotic adjustment (OA) in this respect. Therefore, RWC analysis is an appropriate estimate of plant water status in terms of cellular hydration under the possible effect of both leaf water potential and OA. The method is simple and measure of water deficit in the leaf. RWC value generally range between 98% in turgid and transpiring leaves to about 40% in severely desiccated and dying leaves. In most crop species the typical RWC at about wilting is around 60% to 70%, with exceptions. A wide range of variation in **S₁**, **S₂** and **S₃** stress (20.55 to 93.26%; 26.43 to 92.11%; 20.06% to 95.84%) was found among all the genotypes relative leaf water content and the overall mean of physiological character was 63.32%, 64.27% and 63.18%. The PCV (24.74%, 23.11% and 25.19%), GCV (24.31, 23.01 and 25.10%), heritability (h^2) (96.59%, 99.16% and 99.26%) and genetic advance over mean of percentage (49.22%, 47.19% and 51.51%) were observed respectively for this parameter under all the three water stress conditions (50%, 25% and 0%). Similar results were also reported by Manickvelu *et al.* (2006) in rice. High heritability for this parameter was also reported by Meena *et al.* (2014) in chickpea, Garg *et al.* (2017) in rice, Wattoo *et al.* (2019) in maize, Shankar *et al.* (2019) in groundnut and Gedam *et al.* (2021) in onion.

Data of variability for root dry weight under drought has been presented in Table **5**, **7** and **9**. While studying of the genotypes for drought tolerance, dry weight of a plant at age is universally considered as more

stable character than other morphological parameters (Dutta and Bera, 2008; Vijay *et al.* 2018). The PCV and GCV obtained for root dry weight were (40.84 and 35.82 per cent; 52.21% and 52.21%; 65.23% and 62.99%) respectively. The values of heritability (76.92%, 100.00% and 93.24%) along with high genetic advance as per cent mean (64.72%, 107.55% and 125.29%) were observed for this trait under water stress conditions **S₁**, **S₂** and **S₃** respectively. High PCV, GCV, heritability and genetic advance were also reported for this trait by Riaz *et al.* (2013) in cotton and Rajkumar and Frakudin (2018) in sorghum; for high PCV, GCV and heritability by Wattoo *et al.* (2018) in maize. The findings on heritability for this trait was according to Irum *et al.* (2011), Handi and Katageri (2016) in cotton; Lalitha *et al.* (2015) in chickpea and Gurumurthy *et al.* (2019) in black gram.

The results of shoot dry weight indicated that out of 60 genotypes, 30 genotypes showed high values in comparison to the grand mean (0.12gm). High phenotypic and genotypic coefficient of variability along with high heritability and genetic advance as per cent mean recorded were (34.25%, 42.39% and 46.87%), (34.25%, 42.39% and 46.52%), (100.00%, 100.00 % and 98.50 %) and (70.55%, 87.33% and 95.10%) respectively. Similar findings of high PCV, GCV, high h^2 along with high genetic advance as percentage of mean was reported by Kumar *et al.* (2023) in wheat. Similar result of high heritability and high GA as percentage of mean for this trait was also found by Irum *et al.* (2011), Riaz *et al.* (2013), Handi and Katageri (2016) in cotton, Lalitha *et al.* (2015) in chickpea, Langat *et al.* (2019) in common beans (*Phaseolus vulgaris* L.), Gurumurthy *et al.* (2019) in black gram, Shankar *et al.* (2019) in groundnut. High PCV and GCV for this trait were also reported by Doumbia *et al.* (2022) in cowpea.

Perusal data of root shoot ratio by length indicated the range of genetic variability (0.10-0.64), (0.10 to 0.47) and (0.01-0.91) with grand mean of (0.22, 0.22 and 0.23). The PCV and GCV observed were (34.06, 33.15, 43.88) and (32.32, 31.66, 42.44) per cent, respectively while high heritability (bs) of (90.08 per cent, 91.21 per cent, 96.14 per cent) coupled with high genetic advance over percentage of mean (63.20 per cent, 62.29 per cent, 85.71 per cent) were noticed under water stress conditions **S₁**, **S₂** and **S₃** respectively. Similarly high PCV and GCV, heritability and genetic advance of root shoot ratio by length were recorded by Mishra and Sharma, (2015) in muskmelon, Gobu *et al.* (2017) in eggplant, Rajkumar and Frakudin (2018) in sorghum and Shankar *et al.* (2019) in groundnut.

The character root shoot ratio by weight expressed ranged from 0.03 to 0.94, 0.03 to 1.64 and 0.02 to 0.75 under all the three water stress conditions (**S₁**, **S₂** and **S₃**). The grand mean of root shoot ratio in weight was 1.39, 0.26 and 0.23. The estimates of PCV (62.87%, 76.96 % and 65.60%), GCV (60.51%, 76.96% and

64.00%), h^2 (92.63%, 97.74% and 95.17%) and G.A. (119.97%, 154.97% and 128.62%) respectively were recorded for this trait under all the three water regimes (S_1 , S_2 and S_3) created for drought. Similar results of high GCV, heritability and high GA as percentage of mean was reported by Handi and Katageri (2016) in cotton.

High heritability along with high genetic advance as percentage of mean was recorded for most of the traits under stress conditions. High heritability ($\geq 95\%$) combined with high genetic advance as percentage of mean of root fresh weight, root dry weight and root shoot ratio by weight than other traits indicating that simple selection would be effective for these traits among superior genotypes than other traits under water stress conditions.

UNDER PEER REVIEW

Table 9: Estimation of mean, range and genetic variability parameters for nine quantitative characters in Soybean under stress S₃ (0% watering) condition

Genotypes	Mean	Min	Max	var (g)	var (p)	Heritability (%)	GA	GA% mean	GCV (%)	PCV (%)
Root length (cm)	10.41	3.40	20.50	8.82	9.52	92.60	5.89	56.52	28.51	29.63
Shoot length (cm)	49.36	14.10	90.50	135.09	141.01	95.81	23.44	47.48	23.55	24.06
Root fresh weight (g)	0.186	0.013	0.789	0.008	0.009	94.99	0.18	98.50	49.06	50.34
Shoot fresh weight (g)	0.854	0.125	2.348	0.162	0.174	93.05	0.80	93.57	47.09	48.82
Rwc %	63.18	20.06	95.84	251.45	253.31	99.26	32.54	51.51	25.10	25.19
Root dry weight (g)	0.026	0.004	0.135	0.0003	0.0003	93.24	0.033	125.29	62.99	65.23
Shoot dry weight (g)	0.144	0.031	0.378	0.0045	0.0045	98.50	0.137	95.10	46.52	46.87
Root shoot ratio by length	0.227	0.093	0.911	0.0093	0.0097	96.14	0.195	85.71	42.44	43.28
Root shoot ratio by weight	0.229	0.023	0.750	0.0214	0.0225	95.17	0.294	128.62	64.00	65.60




Figure 4: Estimation of mean, range and genetic variability parameters for nine quantitative characters in Soybean under stress S₃ (0% watering) condition



Figure-5: Experimental View at Net house

Conclusion:

It may be concluded that ample scope of direct selection for most of the traits because this exhibited high genetic variability. Under stress conditions, high broad sense heritability ($\geq 95\%$) together with high genetic advance of root fresh weight, root dry weight and root shoot ratio by weight than other traits demonstrated that these traits would grant more superior genotypes through phenotypic based selection than other traits in future soybean crop improvement in relation to drought stress.

Acknowledgements:

Authors express deep sense of gratitude of Madhya Pradesh Council of Science and Technology (MPCOST) for providing financial assistance (Grant No. 2793). Authors are extremely grateful of Director, ICAR-Indian Institute of Soybean Research (ICAR-IISR) Indore, who provided germplasm of soybean for the experimentation. All the assistance for conduct the experiment provided by Head, Faculty of Agriculture Science, is gratefully acknowledged.

Disclaimer (Artificial intelligence):

Authors hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

REFERENCES

1. Adiku, S., Braddock, R. and Rose, C. (1996). Simulating growth dynamics. *Environml. Software*, **11**: 99-103.
2. Anjum, S.A., Ashraf, U., Zohaib A. and Tanveer, M. (2017). Growth and developmental responses of crop plants under drought stress: A review. *Zemdirbyste-Agriculture.*, 104(3):267-276.
3. Ajayi, A. T.. Screening for drought tolerance in cowpea at the flowering stage. *International Journal of Science Letters*. 2022; 4(2): 236-268.
4. Akram, R.M., Fares, W. M., Fateh, H.S.A. and Rizk, A.M.A. (2011). Genetic variability, correlation and path analysis in soybean. *Egypt. Journal Plant Breeding*, 15(1):89-102.
5. Amarapalli G. (2022). Studies on the effect of water stress on root traits in green gram cultivars. *Legume Research*, 45(4), 422-428.
6. Ashfaq W., Brodie G., Fuentes S. and Gupta D. (2022). Infrared thermal imaging and morpho-physiological indices used for wheat genotypes screening under drought and heat stress. *Plants*, 2022;11:3269. <https://doi.org/10.3390/plants11233269>.
7. Aski M.S., Rai N., Reddy VRP, Gayacharan, Dikshit H.K., Mishra G.P., Singh D., Kumar A., Pandey R., Singh M.P., Pratap A., Nair R.M., Schafleitner R. (2022). Assessment of root phenotypes in mungbean mini-core

collection (MMC) from the world vegetable centre (AVRDC) Taiwan. *PLOS ONE*, 16(3): e0247810. <https://doi.org/10.1371/journal.pone.0247810>.

8. Bairagi V., Mishra S., Sen Ramraj, Dixit S., and Tyagi D.B. (2023). Assessment of genetic variability in soybean (*Glycine max* L., Merrill). *Biological Forum-An Int. Journal*, 15(5):258-263.
9. Bayoumi T.Y., Manal H.E. and Metwali E.M. (2008). Application of physiological and biochemical indices as a screening technique for drought tolerance in wheat genotypes. *African J. of Biotechnology*, 7(14):2341-2352.
10. Bhuva B., Giovarruscio M., Rahim N., Bitter K., Mannocci F. (2020). The restoration of root filled teeth: a review of the clinical literature. *International Endodontic J.*, 54(4):509-535.
11. Burton G. W. (1952). Quantitative inheritance of grasses. Proc. 6 th International Grass Land Congress, *Ag. Sciences*, 1: 277- 282.
12. Dai, A. (2013), Increasing Drought under Global Warming in Observations and Models. *Nature Climate Change*, 3:52-58.
13. Doumbia I.Z., Siddiqui M.N., Diallo M., Dembele S., Dolo A. and Mulaudzi T. (2022). Performance of Malian cowpea (*Vigna unguiculata* (L.) Walp) genotypes to drought stress under field conditions. *J. of Agricultural and Crop Research*, 10(8):154-166.
14. Dutta P. and Borua P.K. (2017). Genetic parameters for drought related traits in rice. *International J. of Engineering Science Invention*, 6(9):32-37.
15. Falk K.G., Jubery T.Z., Jamie A., O'Rourke, Singh, A., Sarkar S. *et. al.* (2020). Soybean root system architecture trait study through genotypic, phenotypic, and shape-based clusters. *Plant Phenomics*, :1-23. Article ID 1925495, <https://doi.org/10.34133/2020/1925495>.
16. Fenta B.A., Beebe, S.E., Kunert, K.J., Burrridge, J.D., Barlow, K.M., Lynch, P.J. *et. al.* (2014). Field phenotyping of soybean roots for drought stress tolerance. *Agronomy*, 4:418–435.
17. Fernandes R.C., Oliveira V. F. D., Schroeder L.D., Jesus A.P.D., Busanello C., Santos M. F. *et al.* (2020). Morphologically accessing wheat genotypes in response to drought on early stages of development. *Communicati. in Plant Sciences*, 10:105-113.
18. Fried H.G., Narayanan S., Fallen B. (2018). Characterization of a soybean (*Glycine max* L. Merr.) germplasm collection for root traits. *Plos One journal.pone*, 13(7):1-19.
19. Garg H.S., Kumari P., Bhattacharya C., Panja S. and Kumar R. (2017). Genetic parameters estimation for yield and yield related traits in rice (*Oryza sativa* L.) with drought tolerance trait under stress condition. *J. of Crop and Weed*, 13(1):83-88.

20. Gedam P.A., Thangasamy A., Shirsat D.V., Ghosh S.M., Bhagat K.P., Sogam O.A. *et al.* (2021). Screening of onion (*Allium cepa* L.) genotypes for drought tolerance using physiological and yield based indices through multivariate analysis. *Frontiers in Plant Sci.*, 12: 600371. doi: 10.3389/fpls.2021.600371.
21. Giordani, W., Gonçalves, L. S. A., Moraes, L. A. C., Ferreira, L. C., Neumaier, N., Farias, J. R. B. *et al.* (2019). Identification of agronomical and morphological traits contributing to drought stress tolerance in soybean. *Australian Journal of Crop Science*, 13(1), 35-44. DOI: <https://doi.org/10.21475/ajcs.19.13.01.p1109>.
22. Gobu R., Babu B.N.H., Chandra K. Shankar M. and Omprakash (2017). Genetic variability, heritability and genetic advance in eggplant (*Solanum melongena* L.) genotypes under normal and osmotic stress in invitro condition. *International J. of Current Microbiology and Applied Sciences*, 6(3):749-760.
23. Gurumurthy, S., Sarkar, B., Vanaja, M., Lakshmi, J., Yadav, S.K., and Maheswari, M. (2019). Morpho-physiological and biochemical changes in black gram [*Vigna mungo* (L.) Hepper] genotypes under drought stress at flowering stage. *Acta Physio. Plantarum*, 41(3): 1-14.
24. Handi S.S. and Katageri I.S. (2016). Genetic variability for root and shoot traits under moisture stress in *G. hirsutum* cotton. *Research in Environ. and Life Sciences*, 9(5):546-549.
25. <http://www.sopa.org>.
26. Irum A., Tabasum A. and Iqbal M.Z. (2011). Variability, correlation and path coefficient analysis of seedling traits and yield in cotton (*Gossypium hirsutum* L.). *African J. of Biotechnology*, 10:18104-18110.
27. Johnson H. W., Robinson H. F. and Comstock E. (1955). Estimates of genetic and environment variability in Soyabean. *American J. of plant sciences*, 47:314-318.
28. Kanavi M.S.P., Nagesha N., Somu G., Krishnaprasad B.T. and Rangaiah S. (2020). Principal component analysis of physiological traits governing drought tolerance in germplasm accessions of green gram (*Vigna radiata* L.). *International J. of Current Microbiology and Applied Sciences*, 9(3):2943-2956.
29. Kashiwagi J., Krishnamurthy L., Crouch J.H., Serraj R. (2006). Variability of root length density and its contributions to seed yield in chickpea (*Cicer arietinum* L.) under terminal drought stress. *Field Crops Resea.*, 95:171–181.
30. Kumar D., Rana V., Rana M. and Guleria P. (2023). Genetic variability, correlation and path analysis studies for grain yield and morpho-physiological traits under moisture stress conditions in bread wheat (*Triticum aestivum* L.) under north-western Himalayan conditions. *J. of Cereal Research*, 15(1):92-102.
31. Kumar H.S., Yerrabala S., Lavanya G.R., Goud G.B., Reddy Y.V.K. and Sharmila V. (2021). Association, direct and indirect effects of yield, biochemical and physiological characters on seed yield in chickpea (*Cicer arietinum* L.). *Environment Conservation J.*, 22 (3):169-180.

32. Kumawat A., Mishra S., Sen R, Sharma R. and Bairagi V.D. (2023). Genetic variability studies and morphological characterization in soybean (*Glycine max* L., Merrill). *The Pharma Inno.*, 12(11):1668-1674.
33. Lalitha N., Upadhyaya, H.D., Krishnamurthy L., Kashiwagi J., Kavikishor P.B. and Singh S. (2015). Assessing genetic variability for root traits and identification of trait-specific germplasm in chickpea reference set. *Crop Sci.*, 55:2034-2045.
34. Langat C., Ombori O., Leley P., Karanja D., Cheruiyot R., Gathaara. M, and Masila B. (2019). Genetic Variability of Agronomic Traits as Potential Indicators of Drought Tolerance in Common Beans (*Phaseolus vulgaris* L.). *International J. of Agronomy* Volume 2019 | Article ID 2360848 | <https://doi.org/10.1155/2019/2360848>.
35. Li R., Zeng Y., Xu J., Wang Q., Wu F., Cao M., Lan H., Liu Y. and Lu Y. (2015). Genetic variation for maize root architecture in response to drought stress at the seedling stage. *Breeding Sci.*, 65:298-307.
36. Manavalan, L.P.; Guttikonda, S.K.; Tran, L.- S.P.; Nguyen, H.T. (2009). Physiological and molecular approaches to improve drought resistance in soybean. *Plant and Cell Physiol.*, 50:1260-1276. DOI: <https://doi.org/10.1093/pcp/pcp082>.
37. Meena H.P., Kumar J. and Ramesh M. (2014). Evaluation of the reaction of chickpea (*Cicer arietinum* L.) genotypes to drought conditions using various stress tolerance indices. *Legume Res.*, 37(5): 453-459.
38. Mehra S., Shrivastava M.K., Amrate P.K. and Yadav R.B. (2020). Studies on variability, correlation coefficient and path analysis for yield associated traits in soybean (*Glycine max* L., Merrill). *J. of Oilseeds Research*, 37(1):56-59.
39. Mishra N., Tripathi M.K., Tiwari S., Tripathi N., Gupta N., Sharma A. (2021). Morphological and physiological performance of indian soybean [*Glycine max* (L.) Merrill] genotypes in respect to drought. *Legume Res.*, DOI: 10.18805/LR-4550.
40. Mishra S. and Patidar D. (2023). Effect of drought stress on growth of soybean under seedling stage. *International J. of Economic Plants*, 10(3):231-245.
41. Mishra S. and Sharma A.K. (2015). Genetic variability and correlation study of muskmelon (*Cucumis melo* L.) genotypes under osmotic stress at early seedling growth stage. *Environ. and Ecology*, 34(2):560-564.
42. Mishra S., Chugh C., Sharma A.K., and Sharma V. (2017). Correlation analysis of muskmelon (*Cucumis melo* L.) genotypes under different water regimes in greenhouse. *Bioscience Biotech. Research Communications*, 10(1):22-27.
43. Morsy, A.R., Fares W.M., Ragheb S.B. and Ibrahim M.A. (2015). Stability analysis of some soybean genotypes using a simplified statistical model. *J. of Plant Production*, 6(12):1975-1990.

44. Nair R.M., Boddepalli V.N., Yan M.R., Kumar V., Gill B., Pan R.S. *et al.* (2023). Global status of vegetable soybean. *Plants* 2023, 12(3):609; <https://doi.org/10.3390/plants12030609>.
45. Narayanan S, Mohan A, Gill K.S., Prasad P.V.V. (2014). Variability of Root Traits in Spring Wheat Germplasm. *Plos One*, 9(6):1-12, e100317. doi:10.1371/journal.pone.0100317.
46. Narayanan S. and Fallen B. (2019). Evaluation of soybean plant introductions for traits that can improve emergence under varied soil moisture levels. *Agron.*, 9:118. doi:10.3390/agronomy9030118.
47. Painawadee M., Jogloy S., Kesmala T., Akkasaeng C. and Patannoithai A. (2009). Heritability and correlation of drought resistance traits and agronomic traits in peanut (*Arachis hypogaea* L.). *Asian J. of Plant Sciences*, (5):325-334.
48. Panse V.G. and Sukhatme PV. 1978. Statistical methods for research workers. New Delhi: ICAR.
49. Parameshwarappa S.G., Salimath P.M., Upadhyaya H.D., Patil S.S., Kajjidoni S.T. and Patil B.C. (2010). Characterization of drought tolerant accessions identified from the minicore of chickpea (*Cicer arietinum* L.). *Indian J. of Genetics*, 70(2): 125-131.
50. Pavitra S., Vengadessan V. and Meena S.K. (2022). Genetic variability analysis in rice genotypes under drought stress. *Biological Forum An International J.*, 14(2):883-890.
51. Prince S.J., Song Li, Qiu D., dos Santos J.V.M., Chai C., Joshi T. *et al.* (2015). Genetic variants in root architecture-related genes in *Glycine soja* accession, a potential resource to improve cultivated soybean. *BMC Genom.*, 16:132. DOI 10.1186/s12864-015-1334-6.
52. Priya, S., Bansal, R., Kumar, G., Dikshit, H.K., Kumari, J., Pandey, R. *et al.* (2021). Root Trait Variation in Lentil (*Lens culinaris* Medikus) Germplasm under Drought Stress. *Plants*, 10, 2410. <https://doi.org/10.3390/plants10112410>.
53. Raina S.K., Rane J., Raskar N., Singh A.K., Govindasamy V., Kumar M. *et al.* (2019). Physiological traits reveal potential for identification of drought tolerant mungbean [*Vigna radiate* (L.) Wilczek] genotypes under moderate soil moisture deficit. *Indian J. of Genetics and Plant Breeding*, 79(2):427-437.
54. Rajarajan, K., Ganesamurthy, K. and Yuvaraja, A. (2018). Genetic variability and diversity for shoot/root parameters under early drought stress condition in sorghum (*Sorghum bicolor* (L.) Moench) genotypes. *Forage Res.*, 43(4):266-269.
55. Rajkuar B.K. and Fakrudin B. (2018). Genetic association of root and yield traits in two recombinant inbred populations of sorghum under terminal drought condition. *Bulletin of Environ., Pharmacology and Life Sci.*, 7(4):65-76.

56. Reddy B.V.R. P., Dikshit H.K., Aski M.S. and Mishra G.P. (2023). Comparative analysis of root system architectural traits in genotypes of *vigna radiata* L. and *vigna mungo* L. *Bangladesh J. Bot.*, 52(1):61-69.
57. Riaz M., Farooq J., Sakhawat G., Mahmood A., Sadiq M A, Yaseen M. (2013) Genotypic variability for root/shoot parameters under water stress in some advanced lines of cotton (*Gossypium hirsutum* L.). *Genetics and Molecular Res.*, 12(1):552-561.
58. Saha I. and Choyal P. (2022). Drought stress responses and inducing tolerance by seed priming approach in plants *Plant Stress Volume 4*, April 2022, 100066.
59. Shankar M., Harish Babu BN., Gobu R. and Sheshaiah (2019). Studies on genetic variability, heritability and genetic advance in groundnut (*Arachis hypogaeae* L.) genotypes under normal and moister stress condition in vegetative stage. *J. of Pharmacognosy and Phytochemistry*, 8 (3).
60. Sivasubramanian S. and Menon M. (1973). Heterosis and inbreeding depression in rice. *Madras Agriculture J.*, 60:1139-1149.
61. Stolf- Moreira R., Medri M.E., Neumaier N., Lemos N.G., Pimenta J.A., Tobita S. *et al.* (2010). Soybean physiology and gene expression during drought. *Genetics and Molecular Research*, 9(4):1946-1956.
62. Sunaryo W., Widoretno W., Nurhasanah, Sudarsono (2016). Drought tolerance selection of soybean lines generated from somatic embryogenesis using osmotic stress simulation of poly-ethylene glycol (PEG). *Nusantara Bioscience*, 8(1): 45-54.
63. Syiem R.M., Chaturvedi H.P., Shah P. and Sharma M.B. (2022). Genetic variability and correlation analysis for seedling vigor traits in soybean [*Glycine max* (L.) Merrill] genotypes. *The Pharma Innovations*, 11(6):1697-1699.
64. Thu, N.B.A. Nguyen Q.T., Hoang X.L.T., Thao N.P., and Tran L.S. P. (2014). Evaluation of Drought Tolerance of the Vietnamese Soybean Cultivars Provides Potential Resources for Soybean Production and Genetic Engineering. *BioMed Res. International*, 2014:1-9.
65. Varshney, R.K., Pazhamala, L., Kashiwagi, J., Gaur, P.M., Krishnamurthy, L., Hoisington, D.A. (2011). Genomics and physiological approaches for root trait breeding to improve drought tolerance in chickpea (*Cicer arietinum* L.). In: de Oliveira A.D., Varshney R.K. (eds) Root genomics. *Springer, Germany*, pp 233–250.
66. Vijay R., Ravichandran V., and Boominathan P. (2018). Assessment of soybean genotypes for PEG induced drought tolerance at germination and seedling level. *Madras Agricultural J.*, 105:1-3.
67. Vollmann J. (2016). Soybean versus other food grain legumes: A critical appraisal of the United Nations international year of pulses 2016. *Die Bodenkultur: J. of Land Management, Food and Environment*, 67(1):17-24.

68. Wattoo, F.M., Rana, R.M., Fiaz, S., Zafar, S.A., Noor, M.A., Hassan, H.M. *et al.* (2018). Identification of drought tolerant maize genotypes and seedling based morpho-physiological selection indices for crop improvement. *Sains Malaysiana* 47(2):295-302.
69. Yan, C., Song, S., Wang, W., Wang, C., Li, H. , Wang, F., Li, S. and Sun X. (2020). Screening diverse soybean genotypes for drought tolerance by membership function value based on multiple traits and drought tolerant coefficient of yield. *BMC Plant Biol.*,20:321.<https://doi.org/10.1186/s12870-020-02519-9>.

UNDER PEER REVIEW