

Perception of Childhood Routine Immunisation among Adults in Selected Communities of Ido Local Government Area, Ibadan, Oyo State, Nigeria

ABSTRACT

Aims: Immunisation is one of the medical achievements of the twentieth century, however some vaccines are receiving less and less support. Increasing people's understanding of vaccination's direct advantages which is actually beyond avoiding infectious diseases may aid in regaining vaccine acceptance for children two years and below. The study determines perception of childhood routine immunization among adults in selected communities of Ido Local Government Area, Ibadan, Oyo State.

Study design: Mixed method study design.

Methodology: A mixed-method study design which adopted a multi-stage sampling technique to select 417 respondents for the study was used. Two instruments were utilized for data collection, a questionnaire and in-depth interview guide. The questionnaire assessed perceived severity and benefits had 36 items each and scores below < 19 and score ≥ 19 were categorized as low and high for each of the variable respectively. Quantitative data was analyzed using descriptive and inferential (Chi-square) statistics at a significance level of 0.05. The qualitative data was analyzed using thematic approach.

Results: There were 41 (9.8%) men and 376 (90.2%) women making a total of 417 adults. The respondents 208 (49.9%) have Secondary School Certificate and this was the highest educational qualification among the respondents. Results also revealed that 172 (41.3%) of respondents had good knowledge about children's routine vaccinations. About 396 (95.0%) had high level of severity to vaccine preventable diseases. Also, 409 (98.1%) had high level of perceived benefits of immunization uptake.

Conclusion: This study showed a high level of perceived benefits of immunization uptake among the respondents. Since respondents' perception on childhood routine immunization and education level were substantially correlated, maternal education should be encouraged and improved by the government.

Keywords: Perception, Childhood routine immunization, adults.

1. INTRODUCTION

One of the most less expensive and successful public response to disease prevention is immunization, also it have significantly improved health and decreased death (1). Although immunization of children has been very successful in low- and middle-income countries, but within these countries vaccine-preventable diseases has been a major health problem

among the children. Each year, immunization prevents 2 to 3 million infant deaths worldwide (2). The World Health Organization estimated that due to the disruption of normal immunization services, 80 million children in 68 countries are at danger of contracting diseases like measles, diphtheria, and polio (3). This is partly attributable to vaccinations' success in preventing contagious disease. However, due to the huge range of illnesses and ailments that exist today, many individuals are simply unaware of the crippling effects of this disease. The average population struggles to have a thorough understanding of all the many diseases, symptoms, and suggestions. It's crucial that people are aware about diseases that can be prevented by vaccines in order to lower their prevalence. Serious illnesses were those that were life-threatening, chronic, or had long-lasting repercussions, according to moms who participated in this study. Most moms considered polio, diphtheria, tetanus, and meningitis to be potentially fatal diseases. All but meningitis were deemed unlikely to affect children.

Kazi et al. (4) describes the benefits of adopting healthy habits as a means to lower your chance of contracting an illness. People tend to adopt healthier behaviors when they believe the new behavior will reduce their chances of developing a disease (4). If they think getting the MMR vaccine will protect their body from the disease, some people might be persuaded to do so. Early immunization would avoid illnesses later in life, which would reduce productivity and increase health care expenses for everyone. People may be aware of the protective effects that comprehensive vaccination can have on those who are vulnerable in addition to their own personal benefits (herd immunity). People may be able to see how their individual actions have an effect on society as a whole. A person's impressions of the advantages of vaccination may be influenced by their awareness of their obligation to those in their immediate environment. However, as misunderstandings about vaccination and VPDs spread and are not addressed by immunization programs, it is probable that this demand could decline in the future (5). Delivering accurate information to enhance public awareness of the advantages and security of vaccinations. In order to provide a child with the greatest protection against diseases that can be prevented by vaccination, it is crucial that they receive all of their immunizations at the proper ages and intervals (6). Similar to earlier research where the majority felt that immunizing children was essential for illness prevention, in several studies almost all participants disputed that vaccines are ineffective in preventing children from contracting diseases (7). According to a different survey, parents of kids between the ages of 0 and 6 were generally in agreement that immunizations shield their kids against sickness and that they run the risk of contracting it otherwise. Parents had a little disagreement on whether their child will experience a major adverse effect from receiving a vaccination. Mothers' illiteracy and lack of knowledge of the advantages of vaccine-preventable diseases may contribute to low immunization rates (8). This study aims to determine the perceived severity of children to vaccine preventable disease and the perceived benefits of immunization uptake of children among adults in selected communities of Ido Local Government Area on childhood routine immunization.

2. METHODOLOGY

Study Design

The study used a mixed-method (i.e. quantitative and qualitative) descriptive approach.

Study setting

Some settlements in Ido local government area in Ibadan, Oyo State, were the subject of the study. Ido is a rural local government area (LGA) in the lesser city of Ibadan, Oyo State. Its corporate offices are located in Ido.

Target population

This comprised of adults or caregivers who had children under the age of two and residents of the chosen communities in Ido Local Government Area (LGA), Ibadan, Oyo State.

Study population

Adults or caregivers of children under the age of two and residents of the chosen communities make up the study population (n=417).

Inclusion Criteria

Adults or caregivers of under-two living in the communities for at least 6 months.
Community members above 18 years living in the selected communities for at least 6 months

Exclusion Criteria

Adults or caregivers of under-two years who were sick and unfit at the time of study. Adults or caregivers of under-two years who were unwilling to participate during the data collection exercise.

Sample size determination

Using the Leslie Kish method, the sample size for this study was established, and it was found that target level of dependability should not exceed 0.05 with a 95% confidence range. Using proportion of children who had taken all basic vaccinations in Ido Local Government Area of 41.3% by Fatiregun et al., (9) cited by Adedire et al., (10).

Using the Leslie Kish formula, the sample size for this investigation was established.

$$n = \frac{Z^2 pq}{d^2}$$

Where, n = Sample size

Z = Standard normal deviation; 1.96

p = Prevalence of women who received post-natal care (Fatiregun et al.,

2013)(9)

q = 1 – Prevalence d = Precision; 0.05

z = 1.96; p = 41.3% i.e., 41.3/100 = 0. 413

q = 1 – 0. 413 = 0.587

d² = 0.05 x 0.05 = 0.0025

$$= \frac{1.96 \times 1.96 \times 0.413 \times 0.587}{0.0025} = \frac{0.9313}{0.0025} = 372.5$$

10% non-response rate = 372.5/ (1-0.10) = 414 (approximately)

Hence, after accounting for the 10% attrition rate or non-response rate, 414 respondents were chosen to take part in the study.

Sampling Technique

For this investigation of 10 wards, a multi-stage sampling technique was used.

Stage 1: Out of ten (10) wards in the Ido Local Government Area, five (5) were chosen using simple random sampling technique.

Stage 2: The community's homes were counted, and homes with children under two were chosen using systematic sampling technique.

Stage 3: In cases where there was more than one qualified respondent in the household, balloting was used to choose one of them. Respondents were chosen from each household.

Validity of instruments

The researcher made sure that each question on the questionnaire correlated with the study's objectives in order to determine the instrument's content validity. The supervisor received a draft copy of the questionnaire, which she modified as needed. In order to

improve the instrument's content validity, expert recommendations were integrated into the questionnaire's final draft.

Reliability of the Instruments

In order to evaluate the reliability of the instrument, 10% of the sample size (41 questionnaires) were administered in Oluyole Local Government Area of Ibadan, a similar population group. This setting was used for the reliability test because it shares traits with the setting used for the primary study population. Before the last version of the instrument was used, any necessary adjustments were performed. Forty-one adults were chosen at random. Using the Cronbach Alpha test, the reliability of the questionnaire was assessed from the pilot study and it was found to be reliable with a reliability coefficient of 0.850.

Method of Data Analysis

Quantitative data Analysis

All completed questionnaires' consistency and accuracy were checked. The surveys were manually sorted and then entered into the Statistical Program for the Social Sciences (SPSS) version 25. Descriptive statistics were used to assess the specific objectives. Pearson chi-square was used to determine the statistical significance of the hypotheses, and logistic regression was then used to forecast for the variable with the most impact.

Qualitative data Analysis

Interviews were verbatim transcribed. To guarantee that the transcription was accurate and thorough, individual transcripts were checked. A computer program for qualitative data analysis was used to sort, categorize, and analyze the data (Nvivo). This included multiple overall surface readings of transcripts to capture context and meaning, followed by coding and categorization of recurring concepts/ideas. A master list of all categories was assembled and examined for common themes.

Ethical Consideration

Before the study began, the study proposal was examined and authorized by the Oyo State Ministry of Health's Ethical Review Committee in Ibadan (AD 13/479/44248B).

3. RESULTS AND DISCUSSION

Socio-demographic characteristics of respondents

Three-hundred and sixty-two (86.8%) of the respondents practice monogamy, 335(80.3%) were married, 208(49.9%) had secondary education, 368(88.2%) are Yorubas, and 274(65.7%) were Muslims (Table 1). According to the study's findings, 376 (90.2%) of the respondents were females (Figure 1).

Table 1: Socio-demographic characteristics of respondents

Variables	N	(%)
*Age (years)		
15-30	108	25.9
31-45	204	48.9
46 and above	104	24.9
Marital Status		
Single	77	18.5
Married	335	80.3
Separated	4	1.0
Divorced	1	0.2
Ethnicity		
Yoruba	368	88.2

Igbo	45	10.8
Hausa	2	0.5
Others	2	0.5
Type of family		
Monogamous	362	86.8
Polygamous	55	13.2
Religion		
Christianity	274	65.7
Islamic	140	33.6
Traditional	3	0.7
Educational Background		
No formal education	28	6.7
Primary	43	10.3
Secondary	208	49.9
Tertiary	138	33.1
Occupation		
Unemployed	36	8.6
Trading	189	45.5
Artisan	68	16.3
Civil servant	19	4.6
Self-employed	105	25.2
Level of Income (#)		
0 – 200,000	409	98.1
201,000 – 400,000	4	1
401,000 – 1,000,000	4	1
Years of marriage		
0 – 20	319	76.5
21 – 30	75	18
31 – 70	23	5.5
Number of children		
0 - 3	105	25.2
4 – 6	201	48.2
7 and above	111	26.2
Sex of last child		
Male	41	9.8
Female	376	90.2
Child's age (years)		
0 – 10	208	49.9
11 – 20	137	32.9
21 – 30	72	17.3

*Mean Age 38.77 \pm 10.8 years




Figure 1: Respondents' Gender

UNDER PEER REVIEW

Perceived severity of vaccine preventable diseases

The level of perceived severity to vaccine preventable diseases was measured on a 36-point perceived severity scale. For each statement, the appropriate response was given a score of strongly agree - 4, agree -3, undecided - 2, disagree - 1, strongly disagree- 0. Perceived severity (PS) of <19 was rated as low perceived severity, PS of 19 – 36 was rated as high perceived severity. The perceived severity of vaccine preventable diseases among the respondents was showed in Table 2. Two-hundred and nine (50.1%) of the respondents strongly agreed that children may be paralyzed if not given oral polio vaccine, 217 (60%) strongly agreed that children may have diphtheria, whooping cough, tetanus or hepatitis B if not given pentavalent vaccine, 214 (51.3%) strongly agreed that children may be seriously sick if vaccine is not accepted, and 214 (51.3%) strongly agreed that disease may later manifest in worse form in adulthood if vaccination is not taken (Table 2). The study's findings showed that 396 (95.0%) of the respondents had a high level of perceived severity of diseases that could have been prevented by vaccination (Figure 2).

Table 2: Perceived severity of vaccine preventable diseases

Statements	Strongly Agreed (%)	Agreed (%)	Strongly disagreed (%)	Disagreed (%)	Undecided (%)
Children may be paralyzed if not given oral polio vaccine	209(50.1)	180(43.1)	6(1.4)	16(3.8)	6(1.4)
Children may have diphtheria, whooping cough, tetanus or hepatitis B if not given pentavalent vaccine	217(60.0)	185(39.6)	4(1.9)	13(3.1)	14(3.4)
Children may have eye deficiency if not given Vitamin A Supplement	219(52.5)	162(38.8)	6(1.4)	14(3.4)	16(3.8)
Children may have small greyish-white spots in the mouth, aches and pains if not given measles vaccine	209(50.1)	161(38.6)	13(3.1)	18(4.3)	16(3.8)
Children may have aching muscles, particularly the back and knees if not given yellow fever vaccine	208(49.9)	156(37.4)	12(2.9)	17(4.1)	27(5.8)
Children may have blood and lung infections and brain damage if not given PCV	193(46.3)	165(39.6)	13(3.1)	18(4.3)	28(6.7)
Children may have bloody cough and /or permanent brain damage if not administered BCG	197(47.2)	173(41.5)	7(1.7)	15(3.6)	25(6.0)
Children may be seriously sick if vaccine is not accepted	214(51.3)	166(39.8)	7(1.7)	22(5.3)	8(1.9)
Diseases may later manifest in worse form in adulthood if vaccination is not taken	214(51.3)	159(38.1)	6(1.4)	24(5.8)	14(3.4)




Figure 2: Level of perceived severity of vaccine preventable diseases

Perceived benefits of immunization uptake

The level of benefits of immunization uptake was measured on a 36-point perceived benefit of immunization uptake scale. For each statement, the appropriate response was given a score of strongly agree- 4, agree -3, undecided - 2, disagree - 1, strongly disagree - 0. Perceived benefits (PB) of <19 was rated as low perceived benefits, PB of 19 – 36 was rated as high perceived benefits. The perceived benefits of immunization uptake among the respondents was showed in Table 3. Two-hundred and sixty one (62.6%) of the respondents strongly agreed that vaccinations may prevent diseases in children, 242 (58.0%) strongly agreed that vaccines may help to prevent paralysis of children, 224 (53.8%) strongly agreed that vaccines may prevents brain damage and increase child cognitive functions, and 214 (51.3%) strongly agreed that vaccines may help prevent lung infection, other details are presented in Table 3. The study's findings showed that 409 (98.1%) of the respondents had high level of perceived benefits of immunization uptake (Figure 3) as above.

Table 3: Respondents' perceived benefits of immunization uptake

Statements	Strongly Agreed (%)	Agreed (%)	Strongly Disagreed (%)	Disagreed (%)	Undecided (%)
Vaccinations may prevent diseases in children	261(62.6)	149(35.7)	1(0.2)	5(1.2)	1(0.2)
Vaccinations may strengthen the baby immune system	252(60.0)	151(36.0)	6(1.4)	8(1.9)	0(0.0)
Vaccines may help to prevent paralysis of children	242(58.0)	164(39.3)	3(0.7)	7(1.7)	1(0.2)
It prevent illness later on in life	236(56.6)	167(40.0)	5(1.2)	6(1.4)	3(0.7)

Vaccines may prevent brain damage and increase child cognitive functions	224(53.8)	64(39.4)	6(1.4)	11(2.6)	11(2.6)
Vaccine may help child to grow and develop well	237(56.8)	169(40.5)	4(1.0)	7(1.7)	0(0.0)
Vaccine may help prevent lung infection	214(51.3)	180(43.2)	6(1.4)	4(1.0)	13(3.1)
Vaccines may help to improve the sight of children	213(51.1)	176(42.2)	6(1.4)	12(2.9)	10(2.4)
The risk of spending money on hospital may be reduced	233(55.9)	163(39.1)	5(1.2)	13(3.1)	3(0.7)




Figure 3: Level of perceived benefits of immunization uptake

Results of In-Depth Interview

Fifteen (15) participants were interviewed. The interviews were completed with participants from five communities in Ido Local Government Area, Oyo State. The age range of the participants was 26 to 44 years. Their educational status ranged from no formal education to tertiary education, and majority had secondary education as their highest educational qualifications. All the IDI participants were married, and over three-quarter were Yoruba. The majority had four children; and were engaged with trading as their occupation. They were Muslims and Christians.

Perceived severity of vaccine preventable diseases

Majority 13(86.7%) of the 15 IDI respondents indicated that the severity of these diseases can be serious and hard on the parents as well. Respondents mentioned different severity to mentioned diseases, among which are shortage of blood for child suffering from malaria, loss of strength and stamina in case of diarrhea and high temperature, disability and paralysis for measles affected children. One of the respondents said;

“The stooling, immediately the child will not have strength, and if it is measles the body will be hot and also if there is catarrh, the parent will not be able to sleep self because the child we be disturbed”(NG, 33 years, primary education)

Benefits of immunization uptake

Prevention of sicknesses and diseases: Almost all 14(93.3%) the 15 IDI respondents also reported that the benefits of routine immunization is that it prevents sickness and diseases like measles, whooping cough, pertussis and paralysis in children. One of the respondents said;

"The benefits there are, for instance, if they get measles vaccine, we were made to understand that it prevents measles for the child. So, if an adult that has measles carries the child, the child may not be easily infected because of the immunization received beforehand."(RS, 29 years, tertiary education)

Another respondent said;

"The benefits there are that, even if a child should want to get sick, once the immunization is already in their body, it will reduce the extent of the disease in their body. When children fall sick, an immunized child will respond to that sickness differently than one that has not been immunized."(BA, 39 years, primary education)

Majority 14(93.3%) of the 15 respondents agreed that the immunization is very beneficial to health as it improves children health, prevents sickness and diseases. One of the respondents said;

"The benefits are much. If the child collected all the immunization injection completed, there is nothing like for the child to be feeling hot temperature when the child wants to grow teeth, the body will not be hot unnecessarily. There will be no form of sickness that is peculiar to a child that will be affecting them"(AK, 36 years, SSCE)

Two (13.3%) of the 15 respondents responded that there might be a negative effect on a child immune system if the child did not take the multiple vaccines given during routine immunization as the child might be prone to different disease and sickness. One of the respondents specifically said;

"If the child did not take the injection, it will have effect on the child, the child might be sick at any time because there is no resistance to stop the disease from entering the body"(OM, 43 years, primary education)

Almost all 14(93.3%) of the 15 respondents said their opinion as regards the multiple vaccine collected by the children during routine immunization is good as it prevents all sort of sickness in children. One of the respondents specifically said;

"I've said it earlier, that immunization is very good for children. And it does not cause any harm. There is nothing else to say other than to advise whoever has not vaccinated their children to go and do so because it does a lot of good work in the body. It fights all sorts of diseases in the body. Maybe there is any disease that wants to show up in the body of the child, the immunization will fight it quickly before it becomes obvious in the child or even affects the mother of the child herself." (OE, 36 years, no formal education)

Almost half 7(46.7%) of the 15 respondents said they are of the opinion that multiple vaccines given to children during immunization prevents future development of chronic diseases and sickness. One of the respondents specifically said;

"In my opinion, when it is time for a child to get multiple injections, one will have to take the child to get it, regardless of the pain it may cause. One must not use the temporary pain and discomfort caused by the injections as a reason to not get the immunization. After all, the immunization will benefit the future of the child. And it is good, I don't think it has any negative effect on their future."(RL, 32 years, SSCE)

Cognitive function: Two other respondents (13.3%) added that not only is immunization good for children health, but it's also beneficial to their academic pursuits. One of the respondents specifically said;

"The benefits of immunization, according to what we were told, are that firstly, it is good for children. Secondly, if they want to travel out or gain admission into schools, it is needed. Lastly, it is very good for their health."(BA, 39 years, Primary education)

Two (13.3%) of the 15 respondents reported that going for multiple vaccines given during routine immunization gives a child good brain and also makes a child sharp and fervent. One of the respondents said;

"It is because immunization prevents future diseases and also makes them smart is why we let them take it."(AK, 36 years, SSCE)

Additionally, one (6.7%) of the 15 respondents reported that immunization vaccination makes a child's brain sharp, and makes them smart in their education. Some 10(66.7%) of the 15 respondents reported that it is educative for the parents as they were also given health talk on how to care for their children during routine immunization. One of the respondents said;

"The benefits there are many. When we go for immunizations, we are usually given health talks, and awareness, and even fun activities. From there, we usually even get other benefits that we receive for free from there."(OM, 43 years, Primary education)

However, a respondent had a negative opinion on of the multiple vaccines given as she said some people believed it affect the children in future (RS, 29 years, tertiary education).

4. DISCUSSION

Evidently from the present study, over two-third of the respondents had high level of perceived severity of. This might be attributed to the fact that infectious diseases such as measles, tetanus, mumps vaccine preventable diseases, rubella, etc were a greater risk for children, so if vaccines were not given to these children, it could result to them being infected by these diseases (11). Finding from this study revealed that children may be paralyzed if not given oral polio vaccine; children may have diphtheria, whooping cough, tetanus and hepatitis B if not given pentavalent vaccine; children may have eye deficiency if not given Vitamin A Supplement; children may have small greyish-white spots in the mouth, aches and pains if not given measles vaccine; children may have aching muscles, particularly the back and knees if not given yellow fever vaccine; children may have blood and lung infections and brain damage if not given PCV; children may have bloody cough and /or permanent brain damage if not administered BCG; children may be seriously sick if vaccine is not accepted; and diseases may later manifest in worse form in adulthood if vaccination is not taken. This finding is similar to that of the IDI in which a respondent added that immunization was given to a child at one month, three months, six months, nine months, and one year. In addition to this, one of the respondents specifically said; *"What I know about it is that it is good for children, right from birth, or by the 8th day. It causes the child not to fall sick. When a child does not get immunized, it can make the child to be coughing, or have many other diseases. It is sha very good for children to be immunized"* This findings concurs with that of Enwonwu, et al. (12) in which it was reported that that vaccines act by stimulating the body's immune system to protect the person against subsequent infections or diseases. Vaccines can

protect more children than most other strategies (12). According to Matta et al. (13), high immunization uptake is critical for protecting people from infectious diseases, but barriers to immunization are complex. Ashbaugh and Brooke (14) highlighted that the benefits from vaccines and vaccination programmes go beyond preventing infections. One of the best and most affordable public health interventions for illness prevention, according to Mantel and Cherian (1), is vaccination and immunization. Also, vaccines have substantially improved health and reduced mortality. World Health Organization (2) reported that the universal BCG vaccination at birth is recommended in countries or settings with a high incidence of TB and/or high leprosy burden. Finding from a study conducted by Williams et al. (15) revealed that less than a quarter of the respondents correctly stated the purpose of immunization which is to prevent infectious disease; while finding from Mvundura et al (16) showed that majority of the respondents mentioned the purpose of immunization correctly. Finding of Holipah and Kuroda (2018)(17) also showed that the majority of them knew that routine vaccination prevents children from some serious infectious diseases and its complication. Habib et al., (18) also reported that more than 85% of the participants knew the role of childhood. Finding from the study revealed that vast majority of the respondents have high level of perceived benefits of immunization uptake. This corroborates the finding of Stone Jr. et al. (19) who reported that taking children for immunization and vaccination uptake provide a golden opportunity to curb and prevent infectious diseases such as diphtheria, pertussis and tetanus.

The study revealed that the perceived benefits of immunization uptake of children include prevention of diseases; strengthen the baby immune system; prevent paralysis of children; prevent illness later on in life; prevention of brain damage and increase child cognitive functions; help child to grow and develop well; help prevent lung infection; help to improve the sight of children; and the risk of spending money on hospital may be reduced. Finding in the IDI agrees with this as significant percentage of respondents said; *“What I know about it is that it is good for children, right from birth, or by the 8th day. It causes the child not to fall sick. When a child does not get immunized, it can make the child to be coughing, or have many other diseases. It is sha very good for children to be immunized.”* These findings corroborate that of Matta et al., (13), who reported that high immunization uptake is critical for protecting people from infectious diseases. Hayden (2014) cited by Kazi et al. (4) observed that early immunization would prevent illness later on in life, which would lead to loss in individual productivity and result in additional health care costs. Besides, immunization will strengthen individual's immune system. Ashbaugh and Brooke (14) highlighted that the benefits from vaccines and vaccination programmes go beyond preventing infections. Mantel, & Cherian, (2020) (1) explained that one of the most effective and affordable public health measures for illness prevention is vaccination and immunization. Also, vaccines have substantially improved health and reduced mortality. Kaufman et al. (6) reiterated that it is of importance that a child should receive all immunization at the appropriate ages and intervals in order to ensure maximal protection from vaccine preventable diseases. Enwonwu et al. (12) submitted that vaccines act by stimulating the body's immune system to protect the person against subsequent infections or diseases. Vaccines can protect more children than most other strategies (12). Kagoné et al. (8) also supported the finding of this present study as the researchers revealed that on general, parents of children aged 0 to 6 agreed that immunizations protect their children from disease and that, in the absence of immunizations, their children may contract a disease. The present study also revealed that vaccinations may prevent diseases in children. The IDI also supported this finding a respondent specifically said; *“What I know about it is that it is good for children, right from birth, or by the 8th day. It causes the child not to fall sick. When a child does not get immunized, it can make the child to be coughing, or have many other diseases. It is sha very good for children to be immunized.”* This finding corresponds with that Matta et al. (13), who reported that high immunization uptake is critical for protecting people from

infectious diseases. Finding from a similar study conducted by Mantel and Cherian, (1) revealed that one of the most effective and affordable public health measures for illness prevention is vaccination and immunization.

The study also revealed that vaccine may help child to grow and develop well. Finding from the IDI also supported this finding as majority of the respondents acknowledged that their opinion as regards the multiple vaccine collected by the children during routine immunization is good as it prevents all sort of sickness in children. This agrees with the finding of Mohapatra et al. (20) in it was reported that few caregivers felt that routine immunization improves the growth and intelligence of children. Mantel and Cherian (1) reported that vaccines have substantially improved health and reduced mortality.

Limitation to study

Only individuals in Ido local government area of Oyo State were used for the study thus the result cannot be generalized. Small sample size of four-hundred and seventeen (417) was used in carrying out this research because only a single Local Government Area was used, hence the result cannot be generalized. It was not easy obtaining the ethical approval from the Oyo State Ethics committee. Therefore, the researcher was constrained to wait and consistently get in touch with workers in the committee's office until final approval was secured.

CONCLUSION

This study looked at how residents (adults) in a few communities in Ido local government area, Ibadan, Oyo State, felt about and accepted routine childhood immunization. In conclusion, the qualitative finding has corroborated with the quantitative finding. More so, respondents' level of education was significantly associated with knowledge on childhood routine immunization ($p < 0.05$). Therefore, maternal education should be improved since it is important for increasing childhood immunization and vaccination uptake.

Recommendations

Since immunization is beneficial to children all stakeholders should work together in order to increase vaccination rates. Also, public health administrators and vaccination providers (paediatricians, nurses, health visitors etc) should work together to ensure children receive all vaccinations at the appropriate times and intervals as specified by the national vaccination schedule.

CONSENT

All authors declare that 'written informed consent was obtained from the patient (or other approved parties) for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editorial office/Chief Editor/Editorial Board members of this journal

ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki

REFERENCES

1. Mantel C, Cherian T. New immunization strategies: adapting to global challenges. *Bundesgesundheitsblatt-Gesundheitsforschung-Gesundheitsschutz*.2020;Jan;63(1):25-31.
2. World Health Organization. Immunization coverage: Fact sheet. Geneva: WHO; 2018. Available at <http://www.who.int/mediacentre/factsheets/fs378/en/>
3. World Health Organization. World health statistics 2023: monitoring health for the SDGs, sustainable development goals. World Health Organization; 2023 May 19.
4. Kazi AM, Ahsan N, Mughis W, Jamal S, Allana R, Raza M, Muneer S, Mughal MA, Kaleemuddin H, Sameen F, Ahmed RM. Usability and acceptability of a mobile app for behavior change and to improve immunization coverage among children in pakistan: A mixed-methods study. *International Journal of Environmental Research and Public Health*. 2021 Sep 9;18(18):9527.
5. Butt M, Mohammed R, Butt E, Butt S, Xiang J. Why have immunization efforts in Pakistan failed to achieve global standards of vaccination uptake and infectious disease control?. *Risk management and healthcare policy*. 2020 Feb 12:111-24.
6. Kaufman J, Tuckerman J, Bonner C, Durrheim DN, Costa D, Trevena L, Thomas S, Danchin M. Parent-level barriers to uptake of childhood vaccination: a global overview of systematic reviews. *BMJ global health*. 2021 Sep 1;6(9):e006860. <https://doi.org/10.2147/RMHP.S211170> *health*, 6(9), e006860.
7. Adefolalu OA, Kanma-Okafor OJ, Balogun MR. Maternal knowledge, attitude and compliance regarding immunisation of under-five children in primary health care centers in Ikorodu local government area, Lagos state. <https://doi.org/10.4103/jcls.jcls>
8. Kagoné M, Yé M, Nébié E, Sié A, Müller O, Beiersmann C. Community perception regarding childhood vaccinations and its implications for effectiveness: a qualitative study in rural Burkina Faso. *BMC public health*. 2018 Dec;18:1-0.
9. Fatiregun AA, Adebowale AS, Ayoka RO, Fagbamigbe AF. Assessing full immunisation coverage using lot quality assurance sampling in urban and rural districts of southwest Nigeria. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 2013 Nov 1;107(11):731-40.

10. Adedire EB, Ajayi I, Fawole OI, Ajumobi O, Kasasa S, Wasswa P, Nguku P. Immunisation coverage and its determinants among children aged 12-23 months in Atakumosa-west district, Osun State Nigeria: a cross-sectional study. *BMC public health*. 2016 Dec;16:1-8.
11. Betsch C, Rossmann C, Pletz MW, Vollmar HC, Freytag A, Wichmann O, Hanke R, Hanke W, Heinemeier D, Schmid P, Eitze S. Increasing influenza and pneumococcal vaccine uptake in the elderly: study protocol for the multi-methods prospective intervention study Vaccination60+. *BMC Public Health*. 2018 Dec;18(1):1-3.
12. Enwonwu KG, Ilika A, Ifeadike C, Aniemena C, Egeonu RO. Perception of childhood immunization among mothers of under-five children in Onitsha, Anambra State. *Afrimed Journal*. 2018 Apr 24;6(1):59-64.
13. Matta P, El Mouallem R, Akel M, Hallit S, Fadous Khalife MC. Parents' knowledge, attitude and practice towards children's vaccination in Lebanon: role of the parent-physician communication. *BMC Public Health*. 2020 Dec;20:1-9.
14. Chevalier-Cottin EP, Ashbaugh H, Brooke N, Gavazzi G, Santillana M, Burlet N, Tin TinHtar M. Communicating benefits from vaccines beyond preventing infectious diseases. *Infectious Diseases and Therapy*. 2020 Sep;9:467-80. <https://doi.org/10.1007/s40121-020-00312-7>
15. Williams S, Endacott I, Ekiri AB, Kichuki M, Dineva M, Galipo E, Alexeenko V, Alafiatayo R, Mijten E, Varga G, Cook AJ. Barriers to vaccine use in small ruminants and poultry in Tanzania. *Onderstepoort Journal of Veterinary Research*. 2022;89(1):1-1.
16. Mvundura M, Frivold C, Osborne AJ, Soni P, Robertson J, Kumar S, Anena J, Gueye A, Menozzi-Arnaud M, Giersing B, Kahn AL. Vaccine innovation prioritisation strategy: Findings from three country-stakeholder consultations on vaccine product innovations. *Vaccine*. 2021 Dec 3;39(49):7195-207.
17. Holipah, Maharani A, Kuroda Y. Determinants of immunization status among 12-to 23-month-old children in Indonesia (2008–2013): a multilevel analysis. *BMC public health*. 2018 Dec;18:1-1.
18. Saleh A, Alrashidi AA, Bukhari MA, Habib RF, Alsubhi RA, Saadawi DW, Hatim RF. Assessment of knowledge, attitude and practice of parents towards immunization of children in Saudi Arabia, 2018. *The Egyptian Journal of Hospital Medicine*. 2018 Apr 1;71(2):2585-9.
19. Stone Jr CA, Rukasin CR, Beachkofsky TM, Phillips EJ. Immune-mediated adverse reactions to vaccines. *British journal of clinical pharmacology*. 2019 Dec;85(12):2694-706
20. Mohapatra I, Kumar A, Mishra K. A study on awareness and utilization of Mission Indradhanush in an urban slum of Bhubaneswar. *Journal of Family Medicine and Primary Care*. 2018 Nov;7(6):1294.
21. Almutairi WM, Alsharif F, Khamis F, Sallam LA, Sharif L, Alsufyani A, Alshulah FN, Alqasimi R. Assessment of mothers' knowledge, attitudes, and practices regarding childhood vaccination during the first five years of life in Saudi Arabia. *Nursing Reports*. 2021 Jul 5;11(3):506-16. <https://doi.org/10.3390/nursrep11030047>