

# Original Research Article

## Determine the socio-economic and demographic characteristics of caregivers of children aged 6-59 months in Kericho County, Kenya.

---

### ABSTRACT

Child growth and well-being are critically dependent on proper nutrition, particularly during the developmental phase between the ages of 6 and 59 months. Poor nutritional status is associated with high proportion of mortality and morbidity in Kenya and globally. Malnutrition during this critical window can predispose children to cognitive impairments, infectious diseases, and an increased risk of chronic conditions, contributing to high mortality rates. Despite the severity of the problem, few studies have focused on the factors affecting the nutritional status of children in this age group in Kenya. This study aimed to identify the factors influencing the nutritional status of children aged 6 months to 5 years in Kericho County, Kenya. Specifically, the study sought to assess the socio-economic and demographic characteristics of caregivers. The research employed a cross-sectional analytical design, targeting children attending outpatient child welfare clinics at Kericho County Referral Hospital and Kapkatet Sub-County Hospital. A total of 172 children were selected through systematic random sampling, and data was collected using a structured questionnaire. Anthropometric data were analyzed using ENA for SMART, and all other variables were processed using the Statistical Package for Social Sciences (SPSS) version 27. Descriptive statistics were used to outline the population characteristics, while regression analysis was employed to identify predictors of nutritional status, with a confidence interval of 95% and significance set at  $p \leq 0.05$ . Key findings revealed that stunting affected 25% of the children, while wasting and underweight were prevalent in 23.8%. Dietary practices showed that only 39.2% of children aged 6 to 23 months met the minimum dietary diversity, and 39.2% received a minimum acceptable diet. Moreover, 45.3% of children experienced illness in the past month. Regression analysis identified several key predictors of malnutrition: children of caregivers without formal education were more likely to be underweight (AOR = 0.70,  $p = 0.01$ ). Evidence from this study highlights the critical role of caregiver education and how this influenced child nutritional outcomes. These findings underscore the need for targeted interventions that enhance caregiver education.

**Keywords** *Dietary practice*: refers to the reported behaviours, actions, or individual choices regarding dietary habit such as meal frequency and dietary diversity.

**Morbidity status**: is the state of having illnesses or symptoms of illnesses.

**Nutritional status:** the health condition of children, influenced by the intake and utilization of nutrients, often assessed through anthropometric indicators such as height-for-age, weight-for-age, and weight-for-height.

**Stunting:** a condition characterized by a child's height-for-age falling below a specified standard, typically expressed as a Z-score, indicating a level of growth retardation.

**Underweight:** to a child's weight-for-age falling below a specified standard, usually expressed as a Z-score, suggesting a deficiency in overall body mass relative to age.

**Wasting:** a condition marked by a child's weight-for-height falling below a specified standard, often expressed as a Z-score.

## INTRODUCTION

According to the World Health Organization (2017), our nutritional status is determined by the nutrient composition of the food we consume, our nutritional demands, and our body's capacity to digest, absorb, and utilize those nutrients (1). Children's nutritional health is especially important between the ages of 6 and 59 months since that's when they're developing and growing the most. Indirectly and directly, the growth of a nation is impacted by the nutritional well-being of its children since it shows that the government, communities, and households are investing in the health of their families (2). In the context of early childhood, chronic and frequent malnutrition can have severe and prolonged negative effects on the mental growth and development of young children (2). Malnutrition can also predispose such children to a number of cognitive impairments and infectious diseases. It has also been linked with chronic diseases, exacerbating high mortality and morbidity rates among young children (2). Regionally, it has been established that East African countries take lead in the prevalence of poor nutritional status among six- to fifty-nine-month-old children globally (3). Kenya had a crude mortality rate of 0.8 per 10,000 children per day between 2013 and 2017, with more than 66% of these deaths being attributed to drought (4). Undernutrition remains a key public health issue in less developed countries like Kenya because of poor sanitation, household food insecurity, infections like diarrhoea, and poor dietary consumption.

According to KDHS 2022, in Kenya, the frequency of stunting, wasting, and underweight was 18%, 5%, and 10%, respectively (5). Multiple international and local studies have examined the nutritional condition of children between the ages of six and fifty-nine months (6, 4,7,8, 9, and 2). However, little has been reported on the determinants of the under-five children's nutrition status. This provides a need to conduct this study to identify these determinants that play a vital role in determining below five children's nutrition status. Hence, the determinants of the nutritional condition of children in Kericho County, Kenya, ranging from 6 to 59 months old, were identified in the research.

## **Problem statement**

. According to recent research by the United Nations Children's Fund (UNICEF), more than 25% of children in Kenya below the age of five have stunted development (2). Severe wasting and wasting have been closely linked to preventable child mortality. Gudu et al. (2020) reported that approximately 35,000 Kenyan children aged below five years die annually due to undernutrition (10).

Deficiency in essential macronutrients like fat, carbohydrates, and protein can trigger protein-energy malnutrition (PEM) (11). A combination of macronutrient and micronutrient deficiencies results in severe health problems for young children (12). Malnourished young children, particularly those suffering from severe malnutrition, are at elevated risk of mortality from illnesses like malaria, pneumonia, and diarrhoea. While poor nutrition status is a leading attributor to diseases and mortality in children under five, limited research has explored the specific factors affecting the nutritional status of children aged six months to five years in Kenya. Moreover, few data exist about the predictors of nutritional outcomes for children in Kericho County. This research aims to fill this gap by identifying the determinants of the nutritional condition of children aged 6 to 59 months in Kericho County. The findings of this research will contribute new knowledge by uncovering context-specific factors influencing child nutrition, thereby informing targeted interventions for improving nutritional outcomes in this vulnerable age group.

## **Justification of the study**

The Big Four Agenda, which includes national health care, diet, and food security, is one of the main goals of the Kenyan government. The major goal of the study on the factors that affect children's nutritional status between the ages of 6 and 59 months in Kericho County, Kenya, is to improve the eating habits and nutritional status of babies, young children, and moms. The government in partnership with international players, who include UNICEF, and the implementation of nutritional services and strategies underscore the importance of addressing nutrition-related issues in the region. The mention of UNICEF's provision of counselling to 1.6 million caregivers on optimal feeding practices for young children and infants in 2020 further emphasizes the ongoing efforts to enhance nutritional outcomes, providing context for the study's examination of factors influencing the nutritional status of children in the specified age group in Kericho County. The research is in support of the Kenyan government's nutritional interventional programs by providing empirical evidence regarding the factors that contributed to malnutrition among children aged below 6-59 months in Kericho County.

While prior literature identified multiple risk factors for malnutrition, individual factors were varied from one country to another and with time. As such, the then characterization of such factors in other counties may have not provided an empirical basis for the nutritional intervention programs in Kericho County. The findings regarding the nutritional status of children aged 5-59 months in Kericho County will enable this county to address malnutrition which is a public health problem.

## **Study limitations and delimitations**

**Limitation 1:** The research was cross-sectional, conducted at a single point in time.

**Delimitation 1:** To overcome this, data was collected from a relatively large sample size to enhance representativeness.

**Limitation 2:** Availability of food due to production and supply at the time may have affected interpretation of participants' dietary characteristics.

**Delimitation 2:** The study accounted for this by employing rigorous data collection methods, considering temporal factors such as food production and supply.

**Limitation 3:** Dietary practices may have been influenced by prevailing high food prices and inflation.

**Delimitation 3:** The study acknowledged the impact of high food prices and inflation, incorporating sophisticated statistical analyses for nuanced insights.

## **Scope of the study**

The study scope was spelt out to setup the confines and parameters within which the research was done. It defined specific aspects, participants, and locations included in the investigation. In the study on the determinants of nutritional status in children aged 6 to 59 months in Kericho County, Kenya, the scope encompassed a defined geographical area, Kericho County. The age comprised 6 to 59 months specified as the target population. The study explored factors that influenced nutrition, including, socioeconomic conditions within this specified demographic. The scope set the context for the study's focus and the extent of its findings.

## **Socio-economic and demographic factors**

A study by Ahmad et al. (2020) sought to find out the outcome that socio-economic factors had on nutritional status amongst the under-fives in Punjab province, Pakistan. Stunted growth was statistically linked to sanitation facility, family size, age, wealth quintile, and maternal education. More male children were wasted and underweight as compared to the girls of the same age. (13). Additionally, wasting was mostly associated with food shortages and children or urban areas. Underweight was also associated with access to treated water and incomplete immunization. The researchers concluded that the increase in malnutrition cases among the under-fives was directly proportional to the population's lower socioeconomic status, especially those originating from rural areas. They suggested multiple policies to combat wasting, including the need for the government to take the necessary social, political, and economic policy measures to improve personal hygiene habits, increase nutritional access, promote healthcare education, and enhance economic activities.

Igbokwe et al. (2017) conducted research to know whether socio-demographic variables affected the nutritional quality of young children in Enugu, Nigeria. They sampled nine private primary schools and five public primary schools within a three-month period of the research using cross-sectional descriptive study design. The papers showed that among the research subjects 3.3%, 3.0%, and 0.8% were underweight wasted, and stunted

correspondingly (14). Comparing students of lower socioeconomic classes with the students of upper socioeconomic classes, the former was more wasted, underweight, and children, while the latter had higher levels of obesity and overweight (14). ; Similarly, in the same study, authors established that diet and feeding of the child were significantly influenced by the main source of income of the family, parents' level of education, sex of the child, and age of the child. Stunting and wasting were significantly higher among poor children in the public schools while obesity and overweight were higher among the rich children who attend private schools as found out by Igbokwe et al.

In their study, Chaudhary and Agrawal (2018) investigated the different dietary habits, patterns of malnutrition among children of six-59 months of age as well as the association between demographic variables and malnutrition in India. The research has reported somewhat higher incidences of stunting than the National Family Health Survey 4 at 43 % (15). That meant 56% of the research subjects were born underweight when recruited into the study. (15). The researchers also found that the people of India were not utilizing the Integrated Child Development Services ICDs in appropriate manner. They laid down recommendations that can help enhance delivery of Anganwadi services.

A hospital-based case-control study researched by Gudu et al. (2020) sought to unveil the components influencing poor nutrition among the under-fives in Western Kenya. These researchers found a number of factors that were associated with undernutrition, including pre-lacteal feeding, deworming, and failure of the mother to attend an antenatal clinic (10). Other factors that were found to be independently correlated with undernutrition included paternal lack of formal education, low birth weight, and delayed developmental milestones (10). The study recommendations included improvement of deworming programs, child feeding practices, and prenatal care in order to combat malnutrition in Western Kenya.

In order to establish the degree of relationship between demographic factors and malnutrition cases among children of one day to twenty-four months in Kwale County, Kenya, the study by Ndemwa et al. (2017) adopted a KWAME County population. In their findings, the sex, underweight, stunting, and age showed some variability they noted this was due to age. They found out that the children had higher rates of global acute malnutrition, being underweight and Growth stunting. The males presented larger numbers of stunted growths than the females in the research done. Not surprisingly, if the children grew stunted or underweight, there was a substantial increase for both growing with an age of the children (16).

Ole-Tankoi et al. (2016) identified antecedents of nutritional status of children aged 6-59 months in Trans- Mara East sub country. Data was gathered employing descriptive cross-sectional research design and a structured self-administered questionnaire. These findings detail indicate that poverty, mother's employment status as a homemaker, and the size of the household were the major determinants of stunting in children. When it came to the gender of the child, the likelihood of males being captured as being obese in comparison to the female children was higher. The study used the prevalence of nutrition and undernutrition to determine that poor nutritional status was twice higher in the country sides of Trans-Mara East sub-county (17).

If at all, there are very few studies that have been conducted in this field: that gap has been filled by our study, as indicated in the literature review above. Further, the research studies undertaken in Kenya appeared to have aimed at identifying the role of specified sociodemographic/socioeconomic predictors on the nutritional status of children in the age

group of six to fifty-nine months. The attempt of the study considers this gap by focusing on the level and magnitude of all the socio-economic and demographic factors on the nutritional status of these children.

## **MATERIAL AND METHODS.**

The research was done at Kericho County Referral Hospital (0.3713° S, 35.2801° E) and Kapkatet Sub-County Hospital (-0.629716, 35.197243). The two hospitals are situated in Kericho County in Kenya.

An institution-based cross-sectional analytical research design was used. The cross-sectional design allowed the researcher to examine data from both the children and the caregivers of 6-59-month-old children at one particular moment in time.

### **Independent variables**

Independent variables included socioeconomic and demographic traits such as farm ownership and size, profession, income, number of children, age of the main carer, and educational attainment. They also included nutritional habits including eating often, varying one's diet, and nursing.

The study included:

- i. Primary caregivers who encompassed any person who were responsible for the child's day-to-day wellbeing and care (17). Specifically, the primary caregivers included biological mothers, aunts, grandparents, and others who sufficed as primary caregivers for children whose biological parents were not available or deceased.
- ii. Only caregivers who gave their informed consent were included in the study.

### **Sampling technique and recruitment process**

The study used systematic random sampling method to get the participants from the target population. Systematic random sampling also involves making a sample from the already defined total population using an equal interval. In this case, the target population was composed of caregivers of children 6 to 59 months old who attended child welfare clinics at Kericho County Hospital and Kapkatet Sub-County Hospital, only on the study day. For this purpose, collaboration with the medical practitioners and clinic managers of the selected sites was initiated. Such collaborations also made it easy to access potential participants by arranging for all the necessary approvals and support from the clinic. Hypothesized mean number of caregivers with eligible children based on health records suggest they were about 40 caregivers per day accessing clinics. To invite participants in a more systematic manner, caregivers were chosen following every fourth consecutive criterion-based eligible caregiver rule, with the inclusion criteria being children aged between 6 to 59 months visiting the clinic with their caregiver for health check-ups or immunization sessions

### **Sample size**

The sample size was determined based on Fischer's formula (2003) as follows:

$$N = \frac{Z^2 Pq}{d^2}$$

Where, N denoted the desired sample size from a sample size of over 10,000

Z denoted the standard deviation at 1.96 as the required confidence level

P= the proportion within the target market that was estimated to be having the characteristic that is being measured. Kericho County prevalence of malnutrition =12.4% (18).

$$q = 1 - P$$

D = the set level of statistical confidence, which is 95% CI (0.005).

$$\text{Therefore: } N = \frac{(1.96^2 \times 0.124 \times (1 - 0.124))}{0.005^2}$$

Using this formula, the initial sample size determined was 167. To account for the possibility of attrition the sample size of the primary caregivers of children aged 6-59 years was increased by 3% (19). Thus, the total sample size determined was 172.

### **Data collection tools**

The researcher-administered structured questionnaire was employed in data collection of the primary caregivers' socio-demographic and economic characteristics.

### **3.9. Data collection tools pretesting**

Pretesting was conducted in Longisa County Referral hospital in Bomet County, Kenya among 17 participants which was about 10% of the sample size expected. Pretesting was done to assess the practical problems associated with the researcher-administered questionnaire. Longisa County Referral hospital had similar characteristics with the study hospitals making it suitable choice.

### **Data collection tool validity**

The data collection tool validity was checked by a panel of experts in the field.

### **Data collection tools reliability**

The tool reliability was determined through test-retest method. This was done by administering the tool to a small group of participants and then repeating the procedure five days later. A correlation coefficient of above 0.80 was considered acceptable (6), the correlation coefficient for the study tool was 0.67 and after validation it increased to 0.84.

### **Data collection procedures**

#### **Socio-economic and demographic data**

The demographic as well as the socio-economic variables of the study children's primary caregivers was assessed. This included size of farmland, ownership of livestock, occupation, education, income, family size, religion, ethnicity, marital status, and head of household. The child characteristics included sickness status, and type of birth, gestation age, place of delivery, birth order, sex, and age.

### **Data analysis and presentation**

The collected data was analysed with the help of ENA for SMART, statistical package for social sciences, that is, SPSS version 27 and Microsoft Excel 2010. All the other variables were assessed using Excel and SPSS. So in the analysis of data; we used the chi square test for bivariate testing and logistic regression analysis for multivariate testing, and descriptive statistics including percentage were also computed. Thus, in order to describe the respondents' characteristics, the results of descriptive analysis of the socio-demographic and economic parameters were considered. The percent score was also offered for the percent of respondents who attained a certain value on all of the evaluated factors. Each nutritional status indicator was ascertained from a percent analysis of nutritional status. The chi-square test was employed to test the relationship between nutritional status and socio-demographic and economic characteristics of caregivers. Between the nutritive condition of children and dietary behaviour of the care giver chi-square tests were used. In this study, children whose regression coefficients were statistically significant for the variables in question, between the ages of 6-59 months had their nutritional status assessed. Chi-square analysis was then followed by regression analysis on the variables that showed significance. With the view of controlling for such factors we also computed the Adjusted Chances Ratio (ACR) and the Crude Chances Ratio (CCR). Using a confidence interval of 95% the level of statistical significance was tested at  $p < 0.05$ .

## **RESULTS AND DISCUSSION**

### **Response rate**

Each of the participants identified and approached to participate in the study agreed to do so which gave a 100% response rate. By all participating increases the validity of the study results obtained in the investigation. Additionally, the fact that none declined to participate also means the results represent the whole sample the investigator was interested in studying.

### **Socio-demographic and economic characteristics**

Tables 1 and 2 display the participants' socio-demographic characteristics as they pertain to children and caregivers, whereas Table 4.3 displays their socio-economic features.

**Child specific socio-demographic characteristics**

As shown in table 2, 75 (43.6%) of the children were aged 24-59 years, followed by 55 (32%) who were aged 9 to 23 months and 42 (24.4%) who had an age range of 6 to 8 months. Additionally, 88 (51.2%) of the children were males and 84 (48.8%) were females. The majority of the children (n = 163, 94.8%) were given birth at the health facility. The mode of delivery for about three-quarters of the children (n = 131, 76.2%) was through the vagina.

**Table 2. Child specific socio-demographic characteristics**

Source: Field data, 2024

Variable	N (172)	% (100)
Age of the child		
6-8	42	24.4
9-23	55	32
4-59	75	43.6
Sex of the child		
Male	88	51.2
Female	84	48.8
Place of childbirth		
At home	9	5.2
Health facility	163	94.8
Mode of delivery		
CS	41	23.8
Vaginal	131	76.2

**Caregiver-specific socio-demographic characteristics**

As shown in Table 3, the age range of the respondents was 14 to 47 years, with 163 (92.4%) of them aged 20 to 47 years. The majority of the caregivers practiced Christianity (n = 169, 98.3%) and were either married or cohabiting (63.4%). Households were majorly headed by males (102, 59.3%), and about an equal proportion of the households had 1 to 3 members (n = 80, 46.5%) and 4 to 6 members (n = 79, 45.9%). Almost equal proportion of the household size was observed for households with members ranging from 1 to 3 (n = 80, 46.5%) and 4 to 6 (n = 79, 45.9%). Additionally, 47 (27.3%) of the caregivers used alcohol.

**Table 3. Caregiver-specific socio-demographic characteristics**

Source: Field data, 2024

<b>Variable</b>	<b>N (172)</b>	<b>% (100)</b>
Age of the caregiver		
14 to 19 years	13	7.6
20 to 47 years	159	92.4
Marital status		
Never married	51	29.7
Married or cohabiting	109	63.4
Separated/divorced/widowed	12	7
Religion		
Christian	169	98.3
Muslim	2	1.2
None	1	0.6
Household head		
Female	70	40.7
Male	102	59.3
Caregivers use alcohol		
Yes	47	27.3
No	125	72.7
Household size		
1-3	80	46.5
4-6	79	45.9
7-9	13	7.6

**Socio-economic characteristics**

The results on the socio-economic characteristics of the caregivers are shown in Table 4. The findings revealed that about two-thirds of the caregivers had secondary (n = 77, 44.8%) and tertiary education (n = 41, 23.8%). Casual labor, with a proportion of 46 (26.7%) of the respondents, followed by 40 (23.3%) of respondents on salaried employment and 35 (23.3%) of respondents on farming, was the leading source of income for the caregivers. Over half of the caregivers (n = 103, 59.9%) owned land, but only a third (n = 58, 33.7%) had a kitchen garden. Though the majority of the caregivers (n = 85, 49.4%) served food first to a child, they served the prime portion to the father (50%) and mother (30.8%), respectively.

UNDER PEER REVIEW

**Table 4. Socio-economic characteristics**

Source: Field data, 2024

Variable	N (172)	% (100)
Educational status		
No formal education	11	6.4
Primary	43	25
Secondary	77	44.8
Tertiary	41	23.8
Occupation		
None	19	11
Casual laborer	46	26.7
Farmer	35	20.3
Salaried employed	40	23.3
Self employed	32	18.6
Household land ownership		
Yes	103	59.9
No	69	40.1
Have kitchen garden		
Yes	58	33.7
No	114	66.3
Served food first		
Child	85	49.4
Father	41	23.8
Mother	43	25
Grandparents	3	1.7
Prime portion		
Child	28	16.3
Father	86	50

---

Mother	53	30.8
Grandparents	5	2.9

---

#### **Association between the study variables**

The correlation between the dependent variable nutritional status and the independent variables were done using logistic regression test.

#### **Association between nutritional status and socio-demographic and economics characteristics of the participants**

The unadjusted logistic regression analysis result of the relationship between socio-demographic characteristics and nutritional status is also shown in table5 . Caesarean section babies were also more likely to be low birth weight compared to the vaginal delivery babies (OR = 2.36, 95% CI: 1.17-3.34,  $p = 0.01$ ). The same trend was observed in children grouped under stunting where odds of being stunted if a child was delivered through caesarian section were higher compared to vaginally borne children (OR = 2.40, 95% CI: 1.83-2.75,  $p = 0.03$ ). Additionally, children with caregivers who had no formal education had higher chances of being underweight compared to those who were with caregivers who had attained tertiary education (OR = 2.61, 95% CI: 1.82-2.99,  $p = 0.01$ ). Children aged 6 to 8 months had reduced chances of being stunted with OR = 0.53 (95% CI: 0.19-1.46,  $p = 0.02$ ). Furthermore, when the mother was served first, there were higher chances of children being wasted compared to when children were served first (OR = 1.30, 95% CI: 0.99-2.18,  $P = <0.01$ ).

**Table 5. Logistic regression analysis of nutritional status and socio-demographic and economic characteristics**

Source: Field data, 2024

<b>Variables</b>	Underweight <sup>a</sup>	Wasting <sup>a</sup>	Stunting <sup>a</sup>	Normal nutritional status
<b>Mode of delivery</b>				
Vaginal	1	1	1	1
Caesarean	2.36 (1.17-3.34) *	0.72 (0.30-1.71)	2.40 (1.83-2.75) *	0.26 (0.21-0.90)
<b>Educational status</b>				
Tertiary education	1	1	1	1
Primary education	0.51 (0.35-0.84)	1.60 (0.58-4.43)	1.52 (1.06-1.89)	1.05 (0.60-3.27)
Secondary education	0.79 (0.26-1.03)	1.35 (0.53-3.43)	0.62 (0.40-0.74)	2.03 (0.76-5.45)
No formal education	2.61 (1.82-2.99) *	0.92 (0.17-5.10)	0.29 (0.21-0.32)	0.48 (0.07-0.93)
<b>Age of child</b>				
24-59 months	1	1	1	1
9-23 months	1.66 (0.76-3.62)	2.20 (1.76-2.95)	1.67 (0.78-3.60)	0.99 (0.62-1.34)
6-8 months	0.57 (0.21-1.58)	1.09 (0.88-1.92)	0.53 (0.19-1.46) *	1.16 (0.35-4.63)
<b>Who is served first</b>				
Child	1	1	1	1
Father	2.29 (0.88-5.96)	0.75 (0.60-0.97)	2.20 (0.85-5.74)	0.57 (0.04-3.22)
Mother	1.33 (0.13-1.83)	1.30 (0.99-2.18) *	1.29 (0.28-2.89)	1.24 (0.22-2.22)
Other relatives	-	0.43 (0.09-1.10)	-	-

Normal nutritional status is the reference category, <sup>a</sup>Crude chances ratio, \* Statistically significant.

UNDER PEER REVIEW

The results of adjusted logistic regression analysis on the association between socio-demographic characteristics and nutritional status presented in table 6 indicated that children born through caesarean section had an association with increased risk of being underweight and stunted. (aOR = 3.39, 95% CI: 1.44-7.97,  $p = 0.03$ ) and (aOR = 3.12, 95% CI: 1.32-7.37,  $p = 0.04$ ), respectively. Children whose caregivers had secondary and tertiary education were also significantly associated with reduced chances of being underweight, with aOR = 0.34 (95% CI: 0.12-0.96,  $p = 0.03$ ) and aOR = 0.25 (95% CI: 0.08-0.67,  $p = 0.02$ ) in that order. Additionally, children aged 9 to 23 months were more likely to be underweight and stunted. Their adjusted chances ratio was aOR = 2.74 (95% CI: 1.12-6.71,  $p = 0.04$ ) for underweight and aOR = 2.70 (95% CI: 1.10-6.61,  $p = 0.02$ ) for stunted growth. In the adjusted model, being served first was not associated with any form of nutritional status.

UNDER PEER REVIEW

**Table 6. Binary logistic regression analysis for socio-demographic and economic factors associated with nutritional status**

Source: Field data, 2024

<b>Variables</b>	Underweight <sup>a</sup>	Wasting <sup>a</sup>	Stunting <sup>a</sup>	Normal nutritional status
<b>Mode of delivery</b>				
Vaginal	1	1	1	1
Caesarean	3.39 (1.44-7.97) *	2.33 (1.40-3.68)	3.12 (1.32-7.37) *	1.28 (1.86-6.23)
<b>Educational status</b>				
No. formal education	1	1	1	1
Primary education	2.04 (0.78-5.33)	1.57 (0.54-4.57)	2.23 (0.85-5.88)	1.91 (0.13-3.25)
Secondary education	0.34 (0.12-0.96) *	1.03 (0.38-2.77)	0.43 (0.16-1.17)	2.92 (1.17-4.02)
Tertiary education	0.25 (0.08-0.67) *	0.98 (0.25-1.54)	0.68 (0.24-1.25)	0.31 (0.07-2.04)
<b>Age of child</b>				
6-8 months	1	1	1	1
9-23 months	2.74 (1.12-6.71) *	1.04 (0.41-2.66)	2.70 (1.10-6.61) *	2.03 (1.01-7.02)
24-59 months	2.25 (1.64-3.82)	1.23 (0.76-3.40)	3.18 (1.25-5.01)	0.95 (0.64-5.21)
<b>Who is served first</b>				
Child	1	1	1	1
Father	3.60 (0.12-10.86)	0.43 (0.03-6.18)	3.40 (1.14-10.14)	2.19 (1.01-7.20)
Mother	2.33 (0.73-3.32)	1.11 (0.08-14.84)	2.18 (0.21-8.27)	1.03 (0.99-5.16)
Other relatives	-	2.21 (1.34-4.55)	-	-

Normal nutritional status is reference category, <sup>a</sup> Model adjusted for variables in the table, \* Statistically significant, - no respondent

UNDER PEER REVIEW

## Discussion

### Socio-demographic and economic characteristics

The findings of this study demonstrated many socio-demographic factors that are precise to children and might influence their nutritional status between 6 and 59 months. The distribution of children across different age groups reveals that a substantial proportion (43.6%) falls within the age range of 24-59 months, indicating a concentration of older children in the sample. This observation aligns with other studies that focused on child nutrition, where older children may face different nutritional challenges compared to their younger counterparts (20). The study also reveals that majority of children (94.8 percent) were born in health facilities and hence exercises every control through antenatal and perinatal care as a major contributor towards early childhood health status (21). The mode of delivery, with 76.2% of children born through the vagina, may have implications for early microbial colonization and subsequent health (22). Furthermore, the gender distribution indicates a slightly higher proportion of male children (51.2%), and gender disparities play a vital role in child nutrition outcomes (23). Comparing these findings with other studies, it is essential to consider contextual factors such as geographic location, cultural practices, and healthcare infrastructure to contextualize and interpret the observed patterns (24).

The predominance of caregivers aged 20 to 47 years highlights the reproductive age group's crucial role in child care and nutrition (25). The majority of caregivers who are married or cohabiting have outlined that the presence of a family support system plays a key role in impacting child feeding practices (26). The educational attainment of caregivers, particularly with approximately 45 percent having secondary and 24 percent having tertiary education, is a positive factor for child nutrition, as maternal education is consistently associated with improved child health outcomes (27, 28). Important factors in allocating resources, such as the size of the family and the gender of the head of the household, may affect the accessibility to healthy food (29). Notably, the presence of alcohol use among 27 percent of caregivers raises concerns, as substance use may impact caregiving practices and household priorities.

While a considerable proportion of caregivers' land is their own, the limited presence of kitchen gardens raises concerns about the accessibility of diverse and nutrient-rich foods. According to Rehman et al. (2019), ownership of land is positively associated with improved child nutritional status, indicating a potential link between household land ownership and enhanced access to resources that positively influence children's well-being (30). The dynamics of food distribution within households are evident, with caregivers often prioritizing serving food to children initially (31).

Besides, the results revealed agree with the cross-sectional studies from other countries on household economic status and child malnutrition. For example, cross-sectional study was conducted in Bangladesh to show that students from higher economic class had less prevalence of stunting and wasting as compared to students from lower economic class (32). This is in agreement with results from the Kericho County where casual labour income was established to be the leading source of income implying that poor families may not be able to feed their children adequately and meet other nutritional needs. However, due to COVID-19 restrictions, there has not been effective construction, existence and utilization of kitchen garden, which are critical in providing food for families, hence escalating the use of purchased foods that in most cases are poor in diet diversity (33). In connection with the experience of Kericho County, such policies as those concerning household food security, which encourage the establishment of kitchen gardens and support non-farm income-generating activities may have high potential for combating malnutrition and improving children's health. The nature of this analysis underscores the fact that child malnutrition is a

complex phenomenon and not just a function of either socio-demographic characteristics or economic status of households.

### **Association between nutritional status and socio-demographic and economic characteristics of the participants**

It was revealed that maternal education status and mode of delivery are significant predictors of underweight. Those children of mothers or caregivers with no education had higher odds of being underweight than those of caregivers with a tertiary education. These findings were consistent with a study conducted in Bangladesh that reported that children of mothers with secondary and higher education had a lower risk (RR: 0.83, 95% CI: 0.78, 0.88) of experiencing underweight compared to children of mothers who had no formal education (34). Additionally, another study described that these children were 23% less likely to be underweight if their mothers had secondary education than the children whose mothers had no education level (32). Maternal education was inversely related to underweight; the findings are endorsed by Woldeamanuel & Tesfaye (2019) (35) and (36). This may be attributed to the fact that educated mothers have a boost of information regarding appropriate feeding practices and general child rearing, which could have a positive impact on child nutrition. It also found that underweight was 2.36 more likely to occur in cesarean compared to those that were a result of vaginal delivery. This result goes against the available studies that portrayed an overall 1.88-fold rise in the likelihood of overweight or obesity in children who were aged between 2- 5 years but born through cesarean section compared with the children born through vaginal delivery (37). However, infants born by CS had a significantly lower risk of being underweight, 0.71 times that of infants born vaginally, according to Abbasi et al. (2018) (38). The above results may be due to health policies and initiatives, whereby more emphasis is placed on policy than on initiatives. The research also found that the order of serving has a relationship with malnutrition in children, with a test result of less than 0.05. The analysis of the results showed that children whose moms were served before their parents had a 1,30 times higher risk of wasting. These findings are similar to the study that reported an odds ratio of 2.18 for acute malnutrition related to families /children who ate meals with their families and those who had children ever.

In addition, the maternal nutritional status was also an important determinant of child health status. Poor nutritional status can lead to adverse births and perpetuate malnutrition in children, and maternal undernutrition disrupts the healthy growth of children (39). Shifting focus to child nutritionally, research carried out in Kenya showed that if the mother has a low BMI, there is high probability that resultant children will be underweight proving that maternal health is determinative of child nutrition and nutritional status. In this regard, people's reciprocal dependency is emphasized as a concern for maternal health indicates the inclusion of this segment in multifaceted solutions regarding child malnutrition. The quality of maternal nutrition should be enhanced through nutrition education and food supplementation, which if achieved, should have dramatic effects on children's nutrition status in future. If government policies put focus in maternal health, alongside child nutrition activities then the idea can blend the efforts towards tackling underweight challenges such as malnutrition.

### **Conclusion.**

The study also supported the center's assertion that socio demographic and economic factors play a huge role in the current state of nutrition. Individuals with a secondary or higher degree were less likely to be underweight. Maternal education influences optimal feeding practices, and the study revealed its pivotal role in fostering children's nutritional well-being. The associations observed between child age, hierarchical food distribution, and wasting echo established knowledge, highlighting the need for tailored interventions that

account for children's evolving dietary needs and household dynamics. Importantly, while some variables in the analysis did not show significant associations, the complex interplay of multifaceted factors influencing child nutritional status is evident. This aligns with the understanding that child nutrition outcomes are shaped by a combination of socio-economic, cultural, and behavioral influences that extend beyond singular demographic factors. The higher chances of underweight and stunting among children born via caesarean sections underscore the need for specialized care for this group. This study also outlines the significance of caregiver education, intra-household dynamics, and early feeding practices. The correlation between maternal education and reduced child underweight underscores the empowering impact of education on child well-being.

. The findings stressed the need for targeted interventions and adherence to evidence-based guidelines to optimize child growth. Further research is warranted to explore potential trends in underweight and the duration of illness, contributing to a broader understanding of integrated health and nutrition programs for child well-being. These findings align with existing research, thus emphasizing the need for integrated health and nutrition programs.

### **Definitions, Acronyms, Abbreviations**

**Dietary diversity score** is a numerical representation of the variety of food groups consumed by a child within a specified period, serving as an indicator to predict the sufficiency of both macro and micro-nutrients in their diet.

**Dietary practice:** refers to the reported behaviours, actions, or individual choices regarding dietary habit such as meal frequency and dietary diversity.

**Morbidity status:** is the state of having illnesses or symptoms of illnesses.

**Nutritional status:** the health condition of children, influenced by the intake and utilization of nutrients, often assessed through anthropometric indicators such as height-for-age, weight-for-age, and weight-for-height.

**Stunting:** a condition characterized by a child's height-for-age falling below a specified standard, typically expressed as a Z-score, indicating a level of growth retardation.

**Underweight:** to a child's weight-for-age falling below a specified standard, usually expressed as a Z-score, suggesting a deficiency in overall body mass relative to age.

**Wasting:** a condition marked by a child's weight-for-height falling below a specified standard, often expressed as a Z-score.

**AOR:** Adjusted Chances Ratio

**BMI:** Body Mass Index

**CWC:** Child Welfare Clinic

**DDS:** Dietary diversity score

**FBP:** Food by prescription

**FFQ:** Food Frequency Questionnaire

**HAZ:** Height-for-age

**KCRH:** Kericho County Referral Hospital

**KEPI:** Kenya Expanded Program on Immunization

**KSCH:** Kapkatet Sub-County Hospital

**MUAC:** Mid Upper Arm Circumference

**PEM:** Protein Energy Malnutrition

**SPSS:** Statistical Package for Social Sciences

**WASH:** Water, Sanitation and Hygiene

**WAZ:** The weight-for-age z- score

**WHZ:** weight-for-length/height z-score

**WHO:** World Health Organization

### **Recommendations.**

The research outcomes resulted in 2 notable recommendations as follows:

1. There is a need to establish nutrition specific intervention program in Kericho County targeting children aged 6 to 59 months. This program should comprehensively address both acute and chronic malnutrition.
2. There is a need to intensify the promotion, awareness and support among caregivers on exclusive breastfeeding for the first six months of life, followed by the introduction of safe complementary foods while continuing breastfeeding up to two years of age or beyond

## **CONSENT**

Study name: Determinants of socio-economic and demographic characteristics of caregivers of children aged 6-59 months in kericho county, kenya

Principal investigator: Daniel Kipngeno Cheruiyot

### Introduction

My name is....., a researcher in this county. I am conducting a study on determinants of socio-economic and demographic characteristics of caregivers of children aged 6-59 months in kericho county, kenya. Your child has been selected as one of the people who can participate in this study to help us find answers to that problem. To be able to enrol, i am seeking for your permission so that you get to know more about the study. Am presenting this information pack for your perusal. Please read this consent document and ask questions for clarifications. At the bottom of the document place your signature. Your signature shall be interpreted as consent to the study. To be able to grant permission you must be 18 years and above therefore a legal guardian who must be above 18 years will sign on behalf of the child.

You may withdraw from the study anytime you wish to without prior notice through verbal or written communication.

### Background of the study

Kenya is only one of several nations throughout the world that are dealing with the public health crisis of malnutrition. Malnutrition may occur when there is an imbalance of vital nutrients, an imbalance in the intake or utilization of nutrients, or both. Health care for everyone and ensuring adequate nutrition and food security are two of the four tenets of kenya's big four agenda, which is in line with the planned study.

The national commission for science, technology & innovation (nacosti), the statutory organization tasked with regulating research in kenya by law, has examined and authorized the study. The mount kenya university scientific and ethical review committee approved the study's ethical component.

### Benefits

#### *Short term*

You will know your nutritional status, how to have a good nutritional status and the available options for improving nutrition at no cost.

#### *Long term*

You will be part of solution on malnutrition problem that part of the people of kericho county are suffering from.

## Risks

You will compromise confidential information during this study. However, that information will be kept confidential.

## Participants' obligation

To sincerely answer the asked questions to the ability and best of their knowledge.

## Confidentiality

A unique number will be assigned you at the recruitment. This number will be used as your name throughout the study and any other time you will be required in reference to this study.

No record of your name will be entered anywhere in the data collection tools or any place. The data will be kept in confidential lockable locker. Am the only person who will keep the key.

The data will be published but your name will not appear anywhere. Should the names be required for whatever reason, your name will not be disclosed without you being contacted and consent sought from you.

## Disclaimer

To participate in this study is voluntary. No charges whatsoever will be charged. There will be no coercion or threats. If you wish to consult someone before giving consent, that will be granted. You can choose to discontinue from the study whenever you want to. Should you decline consent to this study, there will be no loss of benefit, penalty or denial of services. If you wish this information, be translated to a language of your choice, a translator will be provided before.

## Consent form

### Purpose of the study

(1) To determine the socio-economic & demographic characteristics of caregivers of children of 6-59 months years old in kericho county, kenya  
Having gone through the content of the research, i consent to the study.

I ..... Do consent to the study titled determinants of nutritional status of children of six to fifty-nine months in kericho county

Signature.....date.....

Mothers below 18 years old: i ..... Do assent to the study titled determinants of nutritional status of children of six to fifty-nine months in kericho county

Signature.....date.....

Researcher  
name.....signature.....date.....

All authors declare that 'written informed consent was obtained from the patient (or other approved parties) for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editorial office/Chief Editor/Editorial Board members of this journal.

### **Ethical approval.**

This study initially sought approval from the Mount Kenya University Institutional Scientific and Ethical Review Committees (ISREC). There is also the need to make a research authorization from NACOSTI (the National Commission for Science, Technology, and Innovation) before conducting any study. The Kericho County Government and the administrative units of the participating health institutions offered their blessings for the research exercise.

Participants' identity was maintained throughout the study by providing them with numerical codes and in the self-completed questionnaire, there were no questions on identifying details of the participants. Each of the child's caregiver was asked for their permission to participate in the study for them to be selected and they were free to pull out at any time without any penalties during the study process. In the case of the underage caregivers, consent was sought from the guardian of the minor, The primary caregivers responded to the questionnaire on behalf of the child they consented.

To assess demographic, socio- economic characteristics of each child, a pre-tested trans-adapted structured questionnaire was completed by the responding primary caregivers. In addition, the pilot testing of the questionnaire was also conducted and the results used to make modifications in the main one. The interviewer had good knowledge of some of the local languages, in this case she was able to interview the primary caregivers some of whom could barely communicate in English or Kiswahili. The mother or primary caregiver of each child was expected to answer the interviewer's questions. Furthermore, all information sources were cited as they should.

### **REFERENCES**

1. World Health Organization. (2017). Guideline: Assessing and managing children at primary health-care facilities to prevent overweight and obesity in the context of the double burden of malnutrition. World Health Organization. <https://iris.who.int/handle/10665/259133>

2. UNICEF. (2022). *Nutrition: Preventing and treating maternal, adolescent and children malnutrition*. Retrieved on April 26, 2022 from <https://www.unicef.org/kenya/nutrition>
3. Yusuf, A. A. (2017). Prevalence of malnutrition among the children under 5 years in East African countries from 2013 to 2017. *International Journal for Scientific Research & Development*, 5(07), 222-222.
4. Kigaru, D. M. D., & Milelu, M. M. (2017). Dietary diversity, water and sanitation practices and nutritional status of children aged 6-59 months in Kitui County, Kenya. *International Journal of Food Science and Nutrition*, 2(5), 2455-4898.
5. KNBS and ICF. (2023). Kenya Demographic and Health Survey 2022. Key Indicators Report. Nairobi, Kenya, and Rockville, Maryland, USA: KNBS and ICF
6. Bayih, M. T., Arega, Z., & Motbainor, A. (2022). Nutritional status of 6–59 months of age children is not significantly varied between households with and without home gardening practices in Zege, North West Ethiopia, 2020: community based comparative study. *BMC pediatrics*, 22(1), 1-11.
7. Mboya, D., Bitok, L. K., Karani, A. K., Osano, B., & Habtu, M. (2015). Outcomes of severely malnourished children aged 6-59 months on outpatient management program in Kitui Hospital, Kenya. *Open Journal of Pediatrics*, 5, 326-333.
8. Kahsay, A., Mulugeta, A., & Seid, O. (2015). Nutritional status of children (6-59 months) from food secure and food insecure households in rural communities of SaesieTsaeda-Emba District, Tigray, North Ethiopia: comparative study. *International Journal of Nutrition and Food Sciences*, 4(1), 51.
9. Kwena, A. M., & Baliddawa, J. B. (2012). Nutritional status of children aged 6 to 59 months in community based education and service centres (COBES) in Western Kenya. *African Journal of Food, Agriculture, Nutrition and Development*, 12(3), 6229-6244

10. Gudu, E., Obonyo, M., Omballa, V., Oyugi, E., Kiilu, C., Githuku, J., ... & Ransom, J. (2020). Factors associated with malnutrition in children < 5 years in western Kenya: a hospital-based unmatched case control study. *BMC nutrition*, 6(1), 1-7.
11. Chakrabarty, K., Chakrabarty, A. S., Chakrabarty, K., & Chakrabarty, A. S. (2019). Nutritional deficiencies and disorders. *Textbook of Nutrition in Health and Disease*, 151-170. [https://link.springer.com/chapter/10.1007/978-981-15-0962-9\\_8](https://link.springer.com/chapter/10.1007/978-981-15-0962-9_8)
12. Kiani, A. K., Dhuli, K., Donato, K., Aquilanti, B., Velluti, V., Matera, G., ... & Bertelli, M. (2022). Main nutritional deficiencies. *Journal of Preventive Medicine and Hygiene*, 63(2 Suppl 3), E93. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9710417/>
13. Ahmad, D., Afzal, M., & Imtiaz, A. (2020). Effect of socioeconomic factors on malnutrition among children in Pakistan. *Future Business Journal*, 6(1), 1-11.
14. Igbokwe, O., Adimorah, G., Ikefuna, A., Ibeziako, N., Ubesie, A., Ekeh, C., & Iloh, K. (2017). Socio-demographic determinants of malnutrition among primary school aged children in Enugu, Nigeria. *Pan African Medical Journal*, 28(1).
15. Chaudhary, P., & Agrawal, M. (2018). Malnutrition and associated factors among children below five years of age residing in slum area of Jaipur City, Rajasthan. *India Asian. J Clin Nutr*, 11(1), 1-8.
16. Ndemwa, M., Wanyua, S., Kaneko, S., Karama, M., & Anselimo, M. (2017). Nutritional status and association of demographic characteristics with malnutrition among children less than 24 months in Kwale County, Kenya. *The Pan African Medical Journal*, 28.
17. Ole Tankoi, E. O., Asito, S. A., & Adoka, S. (2016). Determinants of Malnutrition among Children Aged 6–59 Months in Trans-Mara East Sub-County, Narok County, Kenya. *International Journal of Public Health & Safety*, 1(116), 2.

18. Kenya National Bureau of Statistics and ICF. Kenya Demographic and Health Survey 2022. Key Indicators Report. Nairobi, Kenya, and Rockville, Maryland, USA: KNBS and ICF. [Internet]. 2023. Available from: [www.DHSprogram.com](http://www.DHSprogram.com).
19. Anino, O. C., Were, G. M., & Khamasi, J. W. (2015). Impact evaluation of positive deviance hearth in Migori County, Kenya. *African Journal of Food, Agriculture, Nutrition and Development*, 15(5), 10578-10596.
20. Akseer, N., Al-Gashm, S., Mehta, S., Mokdad, A., & Bhutta, Z. A. (2017). Global and regional trends in the nutritional status of young people: A critical assessment of progress towards the SDGs. *Nature*, 555(7694), 41–47. <https://doi.org/10.1038/nature25770>
21. UNICEF. (2020). Strategy for improved nutrition of children and women in developing countries. Retrieved from <https://www.unicef.org/about/execboard/files/2019-PL3-EN.pdf>
22. Levin, B. R., Antia, R., Paul, M. R., & Lipsitch, M. (2016). The population genetics of antibiotic resistance. *Clinical Infectious Diseases*, 62(Suppl 2), S59–S64. <https://doi.org/10.1093/cid/civ961>
23. Santoso, M. V., Kerr, R. B., Hoddinott, J., Garlick, C., Olmos, S., & Young, S. L. (2019). Role of gender in determining child nutrition outcomes in Tanzania. *Journal of Nutrition*, 149(11), 1995–2006. <https://doi.org/10.1093/jn/nxz183>
24. Jones, A. D., Hoey, L., Blesh, J., Miller, L., Green, A., & Shapiro, L. F. (2019). A systematic review of the measurement of sustainable diets. *Advances in Nutrition*, 10(4), 748–769. <https://doi.org/10.1093/advances/nmz004>
25. Nyamasege, C. K. (2021). Maternal and caregiver demographic characteristics and their association with child nutrition in Kenya. *Maternal & Child Nutrition*, 17(4), e13150. <https://doi.org/10.1111/mcn.13150>
26. Mahmood, T., Rizwan, M., & Maqsood, R. (2021). Role of family support in shaping early child nutrition outcomes. *BMC Public Health*, 21(1), 1234. <https://doi.org/10.1186/s12889-021-11234-z>

27. Chakona, G. (2020). Social and cultural aspects of food security in rural South Africa. *Food Security*, 12(1), 217-232. <https://doi.org/10.1007/s12571-019-00986-y>
28. Mensch, B. S., Chuang, E. K., Melnikas, A. J., & Psaki, S. R. (2019). Evidence for causal links between education and maternal and child health: Systematic review. *Tropical Medicine & International Health*, 24(5), 504-522. <https://doi.org/10.1111/tmi.13218>
29. Drammeh, W., Hamid, N. A., & Rohana, A. J. (2019). Determinants of household food insecurity and its association with child malnutrition in Sub-Saharan Africa: A review of the literature. *Current Research in Nutrition and Food Science Journal*, 7(3), 610-623. <http://www.foodandnutritionjournal.org/volume7number3/determinants-of-household-food-insecurity-and-its-association-with-child-malnutrition-in-sub-saharan-africa-a-review-of-the-literature/>
30. Rehman, A., Ping, Q., & Razzaq, A. (2019). Pathways and associations between women's land ownership and child food and nutrition security in Pakistan. *International Journal of Environmental Research and Public Health*, 16(18), 3360. <https://www.mdpi.com/1660-4601/16/18/3360/pdf>
31. Raza, A., Fox, E. L., Morris, S. S., Kupka, R., Timmer, A., Dalmiya, N., & Fanzo, J. (2020). Conceptual framework of food systems for children and adolescents. *Global Food Security*, 27, 100436. <https://doi.org/10.1016/j.gfs.2020.100436>
32. Chowdhury, T. R., Chakrabarty, S., Rakib, M., Afrin, S., Saltmarsh, S., & Winn, S. (2020). Factors associated with stunting and wasting in children under 2 years in Bangladesh. *Heliyon*, 6(9). [https://www.cell.com/heliyon/pdf/S2405-8440\(20\)31692-3.pdf](https://www.cell.com/heliyon/pdf/S2405-8440(20)31692-3.pdf)
33. Mwangi, I. P. (2022). *Potential For Periodic Markets As Nodes Of Food Trade And Territorial Development: A Case Of Machakos-Kitui Road, Machakos County, Kenya* (Doctoral dissertation, University of Nairobi).

34. Hasib, M., Hassan, M. N., Hasan, M., & Khan, M. S. I. (2020). Effect of nutritional status on Rohingya under-five children in Bangladesh. *IJPHS*, 9, 358.  
[https://www.academia.edu/download/65240150/13\\_L20546\\_38782\\_1\\_ED\\_EDIT\\_C\\_b.pdf](https://www.academia.edu/download/65240150/13_L20546_38782_1_ED_EDIT_C_b.pdf)
35. Woldeamanuel, B. T., & Tesfaye, T. T. (2019). Risk factors associated with under-five stunting, wasting, and underweight based on Ethiopian demographic health survey datasets in Tigray region, Ethiopia. *Journal of nutrition and metabolism*, 2019.  
<https://www.hindawi.com/journals/jnme/2019/6967170/abs/>
36. Mtongwa, R. H., Festo, C., & Elisaria, E. (2021). A comparative analysis of determinants of low birth weight and stunting among under five children of adolescent and non-adolescent mothers using 2015/16 Tanzania Demographic and Health Survey (TDHS). *BMC nutrition*, 7, 1-10.  
<https://link.springer.com/article/10.1186/s40795-021-00468-6>
37. Papadopoulou, S. K., Mentzelou, M., Pavlidou, E., Vasios, G. K., Spanoudaki, M., Antasouras, G., ... & Giaginis, C. (2023). Caesarean Section Delivery Is Associated with Childhood Overweight and Obesity, Low Childbirth Weight and Postnatal Complications: A Cross-Sectional Study. *Medicina*, 59(4), 664.  
<https://doi.org/10.3390/medicina59040664>
38. Abbasi, S., Mahmood, H., Zaman, A., Farooq, B., Malik, A., & Saga, Z. (2018). Indicators of malnutrition in under 5 Pakistani children: a DHS data secondary analysis. *Journal of Medical research and Health education*, 2(3), 1-10. <http://www.imedpub.com/medical-research-and-health-education/>
39. [Kumar, M., & Mohanty, P. \(2022\). Does maternal overnutrition carry child undernutrition in India?. \*Plos one\*, 17\(6\), e0265788.](#)