

Numerical solution of fractional diffusion equation by shifted Legendre operational matrix method and fractional linear multi-step methods.

Abstract:

The paper deals with an efficient scheme to solve fractional diffusion equation including both time and spatial fractional derivative in Caputo sense. Firstly, the operational matrix of fractional derivative on shifted Legendre orthogonal polynomials is computed and used with spectral Tau method to reduce the problem to the system of fractional ordinary differential equation (FODE). Then the fractional linear multi-step method (FLMMS) with the help of the initial conditions permit to obtain the approximate solution. In order to show the validity of the proposed method, two numerical examples are reported.

Keywords: fractional diffusion equation, operational matrix of shifted Legendre polynomials, spectral Tau method, fractional ordinary differential equation, Linear multi-step methods.

Mathematics Subject Classification: 34A05, 34A08, 42A10, 40A30, 65B10.

1 Introduction

The diffusion equation models many engineering problems such as heat transfer, chemical diffusion, mass transfer, refrigeration and so on. Most of these problems can be well explained by fractional order partial differential equations (FDEs) [1, 3, 4, 5, 7, 8, 23]. In the present paper we consider the problem:

$$\begin{cases} \frac{\partial^\alpha u(x, t)}{\partial t^\alpha} = a \frac{\partial^\sigma u(x, t)}{\partial x^\sigma} + K(x, t) \\ u(0, t) = p(t) \\ u(1, t) = q(t) \\ u(x, 0) = g(x), \end{cases} \quad (1.1)$$

where $0 < \alpha \leq 1$; $0 < \sigma \leq 2$; $0 \leq x, t \leq 1$, a is the coefficient of diffusibility, $K(x, t)$ is the source function. It is not easy to obtain the analytic solution of (1.1). As result for the solution of (1.1), the numerical techniques such as: homotopy perturbation method [25, 26, 27, 28], Tau method

2 Preliminaries

[22, 29, 30, 31, 32, 33, 34, 35], sinc-Legendre method and so on are employed. These methods are very efficient but due to the high computational complexities they are hard to be applied to fractional partial differential equation. We need therefore an easy and efficient method to solve such equations. The motivation of this study is to built the numerical solution wich is a combination of operational matrix [2] method based on the fractional derivative of shifted Legendre polynomial [35, 36] and the FLMMs Method [15, 19, 21]. The application of the orthogonal shifted Legendre operational matrix and the spectral Tau method lead to system of FODE [10, 24] wich can be solved by FLMMs method. In the first step, the shifted Legendre operational matrix method is used to treat space variable x wich come from fractional part of the problem, and in the second step the spectral Tau method with the aid of the residual equation permit to obtain a system of FODE, then FLMM method is used to solve the equation, including time variable t wich come from fractional part of the problem. The paper is organized as follows: In section 2, mathematical preliminaries used in this paper are given. In section 3, shifted Legendre polynomial and the operational matrix are introduced. In section 4, application of the proposed method for solving space and fractional diffusion problem is presented. In section 5, the proposed method are used to demonstrate his reliability and his accuracy on two illustratives examples. In section 6, the concluded results are presented.

2 Preliminaries

In this section, we recall the definitions of fractional Riemann-Liouville integral and his derivative, the fractional derivative in Caputo sense, the fractional ordinary differential equation and the linear multi-step method for fractional equation [1, 2, 3, 4, 5, 6, 7, 8].

2.1 The fractional derivatives and property

Definition 2.1 : *The Riemann–Liouville (RL) fractional integral of order $\alpha > 0$ and origin at to for a function $y(t) \in L^1[0, \pi]$ is defined as:*

$$J_{t_0}^\alpha y(\tau) = \frac{1}{\Gamma(\alpha)} \int_{t_0}^t (t - \tau)^{\alpha-1} y(\tau) d\tau. \tag{2.1}$$

Definition 2.2 : *The RL fractional derivative is defined as:*

$$\widehat{D}_{t_0}^\alpha y(\tau) := D^m J_{t_0}^{m-\alpha} y(t) = \frac{1}{\Gamma(m - \alpha)} \frac{d^m}{dt^m} \int_{t_0}^t (t - \tau)^{m-\alpha-1} y(\tau) d\tau, \tag{2.2}$$

where $m = \lceil \alpha \rceil$ is the smallest integer greater or equal to α and $D^m, y^{(m)}, \frac{d^m}{dt^m}$ denotes the standard integer-order derivative.

Definition 2.3 : *For $y \in A^m([t_0, T])$ (i.e., $y^{(m)}$ absolutely continuous), the Caputo derivative is given by:*

$$D_{t_0}^\alpha y(\tau) := J_{t_0}^{m-\alpha} D^m y(t) = \frac{1}{\Gamma(m - \alpha)} \int_{t_0}^t (t - \tau)^{m-\alpha-1} y^{(m)}(\tau) d\tau. \tag{2.3}$$

If $t_0 = 0$, then $D_{t_0}^\alpha y(\tau) = D^\alpha y(t)$ and we have

$$D^\alpha y(t) = \frac{1}{\Gamma(m - \alpha)} \int_0^t \frac{y^{(m)}(\tau)}{(t - \tau)^{\alpha+1-m}} d\tau, \quad m - 1 < \alpha \leq m, \quad m \in \mathbb{N}. \tag{2.4}$$

For the Caputo derivative:

$$D^\alpha C = 0, \quad (C \text{ is a constant}), \tag{2.5}$$

2.2 Fractional ordinary differential equation (FODE)

$$D^\alpha t^\beta = \begin{cases} 0, & \text{for } \beta \in \mathbb{N}_0 \text{ and } \beta < [\alpha], \mathbb{N}_0 = \{0, 1, \dots\} \\ \frac{\Gamma(\beta + 1)}{\Gamma(\beta + 1 - \alpha)} t^{\beta - \alpha}, & \text{for } \beta \in \mathbb{N}_0 \text{ and } \beta \geq [\alpha] \text{ or } \beta \notin \mathbb{N} \text{ and } \beta > [\alpha], \end{cases} \quad (2.6)$$

where $[\alpha]$ is the smallest integer greater than or equal to α and $\lfloor \alpha \rfloor$ is the largest integer less than or equal to α . The Caputo fractional differentiation is a linear, then we have:

$$D^\alpha (\lambda f(x) + \mu g(x)) = \lambda D^\alpha f(x) + \mu D^\alpha g(x), \text{ with } \lambda \text{ and } \mu \text{ constants.} \quad (2.7)$$

We have also [5]

$$J_{t_0}^\alpha D_{t_0}^\alpha y(t) = y(t) - T_{m-1}[y; t_0](t), \quad (2.8)$$

where $T_{m-1}[y; t_0](t) = \sum_{k=0}^{m-1} \frac{(t - t_0)^k}{k!} y^{(k)}(t)$ is the polynomial.

The relation between Riemann-Liouville and the Caputo derivative is of the form

$$\widehat{D}_{t_0}^\alpha y(t) = D_{t_0}^\alpha y(t) + \sum_{k=0}^{m-1} \frac{(t - t_0)^k}{k!} y^{(k)}(t). \quad (2.9)$$

2.2 Fractional ordinary differential equation (FODE)

The most important application of the fractional Caputo's derivative is the fractional ordinary differential equation (FODE) [9] of the form

$$\begin{cases} D_{t_0}^\alpha y(t) = f(t, y(t)) \\ y(t_0) = y_0, y'(t_0) = y_0^{(1)}, \dots, y^{(m-1)}(t_0) = y_0^{(m-1)} \end{cases} \quad (2.10)$$

where $m - 1 < \alpha \leq m$, $0 \leq t \leq T$, $f(t, y(t))$ is continuous and $y_0^{(1)}, \dots, y_0^{(m-1)}$ are the value of derivative at t_0 . The application to both sides of Equation (2.10) of the RL integration $J_{t_0}^\alpha$, together with Equation (2.8), leads to the reformulation of FDE in terms of the weakly-singular Volterra Integration Equation (VIE) [15, 16, 19]

$$y(t) = T_{m-1}[y; t_0](t) + \frac{1}{\Gamma(\alpha)} \int_{t_0}^t (t - \tau)^{\alpha-1} f(\tau, y(\tau)) d\tau. \quad (2.11)$$

The integral formulation (2.11) is surely useful since it allows exploiting theoretical and numerical results already available for this class of VIEs in order to study and solve FODEs.

2.3 Convolution quadrature rules

One way to obtain solution of (2.10) is to approximate (2.11) by convolution quadrature [11, 12, 13, 14, 18, 20] formulas, which can be written in the general form:

$$y_n = \varphi_n + \sum_{j=0}^n c_{n-j} f_j, \quad f_j = f(t_j, y_j), \quad (2.12)$$

where φ_n and c_n are known coefficient $t_n = t_0 + nh$ is an assigned grid. With a constant step-size $h > 0$ and $y_n = y(t_n)$. The way in which the coefficients are derived depends on the specific method. In particular, the following two classes of convolution quadrature rules for FODEs are mainly studied in the literature: Product-Integration (PI) rules [10, 17, 20, 24] and Fractional Linear Multi-Step methods (FLMMs) [15, 21, 24].

2.3 Convolution quadrature rules

2.3.1 Product-Integration (PI) rules

Given a grid $t_n = t_0 + nh$, with constant step-size $h > 0$, in PI rules, the solution of the VIE (2.11) at t_n is first written in a piece-wise way:

$$y(t_n) = T_{m-1}[y; t_0](t_n) + \frac{1}{\Gamma(\alpha)} \int_{t_0}^{t_n} (t_n - \tau)^{\alpha-1} f(\tau, y(\tau)) d\tau, \quad (2.13)$$

where $f(\tau, y(\tau))$ can be approximated, in each subinterval $[t_j, t_{j+1}]$ by means of some interpolant polynomial. If $f(\tau, y(\tau))$ is approximated by $f(t_j, y_j)$, we obtain the explicit PI rectangular:

$$y_n = T_{m-1}[y; t_0](t_n) + h^\alpha \sum_{j=0}^{n-1} b_{n-j-1}^{(\alpha)} f(j, y(t_j)). \quad (2.14)$$

If $f(\tau, y(\tau))$ is approximated by $f(t_{j+1}, y(t_{j+1}))$, we obtain the implicit PI rectangular:

$$y_n = T_{m-1}[y; t_0](t_n) + h^\alpha \sum_{j=1}^{n-1} b_{n-j}^{(\alpha)} f(j, y(t_j)), \quad (2.15)$$

with $b_n^{(\alpha)} = \frac{((n+1)^\alpha - n^\alpha)}{\Gamma(\alpha+1)}$. If $f(\tau, y(\tau))$ is approximated by $f(t_{j+1}, y_{j+1}) + \frac{s-t_{j+1}}{h} (f(t_{j+1}, y_{j+1}) + f(t_j, y_j))$, we obtain implicit Trapezoidal rule [17]

$$y_n = T_{m-1}[y; t_0](t_n) + h^\alpha \left(\hat{a}_n^{(\alpha)} f_0 + \sum_{j=1}^{n-1} a_{n-j}^{(\alpha)} f(t_j, y(t_j)) \right), \quad (2.16)$$

with

$$\hat{a}_n^{(\alpha)} = \frac{(n-1)^{\alpha+1} - n^\alpha (n-\alpha-1)}{\Gamma(\alpha+2)}, \quad a_n^{(\alpha)} = \begin{cases} \frac{1}{\Gamma(\alpha+2)}, & \text{for } n=0, \\ \frac{(n-1)^{\alpha+1} - 2n^{\alpha+1} + (n+1)^{\alpha+1}}{\Gamma(\alpha+2)}, & \text{for } n=1, 2, \dots \end{cases}$$

2.3.2 Fractional Linear Multi-Step Methods (FLMMs)

Another way to approximate FODE is to use the FLMMs Method. Given the initial value problem:

$$y'(t) = f(t), \quad y(t_0) = y_0, \quad (2.17)$$

its solution can be approximated by means of an linear multi-step method (LMM) given by:

$$\sum_{j=0}^k \rho_j y_{n-j} = h \sum_{j=0}^k \sigma_j f(t_{n-j}),$$

where $\rho(z) = \rho_0 z^k + \rho_1 z^{k-1} + \dots + \rho_k$ and $\sigma(z) = \sigma_0 z^k + \sigma_1 z^{k-1} + \dots + \sigma_k$ are the first and second characteristic polynomial of the LMM. Problem (2.17) can be rewritten in the integral from:

$$y(t) = y_0 + \int_{t_0}^t f(\tau) d\tau.$$

The solution $y(t)$ can be approximated by using LMMs reformulated in terms of convolution quadrature formulas:

$$y_n = h \sum_{j=0}^n \omega_{n-j} f(t_j), \quad n \geq k$$

2.3 Convolution quadrature rules

where the weights ω_n depend on the characteristic polynomials $\rho(z)$ and $\sigma(z)$, but not on h .

$$\delta(\zeta) = \sum_{n=0}^{\infty} \omega_n \zeta^n, \quad \delta(\zeta) = \frac{\rho(1/\zeta)}{\sigma(1/\zeta)}.$$

The idea underlying FLMMs is to derive convolution quadratures for the RL integral (2.1) with convolution weights given by the function:

$$F\left(\frac{\delta(\zeta)}{h}\right) = \left(\frac{\delta(\zeta)}{h}\right)^{-\alpha} = h^\alpha \left(\frac{\rho(1/\zeta)}{\sigma(1/\zeta)}\right), \quad (2.18)$$

where $F(s) = s^{-\alpha}$ the Laplace transform of the kernel $\frac{t^{\alpha-1}}{\Gamma(\alpha)}$ in (2.1). The assumptions that make possible this generalization of LMMs are that the generating function $\delta(\zeta)$ has no zeros in the closed unit disc $|\zeta| \leq 1$, except for $\zeta = 1$, and $|\arg \delta(\zeta)| < \pi$ for $|\zeta| < 1$.

When an LMM is generalized to Equation (2.1) in the above Lubich sense, the resulting FLMM reads as:

$$hJ_{t_0}^\alpha f(t_n) = h^\alpha \sum_{j=0}^n \omega_{n-j}^{(\alpha)} f(t_j), \quad (2.19)$$

where the convolution quadrature weights $\omega_n^{(\alpha)}$ are obtained from:

$$\sum_{n=0}^{\infty} \omega_n^{(\alpha)} \zeta^n = \omega^{(\alpha)}(\zeta), \quad \omega^{(\alpha)}(\zeta) = (\delta(\zeta))^{-\alpha}.$$

Thus, to handle non-smooth functions (as happens in the solution of fractional-order problems), it is necessary to introduce a correction term:

$$hJ_{t_0}^\alpha f(t_n) = h^\alpha \sum_{j=0}^n \omega_{n,j}^{(\alpha)} f(t_j) + h^\alpha \sum_{j=0}^n \omega_{n-j}^{(\alpha)} f(t_j). \quad (2.20)$$

From the application of the discretized convolution quadrature rule (2.20) to integral Equation (2.11), we are able to derive FLMMs for the approximation of the solution of FDEs:

$$y_n = T_{m-1}[y; t_0](t_n) + h^\alpha \sum_{j=0}^s \omega_{n,j}^{(\alpha)} f(t_j, y_j) + h^\alpha \sum_{j=0}^n \omega_{n-j}^{(\alpha)} f(t_j, y_j). \quad (2.21)$$

The starting weights $\omega_{n,j}$ are chosen by imposing that the quadrature rule (2.20) is exact when applied to $f(t) = t^\gamma$, with γ assuming all the possible fractional values expected in the expansion of the true solution and, hence, by solving at each step the algebraic linear system:

$$\sum_{j=0}^s \omega_{n,j}^{(\alpha)} j^\gamma = - \sum_{j=0}^s \omega_{n-j}^{(\alpha)} j^\gamma + \frac{\Gamma(\gamma+1)}{\Gamma(1+\gamma+\alpha)} n^{\gamma+\alpha}. \quad (2.22)$$

One of the simplest FLMMs is obtained from the implicit Euler method (or BDF1). No starting weights are necessary in this case, and since the generating function is $\delta(\zeta) = 1 - \zeta$, we see that $\omega_n^{(\alpha)}$, $n = 0, 1, 2, \dots$, are the coefficients of the generalized binomial series $(1 - \zeta)^{-\alpha}$, namely:

$$\omega_n^{(\alpha)} = (-1)^n \binom{-\alpha}{n} = (-1)^n \frac{\Gamma(1-\alpha)}{\Gamma(n+1)\Gamma(-\alpha-n+1)},$$

3 Properties of shifted Legendre polynomials

which can be also evaluated by the recurrence $\omega_n^{(\alpha)} = \left(\frac{1 - (1 - \alpha)}{n}\right) \omega_{n-1}^{(\alpha)}$ with $\omega_0^{(\alpha)} = 1$. The corresponding method:

$$y_n = T_{m-1}[y; t_0](t_n) + h^\alpha \sum_{j=0}^n (-1)^n \binom{-\alpha}{n-j} f(t_j, y_j),$$

is commonly referred to as the Grünwald–Letnikov scheme [23].

3 Properties of shifted Legendre polynomials

The well-known Legendre polynomials [35] are defined on the interval $[-1, 1]$ and can be determined with the aid of the following recurrence formulae:

$$P_{i+1}(t) = \frac{2i+1}{i+1} t P_i(t) - \frac{i}{i+1} P_{i-1}(t), \quad i = 1, 2, 3, \dots,$$

where $P_0(t) = 1$ and $P_1(t) = t$. In order to use these polynomials on the interval $x \in [0, 1]$ we define the so-called shifted Legendre polynomials by introducing the change of variable $t = 2x - 1$. Let the shifted Legendre polynomials $P_i(2x - 1)$ be denoted by $G_i(x)$. Then $G_i(x)$ can be obtained as follows:

$$G_{i+1}(x) = \frac{(2i+1)(2x-1)}{i+1} G_i(x) - \frac{i}{i+1} G_{i-1}(x), \quad i = 1, 2, 3, \dots, \quad (3.1)$$

where $G_0(x) = 1$ and $G_1(x) = 2x - 1$. The analytic form of the shifted Legendre polynomial $G_i(x)$ of degree i given by:

$$G_i(x) = \sum_{k=0}^i (-1)^{i+k} \frac{(i+k)! x^k}{(i-k)! (k!)^2}. \quad (3.2)$$

Note that $G_i(0) = (-1)^i$ and $G_i(1) = 1$. The orthogonality condition is

$$\int_0^1 G_i(x) G_j(x) dx = \begin{cases} \frac{1}{2i+1}, & \text{for } i = j, \\ 0, & \text{for } i \neq j. \end{cases} \quad (3.3)$$

A function $z(x)$, square integrable in $[0, 1]$, may be expressed in terms of shifted Legendre polynomials as

$$z(x) = \sum_{j=0}^{\infty} c_j G_j(x),$$

where the coefficients c_j are given by:

$$c_j = (2j+1) \int_0^1 z(x) G_j(x) dx, \quad i = 1, 2, 3, \dots$$

In practice, only the first $(m+1)$ -terms shifted Legendre polynomials are considered. Then we have

$$z(x) = \sum_{j=0}^m c_j G_j(x) = C^T \Phi(x),$$

where the shifted Legendre coefficient vector B and the shifted Legendre vector $\Phi(x)$ are given by

$$C^T = [c_0, c_1, \dots, c_m],$$

4 Generalized Legendre operational matrix to fractional calculus

$$\Phi(x) = [G_0(x), G_1(x), \dots, G_m(x)]^T. \tag{3.4}$$

The derivative of the vector $\Phi(x)$ can be expressed by

$$\frac{d\Phi(x)}{dx} = D^{(1)}\Phi(x), \tag{3.5}$$

where $D^{(1)}$ is the $(m + 1) \times (m + 1)$ operational matrix of derivative given by

$$D^{(1)} = (d_{ij}) = \begin{cases} 2(2j + 1), & \text{for } j = i - k, \\ 0, & \text{otherwise,} \end{cases} \begin{cases} k = 1, 3, \dots, m, & \text{if } m \text{ odd,} \\ k = 1, 3, \dots, m - 1, & \text{if } m \text{ even,} \end{cases}$$

for example for even m we have

$$D^{(1)} = 2 \begin{bmatrix} 0 & 0 & 0 & 0 & \dots & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & \dots & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 & \dots & 0 & 0 & 0 \\ 1 & 0 & 5 & 0 & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 1 & 0 & 5 & 0 & \dots & 2m - 3 & 0 & 0 \\ 0 & 3 & 0 & 7 & \dots & 0 & 2m - 1 & 0 \end{bmatrix}$$

4 Generalized Legendre operational matrix to fractional calculus

By using Eq.(3.5),it is clear that

$$\frac{d^n \Phi(x)}{dx^n} = (D^{(1)})^n \Phi(x), \tag{4.1}$$

where $n \in \mathbb{N}$ and the superscript, in $D^{(1)}$, denotes matrix powers. Thus

$$D^{(n)} = (D^{(1)})^n, \quad n = 1, 2, 3, \dots \tag{4.2}$$

Lemma 4.1 *Let $G_i(x)$ be a shifted Legendre polynomial then*

$$D^\alpha G_i(x) = 0, \quad i = 0, 1, 2, \dots, [\alpha] - 1, \quad \alpha > 0. \tag{4.3}$$

Proof 4.1 *Using Eqs.(2.5) – (2.6) in Eq. (3.1) the lemma can be proved.*

In the following theorem we generalize the operational matrix of derivative of shifted Legendre polynomials given in (3.5) for fractional derivative.

Theorem 4.1 *Let $\Phi(x)$ be shifted Legendre vector defined in (3.4) and also suppose $\alpha > 0$ then*

$$D^\alpha \Phi(x) \simeq D^{(\alpha)} \Phi(x), \tag{4.4}$$

where $D^{(\alpha)}$ is the $(m + 1) \times (m + 1)$ operational matrix of fractional derivative of order α in the Caputo

4 Generalized Legendre operational matrix to fractional calculus

sense and is defined as follows:

$$D^{(\alpha)} = \begin{bmatrix} 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & \cdots & 0 \\ \sum_{k=\lceil\alpha\rceil}^{\lceil\alpha\rceil} \delta_{\lceil\alpha\rceil,0,k} & \sum_{k=\lceil\alpha\rceil}^{\lceil\alpha\rceil} \delta_{\lceil\alpha\rceil,1,k} & \cdots & \sum_{k=\lceil\alpha\rceil}^{\lceil\alpha\rceil} \delta_{\lceil\alpha\rceil,m,k} \\ \vdots & \vdots & \cdots & \vdots \\ \sum_{k=\lceil\alpha\rceil}^i \delta_{i,0,k} & \sum_{k=\lceil\alpha\rceil}^i \delta_{i,1,k} & \cdots & \sum_{k=\lceil\alpha\rceil}^i \delta_{i,m,k} \\ \vdots & \vdots & \cdots & \vdots \\ \sum_{k=\lceil\alpha\rceil}^m \delta_{m,0,k} & \sum_{k=\lceil\alpha\rceil}^m \delta_{m,1,k} & \cdots & \sum_{k=\lceil\alpha\rceil}^m \delta_{m,m,k} \end{bmatrix},$$

where $\delta_{i,j,k}$ is given by:

$$\delta_{i,j,k} = (2j + 1) \sum_{l=0}^j \frac{(-1)^{i+j+k+l} (i + k)! (j + l)!}{(i - k)! k! \Gamma(k - \alpha + 1) (j - l)! (l!)^2 (k + l - \alpha + 1)}. \tag{4.5}$$

Note that in D^α , the first $\lceil\alpha\rceil$ rows, are all zero.

Proof 4.2 Using Eqs. (2.6), (2.7) and (3.2) we have:

$$\begin{aligned} D^\alpha G_i(x) &= \sum_{k=0}^i (-1)^{i+k} \frac{(i + k)!}{(i - k)! (k!)^2} D^\alpha x^k \\ &= \sum_{k=\lceil\alpha\rceil}^i (-1)^{i+k} \frac{(i + k)!}{(i - k)! (k!) \Gamma(k - \alpha + 1)} x^{k-\alpha}, \quad i = \lceil\alpha\rceil, \dots, m. \end{aligned} \tag{4.6}$$

Now, approximate $x^{k-\alpha}$ by $(m + 1)$ terms of shifted Legendre series, we have

$$x^{k-\alpha} \simeq \sum_{j=0}^m b_{k,j} G_j(x), \tag{4.7}$$

where

$$\begin{aligned} b_{k,j} &= (2j + 1) \int_0^1 x^{k-\alpha} G_j(x) dx = (2j + 1) \sum_{l=0}^j (-1)^{j+l} \frac{(j + l)!}{(j - l)! (l!)^2} \int_0^1 x^{k+l-\alpha} dx \\ &= (2j + 1) \sum_{l=0}^j (-1)^{j+l} \frac{(j + l)!}{(j - l)! (l!)^2 (k + l - \alpha + 1)}. \end{aligned} \tag{4.8}$$

Employing Eqs. (4.6) – (4.8) we get

$$\begin{aligned} D^\alpha G_i(x) &\simeq \sum_{k=\lceil\alpha\rceil}^i \sum_{j=0}^m \frac{(-1)^{i+k} (i + k)!}{(i - k)! (k!) \Gamma(k - \alpha + 1)} b_{k,j} G_j(x) \\ &= \sum_{j=0}^m \left(\sum_{k=\lceil\alpha\rceil}^i \delta_{i,j,k} \right) G_j(x), \quad i = \lceil\alpha\rceil, \dots, m, \end{aligned} \tag{4.9}$$

5 Applications of the operational matrix of fractional derivative

where $\delta_{i,j,k}$ is given in Eq. (4.5). Rewrite Eq. (4.9) as a vector form we have

$$D^\alpha G_i(x) \simeq \left[\sum_{k=\lceil \alpha \rceil}^i \delta_{i,0,k}, \sum_{k=\lceil \alpha \rceil}^i \delta_{i,1,k}, \dots, \sum_{k=\lceil \alpha \rceil}^i \delta_{i,m,k} \right] \Phi(x), \quad i = \lceil \alpha \rceil, \dots, m. \quad (4.10)$$

Also according to Lemma 4.1, we can write

$$D^\alpha G_i(x) = [0, 0, \dots, 0] \Phi(x), \quad i = 0, 1, \dots, \lceil \alpha \rceil - 1. \quad (4.11)$$

A combination of Eqs.(4.10) and (4.11) leads to the desired result.

Remark 4.1 If $\alpha = n \in \mathbb{N}$, then theorem 4.1 gives the same result as Eq. (4.2).

5 Applications of the operational matrix of fractional derivative

In this section, we apply the operational matrix of fractional derivative relative to shifted Legendre polynomial and the spectral Tau method in order to reduce the problem to a system of fractional ordinary differential equation.

5.1 Linear fractional diffusion differential equation with integer order in space

Consider the Linear diffusion fractional differential equation in time with non homogeneous Dirichlet boundary conditions

$$\begin{cases} \frac{\partial^\alpha u(x,t)}{\partial t^\alpha} = a \frac{\partial^2 u(x,t)}{\partial x^2} + f(x,t) \\ u(0,t) = p(t) \\ u(1,t) = q(t) \\ u(x,0) = g(x), \end{cases} \quad (5.1)$$

where $0 < \alpha \leq 1$, $0 \leq t \leq T$, $0 \leq x \leq 1$; p and q are given functions. We approximate $\frac{\partial^2 u(x,t)}{\partial x^2}$, $f(x,t)$ and $g(x)$ as follows using operational matrix $D^{(2)}$

$$\frac{\partial^2 u(x,t)}{\partial x^2} \simeq C^T(t) D^{(2)} \Phi(x), \quad (5.2)$$

$$f(x,t) \simeq F^T(t) \Phi(x), \quad F_i(t) = (2i+1) \int_0^1 f(x,t) G_i(x) dx, \quad i = 1, 2, 3, \dots \quad (5.3)$$

$$g(x) \simeq g^T \Phi(x), \quad g_i = (2i+1) \int_0^1 g(x) G_i(x) dx. \quad (5.4)$$

Then Eq. (5.1) becomes:

$$\begin{cases} D_t^\alpha C^T(t) \Phi(x) = a C^T(t) D^{(2)} \Phi(x) + F^T(t) \Phi(x) \\ C^T(t) \Phi(0) = p(t) \\ C^T(t) \Phi(1) = q(t) \\ C^T(0) \Phi(x) = g^T \Phi(x), \end{cases} \quad (5.5)$$

where $c_i(0) = g_i, \quad i = 0, \dots, N, \quad F(t) = \begin{bmatrix} F_1(t) \\ \vdots \\ F_N(t) \end{bmatrix}, \quad C(0) = \begin{bmatrix} c_0(0) \\ \vdots \\ c_N(0) \end{bmatrix}.$

5.2 Linear fractional diffusion differential equation with fractional order in space

From (5.5), let $R_m(x)$ defined as:

$$R_m(x) = \left[D_t^\alpha C^T(t) - aC^T(t) D^{(2)} - F(t) \right] \Phi(x) = 0, \quad x \in [0, 1], \quad (5.6)$$

application the spectral Tau method [22, 29], we have:

$$\int_0^1 R_m(x) G_j(x) dx = 0, \quad j = 0, 1, \dots, N - 2. \quad (5.7)$$

By the properties of (3.3), we then have

$$\left\{ \begin{array}{l} D_t^\alpha c_i(t) = ac_i(t) + F_i(t), \quad i = 0, \dots, N - 2 \\ \sum_{i=0}^N (-1)^i c_i(t) = p(t) \\ \sum_{i=0}^N c_i(t) = q(t) \\ c(0) = g_i, \quad i = 0, 1, \dots, N. \end{array} \right. \quad (5.8)$$

The problem (5.8) can be written in the vectorial form as:

$$\left\{ \begin{array}{l} D_t^\alpha C(t) = \wedge C(t) + F(t) + V(t) \\ C(0) = G \\ \sum_{i=0}^N (-1)^i c_i(t) = p(t) \\ \sum_{i=0}^N c_i(t) = q(t) \end{array} \right. \quad (5.9)$$

$$\text{where } \wedge = \begin{bmatrix} a & 0 & \cdots & 0 & 0 \\ 0 & a & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 0 \\ 0 & 0 & \cdots & 0 & 0 \end{bmatrix}, \quad V(t) = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ p(t) \\ q(t) \end{bmatrix}.$$

The equation (5.9) can be as well of the form:

$$\left\{ \begin{array}{l} D_t^\alpha C(t) = Q(t, C(t)) \\ C(0) = g, \end{array} \right. \quad (5.10)$$

where $Q(t, C(t)) = \wedge C(t) + F(t) + V(t)$, with $Q_i(t, C(t)) = DC_i(t) + F_i(t) + V_i(t)$, $i = 0 \dots N$.

5.2 Linear fractional diffusion differential equation with fractional order in space

Consider the linear fractional diffusion equation in time and in space with non homogeneous Dirichlet boundary conditions

$$\left\{ \begin{array}{l} \frac{\partial^\alpha u(x, t)}{\partial t^\alpha} = a \frac{\partial^\sigma u(x, t)}{\partial x^\sigma} + k(x, t) \\ u(0, t) = r(t) \\ u(1, t) = s(t) \\ u(x, 0) = h(x), \end{array} \right. \quad (5.11)$$

6 Numerical experiment

where $0 < \alpha \leq 1$, $1 < \sigma \leq 2$, $0 \leq t \leq T$, $0 \leq x \leq 1$; r and s are two given functions. We approximate $h(x)$ and $k(x, t)$ as follows:

$$k(x, t) \simeq K^T(t) \Phi(x), \quad k_i(t) = (2i + 1) \int_0^1 k(x) G_i(x) dx. \quad (5.12)$$

$$h(x) \simeq H^T \Phi(x), \quad H_i = (2i + 1) \int_0^1 h(x) G_i(x) dx, \quad (5.13)$$

The Eq. (5.11) becomes:

$$\begin{cases} D_t^\alpha C^T(t) \Phi(x) = a C^T(t) D^{(\sigma)} \Phi(x) + K^T(t) \Phi(x) \\ C^T(t) \Phi(0) = r(t) \\ C^T(t) \Phi(1) = s(t) \\ C^T(0) \Phi(x) = H^T \Phi(x) \end{cases} \quad (5.14)$$

Where $C_i(0) = H_i$, $i = 0, \dots, N$. Define the residual function $R_m(x)$ as:

$$R_m(x) = \left[D_t^\alpha C^T(t) - a C^T(t) D^{(\sigma)} - K(t) \right] \Phi(x) = 0, \quad x \in [0, 1]. \quad (5.15)$$

The application of the Galerkin-Tau method give

$$\int_0^1 R_m(x) G_j(x) dx = 0, \quad j = 0, 1, \dots, N - 2. \quad (5.16)$$

6 Numerical experiment

We present numerical test using shifted Legendre operational method followed by multi-step method described in the previous section.

Example 6.1 Consider the following linear one-dimensional fractional diffusion equation in time and integer in space:

$$\begin{cases} \frac{\partial^{\frac{1}{2}} u(x, t)}{\partial t^{\frac{1}{2}}} - \frac{\partial^2 u(x, t)}{\partial x^2} = \frac{2t^{\frac{3}{2}} \sin(2\pi x)}{\Gamma\left(\frac{3}{2}\right)} + 4\pi^2 \sin(2\pi x) \\ u(0, t) = u(1, t) = 0 \\ u(x, 0) = 0, \quad (x, t) \in [0, 1] \times [0, 1] \end{cases} \quad (6.1)$$

The exact solution of the problem (6.1) is $u(x, t) = 2t^2 \sin(2\pi x)$. By applying the technique described in section 5.1 with $m = 3$, we approximate solution as

$$u_{app}(x, t) = c_0 G_0(x) + c_1 G_1(x) + c_2 G_2(x) + c_3 G_3(x) = C^T \Phi(x),$$

where c_0, c_1, c_2, c_3 are found from

$$\begin{cases} D_t^{\frac{1}{2}} c_0(t) = 0 \\ D_t^{\frac{1}{2}} c_1(t) = -\frac{12}{\pi} \left(\frac{t^{\frac{3}{2}}}{\sqrt{\pi}} + \pi^2 t^2 \right) \\ c_0(0) = 0 \\ c_1(0) = 0 \end{cases}$$

6 Numerical experiment

and the boundary condition

$$\begin{cases} c_0(t) - c_1(t) + c_2(t) - c_3(t) = 0 \\ c_0(t) + c_1(t) + c_2(t) + c_3(t) = 0, \end{cases}$$

using the operational matrix

$$D^{(2)} = 2 \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 6 & 0 & 0 & 0 \\ 0 & 30 & 0 & 0 \end{bmatrix}, \quad F(t) = \begin{bmatrix} 0 \\ -\frac{12}{\pi} \left(\frac{t^2}{\sqrt{\pi}} + \pi^2 t^2 \right) \\ 0 \\ 28 \left(\frac{t^2}{\sqrt{\pi}} + \pi^2 t^2 \right) \left(\frac{19}{\pi} - \frac{15}{\pi^3} \right) \end{bmatrix}.$$

Setting $N = 101$ in space and $\Delta t = \frac{1}{2^5}$ for step time, Figure 1 displays the profil of exact and approximate solution and show the profil of error $E = u - u_{app}$.

6 Numerical experiment




Figure 1

The results show an excellent agreement between the exact and approximate solution, which confirms the accuracy of the method.

Example 6.2 Consider the following linear one-dimensional fractional diffusion equation in time and in space:

$$\begin{cases} \frac{\partial^\alpha u(x,t)}{\partial t^\alpha} + \frac{\partial^\sigma u(x,t)}{\partial x^\sigma} = f(x,t), & 0 \leq x, t \leq 1, \quad 0 < \alpha \leq 1, \quad 1 < \sigma \leq 2 \\ u(0,t) = t^2 + 1 \\ u(1,t) = 2(t^2 + 1) \\ u(x,0) = x^2 + 1 \end{cases} \quad (6.2)$$

where $f(x,t) = \frac{2\Gamma(3)x^{2-\alpha}(t^2+1)}{\Gamma(3-\alpha)} + \frac{2\Gamma(3)t^{2-\sigma}(x^2+1)}{\Gamma(3-\sigma)}$ and $\alpha = \sigma = \frac{1}{2}$. The exact solution of the problem (6.2) is $u(x,t) = (x^2 + 1)(t^2 + 1)$. By applying the technique described in section 5.1 with $m = 2$, we approximate solution as

$$u_{app}(x,t) = c_0 G_0(x) + c_1 G_1(x) + c_2 G_2(x) = C^T \Phi(x).$$

where c_0, c_1, c_2 are found by solving

$$\begin{cases} D_t^{\frac{1}{2}} c_0(t) = \frac{8}{\sqrt{\pi}} \left(\frac{4}{15}(t^2+1) + \frac{1}{9}t^{\frac{3}{2}} \right) \\ c_0(0) = \frac{4}{3} \end{cases}$$

and the boundary condition

$$\begin{cases} c_0(t) - c_1(t) + c_2(t) = t^2 + 1 \\ c_0(t) + c_1(t) + c_2(t) = 2(t^2 + 1), \end{cases}$$

using the operational matrix

$$D\left(\frac{1}{2}\right) = \frac{8}{\sqrt{\pi}} \begin{bmatrix} 0 & 0 & 0 \\ \frac{1}{3} & \frac{1}{5} & \frac{-1}{21} \\ \frac{3}{5} & \frac{7}{7} & \frac{1}{3} \end{bmatrix}, \quad F(t) = \frac{8}{\sqrt{\pi}} \begin{bmatrix} \frac{4}{15}(t^2+1) + \frac{1}{9}t^{\frac{3}{2}} \\ \frac{12}{35}(t^2+1) + \frac{1}{3}t^{\frac{3}{2}} \\ \frac{4}{63}(t^2+1) + \frac{1}{9}t^{\frac{3}{2}} \end{bmatrix}.$$

6 Numerical experiment

As in the previous problem, by setting $N = 101$ and $\Delta t = \frac{1}{2^5}$ the Figure 2 displays the profil exact and approximate solution and show the error $E = u - u_{app}$.




Figure 2

The results show an excellent agreement between the exact and approximate solution, which confirm the accuracy of the method.

7 Conclusion

In this paper, we have presented a numerical approach for solving fractional linear diffusion equation one in time only and one in time and space, using the shifted Legendre operational matrix. The operational matrix of fractional differentiation in Caputo sense $D^{(\sigma)}$ have been used via the spectral Tau method for reducing the space fractional diffusion equation into fractional ordinary differential equation (FODE) in time that can be solved by fractional linear multi-step method (FLMM). Two numerical example reveals that the proposed method is very accurate and efficace for solving this problem when compared with exact solution.

References

- [1] **A.A.Kilbas, H.M.Srivastava, J.J.Trujillo**, *Theory and Applications of Fractional Differential Equations*, Elsevier, San Diego, 2006.
- [2] **A.Saadatmandia, M.Deaghan**, *A new operational matrix for solving fractional-order differential equations*, Elsevier, San Diego, 59 (2010) 1326-1336, 2009.
- [3] **K.B.Oldham, J.Spanier**, *The Fractional Calculus*, Academic Press, New York, 1974.
- [4] **K.S.Miller, B.Ross**, *An Introduction to The Fractional Calculus and Fractional Differential Equations*, Wiley, New York, 1993.
- [5] **I.Podlubny**, *Fractional Differential Equations*, Academic Press, San Diego, 1999.
- [6] **J.H.He**, *Some applications of nonlinear fractional differential equations and their approximations*, *Bull.Sci.Technol.*15(2)(1999)8690.
- [7] **R.L.Bagley, P.J.Torvik**, *A theoretical basis for the application of fractional calculus to viscoelasticity*, *J.Rheology* 27(3) (1983) 201-210.
- [8] **F.Mainardi**, *Fractional calculus: Some basic problems in continuum and statistical mechanics*, in: *A. Carpinteri, F.Mainardi (Eds.), Fractals and Fractional Calculus in Continuum Mechanics*, Springer, Verlag, New York, 1997, pp.291-348.
- [9] **Xue, D.; Bai, L**, *Benchmark problems for Caputo fractional-order ordinary differential equations*. *Fract. Calc. Appl. Anal.* 2017, 20, 1305–1312.
- [10] **Garrappa.R**, *Numerical Solution of Fractional Differential Equations: A Survey and a Software Tutorial*, www.mdpi.com/journal/mathematics. *Mathematics* 2018, 6, 16; doi:10.3390/math6020016. Received: 8 December 2017; Accepted: 14 January 2018; Published: 23 January 2018.
- [11] **Lubich, C**, *Discretized fractional calculus*. *SIAM J. Math. Anal.* 1986, 17, 704–719.
- [12] **Lubich, C**, *Convolution quadrature and discretized operational calculus. I*. *Numer. Math.* 1988, 52, 129–145.
- [13] **Lubich, C**, *Convolution quadrature and discretized operational calculus. II*. *Numer. Math.* 1988, 52, 413–425.
- [14] **Lubich, C**, *Convolution quadrature revisited*. *BIT* 2004, 44, 503–514.
- [15] **Wolkenfelt, P.H.M**, *Linear Multistep Methods and the Construction of Quadrature Formulae for Volterra Integral and Integro-Differential Equations; Technical Report NW 76/79; Mathematisch Centrum: Amsterdam, The Netherlands, 1979*.
- [16] **Wolkenfelt, P.H.M**, *The construction of reducible quadrature rules for Volterra integral and integro-differential equations*. *IMA J. Numer. Anal.* 1982, 2, 131–152.
- [17] **Garrappa, R**, *Trapezoidal methods for fractional differential equations: theoretical and computational aspects*. *Math. Comput. Simul.* 2015, 110, 96–112.
- [18] **Schädle, A.; López-Fernández, M.A.; Lubich, C**, *Fast and oblivious convolution quadrature*. *SIAM J.Sci.Comput.* 2006, 28, 421–438.

-
- [19] **Lubich, C.**, *On the stability of linear multi-step methods for Volterra convolution equations.* *IMA J. Numer. Anal.* 1983, 3, 439–465.
- [20] **Young, A.**, *Approximate product-integration.* *Proc. R. Soc. Lond. Ser. A* 1954, 224, 552–561.
- [21] **Aceto, L.; Magherini, C.; Novati, P.** *On the construction and properties of m -step methods for FDEs.* *SIAM J.Sci. Comput.* 2015, 37, A653–A675.
- [22] **A. Saadatmandi, M. Dehghan,** *A Tau method for the one-dimensional parabolic inverse problem subject to temperature overspecification,* *Comput. Math. Appl.* 52 (2006) 933–940.
- [23] **Samko, S.G.; Kilbas, A.A.; Marichev, O.I.,** *Fractional Integrals and Derivatives;* Gordon and Breach Science Publishers: Yverdon, Switzerland, 1993; p. 976.
- [24] **Diethelm, K.,** *The Analysis of Fractional Differential Equations; Lecture Notes in Mathematics;* Springer: Berlin, Germany, 2010; Volume 2004, p. 247.
- [25] **S.Momani, Z.Odibat,** *Homotopy perturbation method for nonlinear partial differential equations of fractional order,* *Phys.Lett.A* 365 (2007) 345–350.
- [26] **N.H. Sweilam, M.M. Khader, R.F. Al-Bar,** *Numerical studies for a multi-order fractional differential equation,* *Phys. Lett. A* 371 (2007) 26–33.
- [27] **M.M. Khadar, N.H. Sweilam, A.M.S. Mahdy,** *An efficient numerical method for solving the fractional diffusion equation,* *J. Appl. Math. Bioinformatics* 1 (2) (2011) 1–12.
- [28] **M.M. Khader,** *On the numerical solutions for the fractional diffusion equation,* *Commun. Nonlin. Sci. Num. Simul.* (2010), <https://doi.org/10.1016/j.cnsns.2010.09.007>.
- [29] **E.H. Doha, A.H. Bhrawy, D. Baleanu, S.S. Ezz-Eldien,** *The operational matrix formulation of the Jacobi Tau approximation for space fractional diffusion equation,* *Adv.Differ. Equ.* 2014 (2014) 231.
- [30] **R. Magin, X. Feng, D. Baleanu,** *Solving fractional order Bloch equation,* *Concept Magn. Res.* 34A (1) (2009) 16–23.
- [31] **H. Singh, H.M. Srivastava,** *Jacobi collocation method for the approximate solution of some fractional-order Riccati differential equations with variable coefficients,* *Physica A* 523 (2019) 1130–1149.
- [32] **D. Baleanu, B. Shiri, H.M. Srivastava, M. Al-Qurashi,** *A Chebyshev spectral method based on operational matrix for fractional differential equations involving non-singular Mittag-Leffler kernel,* *Adv. Differ. Equ.* 2018 (353) (2018) 1–23.
- [33] **M.M. Khader, K.M. Saad,** *Numerical studies of the fractional Korteweg-de Vries-Burgers' and Burgers' equations,* *Proceed. Nat. Acad. Sci. India Sec. A: Phys. Sci.* (2020).
- [34] **H. Singh, C.S. Singh,** *A reliable method based on second kind Chebyshev polynomial for the fractional model of Bloch equation,* *Alexandria Eng. J.* 57 (2018) 1425–1432.
- [35] **H. Singh, F.A. Ghassabzadeh, E. Tohidi, C. Cattani,** *Legendre spectral method for the fractional Bratu problem,* *Math. Meth. Appl. Sci.* 43 (9) (2020) 5941–5952.
- [36] **Khader, M.M. and Hendy, A.S.,** (2012) *The Approximate and Exact Solutions of the Fractional-Order Delay Differential Equations Using Legendre Pseudo-Spectral Method.* *International Journal of Pure and Applied Mathematics,* 74, 287–297.

-
- [37] **D.Mbainguesse, B Abbo and Youssef.P**, *SOLVING NONLINEAR FRACTIONAL INTEGRO-DIFFERENTIAL EQUATION OF FREDHOLM SECOND KIND BY AN APPROXIMATION TECHNIQUE OF LAGUERRE POLYNOMIALS AND NEWTON ITERATION METHOD*. *Universal Journal of Mathematics and Mathematical Sciences*, Volume 19, Number 1, 2023, Pages 1-19.
- [38] **D.Mbainguesse, D.Luc, B.Abbo and Youssef.P**, *NUMERICAL SOLUTIONS OF NON-LINEAR HEAT TRANSFER MODELLING IN A TWO-DIMENSIONAL SPACE*. *International Journal of Numerical Methods and Applications*, Volume 24, Number 1, 2024, Pages 63-77.