

Assessment of association of Tephritids with different fruits and vegetables in Terai Agro Ecological Region of West Bengal

ABSTRACT

The flies belong to the family Tephritidae of Insect's order Diptera are referred as Fruit flies because they infest a wide variety of fruits and vegetables. Globally, Dacinae fruit flies of the genus *Bactrocera* Macquart are one of the most important pests of fruits and vegetables. A study was conducted during 2021, to assess the association of tephritids with different fruits and vegetables like Ber, Citrus, Guava, Mango, Bottlegourd, Cucumber and Pumpkin. Total Nine species were collected by installing traps comprising of sex attractants such as Cue-lure, Methyl Eugenol and by rearing from Infested fruits. Among Nine, six species were collected through cue-lure traps namely *Zeugodacus tau* (Walker), *Zeugodacus cucurbitae* (Coq.), *Bactrocera rubigina* (Wang and Zhao), *Zeugodacus caudatus* (Fabricus), *Bactrocera divendri* (Maneesh, Hancock and Prabhakar) and *Dacus longicornis* (Wiedemann). Two species were collected through methyl eugenol traps namely *Bactrocera dorsalis* (Hendel) and *Bactrocera zonata* (Saunders). Four species namely *Zeugodacus tau* (Walker), *Zeugodacus cucurbitae* (Coq.), *Bactrocera dorsalis* (Hendel) and *Bactrocera minax* (Enderlein) were recovered from infested fruits of ber, citrus, bottle gourd, cucumber and pumpkin. Species richness in different fields were maximum in citrus and from recovery of infested fruits maximum in ber. Species composition of *Z. tau* were maximum in pumpkin, bottle gourd and *B. dorsalis* were maximum in mango, guava, ber and citrus. Species composition of *Z. tau* were recovered maximum from infested pumpkin and cucumber, *Z. cucurbitae* were recovered maximum from infested bottle gourd and *B. dorsalis* were recovered maximum from infested ber and citrus fruits. Relative abundance of *Z. tau* were recorded maximum in pumpkin and bottle gourd and *B. dorsalis* were recorded maximum in mango, guava, ber and citrus. Relative abundance of *Z. tau* were recovered maximum in pumpkin and cucumber infested fruits, *Z. cucurbitae* were recovered maximum from bottle gourd infested fruits and *B. dorsalis* were recovered maximum from ber and citrus infested fruits. In different fields, Species evenness was observed maximum in citrus and from recovery of infested fruits maximum in ber.

Keywords: Fruit flies, Tephritid, Bactrocera, Species Richness, Composition, Relative abundance, Evenness

1. INTRODUCTION

Fruits and vegetables are grown all over the country along with other agricultural crops. In the world, India ranks 2nd position for the production of fruits and vegetables. It provides additional income to the country. India is a wide country with diverse climatic conditions where all varieties of fruits and vegetables are grown. Vegetables like cucurbits (pumpkin, Bottle gourd, Ridge gourd, cucumber), onion, Ginger, Solanaceous (tomato, okra, chilli, brinjal), crucifers (cabbage, cauliflower, broccoli) are grown. Fruits like apple, pear, plum, mango, guava, ber, papaya, citrus, banana, pomegranate and many more are cultivated. One of the most important challenging in production of fruits and vegetables is ideal yield and quality. It is reduced by the infestation of Insect pests and diseases. Many insect pests like lepidopteran, coleopteran, dipteran and many more attack the fruits and vegetables. Among them, the most destructive insect pest of fruits and vegetables in the world are fruit flies (Vayssieres *et al.*, 2008; Ekiesiet *et al.*, 2009). The Fruit flies belongs to the family Tephritidae of Insect's order Diptera. Due to its cosmopolitan distribution, polyphagous nature, nature of damage, losses and quarantine status have an economic importance (Drew *et al.*, 2007; Kapoor 2006; Ukeyet *et al.*, 2013). The genera *Anastrepha*, *Bactrocera*, *Ceratitis*, *Dacus*, *Rhagoletis* contain the most important species of economic importance on horticultural crops

worldwide (White and Elson-Harris 1992). They attack wide varieties of wild and commercial vegetables and fruits causing considerable damage to the crops (De Meyer *et al.*, 2012).Tephritids causes fruit dropping and making them inedible and associated with quarantine restrictions that are imposed by importing countries to prevent the entry and establishment of exotic fruit fly species (Ekesi, 2012).

In India 200 species of tephritids (Fruit flies) are recorded which is about 5% of world's tephritid fauna (Verghese *et al.*, 2002).Fruit flies species attack many families belong to fruits and vegetables. About 36% of the tephritid species infest soft fruits, which consist of various commercially cultivated vegetables and fruits (White and Elson-Harris 1992). *Z. cucurbitae*, *B. dorsalis*, *Z. tau* are the serious pests of fruits and vegetables which destroys the growth and development.

Bactrocera dorsalis infests more than 46 hosts including many fruit crops grown commercially like cucurbits, guava, mango, papaya, avocado, papaya (De Meyer 2010; Cugalaet *al.*, 2014; Mwatawalaet *al.*, 2006; Jose and Santos, 2013). *Bactrocera dorsalis* complex has various host range belongs to 38 families, 75 genera, 118 species (Allwood *et al.*, 1999).It attacks many families of vegetables and fruits like Cucurbitaceae, Caricaceae, Rutaceae, Fabaceae, Malvaceae, Myrtaceae, Sapotaceae, Rhamnaceae, Solanaceae, Moraceae, Vitaceae (Tsuruta *et al.*, 1997).*Bactrocera tau* infested fruit species belong to families Leguminosae, Rutaceae, Sapotaceae, Myrtaceae, Solanaceae, Moraceae (Allwood *et al.*, 1999; Boopathi *et al.*, 2013a). *B. tau* infests tomato fruits causing quality reduction and fruit abortion (Gupta *et al.*, 1990; Boopathi *et al.*, 2013a). *B.cucurbitae*, *Dacus punctatifrons* are also attack tomato fruits. *Bactrocera tau*, *Bactrocera latifrons* infests solanaceous fruits like brinjal, chilli, tomato and *Bactroceradiversa* attacks cucurbits.

Due to wide host range, high dispersal capacity, wide tolerance to climate, polyphagous nature and quick proliferation of fruit flies was difficult to control by using insecticides which cause huge yield losses in vegetables and fruit crops (Baral *et al.*, 2006; Muhammad *et al.*, 2004; Ndiayaet *al.*, 2008).Tephritids attack a various variety of fleshy, soft vegetable and fruit crops and lead to 35 – 81% direct loss all over the season depending on variety and locality (Lux *et al.*, 2003a; Lux *et al.*, 2003b; Ekesiet *al.*, 2009; Otieno, 2009). Infestation caused by tephritid fruit flies outcomes in severe losses in trade value and export opportunity because of strict quarantine regulations enforced by many importing countries (Adhikari *et al.*, 2016; Poudel and Regmi 2008). Crop loss occurs in vegetable crops in India was reported as 30 – 40% (Rai *et al.*, 2014).

Forty three species are known in genera, *Bactrocera* and *Dacus*. Out of thirty six species in 11 sub genera, 17 species of *Bactrocera* are economically significant causing heavy losses to wide varieties of vegetables and fruits in India (Kapoor, 2002).Tephritids cause 100% yield loss in cucurbits as they are quarantine pests (Dhillon *et al.*, 2005).In India, melon fruit fly (*Bactroceracucurbitae*) is a major threat to cucurbits (Shah *et al.*, 1948; Senior – white 1924).Tephritid fruit flies also infest many fruit crops like Mango, Guava, Ber, Papaya, Citrus and many more.About 15-85% loss in mango production by fruit flies has been noted (Anonymous 2013).Loss in guava yield is about 3-40% as reported by Haseeb and Sharma (2007).

More than 75% losses are recorded in ber field due to fruit fly infestation (Nizamani *et al.*, 2015; Saxena and Rawat, 1968) and fruit damage about 55-65% (Sharma *et al.*, 1998) and 90-100% (Vadivlu, 2014; Singh and Vashista, 1985).More than 35% post-harvest losses caused by fruit flies (Musasa *et al.*, 2015).In Northern India, *Bactroceraminax* causes a significant damage in tangerine (Nath, 1973).

2. MATERIAL AND METHOD

2.1 Experimental Site

During 2020 and 2021 the field studies were conducted at the Instructional Farm of Uttar Banga Krishi Viswavidyalaya, Pundibari, Cooch Behar, West Bengal, India. During respective years of studythe laboratory investigations were carried out in the Department of Entomology, UBKV, Pundibari, Cooch Behar.The Instructional Farm located at 26° 19' N latitude and 89°

23° E longitude at an altitude of 43 meter above the Mean Sea Level (MSL). Top soils are dark brown in colour, about 1-3 feet deep, sandy loam in texture, high in raw humus content, porous, low water retention capacity, medium to high nitrogen content, low to medium potassium content but poor in secondary micronutrients. Soil of the region is acidic in reaction and pH of soil ranged from 4.2 to 6.8. About 80 per cent of total rainfall received from the South-West monsoon during the rainy months from June to September. Average maximum and minimum temperatures varied from 24°C to 33.2°C. The whole area is humid and warm except having a short spell of winter during the months of December-February. However, the pattern of rainfall is erratic and not uniformly distributed round the year.

2.2 Collection and Rearing of fruit flies

From different fields tephritid fruit flies are collected by installing traps comprising of sex attractants such as Cue-lure and Methyl Eugenol. Cue-lure and Methyl eugenol lure are prepared in laboratory by soaking cotton wick in a solution containing either cue-lure or methyl eugenol chemical, Cypermethrin and water. Then cotton wick is wrapped by using aluminium foil and fixed in traps. The traps containing para-pheromone about four per each field at a height of 1.5 – 2m with a distance of 20m were placed. From each trap, every day fruit flies were recorded. Tephritids infested fruits were collected from Instructional farm of Uttar Banga Krishi Vishwavidyalaya, Pundibari. Damaged fruits collected from Bottle gourd, Cucumber, Pumpkin, Ber, Citrus field were placed in a plastic or glass tray (42x35x7 cm) containing a dry medium such as loose soil and covered with muslin cloth on top to prevent infestation of another flies which cause secondary infection and trays were kept for rearing in an insect rearing cage. Every two days samples were checked for puparia and fruits from which larva emerged was discarded. After pupation, soil was sieved for collecting pupae. Again pupae were placed in a tray containing soil for adult emergence at least 3 – 4 weeks. Sugar solution as a food was placed in a tray which was used by adult fruit flies for development of full body and coloration which was helpful for identification.

2.3 To assess the association of tephritids with different fruits and vegetables

Fruit fly species collected from traps installed in field and rearing of infested fruits in different crops such as bottle gourd, cucumber, pumpkin, ber, citrus, mango and guava used to analyse the tephritids associated with different vegetables and fruits.

2.3.1 Species richness in different fruits and vegetables

Measure of number of species in a sample (Kishore *et al.*, 2018). More species were found when the sample size is large. Species richness was determined by using Menhinick Index.

$$S$$

$$\sqrt{n}$$

$$\text{Species richness} = \frac{S}{\sqrt{n}}$$

Where S = Number of species in a sample

n = Total number of individuals in a sample

2.3.2 Species composition in different fruits and vegetables:

Species composition was calculated to recognize the similar species present in a locality. It is expressed as percent and all species were added up to 100% (Ndah *et al.*, 2013). It was determined by using density of the species in a locality.

$$\text{Species composition} = \frac{\text{Number of individuals of one species}}{\text{Total number individuals of all species}} \times 100$$

2.3.3 Relative abundance of species in different cucurbits and fruits:

After collection of tephritid fruit flies from para pheromone traps and rearing of infested fruits, 100 fruit flies were collected randomly from each crop. Insects were separated based on species and their relative abundance was calculated based on the formula (Kishore *et al.*, 2018).

$$\text{Relative abundance} = \frac{\text{Number of individuals of one species}}{\text{Total number individuals of all species}} \times 100$$

Total number of individuals of all species

2.3.4 Species evenness of tephritids determined in different fruits and vegetables:

Species evenness was calculated to know the equitability of species in a community. It was estimated by using Pielou's Evenness Index (J) (Pielou 1966).

$$\text{Species evenness (J)} = \frac{H'}{\ln(S)}$$

where H' = Shannor – Weiner Index

S = Number of species in a sample

\ln = Natural logarithm of a number.

The findings of the study have been presented in tabular as well as graphical form as applicable in appropriate manner.

3. RESULTS& DISCUSSION

3.1 To assess the association of tephritids with different fruits and vegetables:

3.1.1 Species richness of fruit flies collected from traps:

Number of species in Bottle gourd, Pumpkin, Ber, Mango, Guavaand Citrus were documented (Kishore *et al.*,2018). If the value is more, then the species richness is more. Observations were presented in the Table 1 and Fig.1.

Table 1: Determiation of species richness in different fields.

Fields Taken Into Consideration	Species Richness (R)
Citrus	0.54
Guava	0.35
Pumpkin	0.29
Bottle gourd	0.29
Mango	0.26
Ber	0.26




Fig. 1. Species Richness in Different Fields

From the undergone research work, data represented in Table 1 and Fig.1. Proves that in different fields such as bottle gourd, pumpkin, ber, mango, guava and citrus, species richness is determined. In citrus, species richness value is 0.54 followed by in guava 'R' is 0.35, in pumpkin and bottle gourd 'R' is 0.29, in mango and ber 'R' is 0.26. As the value is more, then species richness is maximum. Species richness is greater in citrus and least in mango and ber.

3.1.1.1 Species richness of fruit flies recovered from infested fruits:

Number of species in Bottle gourd, Cucumber, Pumpkin, Ber and Citrus were documented (Kishore *et al.*,2018). If the obtained value is more, then the species richness is more. Observations were presented in the Table 2 and Fig.2.

Table 2: Determination of species richness from recovery of infested fruits.

Fruit Flies Infested Fruits	Species Richness (R)
Ber	0.38
Cucumber	0.30
Citrus	0.27
Bottle gourd	0.18
Pumpkin	0.14




Fig. 2. Species Richness in Infested Fruits

From the undergone research work, data represented in Table 2 and Fig.2. Proves that from recovery of infested fruits such as bottle gourd, cucumber, pumpkin, ber, and citrus, species richness is determined. In ber, species richness value is 0.38 followed by in cucumber 'R' is 0.30, in citrus 'R' is 0.27, in bottle gourd 'R' is 0.18 and in pumpkin 'R' is 0.14. As the value is more, then species richness is maximum. Species richness is greater in ber and least in pumpkin.

3.1.2 Species composition of fruit flies collected from traps:

Number of similar individuals recorded in different fields were documented. It was expressed in percentage (Ndah *et al.*,2013). If the obtained value is more, then the composition of that species is more. Observations were presented for different fields in the tables and figures.

3.1.2.1 Species composition of fruit flies in pumpkin:

Table 3: Determination of species composition of tephritids in pumpkin collected by traps.

Name of the Species	Species Composition (%)
<i>Zeugodacus tau</i> (Walker)	48.1
<i>Bactrocera dorsalis</i> (Hendel)	19.6
<i>Zeugodacuscucurbitae</i> (Coq.)	17.7
<i>Bactrocerarubigina</i> (Wang and Zhao)	6.3

<i>Zeugodacus caudatus</i> (Fabricus)	3.1
<i>Bactrocerazonata</i> (Saunders)	2.2
<i>Bactroceradivenderi</i> (Maneesh, Hancock and Prabhakar)	1.3
<i>Dacus longicornis</i> (Weidemann)	1.2




Fig. 3. Species Composition (%) in Pumpkin

Overall information from an experiment Table 3 and Fig.3. reveals as species composition in pumpkin is determined in percent. In pumpkin, *Zeugodacus tau* recorded about 48.1% followed by *Bactrocera dorsalis* (19.6%), *Zeugodacuscucurbitae* (17.7%), *Bactrocerarubigina* (6.3%), *Zeugodacuscaudatus* (3.1%), *Bactrocerazonata* (2.2%), *Bactroceradivenderi*(1.3%) and *Dacus longicornis* (1.2%). As the percentage is more, then the species composition of that particular species is more. So, Species composition of *Zeugodacustau* in pumpkin is more and minimum is *Dacus longicornis*.

3.1.2.2 Species composition of fruit flies in Bottle gourd:

Table 4: Determination of species composition of tephritids in Bottle gourd collected by traps.

Name of the Species	Species Composition (%)
<i>Zeugodcus tau</i> (Walker)	36.6
<i>Zeugodacuscucurbitae</i> (Coq.)	26.9
<i>Bactrocera dorsalis</i> (Hendel)	25.7
<i>Bactrocerarubigina</i> (Wang and Zhao)	4.5
<i>Bactrocerazonata</i> (Saunders)	2.4
<i>Zeugodacus caudatus</i> (Fabricus)	1.6
<i>Bactroceradivenderi</i> (Maneesh, Hancock and Prabhakar)	1.2
<i>Dacus longicornis</i> (Weidemann)	0.9




Fig 4: Species Composition (%) in Bottle gourd

Inclusive information from an research is represented in Table 4 and Fig.4. shown as species composition in bottle gourd is determined in percent. In bottle gourd, *Zeugodacus tau* recorded about 36.6% followed by *Zeugodacuscucurbitae* (26.9%), *Bactrocera dorsalis* (25.7%), *Bactrocerarubigina* (4.5%), *Bactrocerazonata*(2.4%), *Zeugodacus caudatus*(1.6%), *Bactroceradivenderi*(1.2%) and *Dacus longicornis* (0.9%). As the percentage is more, then the species composition of that particular species is more. So, Species composition of *Zeugodacustau* in bottle gourd is more and minimum is *Dacus longicornis*.

3.1.2.3 Species composition of fruit flies in Mango:

Table 5: Determination of species composition of tephritids in Mango collected by traps.

Name of the Species	Species Composition (%)
<i>Bactrocera dorsalis</i> (Hendel)	34.8
<i>Zeugodacustau</i> (Walker)	33.3
<i>Zeugodacuscucurbitae</i> (Coq.)	16.3
<i>Zeugodacus caudatus</i> (Fabricus)	5.7
<i>Bactrocerarubigina</i> (Wang and Zhao)	4.7
<i>Bactrocera zonata</i> (Saunders)	3.2
<i>Bactroceradivenderi</i> (Maneesh, Hancock and Prabhakar)	1.6




Fig. 5. Species Composition (%) inMango

Inclusive information from a research is represented in Table 5 and Fig.5. shown as species composition in mango is determined in percent. In mango, *Bactrocera dorsalis* recorded about 34.8% followed by *Zeugodacus tau* (33.3%), *Zeugodacuscucurbitae* (16.3%), *Zeugodacuscudatus*(5.7%), *Bactrocerarubigina*(4.7%), *Bactrocerazonata*(3.2%), *Bactroceradivenderi*(1.6%). As the percentage is more, then the species composition of that

particular species is greater. So, Species composition of *Bactrocera dorsalis* in mango is maximum and minimum is *Bactrocera divendri*.

3.1.2.4 Species composition of fruit flies in Guava:

Table 6: Determination of species composition of tephritids in Guava collected by traps.

Name of the Species	Species Composition (%)
<i>Bactrocera dorsalis</i> (Hendel)	50.3
<i>Zeugodacus tau</i> (Walker)	18.9
<i>Bactrocera rubigina</i> (Wang and Zhao)	7.9
<i>Bactrocera zonata</i> (Saunders)	7.3
<i>Zeugodacus caudatus</i> (Fabricus)	6.8
<i>Zeugodacus cucurbitae</i> (Coq.)	6.5
<i>Bactrocera divendri</i> (Maneesh, Hancock and Prabhakar)	1.8




Fig. 6. Species Composition (%) in Guava

After an experiment, data is represented in Table 6 and Fig. 6. shown as species composition in guava is determined in percent. In guava, *Bactrocera dorsalis* recorded about 50.3% followed by *Zeugodacus tau* (18.9%), *Bactrocera rubigina* (7.9%), *Bactrocera zonata* (7.3%), *Zeugodacus caudatus* (6.8%), *Zeugodacus cucurbitae* (6.5%), *Bactrocera divendri* (1.8%). As the percentage is more, then the species composition of that particular species is greater. So, Species composition of *Bactrocera dorsalis* in guava is maximum and minimum is *Bactrocera divendri*.

3.1.2.5 Species composition of fruit flies in Ber:

Table 7: Determination of species composition of tephritids in Ber collected by traps.

Name of the Species	Species Composition (%)
<i>Bactrocera dorsalis</i> (Hendel)	39.0
<i>Zeugodacus tau</i> (Walker)	23.9
<i>Zeugodacus cucurbitae</i> (Coq.)	20.4
<i>Zeugodacus caudatus</i> (Fabricus)	5.2
<i>Bactrocera zonata</i> (Saunders)	4.7
<i>Bactrocera rubigina</i> (Wang and Zhao)	4.6
<i>Bactrocera divendri</i> (Maneesh, Hancock and Prabhakar)	1.9




Fig. 7. Species Composition (%) in Ber

Overall information from an experiment Table 7 and Fig.7. reveals as species composition in ber is determined in percent. In ber, *Bactrocera dorsalis* recorded about 39.0% followed by *Zeugodacustau* (23.9%), *Zeugodacuscucurbitae* (20.4%), *Zeugodacus caudatus* (5.2%), *Bactrocerazonata* (4.7%), *Bactrocerarubigina* (4.6%) *Bactroceradivenderi*(1.3%) As the percentage is more, then the species composition of that particular species is more. So, Species composition of *Bactroceradorsalis* in ber is more and minimum is *Bactroceradivenderi*.

3.1.2.6 Species composition of fruit flies in Citrus

Table 8: Determination of species composition of tephritids in Citrus collected by traps.

Name of the Species	Species Composition (%)
<i>Bactrocera dorsalis</i> (Hendel)	45.7
<i>Zeugodacus tau</i> (Walker)	22.2
<i>Zeugodacuscucurbitae</i> (Coq.)	19.1
<i>Zeugodacus caudatus</i> (Fabricius)	7.3
<i>Bactrocerazonata</i> (Saunders)	5.7




Fig. 8. Species Composition (%) in Citrus

Inclusive information from a research is represented in Table 8 and Fig.8. shown as species composition in citrus is determined in percent. In citrus, *Bactrocera dorsalis* recorded about 45.7% followed by *Zeugodacus tau* (22.2%), *Bactroceracucurbitae*(19.1%), *Zeugodacuscudatus*(7.3%), *Bactrocerazonata*(5.7%). As the percentage is more, then the species composition of that particular species is greater. So, Species composition of *Bactroceradorsalis* in citrus is maximum and least recorded is *Bactrocerazonata*.

3.1.3 Species composition of fruit flies recovered from infested fruits:

Number of similar individuals recorded from recovery of infested fruits were documented. It was expressed in percentage (Ndah *et al.*,2013). If the obtained value is more, then the composition of that species is more. Observations were presented in the tables and figures.

3.1.3.1 Species composition of fruit flies in Pumpkin:

Table 9: Determination of species composition of tephritids recovered from pumpkin.

Name of the Species	Species Composition (%)
<i>Zeugodacus tau</i> (Walker)	49.8
<i>Zeugodacuscucurbitae</i> (Coq.)	46.5
<i>Bactrocera dorsalis</i> (Hendel)	3.5




Fig. 9. Species Composition (%) inPumpkin

Through the experiment done and result represented in Table 9 and Fig.9. concludes as species composition of tephritids recovered from pumpkin infested fruit is determined in percent. From infested fruits of pumpkin, *Zeugodacus tau* about 49.8% is recorded followed by *Zeugodacuscucurbitae* (46.5%) and *Bactrocera dorsalis* (3.5%). As the value is more, then the species composition in a particular infested fruit is maximum. Here, species composition is greater in *Zeugodacus tau* and least in *Bactrocera dorsalis*.

3.1.3.2 Species composition of fruit flies in Bottle gourd:

Table 10: Determination of species composition of tephritids recovered from bottle gourd.

Name of the Species	Species Composition (%)
<i>Zeugodacuscucurbitae</i> (Coq.)	48.2
<i>Zeugodacus tau</i> (Walker)	47.4
<i>Bactrocera dorsalis</i> (Hendel)	4.3




Fig. 10. Species Composition (%) in Bottle Gourd

Through the research work and data presented in Table 10 and Fig.10. explains as species composition of tephritids recovered from bottle gourd infested fruit is determined in percent. From infested fruits of bottle gourd, *Zeugodacuscucurbitae* about 48.2% is recorded followed by *Zeugodacus tau* (47.4%) and *Bactrocera dorsalis* (4.3%). As the value is more, then the species composition in a particular infested fruit is maximum. Here, species composition is greater in *Zeugodacuscucurbitae* and least in *Bactrocera dorsalis*.

3.1.3.3 Species composition of fruit flies in Cucumber:

Table 11: Determination of species composition of tephritids recovered from cucumber.

Name of the Species	Species Composition (%)
<i>Zeugodacus tau</i> (Walker)	40.4
<i>Zeugodacuscucurbitae</i> (Coq.)	39.3
<i>Bactrocera dorsalis</i> (Hendel)	20.2




Fig. 11. Species Composition (%) in Cucumber

Data recorded from experiment is represented in Table 11 and Fig.11 concludes as species composition of tephritids recovered from cucumber infested fruit is determined in percent. From infested fruits of cucumber, *Zeugodacus tau* about 40.4% is recorded followed by *Zeugodacuscucurbitae*(39.3%) and *Bactrocera dorsalis* (20.2%). As the value is more, then the species composition in a particular infested fruit is maximum. Here, species composition is greater in *Zeugodacus tau* and minimum in *Bactrocera dorsalis*.

3.1.3.4 Species composition of fruit flies in Ber:

Table 12: Determination of species composition of tephritids recovered from Ber.

Name of the Species	Species Composition (%)
<i>Bactrocera dorsalis</i> (Hendel)	35.4
<i>Zeugodacus tau</i> (Walker)	33.8
<i>Zeugodacuscucurbitae</i> (Coq.)	30.6




Fig. 12. Species Composition (%) inBer

Through the research work and data presented in Table 12 and Fig.12. explains as species composition of tephritids recovered from ber infested fruit is determined in percent. From infested fruits of ber, *Bactrocera dorsalis* about 35.4% is recorded followed by *Zeugodacus tau* (33.8%) and *Zeugodacuscucurbitae* (30.6%). As the value is more, then the species composition in a particular infested fruit is extreme. Here, species composition is greater in *Bactrocera dorsalis* and least in *Zeugodacuscucurbitae*.

3.1.3.5 Species composition of fruit flies in Citrus:

Table 13: Determination of species composition of tephritids recovered from Citrus.

Name of the Species	Species Composition (%)
<i>Bactrocera dorsalis</i> (Hendel)	55.7
<i>Bactroceraminax</i> (Enderlein)	44.2




Fig. 13. Species Composition (%) inCitrus

Data recorded from research work and is represented in Table 13 and Fig.13. describes as species composition of tephritids recovered from citrus infested fruit is determined in percent. From infested fruits of citrus, *Bactrocera dorsalis* about 55.7% is recorded followed by

Bactroceraminax(44.2%) As the value is more, then the species composition in a particular infested fruit is greater. Here, species composition is greater in *Bactrocera dorsalis* and minimum recorded in *Bactroceraminax*.

3.1.4. Relative abundance of fruit flies collected from traps:

Abundance of individuals in a species in different fruits and vegetables were documented (Kishore *et al.*,2018). 100 individuals were selected randomly from each crop and recorded the abundance. It is expressed in percent and if the value is more, then the abundance of that species is more. Observations were presented in the Tables and Figures.

3.1.4.1 Relative abundance of fruit flies in Pumpkin:

Table 14: Determination of relative abundance of tephritids in Pumpkin collected by traps.

Name of the Species	Relative Abundance (%)
<i>Zeugodacus tau</i> (Walker)	33
<i>Bactrocera dorsalis</i> (Hendel)	23
<i>Zeugodacuscucurbitae</i> (Coq.)	15
<i>Bactrocerarubigina</i> (Wang and Zhao)	9
<i>Zeugodacus caudatus</i> (Fabricus)	7
<i>Bactrocerazonata</i> (Saunders)	6
<i>Bactroceradivenderi</i> (Maneesh, Hancock and Prabhakar)	5
<i>Dacus longicornis</i> (Weidemann)	2




Fig. 14. Relative Abundance (%) in Pumpkin

Overall information from an experiment Table 14 and Fig.14. declared as relative abundance in pumpkin is determined in percent. About 100 fruit flies randomly taken and separated the species by this relative abundance is estimated. In pumpkin, *Zeugodacus tau* documented about 33% followed by *Bactrocera dorsalis* (23%), *Zeugodacuscucurbitae* (15%), *Bactrocerarubigina* (9%), *Zeugodacus caudatus* (7%), *Bactrocerazonata* (6%), *Bactroceradivenderi*(5%) and *Dacus longicornis* (2%) As the percentage is maximum then the relative abundance of that particular species is more. So, relative abundance of *Zeugodacustau* in pumpkin is more and least documented is *Dacus longicornis*.

3.1.4.2 Relative abundance of fruit flies in Bottle gourd

Table 15: Determination of relative abundance of tephritids in Bottle gourd collected by traps.

Name of the Species	Relative Abundance (%)
<i>Zeugodacus tau</i> (Walker)	29
<i>Zeugodacuscucurbitae</i> (Coq.)	23
<i>Bactrocera dorsalis</i> (Hendel)	15
<i>Bactrocerarubigina</i> (Wang and Zhao)	11
<i>Bactrocerazonata</i> (Saunders)	9
<i>Zeugodacus caudatus</i> (Fabricus)	6
<i>Bactroceradivenderi</i> (Maneesh, Hancock and Prabhakar)	4
<i>Dacus longicornis</i> (Weidemann)	3




Fig. 15. Relative Abundance (%) in Bottle Gourd

From the Table 15 and Fig.15. based on data of an experiment performed concluded as relative abundance in bottle gourd is determined in percent. About 100 fruit flies randomly taken and separated the species by this relative abundance is estimated. In bottle gourd, *Zeugodacus tau* documented about 29% followed by *Zeugodacuscucurbitae*(23%), *Bactrocera dorsalis* (15%), *Bactrocerarubigina* (11%), *Bactrocerazonata*(9%), *Zeugodacus caudatus*(6%), *Bactroceradivenderi*(4%) and *Dacus longicornis* (3%) As the percentage is maximum then the relative abundance of that particular species is more. So, relative abundance of *Zeugodacustau* in bottle gourd is more and least documented is *Dacus longicornis*.

3.1.4.3 Relative abundance of fruit flies in Mango:

Table 16: Determination of relative abundance of tephritids in Mango collected by traps.

Name of the Species	Relative Abundance (%)
<i>Bactrocera dorsalis</i> (Hendel)	39
<i>Zeugodacus tau</i> (Walker)	30
<i>Zeugodacuscucurbitae</i> (Coq.)	12
<i>Zeugodacus caudatus</i> (Fabricus)	9
<i>Bactrocerarubigina</i> (Wang and Zhao)	5
<i>Bactrocerazonata</i> (Saunders)	3
<i>Bactroceradivenderi</i> (Maneesh, Hancock and Prabhakar)	2




Fig. 16. Relative Abundance (%) in Mango

All-encompassing information from a research is represented in Table 16 and Fig.16. shown as relative abundance in mango is determined in percent. About 100 fruit flies randomly taken and separated the species by this relative abundance is estimated. In mango, *Bactrocera dorsalis* recorded about 39% followed by *Zeugodacus tau* (30%), *Zeugodacuscucurbitae* (12%), *Zeugodacuscaudatus*(9%), *Bactrocerarubigina*(5%), *Bactrocerazonata*(3%),*Bactroceradivenderi*(2%). As the percentage is more, then the relative abundance of that particular species is greater. So, relative abundance of *Bactroceradorsalis* in mango is extreme and least documented is *Bactroceradivenderi*.

3.1.4.4 Relative abundance of fruit flies in Guava:

Table 17: Determination of relative abundance of tephritids in Guava collected by traps.

Name of the Species	Relative Abundance (%)
<i>Bactrocera dorsalis</i> (Hendel)	32
<i>Zeugodacus tau</i> (Walker)	19
<i>Bactrocerarubigina</i> (Wang and Zhao)	13
<i>Bactrocerazonata</i> (Saunders)	12
<i>Zeugodacus caudatus</i> (Fabricus)	10
<i>Zeugodacuscucurbitae</i> (Coq.)	8
<i>Bactroceradivenderi</i> (Maneesh, Hancock and Prabhakar)	6




Fig. 17. Relative Abundance (%) in Guava

After an investigation, data is represented in Table 17 and Fig.17. shown as relative abundance in guava is determined in percent. About 100 fruit flies randomly taken and

separated the species by this relative abundance is estimated. In guava, *Bactrocera dorsalis* documented about 32% followed by *Zeugodacus tau* (19%), *Bactrocerarubigina*(13%), *Bactrocerazonata*(12%), *Zeugodacus caudatus*(10%), *Zeugodacuscucurbitae*(8%), *Bactroceradivenderi*(6%). As the percentage is maximum then the relative abundance of that particular species is greater. So, relative abundance of *Bactroceradorsalis* in guava is maximum and minimum is *Bactroceradivenderi*.

3.1.4.5 Relative abundance of fruit flies in Ber:

Table 18: Determination of relative abundance of tephritids in Ber collected by traps.

Name of the Species	Relative Abundance (%)
<i>Bactrocera dorsalis</i> (Hendel)	35
<i>Zeugodacus tau</i> (Walker)	22
<i>Zeugodacuscucurbitae</i> (Coq.)	18
<i>Zeugodacus caudatus</i> (Fabricus)	9
<i>Bactrocerazonata</i> (Saunders)	8
<i>Bactrocerarubigina</i> (Wang and Zhao)	5
<i>Bactroceradivenderi</i> (Maneesh, Hancock and Prabhakar)	3




Fig. 18. Relative Abundance (%) in Ber

Data from an experiment Table 18 and Fig.18. reported as relative abundance in ber is determined in percent. About 100 fruit flies randomly taken and separated the species by this relative abundance is estimated. In ber, *Bactrocera dorsalis* recorded about 35% followed by *Zeugodacus tau* (22%), *Zeugodacuscucurbitae* (18%), *Zeugodacus caudatus* (9%), *Bactrocerazonata* (8%), *Bactrocerarubigina* (5%) *Bactroceradivenderi*(3%) As the percentage is greater, then the relative abundance of that particular species is more. So, relative abundance of *Bactroceradorsalis* in ber is more and least recorded is *Bactroceradivenderi*.

3.1.4.6 Relative abundance of fruit flies in Citrus:

Table 19: Determination of relative abundance of tephritids in Citrus collected by traps.

Name of the Species	Relative Abundance (%)
<i>Bactrocera dorsalis</i> (Hendel)	43

<i>Zeugodacus tau</i> (Walker)	24
<i>Zeugodacuscucurbitae</i> (Coq.)	20
<i>Zeugodacus caudatus</i> (Fabricus)	8
<i>Bactrocera zonata</i> (Saunders)	5




Fig. 19. Relative Abundance (%) in Citrus

Inclusive information from a research is represented in Table 19 and Fig.19. shown as relative abundance in citrus is determined in percent. About 100 fruit flies randomly taken and separated the species by this relative abundance is estimated. In citrus, *Bactrocera dorsalis* recorded about 43% followed by *Zeugodacus tau* (24%), *Zeugodacuscucurbitae*(20%), *Zeugodacus caudatus*(8%), *Bactrocera zonata*(5%). As the percentage is more, then the species composition of that particular species is greater. So, relative abundance of *Bactrocera dorsalis* in citrus is maximum and least recorded is *Bactrocera zonata*.

3.1.5 Relative abundance of fruit flies recovered from infested fruits:

Abundance of individuals in a species recovered from infested fruits were documented (Kishore *et al.*,2018). It is expressed in percent and the abundance of species is more if the value is more. Observations were presented in the Tables and Figures.

3.1.5.1 Relative abundance of fruit flies in Pumpkin:

Table 20: Determination of relative abundance of tephritids recovered from pumpkin infested fruit

Name of the Species	Relative Abundance (%)
<i>Zeugodacus tau</i> (Walker)	21
<i>Zeugodacuscucurbitae</i> (Coq.)	18
<i>Bactrocera dorsalis</i> (Hendel)	11




Fig. 20. Relative Abundance (%) in Pumpkin

Through the experiment done and result represented in Table 20 and Fig.20.concludes as relative abundance of tephritids recovered from pumpkin infested fruit is determined in percent. About 50 fruit flies are randomly collected and segregated different species by this relative abundance is estimated. From infested fruits of pumpkin, *Zeugodacus tau* about 21% is recorded followed by *Zeugodacuscucurbitae* (18%) and *Bactrocera dorsalis* (11%). As the value is more, then the relative abundance in a particular infested fruit is maximum. Here, relative abundance is greater in *Zeugodacus tau* and least in *Bactrocera dorsalis*.

3.1.5.2 Relative abundance of fruit flies in Bottle gourd:

Table 21: Determination of relative abundance of tephritids recovered from Bottle gourd infested fruit

Name of the Species	Relative Abundance (%)
<i>Zeugodacuscucurbitae</i> (Coq.)	24
<i>Zeugodacus tau</i> (Walker)	21
<i>Bactrocera dorsalis</i> (Hendel)	5




Fig. 21. Relative Abundance (%) in Bottle gourd

Through the research work and data presented in Table 21 and Fig.21. explains as relative abundance of tephritids recovered from bottle gourd infested fruit is determined in percent. About 50 fruit flies are randomly collected and segregated different species by this relative abundance is estimated. From infested fruits of bottle gourd, *Zeugodacuscucurbitae* about 24% is recorded followed by *Zeugodacus tau* (21%) and *Bactrocera dorsalis* (5%). As the

value is more, then the relative abundance in a particular infested fruit is maximum. Here, relative abundance is greater in *Zeugodacuscucurbitae* and least documented in *Bactrocera dorsalis*.

3.1.5.3 Relative abundance of fruit flies in Cucumber:

Table 22: Determination of relative abundance of tephritids in recovered from Cucumber infested fruit

Name of the Species	Relative Abundance (%)
<i>Zeugodacus tau</i> (Walker)	22
<i>Zeugodacuscucurbitae</i> (Coq.)	18
<i>Bactrocera dorsalis</i> (Hendel)	10




Fig. 22. Relative Abundance (%) in Cucumber

Data recorded from experiment is represented in Table 22 and Fig.22. concludes as relative abundance of tephritids recovered from cucumber infested fruit is determined in percent. About 50 fruit flies are randomly collected and segregated different species by this relative abundance is estimated. From infested fruits of cucumber, *Zeugodacus tau* about 22% is recorded followed by *Zeugodacuscucurbitae*(18%) and *Bactrocera dorsalis* (10%). As the value is more, then the relative abundance in a particular infested fruit is maximum. Here, relative abundance is greater in *Zeugodacus tau* and minimum in *Bactrocera dorsalis*.

3.1.5.4 Relative abundance of fruit flies in Ber:

Table 23: Determination of relative abundance of tephritids recovered from Ber infested fruit.

Name of the Species	Relative Abundance (%)
<i>Bactrocera dorsalis</i> (Hendel)	20
<i>Zeugodacus tau</i> (Walker)	19
<i>Zeugodacuscucurbitae</i> (Coq.)	11




Fig. 23. Relative Abundance (%) in Ber

Through the research work and data presented in Table 23 and Fig.23, explains as relative abundance of tephritids recovered from ber infested fruit is determined in percent. About 50 fruit flies are randomly collected and segregated different species by this relative abundance is estimated. From infested fruits of ber, *Bactrocera dorsalis* about 20% is recorded followed by *Zeugodacus tau* (19%) and *Zeugodacuscucurbitae* (11%). As the value is more, then the relative abundance in a particular infested fruit is extreme. Here, relative abundance is greater in *Bactrocera dorsalis* and least in *Zeugodacuscucurbitae*.

3.1.5.5 Relative abundance of fruit flies in Citrus:

Table 24: Determination of relative abundance of tephritids recovered from Citrus infested fruit

Name of the Species	Relative Abundance (%)
<i>Bactrocera dorsalis</i> (Hendel)	28
<i>Bactroceraminax</i> (Enderlein)	22




Fig. 24. Relative Abundance (%) in Citrus

Data recorded from research work and is represented in Table 24 and Fig.24. describes as relative abundance of tephritids recovered from citrus infested fruit is determined in percent. About 50 fruit flies are randomly collected and segregated different species by this relative abundance is estimated. From infested fruits of citrus, *Bactrocera dorsalis* about 28% is documented followed by *Bactroceraminax*(22%) As the value is maximum then the relative abundance in a particular infested fruit is greater. Here, relative abundance is greater in *Bactrocera dorsalis* and minimum recorded in *Bactroceraminax*.

3.1.6 Species evenness of tephritids determined in different fruits and vegetables:

3.1.6.1 Species evenness of fruit flies collected from traps:

Evenness of species in different fields were documented (Pielou 1966) and the species evenness is maximum when the value obtained is one and no species evenness observed when the value is zero. Observations were presented in the Table 25 and Fig.25.

Table 25: Estimation of Species evenness of tephritids collected by traps.

Field taken into consideration	Species Evenness (J)
Citrus	0.901
Ber	0.793
Pumpkin	0.792
Guava	0.761
Mango	0.726
Bottle gourd	0.701




Fig. 25. Species Evenness (J) in different fields

After the experiment conducted, the data is recorded in Table 25 and Fig.25. concludes that species evenness in different fields such as pumpkin, bottle gourd, mango, ber, guava and citrus is determined. In citrus species evenness is 0.901 followed by in ber 'J' is 0.793, in pumpkin 'J' is 0.792, in guava 'J' is 0.761, in mango 'J' is 0.726 and in bottle gourd 'J' is 0.701. As the value obtained is 1 then there is species evenness in particular location. Here, Species evenness or equitability is seen in citrus as near to one and no evenness in ber, pumpkin, guava, mango and bottle gourd as the value is not equal to one.

3.1.6.2 Species evenness of fruit flies recovered from infested fruits:

Evenness of species recovered from infested fruits were documented (Pielou 1966) and If the value obtained is one then there is species evenness and no species evenness observed when the value is zero. Observations were presented in the Table 26 and Fig.26.

Table 26: Estimation of Species evenness of tephritids recovered from infested fruits.

Field taken into consideration	Species Evenness (J)
Ber	1.004
Citrus	0.988
Cucumber	0.960
Bottle gourd	0.769
Pumpkin	0.751




Fig. 26. Species Evenness (J) in Infested Fruits

After the experiment conducted, the data is recorded in Table 26 and Fig.26. concludes that species evenness from recovered infested fruits such as pumpkin, bottle gourd, cucumber, ber and citrus is determined. In ber species evenness is 1.004 followed by in citrus 'J' is 0.988, in cucumber 'J' is 0.960, in bottle gourd 'J' is 0.769 and in pumpkin 'J' is 0.751. As the value obtained is 1 then there is species evenness in particular location. Here, species evenness or equitability is seen in ber as it is equal to one and no species evenness or equitability in citrus, pumpkin, cucumber and bottle gourd.

4. CONCLUSION

Among selected fields, Species richness is maximum in citrus and minimum in mango and ber. From the recovery of fruits, Species richness is maximum in ber and least in pumpkin. Species composition of *Z. tau* is maximum and *D. longicornis* is minimum in pumpkin and bottle gourd and *B. dorsalis* is maximum and *B. divenderi* is minimum in mango, ber and guava. In citrus *B. dorsalis* is maximum and *B. zonata* is minimum. Species composition of *Z. tau* from recovery of pumpkin and cucumber, *Z. cucurbitae* from recovery of bottle gourd and *B. dorsalis* from recovery of ber and citrus are maximum. In pumpkin, cucumber and bottle gourd *B. dorsalis* is minimum and in ber *Z. cucurbitae* is minimum and in citrus *B. minax* is minimum. Relative abundance of *Z. tau* is greater and *D. longicornis* is least in pumpkin and bottle gourd, *B. dorsalis* is greater and *B. divenderi* is least in mango, guava and ber. In citrus *B. dorsalis* is maximum and *B. zonata* is minimum. From the recovery of fruits, relative abundance of *Z. tau* in pumpkin and cucumber, *Z. cucurbitae* in bottle gourd, *B. dorsalis* in ber and citrus is maximum. In pumpkin, cucumber, bottle gourd *B. dorsalis* is minimum and *Z. cucurbitae* and *B. minax* is minimum in ber and citrus respectively. Species evenness or equitability is observed in citrus and no evenness in pumpkin, bottle gourd, ber, mango and citrus. From the infested fruits, Species evenness or equitability is observed in ber and no evenness in citrus, bottle gourd, pumpkin and cucumber. Thus, from the aforementioned summarised findings of the present investigation it may be inferred that a good number of tephritid species are available in the agro-ecosystem of northern tract of West Bengal. In cucurbitaceous vegetables the predominant tephritid species found to infest is *Z. tau*. Relative abundance of a number of species differed from vegetation to vegetation and type of parapheromone used for collection. Other population indices are also indicating that trend. Hence,

for efficient management of tephritid flies, use of pheromone traps may be accommodated depending on the crop on which the tephritid flies caused infestation on fruits.

REFERENCES

1. Adhikari, D., Dougoud, J. W., & Schaffner, U. (2016). Assessment of performance of Plant Clinic in Nepal (A case of fruit fly management). MAS ICM thesis, Universite de Neuchate, Switzerland.
2. Allwood, A. J., A. Chinajariyawong, R. A. I. Drew, E. L. Hamacek, D. L. Hancock, C. Hengsawad, J. C. Jinapin, M. Jirasurat, C. K. Krong, S. Kritsaneepaiboon, C. T. S. Leong and S. Vijaysegaran (1999) Host plant records for fruit flies (Diptera: Tephritidae) in South East Asia. *Raffles Bull. Zool. Suppl.* 7: 1–92.
3. Anonymous (2013) CTA practical guide series, No. 14. Available via http://publications.cta.int/media/publications/downloads/1770_pdf.pdf. Accessed 23 June 2016.
4. Baral, K., Roy, B.C., Rahim, K.M.B., Chatterjee, H., Mondal, P., Mondal, D., Ghosh, D. and Talekar, N.S. (2006). Socio-economic parameters of pesticide use and assessment of impact of an IPM strategy for the control of egg plant fruit and shoot borer in West Bengal, India. Technical Bulletin No. 37. AVRDC publication number 06-673. AVRDC – The world Vegetable Centre, Shanha, Taiwan. 36 pp.
5. Boopathi, T., Singh, S.B., Sithanatham S. (2013a). Exploratory bio-ecology studies on fruit flies, *Bactrocera* spp. in Mizoram. *Hexapoda* 20:83-87.
6. Cugula, D., Ekesi, S., Ambasse, D., Adamu, R. and Mohamed, S. (2014). Assessment of ripening stages of Cavendish dwarf bananas as host or non-host to *Bactrocera invadens*. *J. Appl. Entomol.* 138, 449-457.
7. De Meyer, M. (2010). Ecological niche and potential geographic distribution of the invasive fruit fly *Bactrocera invadens* (Diptera, Tephritidae). *Bull. Entomol. Res.* 100, 35-48.
8. De Meyer, M., Mohamed, S. & White, I.M. (2012). Invasive fruit fly pests in Africa: a diagnostic tool and information reference for the four Asian species of fruit fly (Diptera, Tephritidae) that have become accidentally established as pests in Africa, including the Indian Ocean Islands. Online at: <http://www.africamuseum.be/fruitfly/AfroAsia.htm> (accessed 19 August 2012).
9. Dhillon M.K., R. Singh, J.S. Naresh and H.C. Sharma, (2005). Evaluation of bitter melon (*Momordica charantia* L.) genotypes to melon fruit fly, *Bactrocera cucurbitae* (Coquillett) *Indian Journal Plant Protection*, 33:55-59.
10. Drew, R. A. I., Romig, M. C., and Dorji, C. (2007). Records of Dacinae fruit flies and new species of *Dacus* (Diptera: Tephritidae) in Bhutan. *The Raffles Bulletin of Zoology*, 55: 1-21.
11. Ekesi, S., Billah, M.K., Nderitu, P.W., Lux, S.A. and Rwomushana, I. (2009). Evidence for competitive displacement of the mango fruit fly, *Ceratitidis cosyra* by the invasive fruit fly, *Bactrocera invadens* (Diptera: Tephritidae) on mango and mechanisms contributing to the displacement. *Journal of Economic Entomology*. June; 102(3):981-91.
12. Ekesi, S. (2012). Combating fruit flies in Eastern and Southern Africa (COFESA): elements of a strategy and action plan for a regional cooperation program. Online at: <http://www.globalhort.org/media/uploads/File/Fruit%20Fly/Fruit%20fly%20Issue%20Paper%2010.05.2010-1.pdf> (accessed 19 August 2012).
13. Gupta, D., Verma, A.k., Bhalla, O.P. (1990). Population of fruit flies (*Dacus zonatus* and *D. dorsalis*) infesting fruit crops in north-western Himalayan region. *Indian Journal of Agricultural Sciences* 60:471-474.
14. Haseeb, M., Sharma, S. (2007). Studies on incidence and crop losses by fruit borer, *Deudorixisocrates* (Lep: Lycaenidae) on guava. In: Singh G, Kishan R, Chandra R (eds)

- 1st International guava symposium. Acta Hort 735, doi: 10.17660/ Acta Hort .2007. 735.66.
15. Jose, L., Cugala, D., Santos, L. (2013). Assessment of invasive fruit fly fruit infestation and damage in Cabo Delgado Province, Northern Mozambique. *African Crop Science Journal* 21: 21-28.
 16. Kapoor, V. C. (2002). Fruit-fly pests and their present status. *Proceedings of 6th International Symposium*, 6-10 May, 2002, Stellenbosch, South Africa, pp.23-33.
 17. Kapoor, V. C. (2006). Taxonomy and biology of economically important fruit flies of India. *Israel Journal of Entomology*, 35:459-475.
 18. Kishore, P., Kempraj, T., Elaiyabharathi, T., and Kavitha, C. (2018) Species Diversity and Relative Abundance of Fruit Flies (Tephritidae: Diptera) in Cucurbitaceous Vegetables Madras Agric. J., 105 (4-6): 196-200.
 19. Lux, S.A., R.S. Copeland, I.M. White, A. Manrakhan and M.K. Billah. (2003a). A New Invasive Fruit Fly Species from the *Bactrocera dorsalis* (Hendel) Group Detected in East Africa. *Insect Science and Its Application*, 23(4), 355-361.
 20. Lux, S.A., S. Ekesi, S. Dimbi, S. Mohamed and M. Billah. (2003b). Mango-infesting fruit flies in Africa—Perspectives and limitations of biological approaches to their management, pp. 277–293. In *Biological Control in Integrated Pest Management Systems in Africa* (Edited by P. Neuenschwander, C. Borgemeister and J. Langewald). CABI, Wallingford.
 21. Muhammad, I., M. Usman, M. Asif and I.A. Khan. (2004). Integrated Pest Management of Mango against Mealy Bug and Fruit Fly. *International Journal of Agriculture & Biology*. Vol 6 (3): 452–454.
 22. Musasa, S.T., Musundire, R., Mashingaidze, A.B. and Makuza, S.M. (2015). A preliminary study of the orange (*Citrus sinensis*) fruitvalue-chain in Chimanimani Rural District, Zimbabwe. *African J. Agric Res.* 10, 3507-3516.
 23. Mwatawala, M.W., M. De Meyer, R.H. Makundi and A. P. Maerere, (2006). Biodiversity of fruit flies (Diptera, Tephritidae) at orchards in different agro-ecological zones of the Morogoro region *Tanzania. Fruits*, 61: 321–332.
 24. Nath, D.K. (1973). *Callantraminax* (End.) (Teph.: Dipt.), a new record of a ceratitid fruit fly on orange fruits (*Citrus reticulata* Blanco) in India. *Indian Journal of Entomology* 34: 246.
 25. Ndah, N.R., Andrew, E.E and Bechem, E. (2013). Species composition, diversity and distribution in a disturbed Takamanda Rainforest, South West, Cameroon. *African Journal of Plant Science: Vol.7(12)*, pp.577-585.
 26. Ndiaye, M., E. O. Dieng and G. Delhove. (2008). Population Dynamics and On-Farm Fruit Fly Integrated Pest Management in Mango Orchards in the Natural Area of Niayes in Senegal. *Pest Management in Horticultural Ecosystems*, Vol. 14, No. 1, pp 1-8.
 27. Nizamani, I.A., Rustamani, M.A., Nizamani, S.M., Nizamani, S.A. and Khaskheli, M.I. (2015). Population Density of Foliage Insect Pest of Jujube, *Ziziphus mauritiana* Lam. Ecosystem. *Journal of Basic and Applied Sciences*. 11, 304-313.
 28. Otieno, W. (2009). *Bactrocera invadens* in Kenya. In: Fighting Fruit Flies Regionally in Sub-Saharan Africa. Information letter No. 6, December. COLEACP, CIRAD.
 29. Pielou E.C. (1966). The measurement of diversity in different types of biological collections *Journal of Theoretical Biology*, 13:131-144.
 30. Poudel, K. and Regmi, C. (2008). Disease and insect pest of Citrus (A booklet in Nepali version). Nepal Agriculture Research Council. Horticulture Research Station. Khumaltar, Lalitpur.
 31. Rai, A.B., Halder, J., Kodandaram, M.H. (2014). Emerging insect pest problems in vegetable crops and their management in India: An appraisal. *Pest Mgmt Hort Ecosyst.* 20: 113-122.

32. Saxena, D.K., Rawat, R.R. (1968). A study on the incidence, varietal performance and biology of the berfruitfly in Madhya Pradesh. *Madras Agric. J.* 55:328-333.
33. Senior-White, R. (1924). Trypetidae. *Catalogue of Indian Insects IV*. Government of India, Calcutta, India. pp-1-33.
34. Shah, M. I., Batra, H. N. and Ranjhen, P.L. (1948). Notes on the biology of *Dacus* (*Strumeta*) *ferrugineus* (Fab). and other fruit flies in the North-West Frontier Province. *Indian Journal of Entomology*, 10:249-266.
35. Sharma, V.P., Lal, O.P., Rohidas, S.B., Pramanick, P.K. (1998). Varietal resistance in ber (*Zizyphus mauritiana* Lamk.) against the fruitfly, *Carpomyia vesuviana* Costa (Diptera: Tephritidae) under field conditions. *J. Entomol. Res.* 22(1):61-67.
36. Singh, M.P., Vashishta, B.B. (1985). Field screening of some ber cultivars for resistance to ber fruit fly, *Carpomyia vesuviana* Costa. *Indian J. Plant Prot.* 12:55-56.
37. Tsuruta, K., White, I.M., Bandara, H.M.J., Rajapakse, H. (1997). A preliminary notes on the host-plants of fruit flies of the Tribe Dacini (Diptera, Tephritidae) in Sri Lanka. *Esakia*. 37, 149–160.
38. Ukey, N.S., Chandele, A.G., Wagh, S.S., Bansode, G.M. (2013). Species composition of fruit flies, *Bactrocera* spp. (Diptera: Tephritidae) infesting guava in Maharashtra. *Pest Management in Horticultural Ecosystems*; 19(2):242-244.
39. Vadivelu, K. (2014). Biology and management of Ber fruit fly, *Carpomyia vesuviana* Costa (Diptera: Tephritidae). *African Journal of Agricultural Research*. 9(16), 1310-1317.
40. Vayssieres, J.F., Carel, Y., Coubes, M. and Duyck, P.F. (2008). Development of immature stages and comparative demography of two cucurbit infesters in reunion island: *Bactrocera cucurbitae* and *Dacus ciliatus* (Diptera: Tephritidae). *Environ. Entomol.* 37(2):307–314.
41. Verghese, A., Madhura, H.S., Jayanthi, P.D.K. and Stonehouse, J.M. (2002). Fruit flies of economic significance in India, with special reference to *Bactrocera dorsalis* (Hendel). In: *Proceedings of Sixth International Symposium of Fruit flies of Economic Importance, Stellenbosch, South Africa*, pp. 317-324.
42. White, I.M. and Elson-Harris, M.M. 1992. Fruit flies of economic significance: their identification and bionomics. C.A.B International, United Kindom, 601 pp.