

## Original Research Article

### An evaluation of the systemic impact and clinical efficacy of sequential vincristine sulfate versus doxorubicin in the post-operative management of Canine Mammary Tumours (CMT)

---

---

#### ABSTRACT

The present study was conducted to evaluate the systemic impact and clinical efficacy of sequential vincristine sulfate compared to doxorubicin in the post-operative management of Canine Mammary Tumours (CMT) on 18 clinical cases of mammary tumour of various breed, irrespective of their age, sex and divided into three groups consisting of 6 animals in each. Group Surgery was treated with surgical removal of tumour only while animals of Group SurgDox and Group SurgVin were treated with surgical excision followed by administration of Doxorubicin (30mg/m<sup>2</sup>) BSA and Vincristine sulfate (0.025 mg/kg) intravenously along with DNS at 7<sup>th</sup> and 14<sup>th</sup> post-operative days respectively. The haemato-biochemical parameters and wound evaluation were done at different time intervals. Systemic impact on haematological and biochemical parameters were briefly altered and chemotherapeutic medications does not induce any adverse reactions or detrimental effects on vital organs, but these changes remained within the normal physiological range. Doxorubicin and vincristine sulfate therapy showed minimum to no reoccurrence of tumour with few adverse reactions such as inappetance, vomition, anaemia and alopecia. However, these conditions were managed by supportive therapy. Therefore, it could be concluded that sequential vincristine therapy is best for the post-operative management of canine mammary tumours as it effectively regressed the tumour without relapse.

**Keywords:-** Canine, CMT, Doxorubicin, Mammary tumour, Chemotherapy, Vincristine sulfate

#### INTRODUCTION

“Canine mammary tumors are second most common neoplasm of unspayed female dog representing approximately 40-50% of all neoplasms” (Khimta *et al.*, 2010). “However, male dogs are also affected but the prevalence is only 1%” (Rutterman *et al.*, 2000). “The last two sets of glands (4<sup>th</sup> and 5<sup>th</sup> pair) are most commonly affected. Appearance of mammary gland tumours in dogs can vary greatly. The tumours can be firm or soft, well-defined lumps or diffuse swellings. Mammary tumours primarily undergo metastasis to the regional lymphnodes or to the lungs” (Moulton, 1990). “Mostly, canine mammary tumours are hormone dependent and can be prevented if ovariohysterectomy (OH) is performed before 1 year of age. Bitches spayed prior to their first estrous cycle have about 0.5% risk of developing mammary tumours, while the risk is about 8% in bitches spayed between their first and second estrous and after the second estrous, the risk of mammary tumour development is 26

percent”(Brodey *et al.*, 1983). “The age group at which mammary tumours occurred most frequently was 8-10 years followed by 10-12 years” (Kishor *et al.*, 2016). “In general, sexually intact female dogs have seven times more risk of developing mammary tumours compared to neutered animals” (MacPhail, 2013).

“Classical modalities of cancer management include surgery, radiation and chemotherapy” (Kumar and Pawaiya, 2010). “Surgery remains the treatment of choice for most canine tumours except diffuse, unapproachable and non-exteriorisable, inflammatory carcinomas with distal metastasis. The extent of surgery should be determined by location, tumour size and the presence of some single or multiple tumours”(Fossum *et al.*, 1997). “Surgical excision alone yields unsatisfactory results in dogs with malignant mammary tumours exhibiting lymphatic or vascular invasion because these tumours have high rates of recurrence and metastasis” (Simonet *et al.*, 2006). “A number of chemotherapeutic drugs like Cyclophosphamide, Chlorambucil, Vincristine, Vinblastine, Carboplatin, Cisplatin, Methotrexate, 5-Fluorouracil and Doxorubicin etc have been used for the treatment of canine mammary tumours. Not a single chemotherapy protocol has been reported to be completely effective. Surgery remains the best way to treat most tumours while some tumours are too large and deeply situated where complete removal is not possible. In such cases, radical surgery is advocated followed by chemotherapeutic measures to arrest the recurrence of cancer cells resulting from remnants. The importance of chemotherapy has been emphasized and was reported that survival could be prolonged after chemotherapy in cancer patients” (Maiti *et al.*, 2011). “Therefore, adjunct chemotherapy i.e. surgical excision along with chemotherapy is a newer approach for the treatment of tumours. Doxorubicin is produced by *Streptomyces peucetius var. caesius* and exerts its cytotoxic effect as a DNA-intercalating agent to inhibit further DNA and RNA biosynthesis” (Lori *et al.*, 2010). “Vincristine sulfate belongs to antimicrotubule agents which are extracted from *Vinca rosea* that cause mitotic arrest by interfering with polymerization or depolymerization of microtubules which plays a critical role in cell function and division” (Gustafson and Page, 2013). Since, the reports regarding the use of doxorubicin and vincristine sulfate in adjunct chemotherapy for treatment of canine malignant mammary tumour in clinical cases are limited. Therefore, looking to the advantages of adjunct chemotherapy, the current study was designed to compare the systemic impact and clinical efficacy of sequential vincristine sulfate versus doxorubicin in the post-operative management of Canine Mammary Tumours (CMT).

## **MATERIALS AND METHODS**

The current investigation was carried out at the Teaching Veterinary Clinical Complex (TVCC) and Department of Veterinary Surgery and Radiology, College of Veterinary Science & A.H., Anjora, Durg (C.G.) from July 2017 to July 2018 on 18 clinically suspected cases of canine mammary tumour (17 female and 1 male). They were aged 5-12 years and weighed 10-20 kg. The breedwise distribution of dogs was as follows: 4 Labrador, 4 German shepherd, 3 Non-descript, 3 Pomeranian, 1 Golden Retriever, 2 Lasa Apso, 1 Cross breed. Owner's consent was also taken into consideration before grouping of these animals for surgico-chemotherapy. All the selected 18 animals were divided randomly into three groups viz., Group Surgery, Group SurgDox and Group SurgVin, each comprising of 6 animals. Mammectomy was performed in animals of all the three groups under general anaesthesia using atropine, xylazine and ketamine. Post-operatively, dogs were given intravenous fluid therapy (DNS 500 ml), antibiotic (inj. Ceftriaxone 500 mg I/V) and analgesic (inj. Meloxicam 2 ml I/M) for five consecutive days. Antiseptic dressing was done with Povidone iodine liquid and Povidine ointment. Sequential chemotherapy drug (Doxorubicin and Vincristine) was given in Group SurgDox and Group SurgVin after 7 days of operation as post-operative management. Group Surgery was treated with only surgical removal of tumour whereas group SurgDox, Doxorubicin ( $30\text{mg}/\text{m}^2$ ) BSA was given intravenously alongwith DNS at 7<sup>th</sup> and 14<sup>th</sup> post-operative days respectively while in group SurgVin, Vincristine sulfate ( $0.025\text{ mg}/\text{kg}$ ) was administered intravenously alongwith DNS at 7<sup>th</sup> and 14<sup>th</sup> post-operative days. Supportive therapy using Tribivet @  $0.05\text{-}0.2\text{ mg}/\text{kg}$  b.wt. I/M was given for 5 days followed by Polybion syrup one tsf bid and Tab Liv-52<sup>®</sup> every day to dogs of Group SurgDox and SurgVin during course of chemotherapy. The present study was approved by the Animal ethical committee of the College.

## **I) Evaluation of Systemic Effect of sequential chemotherapy**

### **a) Haematological parameters**

Using an automated haematology blood cell counter (B. C. 2800 Vet. Mindray), blood samples (approximately 1 ml) were drawn from dogs' cephalic or saphenous veins in vials containing ethylene diamine tetra acetic acid (EDTA) before and on 10<sup>th</sup>, 30<sup>th</sup> and 60<sup>th</sup> day post surgery for the estimation of Haemoglobin (gm/dl), Packed Cell Volume (%), Total Erythrocyte Count (millions / $\text{mm}^3$ ), Total Leukocyte Count (thousand / $\text{mm}^3$ ) and Total Platelet Count (lac / $\text{mm}^3$ ). The Differential Leukocyte Count (DLC) was performed manually by making blood smears from the drawn blood and after methanol fixation; the smear was stained with Geimsa stain. A percentage was given for each count.

### **b) Biochemical parameters**

For estimation of biochemical parameters, 3 ml of peripheral venous blood were drawn from the dog in a sterile, clean vial before and at 10<sup>th</sup>, 30<sup>th</sup> and 60<sup>th</sup> day post surgery. Serum was separated

for estimation of serum glucose (mg/dl), total serum proteins (gm/dl), serum urea nitrogen (mg/dl), serum creatinine (mg/dl), aspartate aminotransferase (U/L), alanine aminotransferase (U/L), and alkaline phosphatase (U/L) using semi-automated biochemistry analyzer (Diasil-100 Systronics make).

## II) Evaluation of Clinical Efficacy of sequential chemotherapy

This was done on the basis of wound examination and recurrence of tumour following 2 weeks of chemotherapy in post-operative management of canine mammary tumour. Wound was assessed on 0, 3<sup>rd</sup> and 10<sup>th</sup> day following surgery to determine the nature of wound (normal/oedematous) and also type of discharge (bloody/serosanguinous/fetid/pussy). Side effects of the chemotherapy if any were also recorded.

## III) Statistical analysis


The data collected was statistically analysed using analysis of variance (ANOVA) and Duncan's multiple range tests (DMRT). A computerized statistical software program SPSSv17.0 was used to analyze all of the data in order to compare the mean value within each group at various time intervals as well as among groups at different time intervals, which were represented as mean  $\pm$  standard error. The level of significance was set at  $P < 0.05$ .

## RESULTS AND DISCUSSION

The goal of the present investigation was to evaluate the systemic impact and clinical efficacy of sequential vincristine sulfate versus doxorubicin in the post-operative management of Canine Mammary Tumours. All 18 cases of mammary tumour affected dogs presented to Department of Veterinary surgery and Radiology were unspayed and uncastrated. On clinical examination, the shape of mammary tumours varied from ovoid, elongated, rounded to irregularly nodular (**fig. 1**). Most of the tumours were solitary and pedunculated. Grossly, tumours were soft to firm in consistency. The present study revealed that mammary tumours were most by affecting the 5<sup>th</sup> gland followed by 4<sup>th</sup> to 5<sup>th</sup> gland, 4<sup>th</sup> gland and 3<sup>rd</sup> to 4<sup>th</sup> gland probably because of their greater size containing more mammary tissue and thereby subjected to a greater range of physiological changes, predisposing them to neoplasms. The incidence of mammary tumour was observed more in female (94.44%) as compared to male (5.55%). In the present study, out of 18 cases of mammary tumour, only one male dog was affected with mammary tumour and rest were female dogs. In fact, canine mammary tumours are specific tumour in females and rare in males and are often associated with hormonal abnormalities (Moulton *et al.*, 1970). This could be due to action of ovarian hormones (estrogens and progesterone)

on mammary gland tissue during different stages of development which could be a risk factor associated with the development of mammary tumours (Petrov *et al.*, 2014).

**Fig. 1. Showing location of canine mammary tumour involving mammary glands**



### **I) Systemic Effect of sequential vincristine sulfate vs doxorubicin in post-operative management of Canine Mammary Tumours.**

**(A) Haematological parameters:** Haematological parameters did not show much variation in canine mammary tumour dogs before, post-surgery and after chemotherapeutic regimens in different groups at various time intervals (Table 1). There was non-significant ( $P > 0.05$ ) decrease in haemoglobin values in group Surgery at 10<sup>th</sup> day after surgical excision which might be due to surgical stress as reported by Benjamin (2010) and due to loss of blood from highly vascular malignant tumour during surgery. Thereafter, values showed increasing trend upto 60<sup>th</sup> day. This could be due to haemo-concentration of red cells because of tissue alteration in acute inflammation and increased viscosity of blood (Vegard, 2019). Whereas in group SurgDox and SurgVin, haemoglobin showed non-significant decrease upto 30<sup>th</sup> day which was less than 0 day value and indicated anaemia. Thereafter,

haemoglobin values increased upto 60<sup>th</sup> day of observation period. The decrease in haemoglobin value after doxorubicin and vincristine administration might have resulted due to chemotherapeutic damage to red blood cell formation in the bone marrow. Similarly, Brar *et al.* (2002) recorded anaemia due to bone marrow depression resulting in progressive decrease in haemoglobin concentration. Non-significant decrease in packed cell volume and TEC was recorded in all the three groups which could be due to cytotoxic drug that suppress the replicating precursor cells of the bone marrow leading to reduced production of erythrocyte. Nak *et al.* (2005) reported that decreased Hb, PCV and TEC may be attributed to the tumoral bleeding and myelosuppression induced by vincristine drug. There was non-significant decrease in TEC in group SurgDox and SurgVin which could be due to chemotherapy induced erythrocytopenia resulting from myelosuppression. Similar observation was also recorded by Brar *et al.* (2002) and Todorova *et al.* (2005). There was significant ( $P < 0.05$ ) decrease in TLC in group SurgDox and SurgVin upto 30<sup>th</sup> day, as these cytotoxic drugs suppress the replicating precursor cells of bone marrow, thus, resulting in reduced production of leucocytes. Todorova *et al.* (2005) reported a dose dependent reversible leukopenia as the predominant manifestation of doxorubicin bone marrow haematologic toxicity. Total platelet count showed non-significant decrease in group Surgery and SurgDox upto 30<sup>th</sup> day but to its contrary in group SurgVin, platelet count increased (thrombocytosis) upto 30<sup>th</sup> day and declined which was statistically non-significant. In group SurgDox (surgical removal alongwith doxorubicin therapy) where two doses of therapy were given in dogs suffering with mammary tumour revealed decreased in TPC (thrombocytopenia) which was statistically non-significant. Thrombocytopenia was observed due to bone marrow suppression. Doxorubicin hydrochloride localizes in the bone marrow of long bones and with progression of chemotherapy, the ability of bone marrow to produce platelets is diminished. Immune mediated thrombocytopenia occurs when antibody attaches to platelets promoting the destruction by the mononuclear phagocytic system. However, in group SurgVin where surgical excision alongwith two doses of vincristine sulfate therapy was given in dogs suffering with mammary tumour revealed increased in TPC (thrombocytosis) which was statistically non-significant. Therefore, it could be stated that chemotherapeutic agent vinca alkaloid vincristine sulfate is a safe drug since thrombocytosis was evident with minimum haematological alterations.

There was significant ( $P < 0.05$ ) decrease in neutrophils count in group SurgDox and SurgVin, after administration of chemotherapeutic agent doxorubicin and vincristine sulfate which usually localized into the bone marrow of long bone and its myelosuppressive action reduced the neutrophil percentage therefore, it could be stated that the intravenous chemotherapy induces neutropenia.

Whereas, in group Surgery, neutrophil decrease was negligible. Similarly, Ravikumar *et al.* (1999) observed a progressive decrease in neutrophils count till the end of vincristine therapy which resulted due to decreased bone marrow production. Whereas, lymphocyte count in group SurgDox and SurgVin showed significant ( $P<0.05$ ) increase at 30<sup>th</sup> day intervals which decreased significantly ( $P<0.05$ ) on 60<sup>th</sup> day whereas in group Surgery there was marginal variation in lymphocyte count during period of observation. The significantly ( $P<0.05$ ) reversible leukopenia and neutropenia observed following chemotherapy (doxorubicin and vincristine) in the present study was probably due to the action of cytotoxic drugs which suppressed the replicating precursor cells of bone marrow and created myeloid toxicity. There was non-significant variation in the values of eosinophils and monocytes before and after surgicochemotherapeutic regimens and all the haematological values remained within normal physiological range. Srivastava *et al.* (2009) reported that during the course of vincristine therapy, reticuloendothelial system depresses which causes the changes in haematological parameters in dogs affected with mammary tumours as observed in the present study. Similarly, Khan *et al.* (2017) also observed a significant ( $P<0.05$ ) decrease in haematological parameters viz., haemoglobin, PCV, TEC, TLC, Platelet count, neutrophils and eosinophils after surgical excision along with doxorubicin treatment in canine affected with mammary tumour which could be due to total suppression of the bone marrow activity, accidental infection and allergic reaction.

**Table 1: Alternation in Haematological parameters after sequential chemotherapy in different groups at various time interval**

Parameters	Groups	0 day	10 <sup>th</sup> day	30 <sup>th</sup> day	60 <sup>th</sup> day
Haemoglobin (gm%)	Surgery	12.40 <sup>aA</sup> ±0.93	12.15 <sup>aA</sup> ±0.44	12.20 <sup>aB</sup> ±0.30	12.50 <sup>aB</sup> ±0.50
	SurgDox	12.63 <sup>bA</sup> ±0.40	11.24 <sup>aA</sup> ±0.43	10.85 <sup>aA</sup> ±0.54	11.20 <sup>aA</sup> ±0.36
	SurgVin	12.10 <sup>bA</sup> ±0.408	11.73 <sup>abA</sup> ±0.71	10.99 <sup>aA</sup> ±0.36	11.50 <sup>abA</sup> ±0.54
Packed Cell Volume (%)	Surgery	37.68 <sup>aAB</sup> ±1.24	36.58 <sup>aB</sup> ±1.50	36.07 <sup>aB</sup> ±1.74	36.43 <sup>aB</sup> ±1.51
	SurgDox	34.16 <sup>bA</sup> ±1.24	30.72 <sup>aA</sup> ±1.50	29.60 <sup>aA</sup> ±1.51	31.13 <sup>aA</sup> ±1.70
	SurgVin	39.56 <sup>aB</sup> ±1.24	37.60 <sup>aB</sup> ±1.50	35.71 <sup>aB</sup> ±1.72	36.60 <sup>aB</sup> ±1.52
Total Erythrocyte Count (TEC) (x 10 <sup>6</sup> cu mm)	Surgery	6.94 <sup>aA</sup> ±0.20	6.89 <sup>aA</sup> ±0.25	6.80 <sup>aA</sup> ±0.20	6.95 <sup>aA</sup> ±0.28
	SurgDox	7.99 <sup>bB</sup> ±0.20	6.80 <sup>aA</sup> ±0.50	6.20 <sup>aA</sup> ±0.25	6.72 <sup>aA</sup> ±0.70
	SurgVin	7.50 <sup>bAB</sup> ±0.50	6.65 <sup>abA</sup> ±0.60	6.06 <sup>aA</sup> ±0.40	6.10 <sup>abA</sup> ±0.20
Total Leukocyte Count (TLC) (x 10 <sup>3</sup> cu mm)	Surgery	11.50 <sup>aB</sup> ±0.40	11.20 <sup>aB</sup> ±0.70	11.68 <sup>aC</sup> ±0.60	11.78 <sup>aB</sup> ±0.50
	SurgDox	10.70 <sup>cA</sup> ±0.46	9.20 <sup>bA</sup> ±0.50	8.73 <sup>aA</sup> ±0.30	9.10 <sup>abA</sup> ±0.60
	SurgVin	12.05 <sup>cC</sup> ±0.30	11.60 <sup>bB</sup> ±0.45	11.08 <sup>aB</sup> ±0.60	11.28 <sup>bB</sup> ±0.16
Total Platelet Count (TPC)	Surgery	3.55 <sup>aB</sup> ±0.20	3.40 <sup>aA</sup> ±1.10	3.10 <sup>aA</sup> ±1.17	3.46 <sup>aA</sup> ±1.19
	SurgDox	3.80 <sup>cB</sup> ±0.20	2.99 <sup>bA</sup> ±0.30	2.30 <sup>aA</sup> ±0.70	2.57 <sup>aA</sup> ±0.14
	SurgVin	3.10 <sup>aA</sup> ±0.70	3.89 <sup>bA</sup> ±0.11	4.10 <sup>bB</sup> ±0.16	3.99 <sup>bB</sup> ±0.19
Neutrophils (%)	Surgery	70.85 <sup>bA</sup> ±0.96	69.95 <sup>abA</sup> ±1.15	69.50 <sup>aC</sup> ±1.10	70.20 <sup>abC</sup> ±1.58
	SurgDox	73.55 <sup>cAB</sup> ±0.67	67.16 <sup>bA</sup> ±1.10	60.43 <sup>aA</sup> ±0.28	62.70 <sup>aA</sup> ±0.50
	SurgVin	74.88 <sup>dB</sup> ±0.96	68.67 <sup>cA</sup> ±1.15	61.90 <sup>aB</sup> ±1.10	65.30 <sup>bB</sup> ±0.58

<b>Lymphocyte (%)</b>	<b>Surgery</b>	20.57 <sup>aA</sup> ±1.30	22.52 <sup>cA</sup> ±1.10	21.20 <sup>bA</sup> ±1.09	20.45 <sup>aA</sup> ±0.30
	<b>SurgDox</b>	21.15 <sup>aA</sup> ±0.08	31.30 <sup>cB</sup> ±0.75	35.85 <sup>dC</sup> ±.33	27.45 <sup>bB</sup> ±0.35
	<b>SurgVin</b>	22.65 <sup>aB</sup> ±0.07	32.42 <sup>cC</sup> ±0.90	34.63 <sup>dB</sup> ±0.30	29.62 <sup>bC</sup> ±0.60
<b>Eosinophil (%)</b>	<b>Surgery</b>	2.98 <sup>bC</sup> ±0.10	2.45 <sup>abA</sup> ±0.90	2.20 <sup>aA</sup> ±0.11	2.40 <sup>abA</sup> ±0.30
	<b>SurgDox</b>	2.65 <sup>aAB</sup> ±0.60	2.22 <sup>aA</sup> ±0.90	2.02 <sup>aA</sup> ±0.54	2.20 <sup>aA</sup> ±0.40
	<b>SurgVin</b>	2.28 <sup>aA</sup> ±0.15	2.05 <sup>aA</sup> ±0.07	1.90 <sup>aA</sup> ±0.21	2.00 <sup>aA</sup> ±0.03
<b>Monocyte (%)</b>	<b>Surgery</b>	3.48 <sup>aA</sup> ±0.10	3.85 <sup>aA</sup> ±0.90	4.05 <sup>aA</sup> ±0.05	3.99 <sup>aA</sup> ±0.20
	<b>SurgDox</b>	3.80 <sup>aA</sup> ±0.20	4.10 <sup>aA</sup> ±0.50	4.50 <sup>aA</sup> ±0.35	4.20 <sup>aA</sup> ±0.11
	<b>SurgVin</b>	3.60 <sup>aA</sup> ±0.55	3.85 <sup>abA</sup> ±0.90	4.25 <sup>bA</sup> ±0.70	4.00 <sup>abA</sup> ±0.30

abcd-value (Mean±S.E.) bearing different superscript differ significantly (P<0.05) within groups

ABC-value (Mean±S.E.) bearing different superscript differ significantly (P<0.05) between groups

**(B) Biochemical parameters:** Serum glucose levels in group SurgDox showed significant (P<0.05) increasing trend upto 60<sup>th</sup> day after doxorubicin but within normal physiological range whereas non-significant decrease upto 60<sup>th</sup> day was recorded in group SurgVin after vincristine therapy (Table 2). The increase in glucose level in group SurgDox might be due to stress created on body by chemotherapeutic agent doxorubicin which caused the release of glucocorticoid and mineralocorticoid due to stimulation of adrenal cortex and epinephrine and non-epinephrine as a result of stimulation of the medulla. But serum glucose level in group Surgery showed irregular variation throughout the observation period and values remained within normal physiological limits. Total serum proteins value in the animals of group SurgDox, showed significant (P<0.05) decrease at 10<sup>th</sup> day which further increased significantly (P<0.05) upto 60<sup>th</sup> day. However, the TSP values in group Surgery and SurgVin showed marginal changes during observation period. The marginal irregular pattern of serum total proteins after chemotherapy could be consequential to gastro-intestinal disorder of cytotoxic drugs (Sandhu and Rampal, 2006). In contrast to our study, Gupta *et al.* (2014) reported a significant increase (P<0.05) in total protein value of surgically treated animals from preoperative 5.87±0.35g/dl to 6.485±0.33g/dl following surgery. Whereas, in surgical excision followed by vincristine treated group, there was non significant change in total protein values during the course of therapy but it remained towards the higher side of normal range. The animals of group SurgDox showed significant (P<0.05) increase in serum urea nitrogen and serum creatinine levels upto 30<sup>th</sup> day which further decreased significantly (P<0.05) at 60<sup>th</sup> day. Whereas group Surgery and SurgVin showed irregular changes throughout the observation period. Increase in BUN may be linked to glomerular filtration rate or increased protein catabolism caused by necrosis of tumour or metabolic side effects of neoplasia and increase in serum creatinine was attributed to the increase in catabolic activity. However, Todorova *et al.* (2005) did not find any significant change in the blood urea nitrogen and serum creatinine level after 2<sup>nd</sup> dose of doxorubicin. Similarly, Srivastava *et al.* (2009) has reported

that the plasma concentration of total protein, urea nitrogen, creatinine and glucose remained within normal range during the entire period of vincristine therapy with surgical excision. Khan *et al.* (2017) also reported significant ( $P<0.05$ ) increase in BUN, serum creatinine in dog treated with surgery followed with doxorubicin therapy but the values were within physiological limits whereas a non-significant change were recorded in surgically treated mammary tumour dog. The animals of group SurgDox and SurgVin showed significantly ( $P<0.05$ ) increasing trend in AST levels upto 30<sup>th</sup> day and thereafter the values decreased on 60<sup>th</sup> day after doxorubicin and vincristine therapy respectively. However, group Surgery showed non-significant variation after surgical excision. Whereas ALT level showed significant ( $P<0.05$ ) increasing trend in group Surgery and SurgDox upto 60<sup>th</sup> day following surgical removal of tumour and doxorubicin treated group respectively but animals of group SurgVin showed non-significant decrease in ALT level at 10<sup>th</sup> day which later on increased at 30<sup>th</sup> day and again decreased at 60<sup>th</sup> day after vincristine therapy. In the present study, the increase in AST and ALT levels following chemotherapy in SurgDox and SurgVin groups may probably be due to the result of detoxification of doxorubicin hydrochloride and vincristine in the liver which gets loaded exclusively for secretion of transaminase (Todorova *et al.*, 2005). In contrast to present study, Palta (2000) reported negligible changes in AST and ALT values after neoadjuvant and adjuvant chemotherapy with vincristine in canine mammary neoplasm cases. But Gupta *et al.* (2014) observed a non-significant increase in ALT values after second cycle of therapy in surgical excision followed by vincristine treated group which might be due to increase in metabolic activity of liver for the detoxification of drugs. Similar significant ( $P<0.05$ ) increase in AST and ALT was also observed in dog treated with surgery and followed sequential doxorubicin therapy whereas as a non-significant change in surgically treated mammary tumour dog but all values were within physiological limits as reported by Khan *et al.* (2017). The values of alkaline phosphatase (ALP) were towards the higher limit of normal range in all the three groups at time of presentation. Similarly, Bala (2005) reported elevated levels of alkaline phosphatase at presentation of canine mammary tumour. Alkaline phosphatase (ALP) levels in group Surgery, SurgDox and SurgVin showed significant ( $P<0.05$ ) decrease upto 30<sup>th</sup> day from  $85.40\pm 15.35$  to  $53.87\pm 9.70$  U/L,  $93.80\pm 18.60$  to  $50.67\pm 4.68$  U/L and  $80.55\pm 12.80$  to  $60.68\pm 18.45$  U/L respectively with a further increase upto the end of observation period. This indicates that increase in the alkaline phosphatase level could be attributed to presence of malignancy and addition of chemotherapy (vincristine and doxorubicin) induced stress, which was intensely reflected by rise in alkaline phosphatase activity during the study. Chauhan and Agarwal (2008)

recorded that alkaline phosphatase activity increases in case of carcinoma. Gupta *et al.* (2014) reported progressive increase in serum ALP value after first cycle and second cycle of vincristine therapy.

**Table 2 : Alternation in Biochemical parameters after sequential chemotherapy in different groups at various time interval**

Parameters	Groups	0 day	10 <sup>th</sup> day	30 <sup>th</sup> day	60 <sup>th</sup> day
Glucose (mg/dl)	Surgery	69.31 <sup>ab</sup> ±1.04	73.53 <sup>bb</sup> ±1.09	72.46 <sup>bb</sup> ±0.10	73.38 <sup>bb</sup> ±0.48
	SurgDox	73.54 <sup>ac</sup> ±0.75	75.69 <sup>bc</sup> ±0.63	78.95 <sup>cc</sup> ±1.10	83.55 <sup>dc</sup> ±1.35
	SurgVin	67.43 <sup>aa</sup> ±1.09	66.59 <sup>aa</sup> ±1.03	66.14 <sup>aa</sup> ±1.45	65.98 <sup>aa</sup> ±1.58
Total Serum Proteins (gm/dl)	Surgery	6.84 <sup>aa</sup> ±2.10	6.43 <sup>ab</sup> ±1.05	6.31 <sup>ab</sup> ±0.75	6.14 <sup>aa</sup> ±1.04
	SurgDox	7.14 <sup>cc</sup> ±0.89	6.74 <sup>ac</sup> ±1.10	7.03 <sup>bb</sup> ±1.32	7.50 <sup>dc</sup> ±0.67
	SurgVin	6.32 <sup>ab</sup> ±0.59	6.25 <sup>aa</sup> ±0.40	6.30 <sup>aa</sup> ±0.10	6.68 <sup>bb</sup> ±0.67
Serum Urea Nitrogen (mg/dl)	Surgery	11.20 <sup>aa</sup> ±2.15	10.47 <sup>aa</sup> ±1.39	10.89 <sup>aa</sup> ±2.80	11.05 <sup>aa</sup> ±1.49
	SurgDox	12.39 <sup>ab</sup> ±0.45	14.18 <sup>bc</sup> ±0.30	17.71 <sup>cb</sup> ±0.85	15.64 <sup>cb</sup> ±0.60
	SurgVin	11.02 <sup>aa</sup> ±1.15	11.59 <sup>abb</sup> ±1.39	11.10 <sup>aa</sup> ±1.20	11.80 <sup>ba</sup> ±1.10
Serum Creatinine (mg/dl)	Surgery	1.12 <sup>ac</sup> ±0.022	1.23 <sup>bb</sup> ±0.038	1.25 <sup>bb</sup> ±0.026	1.30 <sup>bb</sup> ±0.30
	SurgDox	1.02 <sup>ab</sup> ±0.22	1.55 <sup>bc</sup> ±0.38	1.89 <sup>cc</sup> ±0.26	1.28 <sup>abb</sup> ±0.60
	SurgVin	0.83 <sup>aa</sup> ±0.56	0.88 <sup>aa</sup> ±0.84	0.92 <sup>aa</sup> ±0.74	0.90 <sup>aa</sup> ±0.45
Aspartate aminotransferase	Surgery	33.63 <sup>ab</sup> ±2.072	34.73 <sup>abA</sup> ±1.90	35.54 <sup>bcA</sup> ±1.67	36.65 <sup>ca</sup> ±1.56
	SurgDox	30.64 <sup>aa</sup> ±1.45	36.76 <sup>bb</sup> ±1.36	42.42 <sup>cb</sup> ±2.70	37.44 <sup>bAB</sup> ±1.40
	SurgVin	34.53 <sup>ab</sup> ±0.50	39.74 <sup>cc</sup> ±0.66	42.84 <sup>db</sup> ±0.89	38.16 <sup>bc</sup> ±0.46
Alanine Aminotransferase	Surgery	33.34 <sup>ab</sup> ±5.40	38.90 <sup>bb</sup> ±4.90	43.03 <sup>bcB</sup> ±6.74	45.48 <sup>db</sup> ±3.64
	SurgDox	32.43 <sup>ab</sup> ±6.56	40.60 <sup>bb</sup> ±5.63	44.70 <sup>cb</sup> ±3.86	46.90 <sup>db</sup> ±4.60
	SurgVin	30.40 <sup>ba</sup> ±4.20	29.12 <sup>aa</sup> ±2.92	29.84 <sup>abA</sup> ±3.54	29.58 <sup>abA</sup> ±2.05
Alkaline phosphatase (U/L)	Surgery	85.40 <sup>db</sup> ±15.35	60.70 <sup>ba</sup> ±8.06	63.87 <sup>ab</sup> ±9.70	75.40 <sup>cc</sup> ±13.39
	SurgDox	93.80 <sup>dc</sup> ±18.60	65.55 <sup>bb</sup> ±11.60	50.67 <sup>aa</sup> ±4.68	69.65 <sup>cb</sup> ±5.40
	SurgVin	80.55 <sup>da</sup> ±12.80	60.68 <sup>ba</sup> ±8.45	58.76 <sup>aa</sup> ±15.44	60.77 <sup>ca</sup> ±8.89

abcd-value (Mean±S.E.) bearing different superscript differ significantly (P<0.05) within groups

ABC-value (Mean±S.E.) bearing different superscript differ significantly (P<0.05) between groups

## **II) Clinical efficacy of sequential vincristine sulfate vs doxorubicin in post-operative management of Canine Mammary Tumours.**

The nature of wound healing process was normal in all the three groups without any complications or discharge. In all the three groups, complete wound healing was observed in 8 to 10 days and was followed for at least three months after surgery and post chemotherapeutic regimens (doxorubicin and vincristine sulfate). The symptoms associated after administration of chemotherapeutic drug were inappetence, vomiting, anaemia and alopecia. These above symptoms were comparatively more in animals of group SurgDox as compared to group SurgVin. However, the condition of the animal was managed by supportive therapy with administration of intravenous fluids, antacids and liver tonics as a palliative measure. There was no recurrence in group SurgVin during the study period while, minimum recurrence (+) was observed in one case in group SurgDox whereas group Surgery showed mild to moderate recurrence (+, ++) in two cases. These findings are in

agreement with Karma (2006) and Srivastava *et al.* (2009) who observed that surgery combined with Vincristine therapy has been proven as excellent therapeutic regimen for canine mammary tumour leading to complete regression of the neoplasm without relapse. However, surgical excision following doxorubicin therapy was also effective in management of canine mammary tumour but showed minimum recurrence as observed in one case. The overall response of post-operative management canine mammary tumour using surgical excision alongwith sequential administration of chemotherapeutic drug (doxorubicin and vincristine) indicates that vincristine sulfate had minimum side effect on systemic impact on haematological and biochemical parameters, which proves its superiority over doxorubicin.

**Table 3: Showing Clinical efficacy after sequential chemotherapy in management of Canine Mammary Tumours**

Animal No.	Group Surgery		Group SurgDox		Group SurgVin	
1	Nil	++	V, Al	+	I	0
2	Nil	0	A, I	0	I	0
3	Nil	0	I, A	0	A, V	0
4	Nil	+	I	0	A, V, Al	0
5	Nil	0	I, V, Al	0	V, Al	0
6	Nil	0	A, V	0	I	0

**0 indicates no recurrence, + indicates minimum recurrence, ++ moderate recurrence, +++ severe recurrence**

**V= Vomition, ; A= Anaemia ; I= Inappetence ; Al= Alopecia**

## CONCLUSION

Systemic impacts on haematological and biochemical parameters were temporarily altered, but these alterations remained within the normal physiological range as chemotherapeutic drugs do not cause any adverse reaction or deleterious effect on vital organs. Sequential vincristine sulfate and Doxorubicin therapy showed minimum to no reoccurrence of tumour with few adverse reactions such as inappetance, vomition, anaemia and alopecia. However, these conditions were managed by supportive therapy. Therefore, it can be concluded that sequential vincristine sulfate therapy is more effective for post-operative treatment of canine malignant tumour as compared to sequential doxorubicin therapy as it effectively suppressed the development of new tumour cells and metastasis.

## Ethical Approval

Animal Ethic committee approval has been collected and preserved by the author(s)

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

## REFERENCES

1. Bala, M. (2005). Clinical studies on the evaluation of doxorubicin as an adjuvant chemotherapy for the management of canine mammary neoplasms. M.V.Sc. Thesis, Punjab Agricultural University, Ludhiana, India
2. Benjamin, M. M. (2010). Haematology, In: Outline of Veterinary Clinical Pathology. 3<sup>rd</sup> Edn., Kalyani Publishers, New Delhi-Ludhiana, Pp:51-56.
3. Brar, R. S., Sandhu, H. S. and Singh, A. (2002). Veterinary Clinical Diagnosis by Laboratory methods. 1<sup>st</sup> Edn. Kalyani Publishers, New Delhi, Pp:28-131.
4. Brodey, R. S., Goldschmidt, M. H. and Roszel, J. R. (1983). Canine mammary gland neoplasms. *J. Am. Animal Hosp. Assoc.*, 19:61-90.
5. Chauhan, R. S. and Agarwal, D. K. (2008). Textbook of Veterinary Clinical Laboratory Diagnosis, 2<sup>nd</sup> Edn., pp:150.
6. Fossum, T. W., Hetlund, C. S., Hulse, D. A., Johnson, A. L., Seim, S. B., Wilard, M. D. and Carroll, G. L. (1997). Small Animal Surgery, Mosby Book Inc. St. Louis, Missouri, pp:539-544.
7. Gupta, K., Sood, N., Uppal, S., Mohindroo, J., Mahajan, S., Raghunath, M. and Singh, K. (2012). Epidemiological studies on canine Mammary tumour and its relevance for breast cancer studies. *Journal of Pharmacy*, 2(2):322-333.

8. Gupta, P., Raghunath, M., Gupta, A. K., Sharma, A. and Kour, K. (2014). Clinical study for diagnosis and treatment of canine mammary neoplasms (CMNs) using different modalities. *Indian Journal of Animal Research*, **48**(1):45-49.
9. Gustafson, D. L. and Page, R. L. (2013). Cancer chemotherapy. In: S.J. Withrow, D. M. Vail and R. L. Page (ed): Withrow & MacEwen's Small Animal Clinical Oncology, 5<sup>th</sup> edn. Saunders Elsevier, St. Louis, Missouri, pp. 157-179.
10. Karma, S. (2006). Studies on the Prevalence and Treatment approach of Mammary Tumours in female dog. M.V.Sc. thesis submitted to Indira Gandhi Krishi Vishwavidyalaya, Raipur Chhattisgarh, India.
11. Khan, G. J., Thorat, M. G., Raulkar, R. V., Upadhye, S. V., Jadhao, P. T., Chepte, S. D. and Deshmukh, S. G. (2017). Surgico-chemotherapeutic management of mammary tumour in canine. *International Journal of Science, Environment and Technology*, **6**(5):3154-3159.
12. Khimta, S., Maiti, S. K., Kumar, N. and Sharma, A. K. (2010). Occurrence of neoplasms in canine-a retrospective study. *Indian Journal of Animal Science*, **80**:7-11
13. Kishor, T. K., Rao, S., Satyanarayana, M. L., Narayanaswamy, H. D., Byregowda, S. M., Nagaraja, B. N., Purushotham, K. M. and Kavya, N. (2016). Classification and staging of canine mammary gland tumours. *Journal of Cell and Tissue Research*, **16**(3):5787-5792.
14. Kumar, P. and Pawaiya, R.V.S. (2010). Advances in cancer diagnostics. *Brazilian Journal of Veterinary Pathology*, **3**(2):142-153.
15. Lori, J. C., Stein, T. J. and Thamn, D. H. (2010). Doxorubicin and cyclophosphamide for treatment of canine lymphoma:a randomized, placebo-controlled study. *Vet. Comp. Oncol.* **8**(3):188-195
16. MacPhail, C. M. (2013). Surgery of the female reproductive tract. In: Fossum, T. W. (ed). *Small Animal Surgery*. 4<sup>th</sup> Edition, St Louis, Mosby Elsevier. USA, pp 809-814.
17. Maiti, S.K., Manikandan, N.,Shivakumar, M.U., Kumar, N., Saikumar, G. and Gupta, O. P. (2011). Therapeutic evaluation of methotrexate with or without Cox-2 inhibitors in the management of canine mammary tumour. *Indian Journal of Canine Practice*, **3**:117-126.
18. Moulton, J. E. (1990). Tumours of the mammary gland. Tumours in Domestic Animals. 3rd edition. University of California Press, London, UK. pp:518- 552.
19. Moulton, J.E., Taylor, D. N., Dorn, C. R. and Anderson, A. C. 1970. Canine mammary tumours. *Pathol. Vet.*, **7**:289-320
20. Nak, D., Nak, Y., Cangul, I. T. and Tuna, B. (2005). A Clinicopathological study on the effect of vincristine on Transmissible Venereal Tumour in dogs. *Journal of Veterinary Medicine Series A: Physiology, Pathology, Clinical Medicine.*, **52**:366-370.
21. Palta, M. (2000). Clinical studies on multimodality in the management of canine mammary neoplasm. M.V.Sc. Thesis, Punjab Agricultural University, Ludhiana, India
22. Petrov, E. A., Ilievska, K., Trojancanec, P., Celeska, I., Nikolovski, G., Gjurovski, I. and Dovenski, T. 2014. Canine mammary tumours - clinical survey. *Mac Vet Rev.*, **37**(2):129-134.
23. Ravikumar, K. P. Vasanth, M. S. and Jayadevappa, S. M. (1999). Comparative efficacy of surgery and surgery with chemotherapy for mammary tumours in canines. *Indian J. Vet.Surg.*,**20**:115
24. Rutterman, G. R., Withrow, S. J. and MacEwen, W. G. (2000). Tumours of the mammary gland. In: *Small Animal Clinical Oncology*. 3rd Edn. Philadelphia WB Saunders Co. pp:450-467.

25. Sandhu, H.S. and Rampal, S. (2006). Essentials of Veterinary Pharmacology and Therapeutics. 1<sup>st</sup> Edn,pp:1359-1381.
26. Simon, D., Schoenrock, D., Baumgartner, W. and Nolte, I. (2006). Postoperative adjuvant treatment of invasive malignant mammary gland tumors in dogs with Doxorubicin and Docetaxel. *Journal of Veterinary Internal Medicine*, **20**: 1184-1190.
27. Srivastava, A. K., Singh, B., Srivastava, A. K., Lal, H. P., Sinha, N. and Sharma, A.K. (2009). A study on chemo-surgical management of mammary tumours in dogs. *Indian J. Vet. Surg.*,**30**(1):27-30.
28. Todorova, I., Simeonova, G., Simeonova, R. and Dinev, D. (2005). Efficacy and toxicity of oxorubicin and cyclophosphamide chemotherapy in dogs with spontaneous mammary tumours. *Trakia Journal of science.*, **3**(5):51-58.
29. Vegard, J. L. (2019). Text book of Veterinary General Pathology, 2<sup>nd</sup> Edn International Book Distributing Co., Lucknow (U.P.) India, pp:250
30. Punchkande N, Dewangan R, Sharda R, Jolhe D, Sahu D, Sidar SK, Yadav D. Treatment of canine transmissible venereal tumour using different surgico-chemotherapeutic protocols. *Indian Journal of Animal Research*. 2022;**56**(7):899-903.