

Antimicrobial Resistance Profile of *Salmonella enterica* Isolated from Improved Poultry Breed Farming Chain, Maharashtra, India

Abstract

With rising demand for poultry meat, globally backyard farms have adopted semi-intensive systems using improved birds. Food safety is a prime issue in as antimicrobial-resistant *Salmonella* spp. found in meat, which has often been traced back to farms. This study was focused on antimicrobial-resistant *Salmonella* spp. in an improved poultry breed farming chain in Maharashtra, India. A total of 364 samples were collected from hatcheries (n=124), improved breed farms (n=110), and households backyard with improved breeds (n=130). The isolation of *Salmonella* spp. was carried out as per IS-5887 (Part 3): 1999 protocol. A total of 15 isolates recovered with the prevalence rates of 4.83, 6.36, and 1.53 percent were found in hatcheries, farms, and backyard households, respectively. Five isolates that were randomly analyzed showed a homologous sequence as *Salmonella enterica*. Antimicrobial susceptibility testing of all isolates revealed higher resistance against Erythromycin (100%), followed by Ceftazidime (40%), while sensitivity (93.33%), was recorded against Ampicillin/Sulbactam, Amoxicillin/Sulbactam, and Enrofloxacin. The average multiple antibiotic resistance (MAR) index, of *Salmonella* isolates was 0.117. Genotypic resistance pattern revealed that all isolates were carrying the *bla*TEM gene, while none were harbouring broad spectrum extended-spectrum beta-lactamases (ESBLs) genes; *bla*SHV, *bla*OXA, and *bla*CTXM. Whereas All isolates were positive for the tetA gene but none of them was positive for *Sul1* genes. The study highlights the low antimicrobial resistance in *Salmonella* isolates which might be due to low usage. However monitoring of multidrug-resistant *Salmonella enterica* in improved poultry breed farming chains is essential to protect human health.

Keywords: improved poultry breed, farming chain, *Salmonella*, antimicrobial resistance, genotypic resistance

Background

Poultry meat consumption is on the rise in world among all animal-derived food items (Waghmare et al. 2021). Meat, due to its high nutritive value containing essential amino acids and various macro- and micronutrients, plays a crucial role in a balanced diet (Van Boeckel et al. 2019). In recent years with the surge in chicken demand, backyard poultry farms have transformed themselves into semi-intensive farming systems with the help of improved native or coloured variety birds and good quality feed (Chaiban et al. 2020). The intervention of backyard poultry farming with improved native or coloured poultry varieties suitable for backyard production can be a source of a supportable food production structure (Singh et al. 2022). Breeder stock and hatcheries can be an important source of *Salmonella* spp. and can be improved through the execution of effective intervention methods (Sivaramalingam et al. 2013).

Salmonellosis stands as a major global foodborne illness, contributing to 93.8 million of gastroenteritis cases and 155,000 deaths per annum globally (Heredia and Gracia 2018). *Salmonella* in food chains is emerging

worldwide and poultry is recognized as a foodborne pathogen reservoir, with several reports highlighting the occurrence of *Salmonella* linked to the backyard and commercial live poultry, production settings (Samanta et al. 2014; Elmonir et al. 2023).

In the past few years, there has been a significant surge in phenotypic and genotypic resistance among non-typhoidal *Salmonella* isolates to β -lactams, tetracycline, and sulphonamides (Egualo et al. 2017). The extensive use of tetracycline in poultry has caused the advance of tetracycline resistance in *Salmonella* spp. due to selective pressure (Waghmare et al. 2018). CDC, 2013, and WHO, 2017 reported that Gram-negative microorganisms that harvest beta-lactamases as one of the world's most insistent threats. Recent food safety studies have shown that *Salmonella* strains isolated from poultry and other foods possess extended-spectrum β -lactamases (ESBLs), which become a potential threat to human health as they may impact treatment regimens for ESBL-producing pathogens (Orabi et al. 2022; Dinh et al. 2023).

Broiler meat harvested from organic and pastured poultry farming production facilities has high demand (Van Loo et al. 2011; Rothrock et al. 2016). The appearance of antimicrobial resistance (AMR) in *Salmonella* from backyard poultry farming systems may pose a significant growing threat to anthropoid and animal health (Van Boeckel et al. 2019; Hedman et al. 2020). Similarly, studies conducted by other researchers reported significantly lower *Salmonella* contamination of fecal matter and bird feed (Siemon et al. 2007; Alali et al. 2010). An additional hypothesis is that improved breed poultry production systems work on reared-antibiotic-free rearing, which might potentially affect the antimicrobial resistance profile and presence of antibiotic resistance genes in the *Salmonella* populations along the hatchery- to- backyard farm continuum.

Therefore, to better understand the prevalence of antimicrobial resistant (AMR) *Salmonella* in these improved native breed poultry farms and backyard poultry management systems, some more work is essential; especially since market demand and production for backyard poultry has been increased. Thus, the present study was carried out to evaluate the antimicrobial resistance profile of *Salmonella* spp. isolated from the backyard poultry production system. This information will provide vital data on the prevalence, and AMR profile of *Salmonella* spp. in improved indigenous poultry breed farming chains which help to improve backyard poultry food safety systems.

Materials and methods

Sample collection

In the current study, a total of 364 samples were collected from improved indigenous poultry breed hatcheries (n=5), improved indigenous breed commercial farms (n=5), and improved breed-rearing backyard poultry setups (n=30) in Marathwada, Maharashtra state, India. The samples from hatcheries (n=124) were comprised of deceased chicks (25), unhatched eggs (25), shell waste (12), and tray swabs (12). A total of 110 samples (10 each of feed, water, litter, feces, walls swab, utensils swab, worker's hands swab, and cages; and 30 cloacal swabs) were selected from five different farms in Pathardi, Maharashtra, India. Similarly, 30 household backyard farms from the Parbhani, Vasmat, and Selu areas of Marathwada, Maharashtra, India were screened by collecting 130 samples (30 each of cloacal swabs, water and feces, 15 each of cage dust swabs and utensils swab and 10 feed samples). All the samples were collected aseptically, using sterilized polypropylene bags, and transported to the laboratory under a cold chain as per standard methods.

Isolation and characterization of *Salmonella* spp.

Isolation of *Salmonella* spp. was performed as per IS-5887 (Part 3): 1999 protocol with slight modifications based on the samples collected. The sample was processed (pre-enrichment) by inoculating a 5 gm sample of dead chick and unhatched egg in 45 ml of buffered peptone water (BPW), swabs and feces (1gm) were pre-enriched with 9 ml buffered peptone water, in sterile test tubes, whereas 25 g of each feed and water was inoculated in 225 ml of buffered peptone water and all samples were incubated at $37^{\circ}\text{C} \pm 1^{\circ}\text{C}$ for 24 hours. Further, enrichment was done by transferring 0.1 ml of inoculum from BPW into 10 ml Rappaport-Vassiliadis (RV) medium and incubated at 42°C for 24 hrs. A loop-full of inoculum from RV broth was streaked on Xylose Lysine Deoxycholate (XLD) Agar and Brilliant Green Sulpha (BGSA) Agar, plates by four quadrant streaking method. The inoculated plates were incubated at $37^{\circ}\text{C} \pm 10^{\circ}\text{C}$ for 24 hrs. for the development of colonies of *Salmonella* spp. The bacterial isolates were further identified by cultural, microscopic morphology, and biochemical characteristics (BAM 2007).

Black center smooth and round or large with black center colonies of *Salmonella* spp. were characterized by polymerase chain reaction (PCR) assays with primers designed for the *invA* gene (Nair et al. 2015). DNA of *Salmonella* isolates was extracted using heat lysis DNA Isolation protocol (Dashti et al. 2023). The details of PCR primers and cycling conditions used in the present study are mentioned in Table 1 and Table 2.

Sequencing of the *invA* gene amplified product

Amplified *invA* gene products of selected isolates from hatchery, farm and backyard households were sequenced from an outsourced agency (Submitted to GeneOmbio Technologies Pvt Ltd, India) for sequencing. The resulting sequences were analyzed, and sequence homology searches were conducted using the BLAST algorithm (https://blast.ncbi.nlm.nih.gov/Blast.cgi?PROGRAM=blastn&BLAST_SPEC=GeoBlast&PAGE_TYP E=BlastSearch).

Antimicrobial susceptibility test

As per the Clinical and Laboratory Standards Institute (CLSI) protocol, the Kirby–Bauer drug-sensitive disk technique was used to test the sensitivity of PCR-positive *Salmonella* isolates to 14 commonly used antimicrobials. Antimicrobial tested were Ampicillin/Sulbactam (A/S) 10/10 μg , Amoxicillin/Sulbactam (AMS) 30/15 μg , Amoxiclav (AMC), Tetracycline (TE) 30 μg , Enrofloxacin (EX) 10 μg , Ceftazidime (CAZ) 30 μg , Levofloxacin (LE) 5 μg , Chloramphenicol (C) 30 μg , Erythromycin (E) 15 μg , Gentamicin (GEN) 10 μg , Amikacin (AK) 10 μg , Ciprofloxacin (CIP) 5 μg , Co-Trimoxazole (COT) 25 μg and Nalidixic Acid (NA) 30 μg . Briefly, bacterial suspensions were achieved from overnight-grown cultures, which adjusted to the 0.5 McFarland turbidity standard. The broth culture was spread on the surface of a Muller Hinton agar plate using a sterile cotton swab. After about 20 min, the disks were applied to the plates and incubated at 37°C for 18 hr. Finally, using a scale (Himedia) the diameter of the inhibition zone was measured to categorize the resistance pattern. The results of phenotypic antimicrobial resistance patterns were analyzed as per the European Committee for Antimicrobial Susceptibility Testing (EUCAST) guidelines (2021).

Multiple Antibiotic Resistance (MAR) Index

The Multiple Antibiotic Resistance (MAR) index of individual isolates was calculated according to the method described by Krumperman (1983) by dividing the number of antibiotics to which the isolates were found resistant to the total number of antibiotics to which the isolate was exposed.

Molecular characterization of AMR genes in *Salmonella* spp.

All the 15 isolates were screened for the presence of 06 antimicrobial resistance genes. The genes that encode resistance to β -lactamases (*blaTEM*, *blaSHV*, *blaCTX-M*, and *blaOXA*), tetracyclines (*tetA*), and sulfonamides (*SulI*) were evaluated.

Genomic DNA of *Salmonella* spp. isolates were extracted by boiling method described by Anejo-Okopiet *et al.* (2016). All *Salmonella* isolates were screened for the presence of antimicrobial resistance genes through singleplex and multiplex PCR protocol described by Ng *et al.* (2001) for *tetA*, Ma *et al.* (2017) for *SulI* and Fang *et al.* (2008) for a group of β -lactamase genes (*blaSHV*, *blaCTXM*, *blaOXA* and *blaTEM*). Primers used are listed in Table 1. The cycling conditions for PCR are mentioned in Table 2.

The singleplex PCR for *invA*, *sulI* and *tetA* was performed in 25 μ l reaction volume containing 12.5 μ l of 2x PCR master mix (HiMedia Laboratories Pvt. Ltd., Mumbai), 1.5 μ l of each primer (10 pmol/ μ l) (Eurofins Genomics India Pvt. Ltd., Bangalore), 1 μ l of genomic DNA and 8.5 μ l molecular biology grade water (HiMedia Laboratories Pvt. Ltd., Mumbai) used to make desired volume. Whereas, multiplex PCR reaction was performed as described by Fang *et al.* (2008) in a 25 μ l volume containing 12.5 μ l 2x PCR Master Mix (Takara Bio Inc., Shiga, Japan) supplied with Taq DNA polymerase, buffer, MgCl₂, and dNTPs. In this PCR 1 μ l (10 pmol/ μ l) each of forward and reverse primer was used. Similarly, the reaction mixture contained 3 μ l DNA template and 7.5 μ l nuclease-free water to make a final volume of 25 μ l.

Table 1. Primers used for genotype characterization of *Salmonella* spp.

Sr.no	Primer	Target	Primer sequence (5'- 3')	Product Size (bp)	Reference
1	<i>invA</i>	Invasion-associated protein	F: TCGTGACTCGCGTAAATGGCGATA R: GCAGGCGCACGCCATAATCAATAA	423	(Nair et al. 2015)
2	<i>tetA</i>	Tetracycline	F: GCT ACA TCC TGC TTG CCT TC R: CAT AGA TCG CCG TGA AGA GG	210	(Ng et al. 2001)
3	<i>SulI</i>	Sulphonamide	F: TTTCTGACCCTGCGCTCTAT R: GTGCGGACGTAGTCAGCGCCA	425	(Ma et al. 2007)
4	<i>blaTEM</i>	Broad Spectrum β -lactamase	F: CGCCGCATACACTATTCTCAGAATGA R: ACGCTCACCGGCTCCAGATTTAT	445	(Fang et al. 2008)
5	<i>blaSHV</i>		F: CTTTATCGGCCCTCACTCAA R: AGG TGC TCA TCA TGG GAA AG	237	
6	<i>blaCTX-M</i>		F: ATGTGCAGYACCAGTAARGTKATG GC R: TGGGTRAARTARGTSACCAGAAAYCAGCGG	593	
7	<i>blaOXA</i>		F: ACA CAA TAC ATA TCA ACT TCG C R: AGT GTG TTT AGA ATG GTG ATC	813	

Table 2. PCR cycling conditions used under study for different primers

PCR steps	PCR conditions	Target genes			Broad Spectrum β -lactamase
		<i>invA</i>	<i>tetA</i>	<i>sulI</i>	<i>blaSHV, blaCTXM, blaOXAblaTEM</i> (Multiplex PCR)
Initial Denaturation	Temperature (°C)	94	95	94	94
	Time (min.)	5	3	3	5
Denaturation	Temperature (°C)	94	95	94	94
	Time (sec.)	30	30	30	45
Annealing	Temperature (°C)	56	60	60	63
	Time (sec.)	60	30	30	60
Extension	Temperature (°C)	72	72	72	72
	Time (sec)	90	60	45	60
Final Extension	Temperature (°C)	72	72	72	72
	Time (min.)	10	8	8	7
Number of cycles		35	35	30	30

Results and discussion

Prevalence of *Salmonella* isolates:

As is shown in Table 3, a total of 15 PCR *invA* gene positive *Salmonella* isolates were recovered from 364 samples, in which *Salmonella* was detected from Dead chick, Unhatched egg, Eggshell waste, Cloacal swabs, Faeces and Litter. All *Salmonella* isolates yielded desired amplified products of 423 bp for the *invA* gene (Fig. 1). The PCR-based method utilizing the *invA* genus-specific primer is a reliable approach due to its rapidity, high specificity, and sensitivity in confirming the isolates (Salehi et al. 2005).

Among the isolates, *Salmonella* strains were observed in 4.83% (6/124) of those from Hatcheries, 6.76% (7/110) of those from commercial farms, and 1.53% (2/130) of those from backyard household farms.

Salmonella can be spread horizontally and through fertilized eggs and it is zoonotic in nature (Zhao et al. 2021). Considering this, in mandate to better assess the tangible infection by *Salmonella* spp., we obtained several samples from hatcheries, farms and backyard farms from various locations by maintaining the supply chain in improved indigenous poultry breed farming.

Results of prevalence from hatcheries are in agreement with earlier studies of Muhammad *et al.* (2010) and Sohail *et al.* (2021) who reported occurrence of 4.4% and 10.34% in hatchery samples, whereas, the present findings are not in contrast with the findings of Mulika *et al.* (2011), Xu *et al.* (2020) and Withenshaw *et al.* (2021) wherein, the prevalence of *Salmonella* spp. in hatchery samples ranging from 16.6 to 33.3% was recorded. Decontamination, sanitation, waste disposal, biosecurity measures, and proper monitoring are necessary to prevent *Salmonella* transmission from hatchery to another setting. The improvement of hatchery hygiene and the application of efficient pathogen detection and disease control strategies would improve *Salmonella* control (Fahmy et al. 2023).

The prevalence of *Salmonella enterica* in Indigenous Poultry farms agrees with previous studies by Samanta *et al.* (2014), who reported a prevalence of 6.1 percent respectively. Abunna *et al.* (2016) stated that the distribution of *Salmonella* spp. varied depending on the sample type, poultry growth stage, and breeds. Soil and fecal matter present on the feathers and feet of birds serve as significant contributors to the microbial contamination observed in poultry carcasses (Orji *et al.* 2005). The results of the Prevalence of *Salmonella enterica* in Household Backyard Poultry Farming are not in agreement with previous reports by Jajereet *et al.* (2019), Koro *et al.* (2022) and Elmoniret *et al.* (2023). In household backyard poultry farming *Salmonella* spp. was observed only in cloacal swab samples (6.67%). The findings are not per the findings of Bhowmick *et al.* (2023) and Eid *et al.* (2023) who reported 30.82 and 31.00% prevalence, respectively in backyard poultry farm cloacal swabs. The presence of *Salmonella* in the backyard poultry might have been the sequel of a long-term persistence of *Salmonella* in these backyards or it could be the summary of laying hens purchased directly from contaminated hatcheries or other households (Trampel *et al.* 2014). The variances in sample types, topographical locations, sampling protocols, or isolation methods resulted in a multifarious and diverse isolation assessment.

Table 3. Prevalence of *Salmonella* in Backyard Poultry Farming System (n = 364)

Category	Type of Samples analyzed	Total Number of Samples	Positive Sample type	Total Number	Overall prevalence (%)
Hatchery	Dead chick (internal organ) Unhatched Egg (Yolk and Embryo) Tray swab Eggshell waste	124	Dead chick (2) Unhatched Egg (3) Shell waste (1)	6	4.83%
Indigenous poultry breed farms	Cloaca swab Utensils swab Wall dust swab Workers hand swab Feed Water Faeces Litter	110	Cloaca swab (3) Faeces (2) Litter (2)	7	6.36%
Household backyard poultry	Cloaca swab Cage dust swab Utensils swab Feed Water Faeces	130	Cloacal swab	2	1.53%

Sequencing of selected *invA* gene PCR Products for serotype confirmation

A total of five selected isolates from hatcheries (A5/S/2 and A5/U/3), farms (B1/CI/2) and household backyard poultry (C8/CI/1 and C11/CI/1) were sequenced using amplified *invA* gene (423bp) product. The sequencing homologous was observed on NCBI as *Salmonella enterica* and showed homology of 99.49%, 99.69% and 99.49% with sequences accession no. AP020332.1, CP074202.1, CP051329.1, respectively which were available on NCBI.

The results are in agreement with Pavon and Rivera (2021), wherein the possible use of the *invA* virulence gene for molecular typing of *Salmonella* through sequencing and phylogenetic analyses was

suggested. Amongst various types, Serovars Enteritidis and Typhimurium were being the most informed serovars. *S. enterica* is responsible for infections in humans and animals (Andino & Hanning., 2015) which makes it a serious concern.




Fig. 1 Agarose gel showing PCR amplified product of *invA* gene of *Salmonella* spp.

Antimicrobial resistance pattern of *Salmonella* isolates

In the present study antimicrobial susceptibility testing discovered upper resistance against Erythromycin (100%), followed by Ceftazidime (40%), while sensitivity was recorded against Ampicillin/Sulbactam (93.33%), Amoxicillin/Sulbactam (93.33%), Enrofloxacin (93.33%), Levofloxacin (93.33%), and Nalidixic Acid (93.33%), whereas, intermediate sensitivity recorded against Ceftazidime (53.34%), followed by Tetracycline (46.67%) and Amoxiclav (26.67%). In the present study, a higher resistance pattern against Erythromycin (100%) is in agreement with the findings of Bhuvaneshwari et al. (2015) and Al Mamum et al. (2017). The resistance pattern of Ceftazidime (40%) was found in agreement with Herrera-Sánchez et al. (2020), wherein the resistance reported was 75.5 percent. The results about Ampicillin/Sulbactam (0%), Gentamicin (0%), Amikacin (0%), Co-Trimoxazole (0%), Nalidixic acid (0%), Amoxicillin/Sulbactam (6.67%), Amoxyclav (6.67%), Tetracycline (6.67%), Enrofloxacin (6.67%), Levofloxacin (6.67%), Chloramphenicol (6.67%), Ciprofloxacin (6.67%) resistance patterns are not in agreement with Akondet et al. (2012), Thakur et al. (2013), Samanta et al. (2014) and Waghmare et al. (2018). Bhuvaneshwari et al. (2015) reported that β -lactam and macrolide antibiotics are used routinely for curbing bacterial infection among chickens in commercial farms compared to backyard farms. The resistance of *Salmonella* Enteritidis to ampicillin decreased in our study this might be due to the reduction in antibiotic selective pressure (Varijakshapanicker et al. 2019). The exercise of using antimicrobial agents for growth, and treatment in livestock decreases their efficacy and is found to be a significant factor in the emergence of multidrug-resistant *Salmonella* (Thakur et al. 2013).

Multiple Antibiotic Index (MAR) for *Salmonella* isolates

In the present study, only 3 *Salmonella* isolates were resistant to 3 or more than 3 antibiotics. The MAR index of A5/S/2, B1/CL/2, and B3/Fe/2 were 0.21, 0.50, and 0.21, respectively (Table 4). The MAR index of the remaining 12 isolates varies from 0.07 to 0.14 which indicates lower selective pressure for antimicrobial resistance. The results of the present study do not agree with the studies carried out by Yoke-Kqueen et al. (2008) and Talukder et al. (2021) who reported that 91.01 and 100 percent *Salmonella* isolates have a MAR index of

more than 0.2. The lower multidrug resistance pattern displayed by *Salmonella* isolates demonstrates that anthropoids consuming meat and meat products from concerned geographical areas are at low risk of multidrug-resistant *Salmonella* infection.

Table 4. Multiple antibiotic resistance [MAR] index and antibiogram of *Salmonella* isolates

Sr. no.	Source of samples	Sample code	Resistance to the number of Antibiotics	Antibiogram	MAR index
1.	Hatchery	A5/S/2	3	CAZ, C, E	0.21
2.		A5/U/3	1	E	0.07
3.		A5/U/4	2	CAZ, E	0.14
4.		A5/U/5	1	E	0.07
5.		A5/C/2	1	E	0.07
6.		A5/C/4	1	E	0.07
1.	Improved breed poultry farm	B2/L/1	1	E	0.07
2.		B4/L/2	1	E	0.07
3.		B1/Cl/2	7	AMS, AMC, EX, CAZ, LE, E, CIP	0.5
4.		B2/Cl/1	2	CAZ, E	0.14
5.		B2/Cl/3	1	E	0.07
6.		B1/Fe/1	2	CAZ, E	0.14
7.		B3/Fe/2	3	CAZ, E, TE	0.21
1.	Household backyard poultry	C8/Cl/1	1	E	0.07
2.		C11/Cl/1	1	E	0.07
				Total average	0.117

Characterization of Antimicrobial Resistance Genes: Extended Spectrum Beta Lactamase genes, tetracycline gene and sulphonamides gene

The carriage status of *Salmonella* strains for 04 antimicrobial Beta Lactamase resistance genes, 01 tetracycline (*tetA*) and 01 sulphonamides (*SulI*) were screened by PCR.

All the *Salmonella* isolates were screened for ESBL genes *blaTEM*, *blaSHV*, *blaOXA* and *blaCTX-M*, with an amplicon size of 445bp, 237bp, 813bp and 593bp respectively. Among the 15 isolates, all (100 %) isolates harboured the *blaTEM* gene (Fig.2), whereas none of them were found to be carrying *blaSHV*, *blaOXA* and *blaCTX-M* genes. However, the *tetA* gene encoding resistance to tetracycline was also detected in all the *Salmonella* isolates (Fig 3). With regard to sulphonamides none of them were found to carry the *SulI* gene encoding resistance to sulphonamides

The results of the *blaTEM* gene are in agreement with reports of Bae et al.(2013) who reported 90.91% of isolates harbouring the *blaTEM* gene. Similarly, Eguale et al. (2017) reported the *blaTEM* gene in 79.1%. Several workers reported a lower prevalence of the *blaTEM* gene in *Salmonella* isolates isolated from poultry (Thakur et al. 2013; Elumalai et al. 2014; Herrera-Sánchez et al. 2020). In our study, none of the *Salmonella* isolates were found to carry *blaSHV*, *blaOXA*, and *blaCTX-M* genes. The study subsequently did not identify MDR *Salmonella* spp. with sequences of the ESBL broad-spectrum β -lactamases (*blaSHV*, *blaOXA*, and *blaCTX-M*) in these isolates. The *blaTEM* gene in the *Salmonella* strains could result in the widespread

prevalence of *bla*TEM positive. *Salmonella* by transfer of gene to *bla*TEM negative strains through plasmid conjugation, transformation, and transduction. (Lai et al. 2023).

Our results for carriage of the *tetA* gene are in agreement with earlier reports of Adesijiet al. (2017), Waghmareet al. (2018), and Soufi et al. (2012) who reported *Salmonella* isolates carrying the *tetA* gene in 100.00, 100.00 and 71.00 percent, respectively. Amplified expression of intrinsic resistance mechanisms for tetracycline resistance in microbes is due to the acquisition of mobile genetic elements, ribosomal binding place mutations, and chromosomal mutations (Pavelquesi et al. 2021).

The *SulI* gene is a sulphonamide-resistant dihydropteroate synthase of Gram-negative microorganisms (Alcock et al. 2023). The *SulI* gene is found related to other resistance genes in class 1 integrons (Sköld 2021). A high percentage of *SulI* gene percent in *Salmonella* isolates was reported by Adesiji et al. (2014) who reported 100 percent presence in *Salmonella* spp. isolated from humans and poultry. Machado et al. (2013) and Ma et al. (2017) reported that no one of the isolates was carrying sulfonamide-resistant *Sul* genes. The absence of *SulI* in all isolates, indicates bacteria are not subjected to selective pressure by sulfonamides, a valuable means for the conservation and further addition of resistance to other antimicrobial elements.

Several factors contribute to the progress of antimicrobial resistance in bacteria, including alterations in bacterial cell permeability, enzymatic modification of drugs, and the exclusion of antimicrobials by membrane-bound efflux pumps (Chen et al. 2004). Antibiotic resistance often stems from genetic changes encoded by chromosomal and plasmid genes, with these genes primarily located on integrons, plasmids, and transposons, which are mobile genetic elements (Thong and Modarressi 2011).




Fig. 2 Agarose gel showing PCR product of *bla*TEM gene of *Salmonella* spp.




Fig. 3 Agarose gel showing PCR product of *tetA* gene of *Salmonella* spp.

Conclusions:

The findings of the existing study showed that an improved native poultry breed supply chain is a potential source of virulent *Salmonella* Enterica uncovering humans to potential zoonotic infections via meat, egg, or direct exposure. But lesser MAR along with the absence of genotypic resistance for ESBL broad-spectrum β -lactamases in *Salmonella* isolates was found under study. Minor use of β -lactam and sulphonamide antibiotics in the production chain of improved breed poultry production chain might have resulted in the non-development of resistance. Overall, results suggested there were no major MDR *Salmonella* spp. circulating in the improved native poultry breed production chain of Maharashtra, India. Judicial use of antimicrobials and biosecurity measures shall be implemented in small-scale farms, backyards, and hatcheries principally in countryside areas of Maharashtra, India.

Declarations:

Ethics approval

Not applicable

Consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

Not applicable

Code availability

Not applicable

References

Abunna F, Bedasa M, Beyene T, Ayana D, Mamo B, and Duguma R (2016) *Salmonella*: Isolation and antimicrobial susceptibility tests on isolates collected from poultry farms in and around Modjo, Central Oromia, and Ethiopia. *J. Anim. Poultry Sci.* 5(2): 21-35.

- Adesiji YO, Deekshit VK, and Karunasagar I (2014). Antimicrobial-resistant genes associated with *Salmonella* spp. isolated from human, poultry, and seafood sources. *Food science and nutrition*, 2(4), 436-442.
- Adesiji YO, Shivakumaraswamy SK, Deekshit VK, Kallappa GS and Karunasagar I (2017) Molecular characterization of antimicrobial multi-drug resistance in non-typhoidal *Salmonellae* from chicken and clam in Mangalore, India. *The Journal of Biomedical Research*, 1(1): 1–8.
- Akond MA, Shirin M, Alam S, Hassan SMR *et al* (2012) Frequency of drug-resistant *Salmonella* spp. isolated from poultry samples in Bangladesh. *Stamford Journal of Microbiology*, 2(1), 15-19.
- Al Mamun MA, Kabir SL, Islam MM, Lubna M *et al* (2017) Molecular identification and characterization of *Salmonella* species isolated from poultry value chains of Gazipur and Tangail districts of Bangladesh. *African Journal of Microbiology Research*, 11(11), 474-481.
- Al-Abadi, and Al- Mayah (2012) Isolation and identification of *Salmonella* spp. from chicken and chicken environment in Basrah province. MSc thesis submitted to Department of Pathology and Poultry Diseases, College of Veterinary Medicine Basrah University, Basrah, Iraq.
- Alali WQ, Thakur S, Berghaus RD, Martin MP & Gebreyes WA (2010) Prevalence and distribution of *Salmonella* in organic and conventional broiler poultry farms. *Foodborne pathogens and disease*, 7(11), 1363-1371.
- Alcock BP, Huynh W, Chalil R, Smith KW, Raphenya AR, Wlodarski MA, ... & McArthur AG (2023) CARD 2023: expanded curation, support for machine learning, and resistome prediction at the Comprehensive Antibiotic Resistance Database. *Nucleic acids research*, 51(D1), D690-D699.
- Andino A & Hanning I (2015) *Salmonella enterica*: survival, colonization, and virulence differences among serovars. *The Scientific World Journal*, 2015(1), 520179.
- Anejo-Okopi JA, Isa SE, Audu O, Fagbamila IO *et al* (2016) Isolation and polymerase chain reaction detection of virulence *invA* gene in *Salmonella* spp. from poultry farms in Jos, Nigeria. *Journal of Medicine in the Tropics*, 18(2), 98.
- Bacteriological Analytical Manual (2007) BAM: *Salmonella*, Chapter 5 *Salmonella*. <http://www.fda.gov/downloads/Food/FoodScienceResearch/UCM309839>.
- Bae DH, Dessie HK, Baek HJ, Kim SG *et al* (2013) Prevalence and characteristics of *Salmonella* spp. isolated from poultry slaughterhouses in Korea. *Journal of Veterinary Medical Science*, 75(9), 1193-1200.
- Bhowmick S, Pal S, Sunder J, Sujatha T *et al* (2023) Exploring broilers and native fowls of Andaman and Nicobar Islands as a source of β -lactamase-producing Enterobacteriaceae even with limited anthropogenic activities and docking-based identification of catalytic domains in novel β -lactamase variants. *Frontiers in veterinary science*, 9, 1075133.
- Bhuvaneshwari M, Shanmughapriya S and Natarajaseenivasan K (2015) Prevalence of Multidrug-Resistant (MDR) *Salmonella enteritidis* in Poultry and Backyard Chicken from Tiruchirappalli, India. *Microbiol. J.* 5(2): 28-35.
- CDC: The Centers for Disease Control and Prevention. Antibiotic Resistance Threats in the United States; CDC: Atlanta, GA, USA, 2013.
- Chaiban C, Robinson TP, Fivres EM, Ogola J *et al* (2020) Early intensification of backyard poultry systems in the tropics: A case study. *Animal*. 14, 2387–2396. <https://doi.org/10.1017/S175173112000110X>
- Chen S, Zhao S, White DG, Schroeder CM *et al* (2004) Characterization of multiple-antimicrobial-resistant *Salmonella* serovars isolated from retail meats. *Applied and Environmental Microbiology*, 70(1), 1-7.
- Dashti AA, Jadaon MM, Abdulsamad AM & Dashti HM (2009) Heat treatment of bacteria: a simple method of DNA extraction for molecular techniques. *Kuwait Med J*, 41(2), 117-122.

- Dinh A, Duran C, Singh S, Tesmoingt C, Bouabdallah L, Hamon A & Michelon H (2023) Real-life temocillin use in Greater Paris area, effectiveness and risk factors for failure in infections caused by ESBL-producing Enterobacterales: a multicentre retrospective study. *JAC-Antimicrobial Resistance*, 5(1), dlac132.
- Eguale T, Birungi J, Asrat D, Njahira MN *et al* (2017) Genetic markers associated with resistance to beta-lactam and quinolone antimicrobials in non-typhoidal *Salmonella* isolates from humans and animals in central Ethiopia. *Antimicrobial Resistance & Infection Control*, 6, 1-10.
- Eid S, Ibrahim HM, Shaltot SH and El Oksh AS (2023) An Overview of the Current Situation of Salmonellosis in Pigeons, Household Chickens, and Commercial Broilers with a Special Reference to a Customized Vaccine Developing Trial. *Journal of Advanced Veterinary Research*, 13(3), 322-332.
- Elmonir W, Abdeltawab D, El-Sharkawy H and Zahran RN (2023) Serotypes Diversity, Virulence, and Antimicrobial Resistance of Non-Typhoidal *Salmonella* Isolates in Commercial and Backyard Egg Production Systems in Egypt.
- Elumalai S, Muthu G, Esther R, Selvam M and Ramesh S (2014) Detection of TEM-, SHV- and CTX-M-type β -lactamase production among clinical isolates of *Salmonella* species. *Journal of Medical Microbiology* 63: 962–967.
- Fahmy SS, Dahshan AH, & El Sawah A (2023) Prevalence of Antimicrobials and Virulence-Related Genes in *Salmonellae* Detected in Local Hatcheries in Northern Upper Egypt. *Journal of Veterinary Medical Research*, 30(2), 71-77.
- Fang H, Ataker F, Hedin G and Dornbusch K (2008) Molecular epidemiology of extended-spectrum β -lactamases among *Escherichia coli* isolates collected in a Swedish hospital and its associated health care facilities from 2001 to 2006. *Journal of Clinical Microbiology*, 46(2), 707-712.
- Hedman H D, Vasco K A and Zhang L (2020) A review of antimicrobial resistance in poultry farming within low-resource settings. *Anim. Open Access J. MDPI* 10, 1264. <https://doi.org/10.3390/ani10081264>.
- Heredia N and Garcia S (2018) Animals as sources of food-borne pathogens: A review. *Animal Nutrition*, 4(3), 250-255.
- Herrera-Sánchez MP, Rodríguez-Hernández R and Rondón-Barragán IS (2020) Molecular characterization of antimicrobial resistance and enterobacterial repetitive intergenic consensus-PCR as a molecular typing tool for *Salmonella* spp. isolated from poultry and humans. *Veterinary World*, 13(9), 1771–1779.
- Igbinosa EO, Beshiru A, Igbinosa IH and Okoh AI (2022) Antimicrobial resistance and genetic characterisation of *Salmonella enterica* from retail poultry meats in Benin City, Nigeria. *LWT*, 169, 114049.
- Jajere SM, Hassan L, Aziz SA, Zakaria Z *et al* (2019) *Salmonella* in native “village” chickens (*Gallus domesticus*): prevalence and risk factors from farms in South-Central Peninsular Malaysia. *Poultry Science*, 98(11), 5961-5970.
- Khanal PR, Satyal D, Bhetwal A, Maharjan A, Shakya S, Tandukar S, Parajuli NP (2017) Renaissance of Conventional First-Line Antibiotics in *Salmonella enterica* Clinical Isolates: Assessment of MICs for Therapeutic Antimicrobials in Enteric Fever Cases from Nepal. *BioMed Research International*, 2017(1), 2868143.
- Koro A, Elezaj I, Hadžiabdić S, Alić A and Rešidbegović E (2022) Occurrence of *Salmonella* spp. in backyard poultry in Bosnia and Herzegovina. *Iranian Journal of Veterinary Research*, 23(1).
- Krumperman PH (1983) Multiple antibiotic resistance indexing of *Escherichia coli* to identify high-risk sources of fecal contamination of foods. *Applied and Environmental Microbiology*, 46(1), 165-170.

- Lai J, Mu H, Zhou B, He J, Cheng X, Gan Y & Ding H (2023) Bla TEM-positive *Salmonella enterica* serovars Agona and Derby are prevalent among food-producing animals in Chongqing, China. *Frontiers in Microbiology*, 14, 1011719.
- Ma M, Wang H, Yu Y, Zhang D and Liu S (2007) Detection of antimicrobial resistance genes of pathogenic *Salmonella* from swine with DNA microarray. *Journal of Veterinary Diagnostic Investigation*, 19(2), 161-167.
- Ma S, Lei C, Kong L, Jiang W *et al* (2017) Prevalence, antimicrobial resistance, and relatedness of *Salmonella* isolated from chickens and pigs on farms, abattoirs, and markets in Sichuan Province, China. *Foodborne Pathog. Dis.* <https://doi.org/10.1089/fpd.2016.2264>.
- Machado E, Coque TM, Cantón R, Sousa JC, Peixe L (2013) Commensal *Enterobacteriaceae* as reservoirs of extended-spectrum beta-lactamases, integrons, and sul genes in Portugal. *Front., Microbiol.* 2013;4:80. <https://doi.org/10.3389/fmicb.2013.00080>.
- Muhammad M, Muhammad LU, Ambali AG, Mani AU, Azard S and Barco L (2010) Prevalence of *Salmonella* associated with chick mortality at hatching and their susceptibility to antimicrobial agents. *Veterinary Microbiology*, 140(1-2), 131-135.
- Mulika L and Yuwapanichsampan S (2011) Prevalence of *Salmonella* spp. and Its Resistance to Antimicrobial Drugs in Poultry Hatchery. *KKU Veterinary Journal*, 18(1), 12-28.
- Nair A, Balasaravanan T, Malik SS, Mohan V *et al* (2015) Isolation and identification of *Salmonella* from diarrheagenic infants and young animals, sewage waste and fresh vegetables. *Veterinary World*, 8(5), 669.
- Ng LK, Martin I, Alfa M and Mulvey M (2001) Multiplex PCR for the detection of tetracycline resistant genes. *Molecular and cellular probes*, 15(4), 209-215.
- Orabi A, Armanious W, Radwan IA, Girh ZM, Hammad E, Diab MS & Elbestawy AR (2022) Genetic correlation of virulent *Salmonella* serovars (Extended Spectrum β -Lactamases) isolated from broiler chickens and human: a public health concern. *Pathogens*, 11(10), 1196.
- Orji MU, Onuigbo HC and Mbata TI (2005) Isolation of *Salmonella* from poultry droppings and other environmental sources in Awka, Nigeria. *Int. J. Infect. Dis.* 9: 86-89.
- Pavelquesi SLS, de Oliveira Ferreira ACA, Rodrigues ARM, de Souza Silva CM, Orsi DC & da Silva ICR (2021) Presence of tetracycline and sulfonamide resistance genes in *Salmonella* spp.: literature review. *Antibiotics*, 10(11), 1314.
- Pavon RDN & Rivera WL (2021) Molecular serotyping by phylogenetic analyses of a 1498bp segment of the *invA* gene of *Salmonella*. *ASM Sci. J*, 14, 1-14.
- Rothrock MJ, Hiatt KL, Guard JY & Jackson CR (2016) Antibiotic resistance patterns of major zoonotic pathogens from all-natural, antibiotic-free, pasture-raised broiler flocks in the Southeastern United States. *Journal of Environmental Quality*, 45(2), 593-603.
- Salehi TZ, Mahzounieh M and Saeedzadeh A (2005) Detection of *invA* gene in isolated *Salmonella* from broilers by PCR method. *Int. J. Poult. Sci*, 4(8), 557-559.
- Samanta I, Joardar SN, Das PK, Sar TK *et al* (2014) Prevalence and antibiotic resistance profiles of *Salmonella* serotypes isolated from backyard poultry flocks in West Bengal, India. *Journal of Applied Poultry Research*, 23(3), 536-545.
- Siemon CE, Bahnson PB & Gebreyes WA (2007) Comparative investigation of prevalence and antimicrobial resistance of *Salmonella* between pasture and conventionally reared poultry. *Avian diseases*, 51(1), 112-117.

- Singh M, Mollier RT, Paton RN, Pongener N *et al* (2022) Backyard poultry farming with improved germplasm: Sustainable food production and nutritional security in fragile ecosystem. *Front. Sustain. Food Syst.* 6:962268.
- Sivaramalingam T, Pearl DL, McEwen SA, Ojkic D & Guerin MT (2013) A temporal study of *Salmonella* serovars from fluff samples from poultry breeder hatcheries in Ontario between 1998 and 2008. *Canadian Journal of Veterinary Research*, 77(1), 12-23.
- Sköld O (2000) Sulfonamide resistance: mechanisms and trends. *Drug resistance updates*, 3(3), 155-160.
- Sohail MN, Rathnamma D, Priya SC, Isloor S *et al* (2021) *Salmonella* from farm to table: isolation, characterization, and antimicrobial resistance of *Salmonella* from commercial broiler supply chain and its environment. *BioMed Research International*, 2021.
- Soufi L, Saenz Y, Toro M, Abbassi MS *et al* (2012) Phenotypic and Genotypic Characterization of *Salmonella enterica* Recovered from Poultry Meat in Tunisia and Identification of New Genetic Traits. *Vector Borne Zoonotic Dis.* 12 (12): 10–16.
- Talukder M, Islam MS, Ievy S, Sobur MA *et al* (2021) Detection of multidrug resistant *Salmonella* spp. from healthy and diseased broilers having potential public health significance. *J. Adv. Biotechnol. Exp. Ther.* 4(2), 248-255.
- Taneja N, Appannanavar SB, Kumar A, Varma G *et al* (2014) Serotype profile and molecular characterization of antimicrobial resistance in non-typhoidal *Salmonella* isolated from gastroenteritis cases over nine years. *Journal of Medical Microbiology*, 63(1), 66-73.
- Thakur S, Brake J, Keelara S, Zou M and Susick E (2013) Farm and environmental distribution of *Campylobacter* and *Salmonella* in broiler flocks. *Research in Veterinary Science*, 94(1), 33–42.
- Thong KL and Modarressi S (2011) Antimicrobial resistant genes associated with *Salmonella* from retail meats and street foods. *Food Research International*, 44(9), 2641-2646.
- Trampel DW, Holder TG & Gast R K (2014) Integrated farm management to prevent *Salmonella* Enteritidis contamination of eggs. *Journal of Applied Poultry Research*, 23(2), 353-365.
- Van Boeckel TP, Pires J, Silvester R, Zhao C *et al* (2019) Global trends in antimicrobial resistance in animals in low- and middle-income countries. *Science* 2019, 365.
- Van Loo EJ, Caputo V, Nayga Jr RM, Meullenet JF & Ricke SC (2011) Consumer's willingness to pay for organic chicken breast: Evidence from choice experiment. *Food quality and preference*, 22(7), 603-613.
- Varijakshapanicker P, Mckune S, Miller L, Hendrickx S *et al* (2019) Sustainable livestock systems to improve human health, nutrition, and economic status. *Animal Frontiers*, 9(4), 39-50.
- Waghmare RN, Paturkar AM, Vaidya VM, Zende RJ *et al* (2018) Phenotypic and genotypic drug resistance profile of *Salmonella* serovars isolated from poultry farm and processing units located in and around Mumbai city, India. *Veterinary world*, 11(12), 1682.
- Waghmare RN, Popalghat HK, Londhe SV, Deshmukh VV and Khobe VV (2021) An Online survey of consumers of Maharashtra concerning the expected change in the meat and meat product business.
- WHO. Prioritization of Pathogens to Guide Discovery, Research and Development of New Antibiotics for Drug-Resistant Bacterial Infections, Including Tuberculosis. 2017. Available online: [www.who.int/medicines/areas/rational_use/PPLreport_2017_09_19 .pdf?ua=1](http://www.who.int/medicines/areas/rational_use/PPLreport_2017_09_19.pdf?ua=1) (accessed on 14 December 2022).

Withenshaw SM, Cawthraw S, Gosling B, Newton K *et al* (2021) Risk factor analysis for *Salmonella* contamination of broiler chicken (*Gallus gallus*) hatcheries in Great Britain. *Preventive Veterinary Medicine*, 196, 105492.

Xu X, Biswas S, Gu G, Elbediwi M, Li Y and Yue M (2020) Characterization of multidrug resistance patterns of emerging *Salmonella enterica* serovar Rissen along the food chain in China. *Antibiotics*. doi: 10.3390/antibiotics9100660.

Yoke Kqueen C, Learn Han L, Noorzaleha AS, Son R, Sabrina S, Jiun Horng S and Chai Hoon K (2008) Characterization of multiple antimicrobial resistant *Salmonella enterica* subsp. *enterica* isolated from indigenous vegetables and poultry in Malaysia. *Letters in applied microbiology*, 46(3), 318-324.

Zdragas A, Mazaraki K, Vafeas G, Giantzi V, Papadopoulos T and Ekateriniadou L (2012) Prevalence, seasonal occurrence and antimicrobial resistance of *Salmonella* in poultry retail products in Greece. *Lett. Applied Microbiol.*, 55: 308-313.

Zhao X, Ju Z, Wang G, Yang J, Wang F, Tang H & Sun S (2021) Prevalence and antimicrobial resistance of *Salmonella* isolated from dead-in-shell chicken embryos in Shandong, China. *Frontiers in Veterinary Science*, 8, 581946.

(Revise by following authors guidelines for the journal)

UNDER PEER REVIEW