

1
2
3
4
5
6
7
8
9
10
11
12
13

Antimicrobial Resistance Profile of *Salmonella enterica* Isolated from Improved Poultry Breed Farming Chain, Maharashtra, India

ABSTRACT

Aims: With rising demand for poultry meat, globally backyard farms have adopted semi-intensive systems using improved birds. Food safety is a prime issue in antimicrobial-resistant *Salmonella* spp. found in meat, which has often been traced back to farms.

Study design: We carried out a cross-sectional prospective study.

Place and Duration of Study: The study was conducted in selected districts of Maharashtra India during October 2023 to March 2024.

Methodology: A total of 364 samples were collected from improved indigenous poultry breed hatcheries (n=5), improved indigenous breed commercial farms (n=5), and improved breed rearing backyard poultry setups (n=30) in Marathwada, Maharashtra state, India. The isolation of *Salmonella* spp. was carried out as per IS-5887 (Part 3): 1999 protocol. The bacterial isolates were further identified by cultural, microscopic morphology and biochemical characteristics (BAM 2007). Confirmation of *Salmonella* spp. was done by polymerase chain reaction (PCR) assays with primers designed for the *invA* gene. All *Salmonella* isolates were subjected for antibiotic susceptibility testing by disk diffusion method against 14 commonly used antimicrobials. Further, isolates were characterized for the presence of *bla*TEM, *bla*SHV, *bla*OXA, *bla*CTXM, *tetA*, and *Sul1* AMR genes.

Results: A total of 15 isolates recovered with prevalence rates of 4.83, 6.36, and 1.53 percent in hatcheries, farms, and backyard households, respectively. Five isolates that were randomly analyzed showed a homologous sequence as *Salmonella enterica*. Antimicrobial susceptibility testing of all isolates revealed higher resistance against Erythromycin (100%), followed by Ceftazidime (40%), while sensitivity (93.33%), was recorded against Ampicillin/Sulbactam, Amoxicillin/Sulbactam, and Enrofloxacin. The average multiple antibiotic resistance (MAR) index, of *Salmonella* isolates was 0.117. Genotypic resistance pattern revealed that all isolates (100%) were carrying the *bla*TEM gene, while none were harbouring broad spectrum extended-spectrum beta-lactamases (*bla*SHV, *bla*OXA, and *bla*CTXM) genes. All isolates were positive for the *tetA* gene (100%) but none of them was positive for *Sul1* genes.

Conclusion: The study highlights the low antimicrobial resistance in *Salmonella* isolates isolated from Improved Poultry Breed Farming Chain, which might be due to low usage of antimicrobials. However, monitoring of multidrug-resistant *Salmonella enterica* in improved poultry breed farming chains is essential to protect human health.

15 *Keywords: improved poultry breed, farming chain, Salmonella, antimicrobial resistance,*
16 *genotypic resistance*

17

18 **1. INTRODUCTION**

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

Poultry meat consumption is on the rise in world among all animal-derived food items (Waghamare et al. 2021). Meat, due to its high nutritive value containing essential amino acids and various macro- and micronutrients, plays a crucial role in a balanced diet (Van Boeckel et al. 2019). In recent years with the surge in chicken demand, backyard poultry farms have transformed themselves into semi-intensive farming systems with the help of improved native or coloured variety birds and good quality feed (Chaiban et al. 2020). The intervention of backyard poultry farming with improved native or coloured poultry varieties suitable for backyard production can be a source of a supportable food production structure (Singh et al., 2022). Breeder stock and hatcheries can be an important source of *Salmonella* spp. and can be improved through the execution of effective intervention methods (Sivaramalingam et al. 2013).

Salmonellosis stands as a major global foodborne illness, contributing to 93.8 million gastroenteritis cases and 155,000 deaths per annum globally (Heredia and Gracia, 2018). *Salmonella* in food chains is emerging worldwide and poultry is recognized as a foodborne pathogen reservoir, with several reports highlighting the occurrence of *Salmonella* linked to the backyard and commercial live poultry, production settings (Samanta et al. 2014, Elmonir et al. 2023).

In the past few years, there has been a significant surge in phenotypic and genotypic resistance among non-typhoidal *Salmonella* isolates to β -lactams, tetracycline, and sulphonamides (Egualde et al. 2017). The extensive use of tetracycline in poultry has caused the advance of tetracycline resistance in *Salmonella* spp. due to selective pressure (Waghamare et al. 2018). CDC, 2013, and WHO, 2017 reported that Gram-negative microorganisms that harvest beta-lactamases as one of the world's most insistent threats. Recent food safety studies have shown that *Salmonella* strains isolated from poultry and other foods possess extended-spectrum β -lactamases (ESBLs), which become a potential threat to human health as they may impact treatment regimens for ESBL-producing pathogens (Orabi et al. 2022; Dinh et al. 2023).

Broiler meat harvested from organic and pastured poultry farming production facilities has high demand (Van Loo et al. 2011; Rothrock et al. 2016). The appearance of antimicrobial resistance (AMR) in *Salmonella* from backyard poultry farming systems may pose a significant growing threat to anthropoid and animal health (Van Boeckel et al. 2019;

52 Hedman et al. 2020). Similarly, studies conducted by other researchers reported significantly
53 lower *Salmonella* contamination of fecal matter and bird feed (Siemon et al. 2007; Alali et al.
54 2010). An additional hypothesis is that improved breed poultry production systems work on
55 antibiotic-free rearing, which might potentially affect the antimicrobial resistance profile and
56 presence of antibiotic resistance genes in the *Salmonella* populations along the hatchery-to-
57 backyard farm continuum.

58 Therefore, to better understand the prevalence of antimicrobial resistant (AMR)
59 *Salmonella* in these improved native breed poultry farms and backyard poultry management
60 systems, some more work is essential; especially since market demand and production for
61 backyard poultry has been increased. Thus, the present study was carried out to evaluate
62 the antimicrobial resistance profile of *Salmonella* spp. isolated from the backyard poultry
63 production system. This information will provide vital data on the prevalence, and AMR
64 profile of *Salmonella* spp. in improved indigenous poultry breed farming chains which help to
65 improve backyard poultry food safety systems.

66

67 **2. MATERIAL AND METHODS**

68 **2.1 Sample collection**

69 In the current study, a total of 364 samples were collected from improved indigenous poultry
70 breed hatcheries (n=5), improved indigenous breed commercial farms (n=5), and improved
71 breed-rearing backyard poultry setups (n=30) in Marathwada, Maharashtra state, India. The
72 samples from hatcheries (n=124) were comprised of deceased chicks (25), unhatched eggs
73 (25), shell waste (12), and tray swabs (12). A total of 110 samples (10 each of feed, water,
74 litter, feces, walls swab, utensils swab, worker's hands swab, and cages; and 30 cloacal
75 swabs) were selected from five different farms in Pathardi, Maharashtra, India. Similarly, 30
76 household backyard farms from the Parbhani, Vasmat, and Selu areas of Marathwada,
77 Maharashtra, India were screened by collecting 130 samples (30 each of cloacal swabs,
78 water and feces, 15 each of cage dust swabs and utensils swab and 10 feed samples). All
79 the samples were collected aseptically, using sterilized polypropylene bags, and transported
80 to the laboratory under a cold chain as per standard methods.

81 **2.2 Isolation and characterization of *Salmonella* spp.**

82 Isolation of *Salmonella* spp. was performed as per IS-5887 (Part 3): 1999 protocol with slight
83 modifications based on the samples collected. The sample was processed (pre-enrichment)
84 by inoculating a 5gm sample of dead chick and unhatched egg in 45 ml of buffered peptone
85 water (BPW), swabs and feces (1gm) were pre-enriched with 9 ml buffered peptone water, in
86 sterile test tubes, whereas 25 g of each feed and water was inoculated in 225 ml of buffered
87 peptone water and all samples were incubated at 37°C ± 1°C for 24 hours. Further,

88 enrichment was done by transferring 0.1 ml of inoculum from BPW into 10 ml Rappaport-
89 Vassiliadis (RV) medium and incubated at 42°C for 24 hrs. A loop-full of inoculum from RV
90 broth was streaked on Xylose Lysine Deoxycholate (XLD) Agar and Brilliant Green Sulpha
91 (BGSA) Agar, plates by four quadrant streaking method. The inoculated plates were
92 incubated at 37°C ± 10°C for 24 hrs. for the development of colonies of *Salmonella* spp. The
93 bacterial isolates were further identified by cultural, microscopic morphology, and
94 biochemical characteristics (BAM 2007).

95 Black center smooth and round or large with black center colonies of *Salmonella* spp. were
96 characterized by polymerase chain reaction (PCR) assays with primers designed for the
97 *invA* gene (Nair et al. 2015). DNA of *Salmonella* isolates was extracted using heat lysis DNA
98 Isolation protocol (Dashti et al. 2023). The details of PCR primers and cycling conditions
99 used in the present study are mentioned in Table 1 and Table 2.

100 **2.3 Sequencing of the *invA* gene amplified product**

101 Amplified *invA* gene products of selected isolates from hatchery, farm, and backyard
102 households were sequenced from an outsourced agency (Submitted to GeneOmbio
103 Technologies Pvt Ltd, India) for sequencing. The resulting sequences were analyzed, and
104 sequence homology searches were conducted using the BLAST algorithm
105 (https://blast.ncbi.nlm.nih.gov/Blast.cgi?PROGRAM=blastn&BLAST_SPEC=GeoBlast&PAGE_TYPE=BlastSearch).
106

107 **2.4 Antimicrobial susceptibility test**

108 As per the Clinical and Laboratory Standards Institute (CLSI) protocol, the Kirby–Bauer drug-
109 sensitive disk technique was used to test the sensitivity of PCR-positive *Salmonella* isolates
110 to 14 commonly used antimicrobials. Antimicrobial tested were Ampicillin/Sulbactam (A/S)
111 10/10 µg, Amoxicillin/Sulbactam (AMS) 30/15 µg, Amoxiclav (AMC), Tetracycline (TE) 30
112 µg, Enrofloxacin (EX) 10 µg, Ceftazidime (CAZ) 30 µg, Levofloxacin (LE) 5 µg,
113 Chloramphenicol (C) 30 µg, Erythromycin (E) 15 µg, Gentamicin (GEN) 10 µg, Amikacin
114 (AK) 10 µg, Ciprofloxacin (CIP) 5 µg, Co-Trimoxazole (COT) 25 µg and Nalidixic Acid (NA)
115 30 µg. Briefly, bacterial suspensions were achieved from overnight-grown cultures, which
116 adjusted to the 0.5 McFarland turbidity standard. The broth culture was spread on the
117 surface of a Muller Hinton agar plate using a sterile cotton swab. After about 20 min, the
118 disks were applied to the plates and incubated at 37°C for 18 hr. Finally, using a scale
119 (Himedia) the diameter of the inhibition zone was measured to categorize the resistance
120 pattern. The results of phenotypic antimicrobial resistance patterns were analyzed as per the
121 European Committee for Antimicrobial Susceptibility Testing (EUCAST) guidelines (2021).

122 **2.5 Multiple Antibiotic Resistance (MAR) Index**

123 The Multiple Antibiotic Resistance (MAR) index of individual isolates was calculated
 124 according to the method described by Krumperman (1983) by dividing the number of
 125 antibiotics to which the isolates were found resistant to the total number of antibiotics to
 126 which the isolate was exposed.

127 **2.6 Molecular characterization of AMR genes in *Salmonella* spp.**

128 All the 15 isolates were screened for the presence of 06 antimicrobial resistance
 129 genes. The genes that encode resistance to β -lactamases (*blaTEM*, *blaSHV*, *blaCTX-M*, and
 130 *blaOXA*), tetracyclines (*tetA*), and sulfonamides (*Sul1*) were evaluated.

131 Genomic DNA of *Salmonella* spp. isolates were extracted by boiling method
 132 described by Anejo-Okopiet *al.* (2016). All *Salmonella* isolates were screened for the
 133 presence of antimicrobial resistance genes through singleplex and multiplex PCR protocol
 134 described by Ng *et al.* (2001) for *tetA*, Ma *et al.* (2017) for *Sul1* and Fang *et al.* (2008) for a
 135 group of β -lactamase genes (*blaSHV*, *blaCTXM*, *blaOXA* and *blaTEM*). Primers used are
 136 listed in Table 1. The cycling conditions for PCR are mentioned in Table 2.

137 The singleplex PCR for *invA*, *sul1* and *tetA* was performed in 25 μ l reaction volume
 138 containing 12.5 μ l of 2x PCR master mix (HiMedia Laboratories Pvt. Ltd., Mumbai), 1.5 μ l of
 139 each primer (10pmol/ μ l) (Eurofins Genomics India Pvt. Ltd., Bangalore), 1 μ l of genomic
 140 DNA and 8.5 μ l molecular biology grade water (HiMedia Laboratories Pvt. Ltd., Mumbai)
 141 used to make desired volume. Whereas, multiplex PCR reaction was performed as
 142 described by Fang *et al.* (2008) in a 25 μ l volume containing 12.5 μ l 2x PCR Master Mix
 143 (Takara Bio Inc., Shiga, Japan) supplied with Taq DNA polymerase, buffer, MgCl₂, and
 144 dNTPs. In this PCR 1 μ l (10 pmol/ μ l) each of forward and reverse primer was used.
 145 Similarly, the reaction mixture contained 3 μ l DNA template and 7.5 μ l nuclease-free water to
 146 make a final volume of 25 μ l.

147 **Table 1. Primers used for genotype characterization of *Salmonella* spp.**

Sr. no	Primer	Target	Primer sequence (5'- 3')	Product Size (bp)	Reference
1	<i>invA</i>	Invasion-associate protein	F:TCGTGACTCGCGTAAATGGCGATA R:GCAGGCGCACGCCATAATCAATAA	423	(Nair et al. 2015)
2	<i>tetA</i>	Tetracycline	F: GCT ACA TCC TGC TTG CCT TC R: CAT AGA TCG CCG TGA AGA GG	210	(Ng et al. 2001)
3	<i>Sul1</i>	Sulphonamide	F: TTTCTGACCCTGCGCTCTAT R:GTGCGGACGTAGTCAGCGCCA	425	(Ma et al. 2007)
4	<i>blaTEM</i>	Broad Spectrum β -lactamase	F:CGCCGCATACACTATTCTCAGAATGA R: ACGCTCACCGGCTCCAGATTTAT	445	(Fang et al. 2008)
5	<i>blaSHV</i>		F: CTTTATCGGCCCTCACTCAA	237	

		R: AGG TGC TCA TCA TGG GAA AG	
6	<i>blaCTX-M</i>	F: ATGTGCAGYACCAGTAARGTKATG GC R:TGGGTRAARTARGTSACCAGAAAYCAGCGG	593
7	<i>blaOXA</i>	F: ACA CAA TAC ATA TCA ACT TCG C R: AGT GTG TTT AGA ATG GTG ATC	813

148

149

Table 2. PCR cycling conditions used under study for different primers

PCR steps	PCR conditions	Target genes			Broad Spectrum β -lactamase
		<i>invA</i>	<i>tetA</i>	<i>sul1</i>	<i>blaSHV, blaCTXM, blaOXAblaTEM</i> (Multiplex PCR)
Initial	Temperature (°C)	94	95	94	94
Denaturation	Time (min.)	5	3	3	5
Denaturation	Temperature (°C)	94	95	94	94
	Time (sec.)	30	30	30	45
Annealing	Temperature (°C)	56	60	60	63
	Time (sec.)	60	30	30	60
Extension	Temperature (°C)	72	72	72	72
	Time (sec)	90	60	45	60
Final	Temperature (°C)	72	72	72	72
	Time (min.)	10	8	8	7
Number of cycles		35	35	30	30

150

151

152

153

3. RESULTS AND DISCUSSION

3.1 Prevalence of *Salmonella* isolates:

154

155

156

157

158

159

160

161

162

163

As is shown in Table 3, a total of 15 PCR *invA* gene positive *Salmonella* isolates were recovered from 364 samples, in which *Salmonella* was detected from Dead chick, Unhatched egg, Eggshell waste, Cloacal swabs, Faeces and Litter. All *Salmonella* isolates yielded desired amplified products of 423 bp for the *invA* gene (Fig. 1). The PCR-based method utilizing the *invA* genus-specific primer is a reliable approach due to its rapidity, high specificity, and sensitivity in confirming the isolates (Salehi et al. 2005).

Among the isolates, *Salmonella* strains were observed in 4.83% (6/124) of those from Hatcheries, 6.76% (7/110) of those from commercial farms, and 1.53% (2/130) of those from backyard household farms.

164 *Salmonella* can be spread horizontally and through fertilized eggs and it is zoonotic
165 in nature (Zhao et al. 2021). Considering this, in obligation to better measure the noticeable
166 infection by *Salmonella* spp., we obtained several samples from hatcheries, farms and
167 backyard farms from various locations by maintaining the supply chain in improved
168 indigenous poultry breed farming.

169 Results of prevalence from hatcheries are in agreement with earlier studies of
170 Muhammad *et al.* (2010) and Sohailet *al.* (2021) who reported occurrence of 4.4% and
171 10.34% in hatchery samples, whereas, similar occurrence was reported by Mulikaet *al.*
172 (2011), Xu *et al.* (2020) and Withenshawet *al.* (2021) wherein, the prevalence of
173 *Salmonella* spp.in hatchery samples ranging from 16.6 to 33.3% was recorded.
174 Decontamination, sanitation, waste disposal, biosecurity measures, and proper monitoring
175 are necessary to prevent *Salmonella* transmission from hatchery to another setting. The
176 improvement of hatchery hygiene and the application of efficient pathogen detection and
177 disease control strategies would improve *Salmonella* control (Fahmy et al. 2023).

178 The prevalence of *Salmonellaenterica* in Indigenous Poultry farms agrees with
179 previous studies by Samantaet *al.* (2014), who reported a prevalence of 6.1 percent
180 respectively. Abunnaet *al.* (2016) stated that the distribution of *Salmonella* spp. varied
181 depending on the sample type, poultry growth stage, and breeds. Soil and fecal matter
182 present on the feathers and feet of birds serve as significant contributors to the microbial
183 contamination observed in poultry carcasses (Orji et al. 2005). The results of the
184 prevalence of *Salmonella enterica* in household backyard poultry farming differ with
185 previous reports by Jajereet *al.* (2019), Koro *et al.* (2022) and Elmoniret *al.* (2023). In
186 household backyard poultry farming *Salmonella* spp. was observed only in cloacal swab
187 samples (6.67%). The findings are not per the findings of Bhowmicket *al.* (2023) and Eid *et*
188 *al.* (2023) who reported 30.82 and 31.00% prevalence, respectively in backyard poultry
189 farm cloacal swabs. The presence of *Salmonella* in the backyard poultry might have been
190 the sequel of a long-term persistence of *Salmonella* in these backyards or it could be the
191 summary of laying hens purchased directly from contaminated hatcheries or other
192 households (Trampel et al. 2014). The variances in sample types, topographical locations,
193 sampling protocols, or isolation methods resulted in a multifarious and diverse isolation
194 assessment.

195

196 **Table 3. Prevalence of *Salmonella* in Backyard Poultry Farming System (n = 364)**

Category	Type of Samples analyzed	Total Number	Positive Sample type	Total Number	Overall prevalence
----------	--------------------------	--------------	----------------------	--------------	--------------------


		of Samples		(%)
Hatchery	Dead chick (internal organ)	124	Dead chick (2)	6 4.83%
	Unhatched Egg (Yolk and Embryo)		Unhatched Egg (3)	
	Tray swab		Shell waste (1)	
	Eggshell waste			
Indigenous poultry breed farms	Cloaca swab	110	Cloaca swab (3)	7 6.36%
	Utensils swab		Faeces (2)	
	Wall dust swab		Litter (2)	
	Workers hand swab			
	Feed			
	Water			
	Faeces			
Household backyard poultry	Cloaca swab	130	Cloacal swab	2 1.53%
	Cage dust swab			
	Utensils swab			
	Feed			
	Water			
	Faeces			

197

198 **3.2 Sequencing of selected *invA* gene PCR Products for serotype confirmation**

199 A total of five selected isolates from hatcheries (A5/S/2 and A5/U/3), farms (B1/CI/2)
200 and household backyard poultry (C8/CI/1 and C11/CI/1) were sequenced using amplified
201 *invA* gene (423bp) product. Pavon and Rivera (2021) suggested the possible use of the *invA*
202 virulence gene for molecular typing of *Salmonella* through sequencing. The sequencing
203 homologous was observed on NCBI as *Salmonellaenterica* and showed homology of
204 99.49%, 99.69% and 99.49% with sequences accession no. AP020332.1, CP074202.1,
205 CP051329.1, respectively which were available on NCBI.

206 Amongst various types, SerovarsEnteritidis and Typhimurium were the most
207 informed serovars. *S. enterica* is responsible for infections in humans and animals
208 (Andino&Hanning., 2015) which makes it a serious concern.



209

210 **Fig. 1 Agarose gel showing PCR amplified product of *invA* gene of *Salmonella* spp.**

211 **3.3 Antimicrobial resistance pattern of *Salmonella* isolates**

212 In the present study antimicrobial susceptibility testing discovered upper resistance against
 213 Erythromycin (100%), followed by Ceftazidime (40%), while sensitivity was recorded against
 214 Ampicillin/Sulbactam (93.33%), Amoxicillin/Sulbactam (93.33%), Enrofloxacin (93.33%),
 215 Levofloxacin (93.33%), and Nalidixic Acid (93.33%), whereas, intermediate sensitivity
 216 recorded against Ceftazidime (53.34%), followed by Tetracycline (46.67%) and Amoxiclav
 217 (26.67%). In the present study, a higher resistance pattern against Erythromycin (100%) is in
 218 agreement with the findings of Bhuvanewa et al. (2015) and Al Mamum et al. (2017). The
 219 resistance pattern of Ceftazidime (40%) was noticed by Herrera-Sánchez et al. (2020),
 220 wherein the resistance reported was 75.5 percent. The results about resistance to
 221 Ampicillin/Sulbactam (0%), Gentamicin (0%), Amikacin (0%), Co-Trimoxazole (0%), Nalidixic
 222 acid (0%), Amoxicillin/Sulbactam (6.67%), Amoxyclav (6.67%), Tetracycline (6.67%),
 223 Enrofloxacin (6.67%), Levofloxacin (6.67%), Chloramphenicol (6.67%), Ciprofloxacin
 224 (6.67%) differs from results of with Akond et al. (2012), Thakur et al. (2013), Samanta et al.
 225 (2014) and Waghmare et al. (2018). Kariuki et al, (2015) reported that recommended
 226 treatment for non-typhoid *Salmonella* include ciprofloxacin as first-line treatment,
 227 whereas alternatives could be levofloxacin, moxifloxacin, cotrimoxazole or the extended
 228 spectrum cephalosporins ceftriaxone or cefotaxime. Bhuvanewa et al. (2015) reported that
 229 β -lactam and macrolide antibiotics are used routinely for curbing bacterial infection among
 230 chickens in commercial farms compared to backyard farms. The resistance of *Salmonella*
 231 Enteritidis to ampicillin decreased in our study this might be due to the reduction in antibiotic
 232 selective pressure (Varijakshapanicker et al. 2019). The exercise of using antimicrobial

233 agents for growth, and treatment in livestock decreases their efficacy and is found to be a
 234 significant factor in the emergence of multidrug-resistant *Salmonella* (Thakur et al. 2013).

235

236 **3.4 Multiple Antibiotic Index (MAR) for *Salmonella* isolates**

237 In the present study, only 3 *Salmonella* isolates were resistant to 3 or more than 3
 238 antibiotics. The MAR index of A5/S/2, B1/CL/2, and B3/Fe/2 were 0.21, 0.50, and 0.21,
 239 respectively (Table 4). The MAR index of the remaining 12 isolates varies from 0.07 to 0.14
 240 which indicates lower selective pressure for antimicrobial resistance. The results of the
 241 present study do not agree with the studies carried out by Yoke-Kqueen et al. (2008) and
 242 Talukder et al. (2021) who reported that 91.01 and 100 percent *Salmonella* isolates have a
 243 MAR index of more than 0.2. The lower multidrug resistance pattern displayed by
 244 *Salmonella* isolates demonstrates that anthropoids consuming meat and meat products from
 245 concerned geographical areas are at low risk of multidrug-resistant *Salmonella* infection. The
 246 isolates further need to be subjected for detection of Minimal inhibitory concentration (MIC)
 247 of antimicrobial under study. MIC has a significant impact on the choice of a therapeutic
 248 strategy and reliable assessment of efficiency of an infection therapy with suitable
 249 antimicrobial (Kowalska-Krochmal and Dudek-Wicher 2021).

250 **Table 4. Multiple antibiotic resistance [MAR] index and antibiogram of *Salmonella***
 251 **isolates**

Sr. no.	Source of samples	Sample code	Resistance to the number of Antibiotics	Antibiogram	MAR index
1.	Hatchery	A5/S/2	3	CAZ, C, E	0.21
2.		A5/U/3	1	E	0.07
3.		A5/U/4	2	CAZ, E	0.14
4.		A5/U/5	1	E	0.07
5.		A5/C/2	1	E	0.07
6.		A5/C/4	1	E	0.07
1.	Improved breed poultry farm	B2/L/1	1	E	0.07
2.		B4/L/2	1	E	0.07
3.		B1/Cl/2	7	AMS, AMC, EX, CAZ, LE, E, CIP	0.5
4.		B2/Cl/1	2	CAZ, E	0.14
5.		B2/Cl/3	1	E	0.07
6.		B1/Fe/1	2	CAZ, E	0.14
7.		B3/Fe/2	3	CAZ, E, TE	0.21
1.	Household backyard poultry	C8/Cl/1	1	E	0.07
2.		C11/Cl/1	1	E	0.07

252

253 **3.5 Characterization of Antimicrobial Resistance Genes: Extended Spectrum** 254 **Beta Lactamase genes, tetracycline gene and sulphonamides gene**

255 The carriage status of *Salmonella* strains for 04 antimicrobial Beta Lactamase
256 resistance genes, 01 tetracycline (*tetA*) and 01 sulphonamides (*Sul1*) were screened by
257 PCR.

258 All the *Salmonella* isolates were screened for ESBL genes *blaTEM*, *blaSHV*,
259 *blaOXA* and *blaCTX-M*, with an amplicon size of 445bp, 237bp, 813bp and 593bp
260 respectively. Among the 15 isolates, all (100 %) isolates harboured the *blaTEM* gene (Fig.
261 2), whereas none of them were found to be carrying *blaSHV*, *blaOXA*, and *blaCTX-M* genes.
262 However, the *tetA* gene encoding resistance to tetracycline was also detected in all the
263 *Salmonella* isolates (Fig 3). With regard to sulphonamides none of them were found to carry
264 *Sul1* gene encoding resistance to sulphonamides


265 The results of the *blaTEM* gene are in agreement with reports of Bae et al. (2013)
266 who reported 90.91% of isolates harbouring the *blaTEM* gene. Similarly, Eguale et al. (2017)
267 reported the *blaTEM* gene in 79.1%, Several workers reported a lower prevalence of the
268 *blaTEM* gene in *Salmonella* isolates isolated from poultry (Thakur et al. 2013; Elumalai et al.
269 2014; Herrera-Sánchez et al. 2020). In our study, none of the *Salmonella* isolates were
270 found to carry *blaSHV*, *blaOXA*, and *blaCTX-M* genes. The study subsequently did not
271 identify MDR *Salmonella* spp. with sequences of the ESBL broad-spectrum β -lactamases
272 (*blaSHV*, *blaOXA*, and *blaCTX-M*) in these isolates. The *blaTEM* gene in the *Salmonella*
273 strains could result in the widespread prevalence of *blaTEM*positive. *Salmonella* by transfer
274 of gene to *blaTEM* negative strains through plasmid conjugation, transformation, and
275 transduction (Lai et al. 2023).

276 Our results for carriage of the *tetA* gene are similar with earlier reports of Adesijiet al.
277 (2018), Waghmare et al.(2018), and Soufi et al. (2012) who reported *Salmonella* isolates
278 carrying the *tetA* gene in 100.00, 100.00 and 71.00 percent, respectively. Amplified
279 expression of intrinsic resistance mechanisms for tetracycline resistance in microbes is due
280 to the acquisition of mobile genetic elements, ribosomal binding place mutations, and
281 chromosomal mutations (Pavelquesi et al. 2021).


282 The *sul1* gene is a sulphonamide-resistant dihydropteroate synthase of Gram-
283 negative microorganisms (Alcock et al. 2023). The *Sul1* gene is found related to other
284 resistance genes in class 1 integrons (Sköld 2021). A high percentage of *Sul1* gene percent in
285 *Salmonella* isolates was reported by Adesijiet al. (2014) who reported 100 percent presence
286 in *Salmonella* spp. isolated from humans and poultry. Machado et al. (2013) and Ma et al.

287 (2017) reported that no one of the isolates was carrying sulfonamide-resistant *Sul* genes.
288 The absence of *Sul1* in all isolates, indicates bacteria are not subjected to selective pressure
289 by sulfonamides, a valuable means for the conservation and further addition of resistance to
290 other antimicrobial elements.

291 Several factors contribute to the progress of antimicrobial resistance in bacteria,
292 including alterations in bacterial cell permeability, enzymatic modification of drugs, and the
293 exclusion of antimicrobials by membrane-bound efflux pumps (Chen et al. 2004). Antibiotic
294 resistance often stems from genetic changes encoded by chromosomal and plasmid genes,
295 with these genes primarily located on integrons, plasmids, and transposons, which are
296 mobile genetic elements (Thong and Modarressi 2011).



297
298 **Fig. 2 Agarose gel showing PCR product of *blaTEM* gene of *Salmonella* spp.**



299
300 **Fig. 3 Agarose gel showing PCR product of *tetA* gene of *Salmonella* spp.**

301
302 **4. CONCLUSION**

303
304 The findings of the existing study showed that an improved native poultry breed
305 supply chain is a potential source of virulent *Salmonella* Enterica uncovering humans to

306 potential zoonotic infections via meat, egg, or direct exposure. But lesser MARs along with
307 the absence of genotypic resistance for ESBL broad-spectrum β -lactamases in *Salmonella*
308 isolates was found under study. Minor use of β -lactam and sulphonamide antibiotics in the
309 production chain of improved breed poultry production chain might have resulted in the non-
310 development of resistance. Overall, results suggested there were no major MDR *Salmonella*
311 spp. circulating in the improved native poultry breed production chain of Maharashtra, India.
312 Judicial use of antimicrobials and biosecurity measures shall be implemented in small-scale
313 farms, backyards, and hatcheries principally in countryside areas of Maharashtra, India.

314

315 **ACKNOWLEDGEMENTS**

316

317 Department of Science and Technology, New Delhi for providing financial support under
318 State Science and Technology Programme, Project entitled “Dissemination of poultry and
319 goat meat processing technology and establishment of meat processing demonstration unit
320 for livelihood upliftment of farmers and butcher communities of Marathwada Region,
321 Maharashtra” [DST No. DST/SSTP/2018-19/30 (G) & (C)] and ‘ICAR -NAHEP’ program for
322 providing all the funds for my research work through selection for the Mapping of P.G/Ph.D.
323 research work under ICAR-NAHEP-CAAST-MAFSU project entitled “Centre of Excellence
324 for Advanced Research on Animal Food Safety”.

325

326 **AUTHORS’ CONTRIBUTIONS**

327

328 All authors contributed to the study conception and design. Designed study:
329 [RupeshWaghmare, RadhaRasve]; Experiment: [RadhaRasve, NandiniKuntawar, M.F.M.F.
330 Siddiqui]; Supervision: [RupeshWaghmare]; Data Analysis: [KakasahebKhose,
331 NandiniKuntawar]; Statistical analysis: [RadhaRasve, NandiniKuntawar]; Manuscript:
332 [RupeshWaghmare, RadhaRasve]; Manuscript review: [KakasahebKhose, M.F.M.F.
333 Siddiqui]. All authors read and approved the final manuscript.

334

335 **Disclaimer (Artificial intelligence)**

336 Option 1:

337 Author(s) hereby declare that NO generative AI technologies such as Large Language
338 Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the
339 writing or editing of this manuscript.

340 Option 2:

341 Author(s) hereby declare that generative AI technologies such as Large Language Models,
342 etc. have been used during the writing or editing of manuscripts. This explanation will

343 include the name, version, model, and source of the generative AI technology and as well as
344 all input prompts provided to the generative AI technology

345 Details of the AI usage are given below:

346 1.

347 2.

348 3.

349

350

351 REFERENCES

352

353 Abunna, F., Bedasa, M., Beyene, T., Ayana, D., Mamo, B., &Duguma, R. (2016).
354 Salmonella: isolation and antimicrobial susceptibility tests on isolates collected
355 from poultry farms in and around Modjo, Central Oromia, and Ethiopia.

356

357 Adesiji, Y. O., Deekshit, V. K., &Karunasagar, I. (2014). Antimicrobial-resistant genes
358 associated with Salmonella spp. isolated from human, poultry, and seafood
359 sources. *Food science & nutrition*, 2(4), 436-442.

360

361 Adesiji, Y. O., Shivakumaraswamy, S. K., Deekshit, V. K., Kallappa, G. S., &Karunasagar, I.
362 (2018). Molecular characterization of antimicrobial multi-drug resistance in non-
363 typhoidal Salmonellae from chicken and clam in Mangalore, India. *Journal of*
364 *Biomedical Research*, 32(3), 237.

365

366 Akond, M. A., Shirin, M., Alam, S., Hassan, S. M. R., Rahman, M. M., &Hoq, M. (2012).
367 Frequency of drugresistant Salmonella spp. isolated from poultry samples in
368 Bangladesh. *Stamford Journal of Microbiology*, 2(1), 15-19.

369

370 Alali, W. Q., Thakur, S., Berghaus, R. D., Martin, M. P., &Gebreyes, W. A. (2010).
371 Prevalence and distribution of Salmonella in organic and conventional broiler
372 poultry farms. *Foodborne pathogens and disease*, 7(11), 1363-1371.

373

374 Alcock, B. P., Huynh, W., Chalil, R., Smith, K. W., Raphenya, A. R., Wlodarski, M. A., et al.
375 (2023). CARD 2023: expanded curation, support for machine learning, and
376 resistome prediction at the Comprehensive Antibiotic Resistance
377 Database. *Nucleic acids research*, 51(D1), D690-D699.

377

378 Andino, A., &Hanning, I. (2015). Salmonella enterica: survival, colonization, and virulence
379 differences among serovars. *The Scientific World Journal*, 2015(1), 520179.

380

381 Anejo-Okopi, J. A., Isa, S. E., Audu, O., Fagbamila, I. O., Iornenge, J. C., & Smith, I. S.
382 (2016). Isolation and polymerase chain reaction detection of virulence invA gene in
383 Salmonella spp. from poultry farms in Jos, Nigeria. *Journal of Medicine in the*
384 *Tropics*, 18(2), 98-102.

385

386 Bacteriological Analytical Manual (2007) BAM: *Salmonella*, Chapter 5
387 *Salmonella*.[http://www.fda.gov/downloads/Food/FoodScienceResearch/UCM](http://www.fda.gov/downloads/Food/FoodScienceResearch/UCM309839)
388 309839.

389
390 Bae, D. H., Dessie, H. K., Baek, H. J., Kim, S. G., Lee, H. S., & Lee, Y. J. (2013). Prevalence
391 and characteristics of Salmonella spp. isolated from poultry slaughterhouses in
392 Korea. *Journal of Veterinary Medical Science*, 75(9), 1193-1200.
393
394 Bhowmick, S., Pal, S., Sunder, J., Sujatha, T., De, A. K., Mondal, T., et al. (2023). Exploring
395 broilers and native fowls of Andaman and Nicobar Islands as a source of β -
396 lactamase-producing Enterobacteriaceae even with limited anthropogenic activities
397 and docking-based identification of catalytic domains in novel β -lactamase
398 variants. *Frontiers in Veterinary Science*, 9, 1075133.
399
400 Bhuvananeswa, M., Shanmughap, S., &Natarajase, K. (2015). Prevalence of multidrug-
401 resistant (MDR) Salmonella enteritidis in poultry and backyard chicken from
402 Tiruchirappalli, India. *Microbiology Journal*, 5(2), 28-35.
403
404 CDC: The Centers for Disease Control and Prevention. Antibiotic Resistance Threats in the
405 United States; CDC: Atlanta, GA, USA, 2013.
406
407 Chaiban, C., Robinson, T. P., Fèvre, E. M., Ogola, J., Akoko, J., Gilbert, M., &Vanwambeke,
408 S. O. (2020). Early intensification of backyard poultry systems in the tropics: a case
409 study. *Animal*, 14(11), 2387-2396.
410
411 Chen, S., Zhao, S., White, D. G., Schroeder, C. M., Lu, R., Yang, H., et al. (2004).
412 Characterization of multiple-antimicrobial-resistant Salmonella serovars isolated
413 from retail meats. *Applied and Environmental Microbiology*, 70(1), 1-7.
414
415 Dashti, A. A., Jadaon, M. M., Abdulsamad, A. M., &Dashti, H. M. (2009). Heat treatment of
416 bacteria: a simple method of DNA extraction for molecular techniques. *Kuwait Med*
417 *J*, 41(2), 117-122.
418
419 Dinh, A., Duran, C., Singh, S., Tesmoingt, C., Bouabdallah, L., Hamon, A., et al. (2023).
420 Real-life temocillin use in Greater Paris area, effectiveness and risk factors for
421 failure in infections caused by ESBL-producing Enterobacterales: a multicentre
422 retrospective study. *JAC-Antimicrobial Resistance*, 5(1), dlac132.
423
424 Eguale, T., Birungi, J., Asrat, D., Njahira, M. N., Njuguna, J., Gebreyes, W. A., et al. (2017).
425 Genetic markers associated with resistance to beta-lactam and quinolone
426 antimicrobials in non-typhoidal Salmonella isolates from humans and animals in
427 central Ethiopia. *Antimicrobial Resistance & Infection Control*, 6, 1-10.
428
429 Eid, S., Ibrahim, H. M., Shaltot, S. H., & El Oksh, A. S. (2023). An Overview of the Current
430 Situation of Salmonellosis in Pigeons, Household Chickens, and Commercial
431 Broilers with a Special Reference to a Customized Vaccine Developing
432 Trial. *Journal of Advanced Veterinary Research*, 13(3), 322-332.
433
434 Elmonir, W., Abdeltawab, D., El-Sharkawy, H., & Zahran, R. N. (2023). Serotypes Diversity,
435 Virulence, and Antimicrobial Resistance of Non-Typhoidal Salmonella Isolates in
436 Commercial and Backyard Egg Production Systems in Egypt. *Pak. J. Zool.*, 1-9.
437
438 Elumalai, S., Muthu, G., Selvam, R. E. M., & Ramesh, S. (2014). Detection of TEM-, SHV-
439 and CTX-M-type β -lactamase production among clinical isolates of Salmonella
440 species. *Journal of medical microbiology*, 63(7), 962-967.
441

442 Fahmy, S. S., Dahshan, A. H., & El Sawah, A. (2023). Prevalence of Antimicrobials and
443 Virulence-Related Genes in Salmonellae Detected in Local Hatcheries in Northern
444 Upper Egypt. *Journal of Veterinary Medical Research*, 30(2), 71-77.
445

446 Fang, H., Ataker, F., Hedin, G., & Dornbusch, K. (2008). Molecular epidemiology of
447 extended-spectrum β -lactamases among *Escherichia coli* isolates collected in a
448 Swedish hospital and its associated health care facilities from 2001 to
449 2006. *Journal of Clinical Microbiology*, 46(2), 707-712.
450

451 Hedman, H. D., Vasco, K. A., & Zhang, L. (2020). A review of antimicrobial resistance in
452 poultry farming within low-resource settings. *Animals*, 10(8), 1264.
453

454 Heredia, N., & García, S. (2018). Animals as sources of food-borne pathogens: A
455 review. *Animal nutrition*, 4(3), 250-255.
456

457 Herrera-Sánchez, M. P., Rodríguez-Hernández, R., & Rondón-Barragán, I. S. (2020).
458 Molecular characterization of antimicrobial resistance and enterobacterial repetitive
459 intergenic consensus-PCR as a molecular typing tool for *Salmonella* spp. isolated
460 from poultry and humans. *Veterinary World*, 13(9), 1771.
461

462 Jajere, S. M., Hassan, L., Aziz, S. A., Zakaria, Z., Abu, J., Nordin, F., & Faiz, N. M. (2019).
463 *Salmonella* in native “village” chickens (*Gallus domesticus*): Prevalence and risk
464 factors from farms in South-Central Peninsular Malaysia. *Poultry science*, 98(11),
465 5961-5970.
466

467 Kariuki S., Gordon MA., Feasey N., Parry CM.. (2015). Antimicrobial resistance and
468 management of invasive *Salmonella* disease. *Vaccine*, 19;33Suppl 3(0 3)
469

470 Koro, A., Elezaj, I., Hadžiabdić, S., Alić, A., & Rešidbegović, E. (2022). Occurrence of
471 *Salmonella* spp. in backyard poultry in Bosnia and Herzegovina. *Iranian Journal of*
472 *Veterinary Research*, 23(1), 1.
473

474 Kowalska-Krochmal B., and Dudek-Wicher R.,(2021) The Minimum Inhibitory Concentration
475 of Antibiotics: Methods, Interpretation, Clinical Relevance. *Pathogens*, 10(2):165
476

477 Krumperman, P. H. (1983). Multiple antibiotic resistance indexing of *Escherichia coli* to
478 identify high-risk sources of fecal contamination of foods. *Applied and*
479 *environmental microbiology*, 46(1), 165-170.
480

481 Lai, J., Mu, H., Zhou, B., He, J., Cheng, X., Gan, Y., et al. (2023). Bla TEM-positive
482 *Salmonella enterica* serovars Agona and Derby are prevalent among food-
483 producing animals in Chongqing, China. *Frontiers in Microbiology*, 14, 1011719.
484

485 Ma, M., Wang, H., Yu, Y., Zhang, D., & Liu, S. (2007). Detection of antimicrobial resistance
486 genes of pathogenic *Salmonella* from swine with DNA microarray. *Journal of*
487 *Veterinary Diagnostic Investigation*, 19(2), 161-167.
488

489 Ma, S., Lei, C., Kong, L., Jiang, W., Liu, B., Men, S., ... & Wang, H. (2017). Prevalence,
490 antimicrobial resistance, and relatedness of *Salmonella* isolated from chickens and
491 pigs on farms, abattoirs, and markets in Sichuan Province, China. *Foodborne*
492 *pathogens and disease*, 14(11), 667-677.
493

- 494 Machado, E., Coque, T. M., Cantón, R., Sousa, J. C., & Peixe, L. (2013). Commensal
495 Enterobacteriaceae as reservoirs of extended-spectrum beta-lactamases,
496 integrons, and sul genes in Portugal. *Frontiers in Microbiology*, 4, 80.
497
- 498 Muhammad, M., Muhammad, L. U., Ambali, A. G., Mani, A. U., Azard, S., & Barco, L. (2010).
499 Prevalence of Salmonella associated with chick mortality at hatching and their
500 susceptibility to antimicrobial agents. *Veterinary Microbiology*, 140(1-2), 131-135.
501
- 502 Mulika, L., & Yuwapanichsampan, S. (2011). Prevalence of Salmonella spp. and Its
503 Resistance to Antimicrobial Drugs in Poultry Hatchery. *KKU Veterinary
504 Journal*, 18(1), 12-28.
505
- 506 Nair, A., Balasaravanan, T., Malik, S. V. S., Mohan, V., Kumar, M., Vergis, J., & Rawool, D.
507 B. (2015). Isolation and identification of Salmonella from diarrheagenic infants and
508 young animals, sewage waste and fresh vegetables. *Veterinary world*, 8(5), 669.
509
- 510 Ng, L. K., Martin, I., Alfa, M., & Mulvey, M. (2001). Multiplex PCR for the detection of
511 tetracycline resistant genes. *Molecular and cellular probes*, 15(4), 209-215.
512
- 513 Orabi, A., Armanious, W., Radwan, I. A., Girh, Z. M., Hammad, E., Diab, M. S., & Elbestawy,
514 A. R. (2022). Genetic correlation of virulent Salmonella serovars (Extended
515 Spectrum β -Lactamases) isolated from broiler chickens and human: a public health
516 concern. *Pathogens*, 11(10), 1196.
517
- 518 Orji, M. U., Onuigbo, H. C., & Mbata, T. I. (2005). Isolation of Salmonella from poultry
519 droppings and other environmental sources in Awka, Nigeria. *International Journal
520 of Infectious Diseases*, 9(2), 86-89.
521
- 522 Pavelquesi, S. L. S., de Oliveira Ferreira, A. C. A., Rodrigues, A. R. M., de Souza Silva, C.
523 M., Orsi, D. C., & da Silva, I. C. R. (2021). Presence of tetracycline and
524 sulfonamide resistance genes in Salmonella spp.: literature
525 review. *Antibiotics*, 10(11), 1314.
526
- 527 Pavon, R. D. N., & Rivera, W. L. (2021). Molecular serotyping by phylogenetic analyses of a
528 1498bp segment of the invA gene of Salmonella. *ASM Sci. J*, 14, 1-14.
- 529 Rothrock Jr, M. J., Hiett, K. L., Guard, J. Y., & Jackson, C. R. (2016). Antibiotic resistance
530 patterns of major zoonotic pathogens from all-natural, antibiotic-free,
531 pasture-raised broiler flocks in the Southeastern United States. *Journal of
532 Environmental Quality*, 45(2), 593-603.
533
- 534 Salehi, T. Z., Mahzounieh, M., & Saeedzadeh, A. (2005). Detection of invA gene in isolated
535 Salmonella from broilers by PCR method. *Int. J. Poult. Sci*, 4(8), 557-559.
536
- 537 Samanta, I., Joardar, S. N., Das, P. K., Sar, T. K., Bandyopadhyay, S., Dutta, T. K., &
538 Sarkar, U. (2014). Prevalence and antibiotic resistance profiles of Salmonella
539 serotypes isolated from backyard poultry flocks in West Bengal, India. *Journal of
540 Applied Poultry Research*, 23(3), 536-545.
541
- 542 Siemon, C. E., Bahnson, P. B., & Gebreyes, W. A. (2007). Comparative investigation of
543 prevalence and antimicrobial resistance of Salmonella between pasture and
544 conventionally reared poultry. *Avian diseases*, 51(1), 112-117.
545

546 Singh, M., Mollier, R. T., Paton, R. N., Pongener, N., Yadav, R., Singh, V., et al. (2022).
547 Backyard poultry farming with improved germplasm: Sustainable food production
548 and nutritional security in fragile ecosystem. *Frontiers in Sustainable Food*
549 *Systems*, 6, 962268.
550
551 Sivaramalingam, T., Pearl, D. L., McEwen, S. A., Ojkic, D., & Guerin, M. T. (2013). A
552 temporal study of Salmonella serovars from fluff samples from poultry breeder
553 hatcheries in Ontario between 1998 and 2008. *Canadian Journal of Veterinary*
554 *Research*, 77(1), 12-23.
555
556 Sköld, O. (2000). Sulfonamide resistance: mechanisms and trends. *Drug resistance*
557 *updates*, 3(3), 155-160.
558
559 Sohail, M. N., Rathnamma, D., Priya, S. C., Isloor, S., Naryanaswamy, H. D., Ruban, S. W.,
560 &Veeregowda, B. M. (2021). Salmonella from farm to table: isolation,
561 characterization, and antimicrobial resistance of Salmonella from commercial
562 broiler supply chain and its environment. *BioMed Research International*, 2021(1),
563 3987111.
564
565 Soufi, L., Sáenz, Y., de Toro, M., Salah Abbassi, M., Rojo-Bezares, B., Vinué, L., et al.
566 (2012). Phenotypic and genotypic characterization of Salmonella enterica
567 recovered from poultry meat in Tunisia and identification of new genetic
568 traits. *Vector-Borne and zoonotic diseases*, 12(1), 10-16.
569
570 Talukder, M., Islam, M. S., Levy, S., Sobur, M. A., Ballah, F. M., Najibullah, M., et al. (2021).
571 Detection of multidrug resistant Salmonella spp. from healthy and diseased broilers
572 having potential public health significance. *J. Adv. Biotechnol. Exp. Ther*, 4(2), 248-
573 255.
574
575 Thakur, S., Brake, J., Keelara, S., Zou, M., &Susick, E. (2013). Farm and environmental
576 distribution of Campylobacter and Salmonella in broiler flocks. *Research in*
577 *Veterinary Science*, 94(1), 33-42.
578
579 Thong, K. L., &Modarressi, S. (2011). Antimicrobial resistant genes associated with
580 Salmonella from retail meats and street foods. *Food Research International*, 44(9),
581 2641-2646.
582
583 Trampel, D. W., Holder, T. G., & Gast, R. K. (2014). Integrated farm management to prevent
584 Salmonella Enteritidis contamination of eggs. *Journal of Applied Poultry*
585 *Research*, 23(2), 353-365.
586
587 Van Boeckel, T. P., Pires, J., Silvester, R., Zhao, C., Song, J., Criscuolo, N. G., et al. (2019).
588 Global trends in antimicrobial resistance in animals in low-and middle-income
589 countries. *Science*, 365(6459), eaaw1944.
590
591 Van Loo, E. J., Caputo, V., Nayga Jr, R. M., Meullenet, J. F., &Ricke, S. C. (2011).
592 Consumers' willingness to pay for organic chicken breast: Evidence from choice
593 experiment. *Food quality and preference*, 22(7), 603-613.
594
595 Varijakshapanicker, P., Mckune, S., Miller, L., Hendrickx, S., Balehegn, M., Dahl, G. E.,
596 &Adesogan, A. T. (2019). Sustainable livestock systems to improve human health,
597 nutrition, and economic status. *Animal Frontiers*, 9(4), 39-50.
598

- 599 Waghamare, R. N., Paturkar, A. M., Vaidya, V. M., Zende, R. J., Dubal, Z. N., Dwivedi, A., &
600 Gaikwad, R. V. (2018). Phenotypic and genotypic drug resistance profile of
601 Salmonella serovars isolated from poultry farm and processing units located in and
602 around Mumbai city, India. *Veterinary world*, 11(12), 1682.
603
- 604 Waghamare, R. N., Popalghat, H. K., Londhe, S. V., Deshmukh, V. V., & Khobe, V. V.
605 (2021). An Online survey of consumers of Maharashtra concerning the expected
606 change in the meat and meat product business.
607
- 608 WHO. Prioritization of Pathogens to Guide Discovery, Research and Development of New
609 Antibiotics for Drug-Resistant Bacterial Infections, Including Tuberculosis. 2017.
610 Available online:
611 [www.who.int/medicines/areas/rational_use/PPLreport_2017_09_19](http://www.who.int/medicines/areas/rational_use/PPLreport_2017_09_19.pdf?ua=1)
612 (accessed on 14 December 2022).
613
- 614 Withenshaw, S. M., Cawthraw, S., Gosling, B., Newton, K., Oastler, C. E., Smith, R. P., &
615 Davies, R. H. (2021). Risk factor analysis for Salmonella contamination of broiler
616 chicken (*Gallus gallus*) hatcheries in Great Britain. *Preventive Veterinary*
617 *Medicine*, 196, 105492.
618
- 619 Xu, X., Biswas, S., Gu, G., Elbediwi, M., Li, Y., & Yue, M. (2020). Characterization of
620 multidrug resistance patterns of emerging Salmonella entericaserovar Rissen along
621 the food chain in China. *Antibiotics*, 9(10), 660.
622
- 623 Zhao, X., Ju, Z., Wang, G., Yang, J., Wang, F., Tang, H., et al . (2021). Prevalence and
624 antimicrobial resistance of Salmonella isolated from dead-in-shell chicken embryos
625 in Shandong, China. *Frontiers in Veterinary Science*, 8, 581946.
626
627
628
629