

Original Research Article

Effect of inorganic, organic and biofertilizers on growth and yield of garlic (*Allium sativum* L.) cv. Yamuna Safed - 4 (G-323)

ABSTRACT

A study on the research entitled “Effect of inorganic, organic and biofertilizers on growth and yield of garlic (*Allium sativum* L.) cv. Yamuna Safed - 4 (G-323)” was conducted to evaluate the effect of different fertilizers on growth and yield of garlic at the Horticultural Research Farm of Doon (P.G.) College of Agriculture Science and Technology, Dehradun, Uttarakhand during 2023-2024. The experiment was laid out in randomized block design. The growth and yield parameters of garlic were significantly influenced by different treatments of inorganic, organic and biofertilizers. The maximum growth in terms of plant height (72.46 cm), number of leaves per plant (6.73), length of leaves (43.80 cm), stem girth (10.89 mm), chlorophyll content of leaves (79.88 mg/g) and yield per hectare (108.00 q/ha) were observed in the treatment T₃ [NPK (100%)]. This study highlights the dual importance of achieving high garlic yields with NPK while advocating for the integration of organic and biofertilizers to enhance soil health, reduce dependency on synthetic inputs, and ensure long-term agricultural sustainability.

Keywords: Inorganic, Organic, Biofertilizers, Growth, Yield, Garlic

1. INTRODUCTION

Garlic (*Allium sativum* L.) is one of the most widely grown *Allium* species after onion belonging to the family Alliaceae (Vavilov, 1926). It has been an important vegetable crop in India since ancient times (Bhosle, 2024). It originated in Central Asia and is used as a vegetable as well as for medicinal purposes (Petrovska and Cekovska, 2010). It is the most nutritious crop among the bulbous vegetables. It is also used as a spice and condiment with high nutritional and medicinal value (Sunanta *et al.*, 2023). It contains high amounts of carbohydrates (29%), proteins (6.3%), minerals (0.3%), essential oils (0.1-0.4%) and appreciable quantities of fat, vitamin C and sulphur (Memane *et al.*, 2008).

Garlic has antiviral, antibacterial, antifungal and anti-protozoal properties (Batiha *et al.*, 2020). It is beneficial to the cardiovascular and immune systems due to its antioxidant and anti-cancerous properties. It is described as a heat stimulant, carminative & anti-rheumatic and its oil is a powerful antiseptic mainly due to the presence of allicin; an important organosulfur compound (Shang *et al.*, 2019). It is used as a vermifuge for expelling roundworms and has been recommended for the cure of several ailments viz., wounds, ulcers, pneumonia, bronchitis and gastro-intestinal disorders (Hazra *et al.*, 2011). The applications of synthetic chemical fertilizers alone can have deleterious effects on the soil health, water resources and ecosystem (Phad, 2023). Plant nutrition is one of the important factors in increasing the growth and yield of crop plants by providing essential nutrients that drive key physiological processes, including photosynthesis, root development, flowering, and stress tolerance, which are crucial for maximizing productivity and quality (White and Brown, 2010, Noulas *et al.*, 2023). It is a well-known fact that long-time inorganic fertilizer application leads to harmful effects on soil fertility, resulting in poor yield and quality of crops (Glendining *et al.*, 1996; Brar *et al.*, 2015, Ma *et al.*, 2023). Therefore, the integration of inorganic fertilizers, organic manures and biofertilizers can maintain good soil health, productivity and fertility status of soil (Priyanshu *et al.*, 2020; Hanif *et al.*, 2022).

Keeping in view the above literature, the present investigation was undertaken to find out the effect of individual application of inorganic [Nitrogen, Phosphorus and Potassium (NPK)], organic [Farm Yard Manure (FYM), Vermicompost, Wood ash] and bio-fertilizers [Azotobacter, Arbuscular Mycorrhiza] for the growth and yield of garlic in the Dehradun region of Uttarakhand.

2. MATERIALS AND METHODS

The experiment was conducted at the Horticultural Research Farm, Department of Horticulture, Doon (P.G) College of Agriculture Science and Technology, Selaqui, Dehradun (UK), India from 2023-2024. The experimental field is located at an altitude of 515 m above mean sea level between 30.35°N latitudes and 77.84°W longitudes. The average annual rainfall is 2073.3 mm; average temperature (15.5°C) and average humidity (75.21%) are recorded during the garlic growth period. The soil of the experimental field was sandy loam with pH ranging from 6.5 to 7.5. The soil was very low in available nitrogen (0.02%), medium in available phosphorus (48.9%) and high in available potassium (2.9%). Electrical conductivity of soil

ranged between 0.30-0.73 dS/m indicating the nature of soil is normal. Moreover, the soil contains organic carbon ranging from 0.31% to 0.80%. Garlic cultivar ‘Yamuna Safed – 4 (G-323)’ was used as planting material for the present investigation. The experiment was laid out in a simple Randomized Block Design (RBD) with three replications and seven treatment plots. The treatment consisted of seven different treatments of inorganic, organic and biofertilizers viz., T₁ [Control], T₂ [FYM (100%)], T₃ [NPK (100%)], T₄ [*Azotobacter* (100%)], T₅ [Vermicompost (100%)], T₆ [Wood ash (100%)] and T₇ [Arbuscular Mycorrhiza (100%)]. Required quantity (considered as 100%) of Farm Yard Manure @ 20 t/ha, NPK @ 100:50:50 kg/ha, *Azotobacter* @ 8 kg/ha, vermicompost @ 10 t/ha, wood ash @ 5 t/ha and Arbuscular Mycorrhiza @ 6 kg/ha were applied directly to the soil before planting of garlic cloves in the respective plots as per scheduled treatments.

Data about growth components were collected from 10 plants in each replication of the 7 treatments. A random sampling was done from each plot to determine plant height, number of leaves per plant, length of leaves, stem girth and chlorophyll content of leaves. Plant height and length of leaves were measured with the help of measuring tape and expressed in centimeters. Number of leaves was counted from randomly selected plants of different treatment. Stem girth was taken in the east-west and north-south directions with the help of digital vernier calipers (Corceptive Pvt. Ltd.) and averaged. Stem girth was expressed in mm. The chlorophyll content is analyzed by extraction of chlorophyll in a solvent followed by in-vitro measurements in a spectrophotometer (Parry *et al.*, 2014). Yield per hectare was calculated by weighing all bulbs from each treatment, replication-wise at the time of harvesting.

$$\text{Yield per hectare (kg)} = \frac{\text{Garlic weight (kg)}}{\text{Plot area (m}^2\text{)}} \times 10,000$$

The obtained data was subjected to statistical analysis using the F test according to the procedure of Gomez and Gomez (1984). The critical difference at 5% was calculated to compare the mean value of the determined criteria of different treatments.

3. RESULTS AND DISCUSSION

The effect of different inorganic (NPK), organic (FYM, Vermicompost, Wood ash) and biofertilizers (*Azotobacter*, Arbuscular Mycorrhiza) on growth and yield attributes has been presented in Table 1 and Figure 1. Data recorded for plant height revealed that the maximum

plant height (72.46 cm) was recorded in treatment T₃ [NPK (100%)] and was significantly different from treatment T₅ [Vermicompost (100%)] (69.66 cm) and T₂ [FYM (100%)] (63.20 cm). The minimum plant height (50.03 cm) was recorded in treatment T₁ [Control] which was significantly lower than treatment T₇ [Arbuscular Mycorrhiza (100%)] which is 53.40 cm. Kumar *et al.* (2019) reported that the plant height of garlic was significantly increased with the combined application of organic and inorganic fertilizers maintaining long-term soil fertility and sustaining a high level of productivity due to increased availability of nutrients to the plant initially through organic and inorganic fertilizers in the cropping season.

The maximum number of leaves per plant (6.73) was reported in treatment T₃ [NPK (100%)] followed by T₅ [Vermicompost (100%)] and a minimum number of leaves per plant (5.13) were found in Control which was statistically at par with treatments T₂ [FYM (100%)]. The maximum number of leaves observed in treatment T₃ [NPK (100%)] might be attributed to the **immediate availability of nutrients in inorganic fertilizers** (Eragegowda *et al.*, 2018). Integrated nutrition management may have promoted vegetative growth, which in turn increased the number of leaves and increased photosynthetic activity (Dhaker *et al.*, 2017). The maximum length of leaves per plant (43.80 cm) was reported in treatment T₃ [NPK (100%)] followed by T₅ [Vermicompost (100%)] and a minimum length of leaves per plant (32.36 cm) was found in the control which was significantly different with all other treatments. Results are also similar to the findings of Naidu *et al.* (2000) who reported that the application of nitrogen 150 kg N/ha caused a significant improvement in all growth parameters. **Dawar *et al.* (2005) reported that the highest leaf length (61.06 cm) in garlic was recorded with the application of 120 kg N per ha.**

The maximum stem girth (10.89 mm) was reported in treatment T₃ [NPK (100%)] followed by T₅ [Vermicompost (100%)] and minimum stem girth (6.55 mm) were found in the control which was significantly different among other treatments. The maximum chlorophyll content of leaves (79.88 mg/g) was reported in treatment T₃ [NPK (100%)] followed by T₅ [Vermicompost (100%)] and minimum chlorophyll content of leaves (67.95 mg/g) was found in control which was significantly different among all other treatments. This result conforms with the findings of Sharangi *et al.* (2003), Abd-El-Moneem *et al.* (2005) and Rohidas *et al.* (2010) on garlic. The maximum yield per hectare (108 q/ha) was reported in treatment T₃ [NPK (100%)] followed by T₅ [Vermicompost (100%)] and the minimum yield (63.00 q/ha) was found in the

control which was significantly different from other treatments. Similar results were reported by Bagali *et al.* (2012) on onion, Bhandari *et al.* (2012) and Damse *et al.* (2014) on garlic as they found significantly lower yield in control when compared to different inorganic and organic fertilizer applications.

Table 1. Effect of inorganic, organic and biofertilizers on plant height, number of leaves, length of leaves, stem girth, chlorophyll content and yield of garlic

Symbols	Treatments Details	Plant Height (cm)	Number of leaves per plant	Length of leaves (cm)	Stem Girth (mm)	Chlorophyll content of leaves (mg/g)	Yield (q/ha)
T ₁	Control	50.03	5.13	32.36	6.55	67.95	63.00
T ₂	FYM (100%)	63.20	6.26	40.06	8.83	73.22	88.00
T ₃	NPK (100%)	72.46	6.73	43.80	10.89	79.88	108.00
T ₄	<i>Azotobacter</i> (100%)	58.73	6.06	38.46	8.43	72.47	81.00
T ₅	Vermicompost (100%)	69.66	6.53	41.65	9.45	75.47	91.66
T ₆	Wood ash (100%)	56.20	5.73	37.25	7.82	71.58	74.00
T ₇	Arbuscular Mycorrhiza (100%)	53.40	5.40	35.40	7.28	70.36	68.00
C.D. @ 5 %		2.624	0.489	0.914	0.207	0.858	10.015
SE(m)		0.842	0.157	0.293	0.067	0.276	3.215
C.V.		2.410	4.547	1.322	1.362	0.654	6.794




Figure 1. Effect of inorganic, organic and biofertilizers on plant height, number of leaves, length of leaves, stem girth, chlorophyll content and yield of garlic



Plate 1. Pictorial view of garlic experimental field

4. CONCLUSION

The present study revealed that the treatment T₃ [NPK (100%)] significantly enhanced the growth and yield parameters of garlic, including plant height, number of leaves, leaf length, stem girth, chlorophyll content and total yield, compared to other treatments. Treatments T₅ [Vermicompost (100%)] and T₂ [FYM (100%)] also showed promising results, emphasizing the importance of organic fertilizers in sustainable farming.

The findings underscore the potential of integrating inorganic and organic fertilizers to improve garlic production in the Dehradun region while maintaining soil health. Future research could focus on optimizing the ratios of organic and inorganic inputs to achieve maximum yield and environmental benefits.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

REFERENCES

1. Abd-El-Moneem, K.M.H., Fawaz, S.B.M., Saeed, F.A. and El-Shehaby, A.I. 2005. Effect of clove size and certain micronutrients on fusarium basal rot of garlic. *Assiut Journal of Agricultural Sciences*, 36(4): 163-175.
2. Bagali, A.N., Patil, H.B., Chimmad, V.P., Patil, P.L. and Patil, R.V. 2012. Effect of inorganics and organics on growth and yield of onion (*Allium cepa* L.). *Karnataka Journal of Agricultural Science*, 25(1): 112-115.
3. Batiha, G.E.S., Beshbishy, A.M., Wasef, L.G., Elewa, Y.H.A., Sagan, A.A.A., Hack, M.E.A.E., Taha, A.E., Elhakim, Y.M.A., Devkota, H.P. 2020. Chemical Constituents and Pharmacological Activities of Garlic (*Allium sativum* L.): A Review. *Nutrients*, 12(872):1-21.
4. Bhandari, S.A., Patel, K.S. and Nehete, D.S. 2012. Effect of integrated nutrient management on growth, yield and quality of garlic (*Allium sativum* L.) cv. Gujrat Garlic-3. *Asian Journal of Horticulture*, 7(1): 48-51.

5. Bhosle, S.S. 2024. Review on garlic cultivation in India: Techniques, challenges, and future prospects. *Modern Phytomorphology*, 18: 133 -143.
6. Brar, B.S., Singh, J., Singh, G., and Kaur, G. 2015. Effects of long term application of inorganic and organic fertilizers on soil organic carbon and physical properties in maize–wheat rotation. *Agronomy*, 5(2): 220-238.
7. Damse, D.N., Bhalekar, M.N. and Pawar, P.K. 2014. Effect of integrated nutrient management on growth and yield of garlic. *The Bioscan*, 9(4): 1557-1560.
8. Dawar, M., Hussain, S.A. and Sajid, M. 2005. Effect of planting density and nitrogen levels on the growth and yield of garlic. *Sarhad Journal of Agriculture*, 21(4): 577-582.
9. Dhaker, B., Chhipa, B.G., Rathore, R.S. and Sharma, R.K. 2017. Effect of different sources of nutrients on growth and yield of onion (*Allium cepa* L.). *Journal of Plant Development Sciences*, 9: 1053-1056.
10. Eragegowda, M., Venkatesha, J., Hiremath, J.S., Mahantesh, P.S., Ravi, Y. and Manjesh, G.N. 2018. Effect of integrated nutrient management on growth and yield of garlic (*Allium sativum* L.) cv. AAS-2. *Journal of Pharmacognosy and Phytochemistry*, 7(3S): 387-389.
11. Glendining, M.J., Powlson, D.S., Poulton, P.R., Bradbury, N.J., Palazzo, D., and Li, X. 1996. The effects of long-term applications of inorganic nitrogen fertilizer on soil nitrogen in the Broadbalk Wheat Experiment. *The Journal of Agricultural Science*, 127(3): 347-363.
12. Gomez, K.A. and Gomez, A.A. 1984. Statistical Procedures for Agricultural Research. John Wiley and Sons, 680.
13. Hanif, M., Khan, R., Kiran, J., Haq, I.U., Ali, S., Niaz, Y., Ali, A., Arshad, M., Ansari, M.J., Alharbi, S.A. and Maqbool, A. 2022. Interactive effect of organic and mineral phosphorus on volatile compounds, morphology, and physiology of garlic (*Allium sativum* L.). *Journal of King Saud University-Science*, 34(8):102351.
14. Hazra, P., Chattopadhyay, A. and Karmakar, K. 2011. Modern technology in vegetable production. *New India Publishing Agency*, 288-289.
15. Kumar, S., Naruka, I.S., Shaktawat, R.P.S., Singh O.P. and Meena, K.C. 2019. Response of garlic (*Allium sativum* L.) to organic and inorganic fertilizers. *International Journal of Agriculture Sciences*, 11(7): 8209-8211.

16. Ma, G., Cheng, S., He, W., Dong, Y., Qi, S., Tu, N., & Tao, W. 2023. Effects of organic and inorganic fertilizers on soil nutrient conditions in rice fields with varying soil fertility. *Land*, 12(5): 1026.
17. Memane, P.G., Tomer, R.S., Kakade, D.K., Kulkarni, G.U. and Chovatia, R.S. 2008. Effect of clove weight and plant growth regulators on growth and yield of garlic cv. Gujrat Garlic-3. *The Asian Journal of Horticulture*, 3(1):82-86.
18. Naidu, A.K., Tiwari, J.P., Dwivedi, S.K. and Saxena, S.K. 2000. Effect of various levels of N, P and K on physiological growth determinants of productivity in garlic (*Allium sativum* L.). *Vegetable Science*, 27(2): 165-167.
19. Noulas, C., Torabian, S., and Qin, R. 2023. Crop Nutrient Requirements and Advanced Fertilizer Management Strategies. *Agronomy*, 13(8): 2017.
20. Parry, C., Blonquist, J.M. and Bugbee, B. 2014. In situ measurement of leaf chlorophyll concentration: Analysis of the optical/absolute relationship. *Plant, Cell and Environment*, 37: 2508-2520.
21. Petrovska, B.B. and Cekovska, S. 2010. Extracts from the history and medical properties of garlic. *Pharmacognosy Review*, 4: 106-110.
22. Phad N.M. 2023. Chemical fertilizers and their impact on soil health. *Journal of Emerging Technologies and Innovative Research*, 10(4): 428-432.
23. Priyanshu, A.B., Singh, M.K., Kumar, M., Kumar, V., Malik, S., Sahahi, U.P. and Lodhi, S.K. 2020. Effect of integrated nutrient management on yield and quality of garlic cv. Yamuna Safed-3. *Journal of Agrisearch*, 7(4): 251-254.
24. Rohidas, S.B., Bharadiya, P.S., Jature, S.D. and Ghate, K.B. 2010. Effect of micronutrient on growth and yield of garlic (*Allium sativum* L.) var. G-41. *Asian Journal of Horticulture*, 5(2): 517-519.
25. Shang, A., Cao, S.Y., Xu, X.Y., Gan, R.Y., Tang, G.Y., Corke, H., Mavumengwana, V. and Li, H.B., 2019. Bioactive compounds and biological functions of garlic (*Allium sativum* L.). *Foods*, 8(7): 246.
26. Sharangi, A.B., Pariari, A., Datta, S., and Chatterjee, R. 2003. Effect of boron and zinc on growth and yield of garlic in New Alluvial Zone of West Bengal. *Crop Research Hisar*, 25(1): 83-85.

27. Sunanta, P., Kontogiorgos, V., Pankasemsuk, T., Jantanasakulwong, K., Rachtanapun, P., Seesuriyachan, P., and Sommano, S.R. 2023. The nutritional value, bioactive availability and functional properties of garlic and its related products during processing. *Frontiers in Nutrition*, 10: 1142784.
28. Vavilov, N.I. 1926. Studies on the origin of cultivated plants. Trudy Byuro Prikladnoy Botaniki 16, 139 pp (in Russian).
29. White, P. J., and Brown, P. 2010. Plant nutrition for sustainable development and global health. *Annals of botany*, 105(7): 1073-1080.

UNDER PEER REVIEW