

Effects of *Lactobacillus sporogenes* supplementation on growth performance, survival, immune parameters, and disease resistance in *Labeo rohita* exposed to *Aeromonas hydrophila*

ABSTRACT

The present study was aimed to investigate the influence of *Lactobacillus sporogenes* on the growth characteristics, immune system, and susceptibility to *Aeromonas hydrophila* infection in *Labeo rohita*. Diet containing different levels of *L. sporogenes* at 10^4 cfu/g (F1), 10^6 cfu/g (F2), 10^8 cfu/g (F3) and without (F0, control) in triplicates for 90 days. Results indicated meaningful ($p < 0.05$) increased weight performance and decreased feed conversion ratio (FCR) at 10^8 cfu/g (F3) compared to control group (F0). Probiotic supplementation (F2, F3) led to a noticeable and statistically significant increase ($p < 0.05$) in the non-specific immunological measures, including superoxide anion production (SAP), total serum protein, and lysozyme function, compared to the group without supplementation (F0). Furthermore, the probiotic supplemented group exhibited a substantial ($p < 0.05$) decline in the cumulative percent of mortality (F1: 23.33 ± 0.5 ; F2: 26.67 ± 1.15 and F3: 30 ± 1.35) and higher relative percent survival (F1: 77.73 ± 1.51 ; F2: 79.69 ± 1.26 and F3: 83.69 ± 0.89) compared to the group without probiotic supplementation (F0: CPM-73.33%) when challenged with *A. hydrophila*. Therefore, overall present study recommended that higher level inclusion of *L. sporogenes* can significantly enhance the growth performance, immune function, and ability to resist diseases in *L. rohita* when exposed to *A. hydrophila* infection.

Keywords: *Labeo rohita*, Lactic Acid Bacteria, *L. sporogenes*, non-specific immune parameters, cfu/g

1. Introduction

Due to rising fish consumption and the possibility that capture fishery have been overfished to their capacity, aquaculture has been growing over the past few decades (FAO 2018). Indian major carps *Catla catla*, *Labeo rohita* and *Cirrhinus mrigala* are the most significant economic fish that are considered as highest market priced fish in India. They contribute over 67% of the total production of freshwater fish (Parthasarathy et al. 2011). Aquatic animal production at large-scale facilities is subject to stress, disease, and deterioration of the environment, which results in significant economic losses (Subasinghe et al. 2005). Development of infectious diseases including abdominal dropsy, exophthalmia, red spot, tail and fin rot, haemorrhagic septicemia and bacterial gill disease are frequently encountered in *L. rohita* (Padala et al. 2021). According to Harikrishnan and Balasundaram (2005), *A. hydrophila* generally affects freshwater fish species. As a conventional method of controlling fish disease in aquaculture systems, antibiotics have been administered to minimize the disease condition (Prasad et al. 2012). The extensive application of antibacterial drugs in the aquaculture system damages aquatic animals and its gastrointestinal systems resulting unfit for human utilization (Verschuere et al. 2000). This lead to the use of beneficial bacteria in aquaculture, which opened the doors to the application of probiotics in aquaculture, which have a positive influence through a variety of actions. Potential benefit of probiotics and its impact on the health and nutrition of some aquaculture organisms have been investigated (Merrifield et al. 2010; Mamun et al. 2019).

The term "probiotic" generally pertains to Gram-positive bacteria from the genus *Bacillus* spp. *Streptococcal* spp. and *Lactobacillus* spp. most commonly used and have been shown to improve the health of aquaculture animal without incurring any noticeable adverse effects (Irianto and Austin 2002). In fish culture, using a particular *Lactobacillus* produced superior outcomes (Barbosa et al., 2011). Probiotic refers to a group of bacteria, including the genus *Lactobacillus*, one of the important probiont that can fermenting a range of carbohydrates mostly into lactate and acetate (Gatesoupe 1999). The environmental function of *Lactobacillus* spp. as a helpful flora in the digestive tract includes, among other things, the synthesis of antimicrobial compounds that improve immune response and increase fish resistance to bacterial pathogens, increase nutrient availability, additionally utilizing certain carbohydrates that are not digestible (Dimitroglou et al. 2011).

The use of probiotics in feed formulation for sustainable aquaculture practices has garnered considerable attention in recent decades. Probiotics that increase fish body enzyme activity can enhance feed digestion and feed utilization, improve fish health and performance

(Mohapatra et al. 2012). Application of *Lactobacillus* spp. in several finfishes increased feed utilization, non-specific immunity, and resistant to virulent pathogen (Harikrishnan et al. 2010). Among the *Lactobacillus* spp. *L. sporogenes* considered one of the important probiont used in aquaculture industry and generated several beneficial impacts in finfishes (Sridhar and Joice 2012) and shellfishes (Gupta and Dhawan 2012; Seenivasan et al. 2012). There is, however, a lack of information on Indian Major Carp, *L. rohita*. Hence, a research investigation was undertaken to investigate the effects of *L. sporogenes* on growth efficiency and feed consumption, carcass composition, non-specific immunity, and its potential for enhancing resilience in the presence of *A. hydrophila*.

2. Materials and Method

2.1 Research diet and structure

The probiotic *L. sporogenes* was obtained from M/S Altech Biotechnology Pvt. Ltd, Bangaluru (Bangalore, Karnataka, India). The dry ingredients were mixed well to make a smooth dough after cooking with required temperature and water. The pellets were formed using a manual pelletizer with a diameter of 3 mm. Subsequently, the pellets were subjected to air drying in a hot air oven set at a temperature of 60°C until they reached a moisture content of approximately 10%. Each diet was packaged differently in thick polythene bags, marked and maintained at 4 °C in a freezer. Experimental diets were evaluated for the proximate composition in accordance with AOAC (1995). The nutrient composition of dietary components and its protein contribution exhibited in Table 1.

The *L. rohita* spawn was obtained from the Bhadra Reservoir Project fish seed farm in Shivamogga, Karnataka, India. It was then raised for 3 to 4 months in the college's research and instructional fish farm to reach fingerling stage. Twelve (12) outdoor cement tanks with a 1×1×1 m bare bottom each were used for the investigation. *L. rohita* fingerlings of uniform size, measuring approximately 1.17±0.45 cm in length and weighing around 0.76±0.15 g, were released in each tanks with 15 fingerlings. Each diet was replicated in triplicate. Three experimental diets, labelled as F1, F2, and F3, were formulated to include 104 cfu/g, 106 cfu/g, and 108 cfu/g of *L. sporogenes*, respectively. The control diet (F0) did not contain any *L. sporogenes*. To achieve the desired texture, the ingredients were mixed and hand-kneaded with an appropriate amount of water (1:0.8 ratio) until a dough was formed. After cooking the dough, the feed was rapidly down to ambient temperature (29±2 °C) putting it out in an aluminium tray. Following the methodology outlined by Nesara et al. (2018), the feed was supplemented with *L. sporogenes*, the probiotic bacterium.

Our viability testing showed that the probiotic bacterium remained viable after incorporation into the diet. The probiotic's survival bacterium, *Lactobacillus sporogenes*, in the fish feed was evaluated using the plate count method. Nutrient agar plates (Himedia) were prepared, and the probiotic-containing feed was poured onto the agar plates after being serially diluted. The plates were then incubated and estimated the number of live bacterium cells. The fish received feed for the duration of the experiment in a quantity equal to 5% of their body weight. Physico-chemical parameters were recorded during fortnightly sampling and accordingly all the parameters were maintained.

2.2 Proximate composition of formulated diets

While formulating test diets, all the feed items underwent proximate composition analysis using accepted techniques (AOAC 1995, Table 2). Samples are heated for 30 minutes at 105 °C, allowing them to cool, and then weighing them at a steady weight, the moisture content was calculated. The FOSS Kjeltex system was used to assess crude protein, the Soxtech system (PELICAN) to determine fat content, and the Fibretex system to determine fiber content (PELICAN). The difference approach (Hastings, 1976) was used to calculate the amount of carbohydrates present in nitrogen free extract (NFE) with the standard formula.

2.3 Growth parameters

To assess the growth's effectiveness and feed utilization, fish samples were taken every 15 days. Each sampling involved the collection of at least 50% of the stocked fish, which were then each individually weighed and measured. After each sampling, the quantity of feed provided was modified in accordance with the recorded growth in fish weight during the assessment. Various growth parameters, such as weight gain, feed conversion ratio (FCR), survival rate, and specific growth rate (SGR), were calculated using standard formulae.

2.4 Sample collection and immunological parameters

2.4.1 Blood sampling

After the completion of a 90-day period, a total of 10 fingerlings from every treatment as well as the control group were selected for physiological investigations. To prepare for blood collection, the fish underwent a 24-hour fasting period during which they were not fed. Using a sterile 2 ml syringe, blood was collected by caudal vein. Subsequently, the obtained blood were carefully transferred into dry EDTA (for NBT assay) and clean 2 ml centrifuge tube. Subsequently, the collected blood samples were centrifuged for ten minutes at 10,000 rpm to divide the serum. After centrifugation, the sera obtained from three fish from each treatment group were combined or pooled together. The pooled sera were then stored at a temperature of -40 °C for

next analysis, specifically for measuring parameters such as total serum protein and lysozyme activity.

2.4.2 Nitroblue tetrazolium assay (superoxide anion production)

The nitroblue tetrazolium assay was assessed according to Anderson and Siwicki (1995). Three fish were sampled, and blood was collected from their caudal veins and transferred to EDTA-coated tubes (EDTAK2). After centrifugation (5000 rpm for 10 minutes) the leukocyte-containing buffy coat was dispensed into a microtitre plate for cell adhesion. Following an hour of incubation, the plate was treated with 0.3% NBT solution for another hour. Upon removal of the NBT solution, the cells were treated with a fixing agent, washed, and air-dried. To dissolve the formazan, a solution consisting of 60 μ l of 2N KOH and 70 μ l of dimethyl sulphoxide (DMSO) was added to each well. Following that, the solution with its distinctive turquoise blue color was analyzed using a microplate reader, measuring the optical density at a wavelength of 620 nm.

2.4.3 Lysozyme activity

Lysozyme activity was assessed applying a customized turbidimetric technique reported by Parry et al. (1965). Two replicates of 50 μ l serum were dispensed into each well of a 96-well plate, along with 50 μ l of Phosphate Buffered Saline (PBS). The serum was subjected to sequential dilutions, gradually decreasing the amount of sample by 50 μ l in each consecutive well. Following that, 125 μ l of a suspension of *Micrococcus luteus* was evenly introduced into all the wells. The plate was placed in an incubator at ambient temperature, and the decrease in absorbance at 450 nm was assessed during a duration of 0 to 15 minutes using an ELISA reader. Lysozyme function was quantified based on the rate of reduction in absorbance per minute (0.001 min^{-1}), with one lysozyme unit defined as the enzyme activity that resulted in a decrease in absorbance at this rate.

2.4.4 Total plasma protein

Serum plasma protein was determined by the Lowry method with the GeNei™ protein analysis kit. In order to complete the technique, 50 μ l of distilled water and 50 μ l of serum were combined and added to each well of a 96-well plate. The Lowry reagent was then applied in 200 μ l well, and placed in an incubator for 10 minutes at ambient temperature. After the initial incubation, the plate was incubated for an additional 30 minutes with 200 μ l of the Folin-Ciocalteu reagent inserted into each well. Then, a spectrophotometer had used to measure the absorbance of each well at 650 nm.

2.5 Determination of LD₅₀

A virulent strain of *A. hydrophila* (ATCC 36562) was purchased from Pune, India's National Collection of Industrial Microorganisms (NCIM). The dried bacterial cells in the ampule were revived in trypton soya broth at 37 °C and cultured further to keep in agar slants at chill temperature (4°C). Ten fish fingerling (6.89±1.55g) per tank were used to determine the LD₅₀. Virulent pathogen, *A. hydrophila* were inoculated ranging from 10² to 10⁸ CFU/ml was injected intraperitoneally to each fish. Physiological saline 0.1 ml injections into the fish group considered as the control. Daily mortality rates were documented for 240 hours. Based on Reed and Muench's (1938) estimations, the degree of pathogenicity (LD₅₀) was determined.

2.6. Challenge study

Following a 90-day period of experimental feeding, a sample size of ten fish were selected from both the control and treatment groups in order to assess their disease resistance. Fishes were challenged (intramuscular) with 0.1 ml of *A. hydrophila* (2.3× 10⁷ CFU/ml. One negative control fish group received a dose of 0.1 ml of phosphate buffered saline. Percent survival, was calculated (Amend, 1981) and utilized as a measure of protection against diseases. Ethical permission were taken before sacrificing the fishes (No. FCM/ AEC/DFK 1501-2015-2018/01).

$$RPS = 100 \times (1 - \% \text{ mortality in } L. \text{ sporogenes} \text{ fed fishes} / \% \text{ mortality in control (F}_0\text{) fishes})$$

2.7 Data analysis

One-Way Analysis of Variance (ANOVA) was performed on the experimental and control groups. To assess the significance the Duncan test was employed. These statistical tests were utilized to evaluate and compare the results obtained from the different treatments.

3. Results and Discussion

3.1 Experimental ingredients and diets

In the current study, Table 1 outlines the proximate analysis of the dietary ingredients used. Higher and lower protein content had recorded in fish meal (67.45 %) and tapioca flour (2.34 %) respectively. Moreover, data showed that maximum level of fat, fibre and ash content was present in groundnut oil cake (16.92 %) wheat bran (10.31%) and in fish meal (25.21%). Crude protein contents in all the experiments (F₀, F₁, F₂ and F₃) were ranging from 27.09 % to 27.14% are shown in Table 2.

Table 1: Ingredients used to make various experimental diets and their approximate makeup

	Feed composition			
	Fish meal	Ground nut oil cake	Wheat bran	Tapioca flour
Moisture (%)	9.75±1.06	9.65±0.02	8.02±0.10	7.20±0.07
Dry matter (%)	90.25±1.07	90.35±0.01	91.98±0.08	92.80±0.08
Crude protein (%)	67.45±0.50	38.30±0.42	18.30±0.27	2.34±0.26
Ether extract (%)	13.55±0.07	16.92±0.56	8.54±0.23	4.60±0.14
Crude fibre (%)	1.55±0.05	3.92±0.13	10.31±0.02	2.31±0.05
Ash (%)	25.21± 0.12	7.32± 0.17	4.50± 0.70	5.27± 0.31
Nitrogen free extract (%)	17.51 ± 0.55	23.89 ± 0.76	50.33 ± 0.28	78.28 ± 0.44
Protein contribution (%)	13.59	7.72	5.18	0.68

Table 2: Proximate composition of formulated diet

	F ₀	F ₁	F ₂	F ₃
Moisture (%)	8.17± 0.03	7.76±0.01	7.58±0.03	7.36±0.02
Dry matter (%)	91.82±0.03	92.23±0.01	92.42±0.03	92.63±0.02
Crude protein (%)	27.09±0.31	27.14±0.17	27.13±0.01	27.14±0.01
Ether extract (%)	6.07±0.10	6.10±0.21	6.37±0.17	6.35±0.10
Crude fibre (%)	6.88±0.02	6.86±0.09	6.34±0.04	7.17±0.04
Ash (%)	10.75±0.35	11.00±0.70	11.25±0.35	11.25±0.35
Nitrogen free extract (%)	41.04±0.33	41.14±0.29	41.33±0.18	40.73±0.24

3.2 Growth metrics and feed utilization

The observed data showed dietary supplementation of probiotics has significantly enhanced fish biomass (weight gain) compared to the non-probiotic group (F₀). In 90 days of trail the growth found higher in F₃ followed by F₁, F₂ and F₀ (Table, 3). However, no notable disparity were noted in the treatment groups. The F₃ group exhibited the lowest but optimal feed conversion ratio (FCR), with the F₁, F₂, and F₀ groups following in that order. The results

show that group F1 and F3 had the highest specific growth rates, with values of 2.35 ± 0.11 and 2.37 ± 0.10 , respectively. Group F2 had an intermediate specific growth rate of 2.23 ± 0.02 , which did not differ from the F1 and F3 groups however differed significantly from the non-supplemented (F0). The control fishes (F0) had the lowest specific growth rate of 2.17 ± 0.11 . Furthermore, the inclusion of *L. sporogenes* in the diet generated significantly higher SGR in comparison to the control group.

Table 3: Comparison of the growth performance of *L. rohita* after 90 days of feeding with a control diet and a diet supplemented with probiotic *L. sporogenes* (mean of 3 replicates \pm standard error)

	F ₀	F ₁	F ₂	F ₃
Initial body weight	0.76 ± 0.02^a	0.76 ± 0.02^a	0.76 ± 0.02^a	0.76 ± 0.02^a
Final body weight	5.45 ± 0.50^b	6.35 ± 0.39^a	6.21 ± 0.35^a	6.52 ± 0.49^a
Weight gain	717.10 ± 72.05^b	835.52 ± 86.22^a	817.10 ± 86.09^a	857 ± 81.99^a
Feed conversion ratio	2.58 ± 0.32^a	2.17 ± 0.11^b	2.20 ± 0.13^b	2.16 ± 0.50^b
Specific growth rate	2.17 ± 0.11^b	2.35 ± 0.11^a	2.23 ± 0.02^{ab}	2.37 ± 0.10^a
Survival (%)	95.55^a	95.35^a	95.55^a	84.44^a

Evidence of previous works done by few authors the optimal dietary supplement of *L. sporogenes* range was decided at levels of 10^4 cfu/g, 10^6 cfu/g, and 10^8 cfu/g. In our study results found that there was no discernible variation in treatment groups (F1, F2) although F3 group fishes showed increase weight gain compared to the fishes fed with 10^4 cfu/g, 10^6 cfu/g of *L. sporogenes* and control group. Results suggested that group F3 with 10^8 cfu/g able to meet fish requirement. However, Eid and Mohamed (2008) opinioned that incorporation of commercial blend of probiotics Bifilac[®] and Pronifer[®] had a beneficial effect on weight gain, feed utilization on monosex tilapia (*Oreochromis niloticus*). Similarly, Kumar et al. (2006) validated that nutritional feeding of *B. subtilis* orally at 1.5×10^3 cfu/ g⁻¹ to *L. rohita* for 60 days improved growth with decrease in FCR. Also, Chaudhary and Qazi (2006) found that the incorporation of *B. circulans* into the feed of *L. rohita* fingerlings positively influenced their growth performance and feeding efficiency. In contrast, Noh et al. (1994) discovered that the utilization of lactic acid bacteria as a dietary component in sea bass did not exhibit significant effectiveness when compared to its potential outcomes on the feeding efficiency of juvenile carp. However, their investigation revealed that higher yield and feed utilization were achieved

through the dietary supplementation of *Streptococcus facium* in Israeli carp. In another study by Kennedy et al. (1998) noted higher survivability and feed utilization in marine fish with probiotic inclusion diets. A 20 days feeding trial with *A. hydrophila* in *Pangasianodon hypophthalmus* showed improved growth performance and diet efficiency (Mamun et al., 2020).

3.3 Non-specific immune parameters

Following the 90-day feeding trial, noteworthy discrepancies ($p < 0.05$) in total serum protein levels and superoxide anion production in the blood between the treatment and control groups. Although significant higher super oxide anion production was marked in the 10^6 cfu/g inclusion of *L. sporogenes* group (F2) however these values were not remarkably different with inclusion level of 10^8 cfu/g group (F3). Highest lysozyme activity was recorded in 10^8 cfu/g incorporated diet (F3), followed by 10^6 cfu/g (F2) and 10^4 cfu/g (F1) diets. Conversely, a meaningful decrease in lysozyme function recorded in the fishes that did not received probiotic supplementation.

The notable results showed that blood respiratory burst is considered as one of vital bactericidal mechanisms in fish (Ellis, 2001) and it also indicates the health status of the fish (Anderson, 1992). Successful use of probiotic bacteria like *Bacillus* sp. which provide higher yield of fish production and better health condition by evoking innate and adaptive immune system (Zhao et al., 2012). Previous findings showed elevated respiratory burst level with incorporation of 5% *Saccharomyces cerevisiae* in the diets of rainbow trout (Siwicki et al., 1994), seabream (Ortuno et al., 2002) and turbot (Toranzo et al., 1995). Similarly, EL-Boshy et al. (2010) documented a notable augmentation in the activity of neutrophils and macrophages in *Bacillus* sp., supplemented group in comparison to the control group. In our study we found significantly higher respiratory burst encountered in *L. sporogenes* supplemented group than the control. Total protein in serum is a non-specific humoral components indicated higher immunological state when fish possessed higher amount in blood (Mamun et al., 2022a). Meanwhile lower level of total protein in serum lead to microbial infections, organ failure and nutritional imbalance in aquaculture organisms (Rathore et. al., 2021).

Table 4: Production of super oxide anion, total serum protein, and lysozyme activity of *L. rohita* under treatments and control group (mean of 3 replicates± standard error)

Treatments	F ₀	F ₁	F ₂	F ₃
Super oxide anion production	0.175±0.070 ^b	0.186±0.033 ^b	0.232±0.057 ^a	0.197±0.035 ^{ab}
Total serum protein(mg/l)	38.34±0.271 ^b	41.69±0.52 ^b	55.40±0.66 ^a	47.07±0.62 ^{ab}
Lysozyme activity (U/ml)	910.33±2.60 ^b	1242.66±1.45 ^a	1253.00±4.48 ^a	1258.00±4.35 ^a

In our study fishes fed with probiotics exhibited stronger immune responses in contrast to the group not supplemented dietary *L. sporogenes*. Notably, major improvements in serum protein levels was noticed in the F₂ group, which had a probiotic count of 10⁶ cfu/g. A consistent pattern of positive effects stemming from probiotic supplementation on fish health and immunity. This trend is particularly evident in the significant enhancement of lysozyme expression and increased populations of beneficial bacteria like *Lactobacillus* sp. resulting from dietary lactic acid (LA) supplementation in common carp (Hoseini et al. 2023). This phenomenon aligns with the findings of Abdel-Latif et al. (2023), who reported elevated serum lysozyme activity and total immunoglobulin levels in *P. hypophthalmus* fingerlings following multispecies probiotic supplementation. Importantly, the positive impacts of *Bacillus* sp. supplementation extend beyond just lysozyme activity. As reported by Ji et al. (2023), dietary supplementation of *B. subtilis* significantly enhanced disease resistance in Chinese perch (*Siniperca chuatsi*), resulting in heightened resistance against the pathogen *A. hydrophila*. Additionally, El-Son et al. (2022) observed that Sanolife® PRO-F probiotic supplementation in Nile tilapia led to positive effects on antioxidant capacity and lysozyme activity. These multifaceted improvements in diverse immune response parameters further underline the comprehensive benefits of probiotics in promoting overall fish health and immune function. It is worth noting that the effects of probiotic interventions can be species-specific, as evidenced by the study conducted by Keereelang et al. (2022). Their research involving *L. plantarum* (LP) supplementation in black shark minnow (*Labeo chrysophekadion*) exhibited species-specific improvements in immune responses and disease resistance. This was reflected in the enhancement of lysozyme and peroxidase activities, alongside increased survival rates against

A. hydrophila infection. These findings underscore the importance of tailoring probiotic interventions to suit the unique requirements of different fish species. Moreover, the combination of probiotics with other dietary additives has showcased promising outcomes. Yousefi et al. (2023) demonstrated that the supplementation of Gum Arabic (GA) and/or *L. helveticus* (LH) yielded multiple improvements, including enhanced superoxide dismutase and catalase activities. These results suggest potential synergistic effects achievable by integrating probiotics with other dietary components, further enhancing fish health and immune responses.

Lysozyme activity, an important component of the nonspecific immune response in fish, is often used as an indicator of fish health status (Biller et al. 2021). According to Liu et al. (2012) feeding fish *E. coioides* with *B. subtilis* at a concentration of 10^8 cfu/g resulted in significantly higher lysozyme activity compared to lower concentrations and control diets. On the contrary, compared to the control fish, rainbow trout, *O. mykiss*, fed with a greater concentration of *L. rhamnosus* (JCM 1136) at 10^{11} cfu/g had considerably increased serum lysozyme activity (Ezabi et al. 2005). Mahmoudzadeh et al. (2016) supplemented *O. mykiss* diets with *B. subtilis* for the period of 44 days and observed enhanced lysozyme activity and total antibody levels, which aligns with the findings of El-Boshy et al. (2010), who supplemented *S. cerevisiae* to *O. niloticus* for around 21 days and observed similar results. In another study by Sîrbu et al. (2022), lysozyme activity significantly increased in the treatments including probiotics, prebiotics and synbiotics groups compared to the control group. Additionally, Opiyo et al. (2019) found that Nile tilapia grown in ponds with minimal input and supplemented with either *Saccharomyces cerevisiae* or *Bacillus subtilis* at different levels exhibited substantially higher lysozyme activity than the untreated group.

3.4 Body composition of *L. rohita*

The influence of *L. sporogenes* on the proximate whole-body composition is depicted in Table 5. Observed data showed no remarkable ($p>0.05$) difference in all the composite analysis (moisture, dry matter, protein, fat, ash and carbohydrate) of fish muscle either in the probiotic fed groups or without probiotic group despite the fact that fish fed probiotics showed greater levels. According to the current data of fish body composition, there is no noticeable difference between both the treatment and control groups. Likewise, Eid and Mohamed (2008) found no statistically significant variation in the proximate compositions of in tilapia fingerlings when they were supplemented with various levels of commercially produced feed additives.

Meanwhile, carcass content were significantly higher in probiotic mixed feed given to the juveniles of common carp, *Cyprinus carpio* (Mohsen et al. 2016).

Table 5: Comprehensive analysis of the proximate composition (%) of fish muscle considering multiple treatment conditions.

Treatments	F ₀	F ₁	F ₂	F ₃
Moisture (%)	77.55±0.052 ^a	77.98±0.037 ^a	77.55±0.037 ^a	78.16±0.037 ^a
Dry matter (%)	22.44±0.052 ^a	22.01±0.037 ^a	22.44±0.037 ^a	21.84±0.037 ^a
Protein (%)	15.08±0.17 ^a	15.01±0.23 ^a	15.52±0.11 ^a	15.30±0.33 ^a
Ether extract (%)	2.73±0.024 ^a	2.75±0.061 ^a	3.02±0.037 ^a	3.14±0.011 ^a
Ash (%)	2.10±0.017 ^a	2.09±0.018 ^a	2.07±0.014 ^a	2.10±0.020 ^a
NFE (%)	2.52±0.04 ^a	2.14±0.05 ^a	1.82±1.11 ^a	1.35±0.22 ^a

Nitrogen free extract (mean of 3 replicates± standard error).

3.5 Disease resistance of *L. rohita* to *Aeromonas hydrophila*

Feeding for around 90 days with the diet which contained *L. sporogenes* challenged with *A. hydrophila* showed significant increase in relative percentage survival (RPS). The RPS value in F₃, F₂ and F₁ group were 83.65%, 79.69% and 77.73% respectively (Figure 1). Significant incremental ($P<0.05$) cumulative mortality pattern were observed during 10 days of post injection in control group (73.33%) compared to the probiotic groups (26.67±3.95%). Challenged fish in the control group showed severe clinical signs including red spots, haemorrhagic lesions, tail and fin rots and scale loss.

In the present study fishes were challenged with *A. hydrophila* as these *Aeromonas* spp. are Gram-negative aquatic bacteria and ubiquitous in nature which causes fatal haemorrhagic septicemia in catfishes (Mamun et al. 2022b) and caused severe economic loss in freshwater cultured cyprinid fishes (Nithin et al. 2021). In fish, first line of protection against invading pathogens is predominantly attributed to the innate immune system (Magnadottir 2006). On that note Balcazar et al. (2007) found that *Cyprinus carpio* supplemented with probiotic diet showed survival rates increased up to 97.8 to 100% when exposed to *A. hydrophila*. According to Newaj-Fyzul et al. (2007), probiotic-fed fishes had higher protection level of 65 to 100% when challenged with *Aeromonas* spp., compared to 5 to 15% for control fish. Furthermore, the application of probiotic *B. subtilis* in *L. rohita* has been demonstrated to be effective in controlling *A. hydrophila* infection (Kumar et al. 2006). Aly et al. (2008) observed similar

results with the probiotics *B. subtilis* and *L. acidophilus*. It is also noted that *O. niloticus* fed with probiotics exhibited better protection against *A. hydrophila*.




Figure 1: The determination of the relative percentage of survival (RPS) in *L. rohita* was conducted across a range of treatments and the control group following a challenge with *A. hydrophila*.




Figure 2: The percentage cumulative mortality (CPM) of *L. rohita* was assessed in the different treatments and control group against *A. hydrophila*.

Conclusion

Dietary inclusion of *L. sporogenes* had a significant impact on growth efficiency, immunity, and exerted a defensive action against *A. hydrophila* pathogen. Different levels of probiotics in the present study acted in one or the other parameters and promoted the higher yield and immunity. Same view was noted by Tilwani et al. (2022) observed growth improvement and significant higher immunological indices such as lysozyme, phagocytic, immunoglobulin and superoxide dismutase level when Indian Major carp, *Cirrhinus mrigala* fed a probiotic supplemented diet. Considering all these findings, it has been strongly recommend that including *L. sporogenes* in the diet can effectively improve growth efficiency, enhance immunity, and offer protection against *A. hydrophila*. Nevertheless, it is crucial to emphasize the need for additional research to understand the intricate mechanisms responsible for the observed benefits of *L. sporogenes*. Furthermore, such research will facilitate the optimization of *L. sporogenes* application in aquaculture practices, ensuring its maximum efficacy and potential for enhancing feed utilization, growth improvements, innate immunity, and strategies for combating *A. hydrophila*.

References

- Abdel-Latif, H. M., Chaklader, M. R., Shukry, M., Ahmed, H. A. & Khallaf, M. A. (2023). A multispecies probiotic modulates growth, digestive enzymes, immunity, hepatic antioxidant activity, and disease resistance of *Pangasianodon hypophthalmus* fingerlings. *Aquaculture*, 563, 738948.
- Aly, S.M., Ahmed, Y., Ghareeb, A.A.A. & Mohamed, M.F. (2008). Studies on *Bacillus subtilis* and *Lactobacillus acidophilus*, as potential probiotics, on the immune response and resistance of *Tilapia nilotica* (*Oreochromis niloticus*) to challenge infections. *Fish and Shellfish Immunology*, **25**(1): 128-136.
- Anderson, D.P. (1992). Immunostimulants, adjuvants and vaccine carriers in fish: applications to aquaculture. *Annual Review of Fish Diseases*, **2**: 281-307.
- Anderson, D.P. & Siwicki, A. (1993). Basic haematology and serology for fish health. In Programs Paper Presented in Second Symposium on Disease in Asia Aquaculture Aquatic Animal Health and The Environment, Phuket, Thailand, pp. 185-202.
- AOAC. 1995. Official methods of analysis, 12th edn, W. Harwitz, (Ed) Association of Analytical Chemists, Washington D.C.

- Balcázar, J.L., Vendrell, D., de Blas, I., Ruiz-Zarzuela, I., Muzquiz, J.L. & Girones, O. (2008). Characterization of probiotic properties of lactic acid bacteria isolated from intestinal microbiota of fish. *Aquaculture*, **278**(1-4): 188-191.
- Barbosa, M., Jatoba, A., Vieira, F., Silva, B., Mourino, J. & Andreatta, E. (2011). Cultivation of juvenile fat snook (*Centropomus parallelus* poey) fed probiotic in laboratory conditions. *Brazilian Archives of Biology and Technology*, **54**(4): 795-801.
- Biller, J.D., Polycarpo, G.D.V., Moromizato, B.S., Sidekerskis, A.P.D., Silva, T.D.D., Reis, I.C.D. & Fierro-Castro, C. (2021). Lysozyme activity as an indicator of innate immunity of tilapia (*Oreochromis niloticus*) when challenged with LPS and *Streptococcus agalactiae*. *Revista Brasileira de Zootecnia*, **50**: e20210053.
- Chaudhary, A. & Qazi, J.I. (2006). Microbiological upgradation of formulated fish feed. *Proceedings of Pakistan Congress of Zoology*, **26**: 43-73.
- Dimitroglou, A., Merrifield, D.L., Carnevali, O., Picchietti, S., Avella, M., Daniels, C. & Davies, S.J. (2011). Microbial manipulations to improve fish health and production—a Mediterranean perspective. *Fish & shellfish immunology*, **30**(1): 1-16.
- Eid, A. & Mohamed, k.A. (2008). Effect of using probiotic as growth promoter in commercial diets for monosex Nile tilapia (*Oreochromis niloticus*) fingerlings. *8th International Symposium on Tilapia in Aquaculture Cairo Egypt 12-14 October*, pp 241-253.
- El-Boshy, M.E., El-Ashram, A.M., Abdel hamid, F.M. & Gadalla, H.A. (2010). Immunomodulatory effect of dietary *Saccharomyces cerevisiae*, β -glucan and laminaran in mercuric chloride treated Nile tilapia (*Oreochromis niloticus*) and experimentally infected with *Aeromonas hydrophila*. *Fish Shellfish Immunology*, **28**: 802-808.
- El-Ezabi, M., El-Serafy, S., Essa, M., Daboor, S. & Esmael, N. (2011). The viability of probiotics as a factor influencing the immune response in the Nile tilapia, *Oreochromis niloticus*. *Egyptian Journal of Aquatic Biology and Fisheries*, **15**(1): 105-124.
- Ellis, A.E. (2001). Innate host defense mechanism of fish against virus and bacteria. *Developmental and Comparative Immunology*, **25**: 827-839.
- El-Son, M. A., Elshopakey, G. E., Rezk, S., Eldessouki, E. A. & Elbahnaswy, S. (2022). Dietary mixed *Bacillus* strains promoted the growth indices, enzymatic profile, intestinal immunity, and liver and intestinal histomorphology of Nile tilapia, *Oreochromis niloticus*. *Aquaculture Reports*, **27**, 101385.

- FAO. (2020). Aquaculture. <http://www.fao.org/fishery/aquaculture/en>, 2020 (accessed on 20 Dec 2020).
- Gatesoupe, F.J. (1999). The use of probiotics in aquaculture. *Aquaculture*, **180**: 147–165.
- Gupta, A. & Dhawan. A. (2012). Effect of dietary probiotic improval (*Lactobacillus sporogenes* and *Saccharomyces cerevisiae*) on growth and feed utilization of *Macrobrachium rosenbergii* post larvae. *Animal Nutrition and Feed Technology*, **12**: 209-217.
- Harikrishnan, R. & Balasundaram, C. (2005). Modern trends in *Aeromonas hydrophila* disease management with fish. *Reviews in Fisheries Science*, **13**: 281-320.
- Harikrishnan, R., Balasundaram, C. & Heo, M.S. (2010). *Lactobacillus sakei* BK19 enriched diet enhances the immunity status and disease resistance to streptococcosis infection in kelp grouper, *Epinephelus bruneus*. *Fish and Shellfish Immunology*, **29**: 1037-1043.
- Hastings, W.H. (1976). Fish nutrition and fish feed manufacture. In FAO, Rome (Italy). *Fishery Resources and Environment Div.* FAO Technical Conference on Aquaculture. Kyoto (Japan).
- Hoseini, S. M., Yousefi, M., Afzali-Kordmahalleh, A., Pagheh, E. & Taheri Mirghaed, A. (2023). Effects of dietary lactic acid supplementation on the activity of digestive and antioxidant enzymes, gene expressions, and bacterial communities in the intestine of common carp, *Cyprinus carpio*. *Animals*, 13(12), 1934.
- Irianto, A. & Austin, B. (2002). Probiotics in aquaculture. *Journal of Fish Disease*, **25**: 633-642.
- Ji, Z., Zhu, C., Zhu, X., Ban, S., Yu, L., Tian, J. & Jiang, M. (2023). Dietary host-associated *Bacillus subtilis* supplementation improves intestinal microbiota, health and disease resistance in Chinese perch (*Siniperca chuatsi*). *Animal Nutrition*, 13, 197-205.
- Keereelang, J., Mangumphan, K., Chitmanat, C., Tongsiri, S., Linh, N. V. & Van Doan, H. (2022). Dietary effect of *Lactobacillus plantarum* (TISTR 912) on digestive enzyme activity, growth performance, immune response, and disease resistance of black shark minnow (*Labeo chrysophekadion*) against *Aeromonas hydrophila* infection. *Aquaculture Reports*, 27, 101409.
- Kennedy, S.B., Tucker, J.W., Thoresen, D. & Sennett, G. (1998). Current methodology for the use of probiotic bacteria in the culture of marine fish larvae. *Aquaculture* 98. *World Aquaculture Society*. Baton Rouge, 286

- Kumar, R., Mukherjee, S.C., Prasad, K.P. & Pal, A.K. (2006). Evaluation of *Bacillus subtilis* as a probiotic to Indian major carp *Labeo rohita* (Ham.). *Aquaculture Research*, **37**: 1215-1221.
- Liu, C.H., Chiu, C.H., Wang, S.W. & Cheng, W. (2012). Dietary administration of the probiotic, *Bacillus subtilis* E20, enhances the growth, innate immune responses, and disease resistance of the grouper, *Epinephelus coioides*. *Fish & shellfish immunology*, **33**: 699-706.
- Magnadottir, B. (2006). Innate immunity of fish (overview). *Fish & Shellfish Immunology*, **20**: 137–151.
- Mahmoudzadeh, L., Meshkini, S., Tukmehchi, A., Motalebi Moghanjoghi, A.A. & Mahmoudzadeh, M. (2016). Effects of dietary *Bacillus subtilis* on growth performance and immune responses, in rainbow trout, *Oncorhynchus mykiss* (Walbaum, 1792). *Iranian Journal of Fisheries Sciences*, **15**: 347-359.
- Mamun, M. A. A., Nasren, S., Abhiman, P. B., Rathore, S. S., Sowndarya, N. S., Ramesh, K. S. & Shankar, K. M. (2020). Effect of biofilm of *Aeromonas hydrophila* oral vaccine on growth performance and histopathological changes in various tissues of striped catfish, *Pangasianodon Hypophthalmus* (Sauvage 1878). *Indian Journal of Animal Research*, **54**(5): 563-569.
- Mamun, M.A.A., Nasren, S., Abhiman, P.B., Rathore, S.S., Rakesh, K., Sowndarya, N.S. & Shankar, K.M. (2022a). Evaluation of feed utilization, immune response and disease resistance in striped catfish, *Pangasianodon hypophthalmus* (Sauvage 1878) fed with a novel *Aeromonas hydrophila* biofilm vaccine. *Fish and Shellfish Immunology Reports*, 100070.
- Mamun, M.A.A., Nasren, S., Rathore, S.S. & Mahbub Alam, M.M. (2022b). Histopathological analysis of striped catfish, *Pangasianodon hypophthalmus* (Sauvage, 1878) spontaneously infected with *Aeromonas hydrophila*. *Jordan Journal of Biological Sciences*, **15**: 93-100.
- Mamun, M.A.A., Nasren, S., Rathore, S.S., Sidiq, M.J., Dharmakar, P. & Anjusha, V.K. (2019). Assessment of probiotic in aquaculture: functional changes and impact on fish gut. *Microbiology Research Journal International*, **29**: 1-10.
- Merrifield, D.L., Bradley, G., Baker, R.T.M. & Davies, S.J. (2010). Probiotic applications for rainbow trout (*Oncorhynchus mykiss* Walbaum) II. Effects on growth performance,

- feed utilization, intestinal microbiota and related health criteria postantibiotic treatment. *Aquaculture nutrition*, **16**: 496-503.
- Mohapatra, S., Chakraborty, T., Prusty, A.K., Das, P, Paniprasad, K. & Mohanta, K.N. (2012). Use of different microbial probiotics in the diet of rohu, *Labeo rohita* fingerlings: effects on growth, nutrient digestibility and retention, digestive enzyme activities and intestinal microflora. *Aquaculture Nutrition*, **18**: 1-11.
- Mohsen, S.H., Ahmed, Z., Hakim, N.F.A.E., Nawsany, M.E. & State, H.A.A. (2016). Effect of different growth promoters on growth performance, feed utilization and body composition of common carp (*Cyprinus carpio*). *Journal of Fisheries and Aquatic Science*, **11**: 370-377.
- Nayak, A.K., Das, B.K., Kohli, M.P.S. & Mukherjee, S.C. (2004). The immune suppressive effect of α -permethrin on Indian major carp, rohu (*Labeo rohita*). *Fish and Shellfish Immunology*, **16**: 41-50.
- Nesara, K. M., Jayaraj, E. G., Amoga, K. R. & Sandeep, C. V. (2018). Effects of dietary probiotic and organic acid alone and in combination on the growth performance of Indian major carp, *Labeo rohita*. *Journal of Experimental Zoology, India*, **21**: 805-812.
- Newaj-Fyzul, A., Adesiyun, A.A., Mutani, A., Ramsubhag, A., Brunt, J. & Austin, B. (2007). *Bacillus subtilis* AB1 controls *Aeromonas* infection in rainbow trout (*Oncorhynchus mykiss*, Walbaum). *Journal of applied microbiology*, **103**: 1699-1706.
- Nithin, M.S., Girisha, S.K., Kushala, K.B., Chandan, D.V., Puneeth, T.G., Kumar, B.T.N., Vinay, T.N., Suresh, T., Lopamudra, S. & Ramesh, K.S. (2021). Novel lytic bacteriophages (AhFM4 & AhFM5) as bio-control measures against multidrug resistant biofilm producing *Aeromonas hydrophila* (AhZ1K). *Aquaculture*, **544**: 737106.
- Noh, S.H., Han, K., Won, T.H. & Choi, Y.J. (1994). Effect of antibiotics, enzymes, yeast culture and probiotics on the growth performance of Israeli carp. *Korean Journal of Animal Science*, **36**: 480-486.
- Opiyo, M. A., Jumbe, J., Ngugi, C. C. & Charo-Karisa, H. (2019). Dietary administration of probiotics modulates non-specific immunity and gut microbiota of Nile tilapia (*Oreochromis niloticus*) cultured in low input ponds. *International journal of veterinary science and medicine*, **7**: 1-9.
- Ortuno, J., Cuesta, A. Rodriguez, A. Esteban, M. A. & Meseguer, J. (2002). Oral administration of yeast, *Saccharomyces cerevisiae*, enhances the cellular innate

- immune response of *gilthead seabream (Sparus aurata L.)*. *Veterinary Immunology and Immunopathology*, **85**: 41-50.
- Padala, D., Ganapathi, M.N., Anjusha, K.V., Prabhakaran, P.L., Mamun, M.A.A. & Ramesh, K.S. (2021). Effect of dietary peppermint (*Mentha piperita*) on growth, survival, disease resistance and haematology on fingerlings of rohu (*Labeo rohita*). *Aquaculture Research*, **52**: 2697-2705.
- Parry, R.M., Chandan, R.C. & Shahani, K.M. (1965). A rapid and sensitive assay of muramidase. *Proceedings of the society for experimental biology and medicine*, **119**: 384-386.
- Parthasarathy, R. & Ravi, D. (2011). Probiotic bacteria as growth promoter and biocontrol agent against *Aeromonas hydrophila* in *Catla catla* (Hamilton, 1822). *Indian Journal of Fisheries*, **58**(3): 87-93.
- Prasad, L., Nayak, B.B., Kohli, M.P.S., Reddy, A.K. & Srivastava, P.P. (2012). Effect of supplemented bacteria (*Lactobacillus sporogenes*) on growth of *Macrobrachium rosenbergii* postlarvae. *Israeli Journal of Aquaculture-Bamidgeh*, **64**: 1-6
- Rathore, S.S., Murthy, H.S., Mamun, M.A.A., Nasren, S., Rakesh, K., Kumar, B.T.N. & Khandagale, A.S. (2021). Nano-selenium supplementation to ameliorate nutrition physiology, immune response, antioxidant system and disease resistance against *Aeromonas hydrophila* in monosex Nile tilapia (*Oreochromis niloticus*). *Biological trace element research*, **199**: 3073-3088.
- Reed, L. J. & Muench, H., 1938. A simple method of estimating fifty per cent endpoints. *American journal of epidemiology*, **27**: 493-497
- Seenivasan, C., Bhavan, P.S., Radhakrishnan, S. & Shanthi, R. (2012). Enrichment of Artemia nauplii with *Lactobacillus sporogenes* for enhancing the survival, growth and levels of biochemical constituents in the post-larvae of the freshwater prawn *Macrobrachium rosenbergii*. *Turkish Journal of Fisheries and Aquatic Sciences*, **12**(1).
- Sîrbu, E., Dima, M. F., Tenciu, M., Cretu, M., Coadă, M. T., Țoțoiu, A., & Patriche, N. (2022). Effects of Dietary Supplementation with Probiotics and Prebiotics on Growth, Physiological Condition, and Resistance to Pathogens Challenge in Nile Tilapia (*Oreochromis niloticus*). *Fishes*, **7**: 273.

- Siwicki, A.K., Anderson, D.P. & Rumsey, G.L. (1994). Dietary intake of immunostimulants by rainbow trout affects non-specific immunity and protection against furunculosis. *Veterinary Immunology and Immunopathology*, **41**: 125-139.
- Sridhar, K. & Joice, P.E. (2012). Efficacy of probiotic bacteria *Lactobacillus sporogenes* on zinc toxicated fresh water fish *Cyprinus carpio*. *International Journal of Advanced Life Sciences*, **3**:72-80.
- Subasinghe, R.P. (2005). Epidemiological approach to aquatic animal health management: opportunities and challenges for developing countries to increase aquatic production through aquaculture. *Preventive veterinary medicine*, **67**: 117-124.
- Tilwani, Y.M., Sivagnanavelmurugan, M., Lakra, A.K., Jha, N. & Arul, V. (2022). Enhancement of growth, innate immunity, and disease resistance by probiotic *Enterococcus faecium* MC-5 against *Aeromonas hydrophila* in Indian Major carp *Cirrhinus mrigala*. *Veterinary Immunology and Immunopathology*, 110503.
- Toranzo, A.E., Devesa, S., Romalde, J.L., Lamas, J., Riaza, A., Leiro, J. & Barja, J.L. (1995). Efficacy of intraperitoneal and immersion vaccination against *Enterococcus* sp. infection in turbot. *Aquaculture*, **134**: 17-27.
- Verschuere, L., Rombaut, G., Sorgeloos, P. & Verstraete, W. (2000). Probiotic bacteria as biological control agents in aquaculture. *Microbiology and molecular biology review*, **64**: 655–671.
- Yousefi, M., Farsani, M. N., Ghafarifarsani, H. & Raeeszadeh, M. (2023). Dietary *Lactobacillus helveticus* and Gum Arabic improves growth indices, digestive enzyme activities, intestinal microbiota, innate immunological parameters, antioxidant capacity, and disease resistance in common carp. *Fish & Shellfish Immunology*, 135, 108652.
- Zhao, Y.C., Zhang, W.B., Xu, W., Mai, K.S., Zhang, Y.J. & Liu, Z.G. (2012). Effects of potential probiotic *Bacillus subtilis* T13 on growth, immunity and disease resistance against *Vibrio splendidus* infection in juvenile sea cucumber *Apostichopus japonicus*. *Fish & Shellfish Immunology*, **32**: 750–755.