

Survey and monitoring of thrips (*Thrips tabaci*) Lindeman in onion crop

Abstract

Thrips (*Thrips tabaci* Lindeman) is a major pest of onion and cause considerable losses in yield. The nymph and adult of thrips feed leaves by piercing and rasping the leaf tissues and causes lengthwise, silvery stippling or blotching on the onion leaves, resulting in the loss of chlorophyll and reduced photosynthetic efficiency. Thrips attack onion at all ~~the stages of crop growth~~ crop growth stages but their count increases from bulb initiation and ~~remain-remains~~ high up to bulb development and maturity. Onion thrips can also transmit several plant pathogens in onion crop that reduced onion bulb size and quality. Therefore, ~~the objective of this survey was undertaken~~ this survey's objective was to evaluate the damage ~~percent-percentage~~ of thrips in onion ~~crop~~ crops. Survey was conducted during Kharif 2022 and Rabi 2022-23. The maximum thrips ~~damge-damage~~ rating scale of 3 (45%) was recorded at 75 DAT in ~~the~~ Kharif season and Rabi season at 75 DAT thrips damage rating scale ~~were-was~~ recorded at 4 (65%). The survey result show that the thrips population and damage ~~percent-percentage~~ increased due to increased temperature and relative humidity.

Key words Keywords: *Allium cepa*, *Thrips tabaci*, survey

Introduction

Onion (*Allium cepa* L.) is an important vegetable or ~~spices-spice~~ crop cultivated in almost all the ~~states of the country~~ country's states. In India onion is cultivated in 3 seasons i.e., *rabi*, *kharif* and *late kharif* seasons and ~~the~~ maximum area under cultivation is ~~being~~ covered in ~~the~~ *rabi* season (about 60-65%). Thrips (*Thrips tabaci* Lindeman) is a regular and potential pest of onion and cause considerable losses as high as 90% in quality and yield (Gupta *et. al.*, 1984). Srinivas and Lawande (2004) reported that *Thrips tabaci* could cause yield loss in the range of 46-87% in onion. Waiganjoet. *al.*, (2008) estimated the foliage damage of ~~crop-crops~~ around 40-60% which can ~~led-lead~~ to yield losses of 10-20% in the crop. Shibru and Negeri. (2014) reported that onion thrips cause damage on yield 23-85%. Onion thrips are an important vector also for several plant viruses such as tomato spotted wilt virus (Kritzman *et. al.*, 2002). Failure to control this pest by timely and effective means causes considerable damage and ~~result-results~~ in immense economic loss by remarkably reduced yield (Anonymous.2000, Jaun, 2002). Insecticides are a major tool for thrips control, but this strategy is inadequate and unsustainable (Maniania *et. al.*, 2003) because thrips have developed resistance ~~towards-to~~ various ~~group-groups~~ of insecticides

(Lebedev *et. al.*, 2013). The farmers have intensively used particularly those insecticides ~~which-that~~ are used to control this pest and repeated application of the same group of ~~insecticide-insecticides~~ could have led to the development of resistance to insecticides (Shitoleet. *al.*, 2002, Alston & Drost 2008). Pathak *et. al.*, (2018) reported that the lowest thrips population and highest onion seed yield ~~was-were~~ recorded with the application of fipronil insecticides. Turkey and Kumar (2017), Kurbett *et. al.*, (2015) reported that thiamethoxam was ~~proved-provento~~ to be the most effective for thrips control. Asgar *et. al.*, (2018) suggested that the insecticides reduced the thrips population compared to the control and the highest yield was obtained by the use of Dimathoate. Kumar and Singh (2011), and Daset. *al.*, (2017) reported that spray of Imidacloprid given at 15 days interval recorded the lowest thrips population and gave the highest gross yield. Pandey *et al.*, (2013) reported that the lowest thrips population and highest bulb yield was obtained by applying fipronil.

Origin of onion thrips

Thrips tabaci was first reported by Russian Entomologist Karl Eduard Lindeman based on specimens collected in Bessarabia, Russia, that caused severe damage to tobacco plants (Lindeman 1889).

Geographic distribution of onion thrips

The onion thrips ~~is-area~~ global pest of onion grown between ~~see-sea~~ level and 2000 m (Lewis 1973). Onion thrips is a native of the Mediterranean region but has become a major pest of agricultural crops throughout most of the world (Mound 1977).

Biology of onion thrips

~~Thrips-Thrips~~ life cycle deepened environmental ~~condition-conditions~~, like temperature, humidity, and nutrient quality of their food source. ~~Stage-The stages~~ in the developmental life cycle ~~is-are~~ the egg, first larval stage, second larval stage, ~~pre-pupapre-pupa~~, pupa, and adult. Because of their small size, this pest species like other thrips ~~can-not~~ readily be identified ~~to-as-a~~ species even with a hand lens. Adult specimens are usually needed to make species identification under high microscope magnification (Morse *et. al.*, 2005). Brain A.N. 2006 mentioned that the biology of onion thrips was as ~~followed-follows~~ the entire life cycle (egg to adult) required about 19 days. Large ~~population-populations~~ are able ~~to-can~~ develop quickly under dry weather ~~condition-conditions~~ where there are many overlapping generations throughout year. Females have a ~~saw-like~~ saw-like structure that help to make an incision in plant tissue for egg laying. Eggs are placed singly just under the epidermis of succulent leaf, flower, stem, or bulb tissue. Eggs are elliptical, approximately 0.2 mm in length. They are whitish at deposition and change to an orange tint as development conditions. Hatching occurs in 4-5 days. Larvae are whitish to yellowish. There are two larval stages and besides the adults, they are the only damaging stages. Larval development is completed in about 9-10 days.

There are two non-feeding stages called the prepupa and pupa does not feed and ~~occur~~ occurs primarily in the soil. Combined prepupal and pupal development ~~is~~ is completed in 4-6 days. ~~Adult~~ The ~~adult~~ are ~~is~~ about 1 mm in length their body ~~colour~~ color ranges from pale yellow to dark brown, wings are ~~un-banded~~ unbanded and dirty gray. The males are wingless and exceedingly rare while the ~~female~~ females have long, narrow fringed wings. ~~Female~~ Females live for 12-30 days and lay 50-60 ~~kidney-shaped~~ kidney-shaped eggs singly inside leaf tissue with a ~~sharpe~~ sharp ovipositor.

Economic importance

Thrips is a serious pest on a wide range of fruit, vegetable, flower, and Agronomics crops. Thrips are members of the order Thysanoptera, which contains ~~a number of~~ several genera and species. Among species of thrips that attack onion are onion thrips (*Thrips tabaci*). Onion thrips incidence was a major problem due to the damage caused by feeding on vegetative parts which caused discoloration, deformities and reduced marketability the crop thrips may also serve as vectors for plant diseases such as tospovirus (Nault LR 1997). These enveloped viruses are considered among some of the most damaging of emerging plant ~~pathogen~~ pathogens around the world. Thrips damage is usually measured as an overall reduction in bulb size and weight of bulb.

Damage symptoms

Thrips tabaci is considered an ~~in-direct~~ indirect pest of dry bulb onion because it feeds on leaves rather than the marketable portion of the crop. Thrips feeding on onion leaves ~~causes~~ causes silvery leaf ~~spot~~ spots that turn ~~in to~~ into white blotches along the leaves due to removal of cellular content followed by the development of silvery patches and curling of leaves (Bailey 1938). *Thrips tabaci* causes significant yield loss despite decades of research on control strategies worldwide (Lewis 1997b). *T. tabaci* feeding can reduce onion bulb weight (Kendall and Capinera 1987, Fournier *et al.*, 1995, Rueda *et al.*, 2007, Diaz-Monato *et al.*, 2010, Waiganjo *et al.*, 2008 and cause up to 60% yield loss. In addition to injury by feeding *T. tabaci* transmits IYSV and is the only confirmed vector of this pathogen (Poizzer *et al.*, 1994, Kritzman *et al.*, 2001). IYSV was first identified on onion in southern Brazil in 1981 (Poizzer *et al.*, 1994) and was confirmed in the United states in 1989 and worldwide (Gent *et al.*, 2006). IYSV symptoms on leaves appear as lesions (i.e., straw colour to white, dry, and sometimes elongate) along the edges (Gent *et al.*, 2004) indicated that IYSV infection can reduced bulb size. The IYSV infects onion plants early in the growing season, onion yield losses may increase (Diaz-Montano *et al.*, 2010).

Weather factors on outbreak of onion Thrips

Relatively high temperatures and lack of rainfall have been associated with increase in onion thrips population, while high relative humidity and rainfall reduce thrips population (Hamdy, M.K. *et al.*, 1994). In addition to their effect on thrips activity, temperature and relative humidity further influence the intrinsic rate of natural increase of the thrips (Murai T. 2000). The rate of development of thrips *tabaci* is positively affected by increased

temperature and decreased by increased relative humidity (Hamdy, M.K. and Salem M.*et al.*, 1994). The hot and dry weather promotes the increase of *T.tabaci* populations (Bailey1934, Rueda *et al.*, 2007) and the severity of thrips injury (Lewis1973) Additionally water stress may impact the nutritional quality of onion plants and also increases the attractiveness of the plants to thrips .

Material and methods

The survey of onion thrips was conducted during *Kharif* 2022 and Rabi 2022-23in Nashik district taluka Niphad ,Maharashtra .During the survey five villages were selected for survey in and five onion fields in each village for observation of thrips damage scale.The survey conducted twice a season at 45 days after transplanting and 75 days after transplanting.The thrips damage rating scales at least 10 plants randomly in each field should be scored using rating scale given below.

List 1 :Thrips damage Rating

Scale:

Scale	Foliage damage rating
1	1-20%
2	21-40%
3	41-60%
4	61-80%
5	81-100%

Results and Discussion

Kharif 2022

The twenty five different onion fields were surveyed in 5 villages during September and October 2022.The survey for thrips infestation in onion crop was done in all the 25 fields in month of September on 29.09.2022 at different Villages likes – Chitegaon,Chandori,Saikheda ,Chatori and Berewadi of Taluka Niphad District Nashik,Maharashtra.The thrips damage rating scale ranged from 1 to 2 scale were recorded during survey. Based on the survey onion thrips infested fields were recorded 25% at 45 DAT in Saikheda village farmers fields. The mostly farmers transplanting was done of kharif onion in 15th August.During the survey period crop is under vegetative stage. At 75 DAT further the survey of thrips infestation was done in same villages and same 25 onion fields at Taluka Niphad District Nashik in month of October on 29.10.2022 .The thrips damage rating ranged from 1 to 3 scale were recorded during survey. Based on the survey onion thrips infested fields were recorded 45% inSaikheda village farmers fields. During the survey period crop is under bulb maturity stage.

Table 1- Thrips damage rating in villages at 45 DAT

Sr.no	Villages	Damage rating
1	Chitegaon	1
2	Chandori	1
3	Saikheda	2
4	Chatori	1
5	Berewadi	1

Table 2- Thrips damage rating in villages at 75 DAT

Sr.no	Villages	Damage rating
1	Chitegaon	1
2	Chandori	2
3	Saikheda	3
4	Chatori	2
5	Berewadi	1

The result show that overall highest thrips damage rating was recorded 2 (25% damage) at 45 DAT in farmers village Saikheda Taluka Niphad. Further the same places survey result show that at 75 DAT the overall highest thrips damage rating was recorded 3 (45% damage) at same village Saikheda Taluka Niphad due to higher temperature of this week. During this period temperature range from 20.24 to 32.42 °C and relative humidity range from 77.90 to 86.87% in village Saikheda.

Rabi 2022-23

The survey of onion thrips Nashik district taluka Niphad was conducted during *Rabi* 2022 on 45 DAT and 75 DAT. During the survey five villages were selected for survey in each taluka and five onion fields in each village for observation of thrips damage scale. The twenty five different onion fields were surveyed in 5 villages during January and February 2023.

The survey for thrips infestation in onion crop was done in all the 25 fields in month of January at different Villages likes –Chitegaon, Chandori, Saikheda, Chatori and Berewadi of Taluka Niphad District Nashik, Maharashtra. The thrips damage rating scale ranged 1-2 scale were recorded during survey. Based on the survey onion thrips infested fields were recorded 30% at 45DAT..

At 75 DAT further the survey of thrips infestation was done in same villages and same 25 onion fields at Taluka Niphad District Nashik. The thrips damage rating ranged from 1-4 scale were recorded during survey. Based on the survey onion thrips infested fields were recorded 65%.

Table 3- Thrips damage rating in villages at 45 DAT

Sr.no	Villages	Damage rating
1	Chitegaon	1
2	Chandori	1
3	Saikheda	2
4	Chatori	1
5	Berewadi	2

Table 4- Thrips damage rating in villages at 75 DAT

Sr.no	Villages	Damage rating
1	Chitegaon	2
2	Chandori	2
3	Saikheda	4
4	Chatori	2
5	Berewadi	3

The result show that overall highest thrips damage rating was recorded 2 (30% damage) at 45 DAT in village Saikheda and Berewadi and minimum thrips damage sacle 1 (20%) were recorded in farmers field Chitegaon village Taluka Niphad. Further the same places survey result show that at 75 DAT the overall highest thrips damage rating was recorded 4 (65% damage) at village Saikheda Taluka Niphad due to higher temperature of this week and minimum thrips damage scale 2 recorded in farmersfields villages Chitegaon, Chandori and Chatori. During this period temperature range from 12.05 to 26.28 °C and relative humidity range from 62.26 to 72.83% in village Saikheda.

The candidate manuscript does not have a robust scientific discussion, I suggest the authors incorporate the suggested paragraphs, in this way it would improve the scientific quality of the manuscript:

The survey and monitoring study of onion thrips (*Thrips tabaci*) in Maharashtra reveals that environmental conditions, particularly temperature and relative humidity, play a crucial role in pest population dynamics and the extent of crop damage. Thrips population and damage levels were observed to be highest during the bulb maturity stage, with a strong positive correlation with temperature increases. These findings align with studies in Colombia and Venezuela, where the incidence of pests like thrips in onions, tomatoes, and other vegetables is exacerbated by warm, dry conditions (Campos, 2023; Campos et al. 2023). For instance, thrips thrive in Colombia's warmer regions with prolonged dry spells, indicating a similar response to environmental conditions across different geographic locations (Zingaretti and Olivares, 2018; Rodriguez et al. 2023). Likewise, in Venezuela, studies have documented that high temperatures coupled with low rainfall contribute to

Formatted: Indent: Left: 0"

Formatted: Font: Not Bold

Formatted: Font: Not Bold, Italic

Formatted: Font: Not Bold

Formatted: Justified

Formatted: Font: Not Bold

Formatted: Font: Not Bold

thrips outbreaks, highlighting the pest's dependency on climatic factors common to the region (Bertorelli and Olivares, 2020; Parra et al. 2012).

Formatted: Font: Not Bold

The influence of humidity on thrips populations in the Maharashtra study also parallels observations in Latin America. Thrips populations in Venezuelan onion fields are notably lower in regions with high rainfall and humidity, as humidity disrupts thrips' ability to feed and reproduce effectively (Viloria et al. 2023). In Colombia, thrips management strategies for onions and other bulb crops often rely on irrigation schedules that enhance moisture levels during critical pest development phases, which has shown to reduce pest prevalence (Olivares and Hernandez, 2019b). This tactic highlights the importance of integrated pest management (IPM) strategies that incorporate environmental control measures to manage pest pressure (Olivares, 2016; Olivares et al. 2021). Both studies demonstrate the need to adapt pest management practices to local climate conditions, emphasizing that temperature and humidity are paramount to thrips population dynamics across these different agricultural systems (Vega et al. 2022; Olivares and Hernandez, 2019a).

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Agronomic practices also influence pest dynamics, as evidenced by the current study's recommendations for intercropping, bio-pesticides, and selective insecticides. Colombian and Venezuelan farmers commonly employ intercropping with crops like carrots and tomatoes to manage pest populations, as intercropping reduces pest prevalence by altering microclimates within crop fields (Zingaretti et al. 2016; Viloria et al. 2023). Additionally, the use of organic solutions such as neem oil and botanical-based pesticides, like in the Maharashtra study, is increasingly popular in Latin America, where small-scale farmers seek cost-effective and ecologically sustainable pest control measures. This approach reduces thrips populations by creating less favorable environments for pests while enhancing plant resilience against pest attacks (Olivares et al. 2017; Olivares et al. 2020).

Formatted: Font: Not Bold

Formatted: Font: Not Bold

The cumulative findings from India, Colombia, and Venezuela highlight that pest management in bulb crops like onions benefits greatly from holistic approaches that integrate environmental (Hernandez et al. 2020), agronomic (Hernandez and Olivares, 2020), and chemical controls (Hernandez et al. 2018). Future research should focus on developing pest-resistant onion varieties suited to warm, humid environments. Studies could further assess the efficacy of entomopathogenic fungi and natural predators in both regions to offer alternative pest control solutions, reducing reliance on chemical insecticides and mitigating resistance issues (Montenegro et al. 2021). This comparison underscores the interconnectedness of environmental conditions and agronomic practices in shaping effective, sustainable pest management strategies across diverse agricultural landscapes.

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Conclusion

Keeping in view the above facts that the present surveys shows that the thrips population and damage rating increased due to increased temperature and relative humidity. Thrips are major insect pest of onion crop, reduce the yield potential. Management of bulb crops pests relies on insecticide use at transplanting. But insecticide resistance can cause control failure that threaten the long-term viability of this strategy. IPM strategies minimize the loss and increase the marketable value of bulb crops. Thrips population was positively correlated with temperature and negatively with relative humidity and rainfall. Cultural practice such as intercropping with carrot, tomato spider plant, mixed cropping, time of planting, spacing, balance nitrogenous fertilizer dose, irrigation application, coloured sticky traps, bio-pesticides, *B. bassiana*, neem oil, basil oil, geranium oil, datura and neem products can also be used for effective control of thrips and some insecticides fipronil, thiamethoxam, dimethoate, imidachlorprid, spinosad, acetamiprid are can also be used for effective control of onion thrips.

Therefore, it needs to focus on the future the management aspects of onion thrips is need attention to the researchers. It's important to develop resistant varieties, use of entomopathogenic fungi, mass rearing and release of natural enemies, use selective different classes of insecticides within a season to avoid resistance and use all available cultural practices. In addition to these it is necessary to develop alternative tactics that are practical to implement.

Disclaimer (Artificial intelligence)

Authors hereby declare that NO generative AI technologies such as large language models and text to image generators have been used during the writing or editing of this manuscript.

References

- Anonymous (2000). Annual Report: National Research Centre for onion and Garlic. pp.63-64.
- Alston D.G., Drost D. (2008). Utah pests fact sheet. Onion thrips (Thrips tabaci). Internet <http://utahpests.edu/IPM/files/uplod/pdfdocs/factsheet-pdf/ENT-117>

- Asghar Muhammad, Mirza Muhammad Qaddeer Baig, Muhammad Afjal, Naeem Faisal (2018) Evaluation of different insecticides for the management of onion thrips (*Thrips tabaci* Lindeman) on onion crop. *Polish journal of entomology* Vol.87:165-176.
- Brain AN (2006) Biology and Ecology of onion thrips on onion fields *PP2. Cornell University*.
- Bailey, S.F. (1938) Thrips economic importance in California, *Univ. Calif. Coll. Agric. Agric. Exp. Stn. Cir. 346*.
- Bertorelli, M., and Olivares, B.O. Population fluctuation of *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) in sorghum cultivation in Southern Anzoategui, Venezuela. *J. Agric. Univ. P. R.* 2020, 104(1):1-16. <https://doi.org/10.46429/jaupr.v104i1.18283>
- Campos, B. O. (2023). *Banana Production in Venezuela: Novel Solutions to Productivity and Plant Health*. Springer Nature. <https://doi.org/10.1007/978-3-031-34475-6>
- Campos, B. O. O., Araya-Alman, M., & Marys, E. E. (2023). *Sustainable Crop Plants Protection: Implications for Pest and Disease Control* (p. 200). MDPI-Multidisciplinary Digital Publishing Institute. <https://doi.org/10.3390/books978-3-0365-9150-6>
- Das Ajay Kumar, Wajid Hasan and Sushil Kumar Singh (2017). Management of onion thrips, *Thrips tabaci* Using Chemical and Bio-pesticides for Quality onion production. *Trends in Biosciences* 10(22)4384-4388.
- Diaz-Montano J, Fuchs M, Nault BA, Fail J, Shelton AM (2010). Evaluation of onion cultivars for resistance to onion thrips (Thysanoptera: Thripidae) and Iris yellow spot virus. *J. Econ. Entomol.* 103, 925-937.
- Fournier F, Guy B, Robin S. (1995). Effect of thrips *tabaci* (Thysanoptera: Thripidae) on yellow onion yields and economic thresholds for its management. *Entomological Society of America*. 88(5):1401-1407.
- Gupta, R.P., Srivastava, V.K., Bhardwaj, B.S. and Pandey U.B. (1984). Chemical control of *Thrips tabaci* L. infesting onion crop. *J. Ent. Res.* 8(2)196-198.
- Gent, D.H., HR. Schwartz, and R. Khosla. (2004). Distribution and incidence of Iris yellow spot virus in Colorado and its relation to onion plant population and yield. *Plant Dis.* 88:446-452.
- Gent, D.H., L.J. du Toit, S.F., Fichtner, S.K., Mohan, H.R. Pappu and H.F. Schwartz (2006) Iris yellow spot virus an emerging threat to onion bulb and seed production. *Plant Dis.* 90:1468-1480.
- Hamdy MK and Salem M (1994) .The effect of plantation dates of onion, temperature and relative humidity on the population density of the onion

Formatted: Font: Not Italic

Formatted: Portuguese (Brazil)

Formatted: Hyperlink, Font: (Default) +Body, 11 pt, Portuguese (Brazil)

Field Code Changed

Formatted: English (United States)

Formatted: English (United States)

Formatted: English (United States)

Formatted: Spanish (Spain, International)

Formatted: Space After: 10 pt, Line spacing: Multiple 1.15 li

Formatted: English (United States)

thrips. *Thrips tabaci* Lind in Egypt. *Ann Agric Sci Univ Ain Shams(Egypt)* 39:417-424.

- [Hernández, R., Olivares, B., Arias, A., Molina, JC. & Pereira, Y. Agroclimatic zoning of corn cultivation for the sustainability of agricultural production in Carabobo, Venezuela. Rev. Univ. de Geog. 2018, 27 \(2\): 139-159. https://n9.cl/i0upn](#)
- [Hernandez, R.; Olivares, B.; Arias, A; Molina, JC., Pereira, Y. Eco-territorial adaptability of tomato crops for sustainable agricultural production in Carabobo, Venezuela. Idesia, 2020, 38\(2\):95-102. http://dx.doi.org/10.4067/S0718-34292020000200095](#)
- [Hernández, R. & Olivares, B., Application of multivariate techniques in the agricultural land's aptitude in Carabobo, Venezuela. Trop. Subtrop. Agroecosystems, 2020, 23\(2\):1-12. https://n9.cl/zeedh](#)
- Juan, Anciso (2002). *Onion world*, 18(3):10.
- Kurbett A (2015). Studies on elite genotype of chilli against pest complex and their management. Master of sciences (Hort.). Thesis, University of Horticultural sciences, Bagalkot, India.
- Kumar, U. and Singh, S.K. (2011). Evaluation of some chemical, botanical and bio-pesticides against onion thrips under north Bihar condition. *Bihar journal of horticulture*. 1:34-35.
- Kritzman, Gera, A., Racaah, B., VanLent, J.W.M and Peters, D. (2002). The route of tomato spotted wilt virus inside the thrips body in relation to transmission efficiency. *Arch. Virol.*, 147:2143-2156.
- Kendall, D.M. and Capinera .JL. (1987). Susceptibility of onion growth stages to onion thrips (Thysanoptera: Thripidae) damage and mechanical defoliation. *Environmental Entomology Journal* 16:859-863.
- Kritzman, A., M. Lampel, B. Raccach and A Gera. (2001) Distribution and transmission of Iris yellow spot virus. *Plant Dis.* 85:838-842.
- Lebedev G., Abo-Moch F., Gafni G., Ben-Yakir D., Ghanim M. (2013). High level of resistance to spinosad, emamectin benzoate and carbosulfan in populations of thrips tabaci collected in Isrel. *Pest management science* 69(2):274-277.
- Lewis (1997). Pest thrips in perspective. In: Lewis, T. (ed): *Thrips as crop pest CAB international, Wallingford, UK: 1-14.*
- Lewis, T. (1973). Chemical control pp567-593. In T. Lewis (ed). *Thrips as crop pests. CAB International New York.*

Formatted: Font: Not Italic

- Lindeman, K.(1889) Die schadlichsten insekten des tabak in Bessarabein .Bull.Soc.Imp.Nat.Moscou2:10-77.
- Maniania,N.K.,Sithanantham,S.,Ekesi,S.,AmpongNyarko,K.Baumgartner,J,Lohr,B. and Matoka,C.M. (2003). Afield trial of the entomopathogenous fungus *Metarhizium anisopliae* for control of onion thrips *Thripstabaci*. *Crop Protect*.22:553-559.
- Murai, T.(2000).Effect of temperature on development reproduction on onion thrips,Thripstabaci Lindeman (Thysanoptera:Thripidae) ,on pollen and honey solution .Appl.Entomol.Zool.35:499-504.
- Montenegro, E; Pitti, J; Olivares, B. Identificacion of the main subsistence crops of the Teribe: a case study based on multivariate techniques. Idesia, 2021, 39(3): 83-94. <http://dx.doi.org/10.4067/S0718-34292021000300083>
- Morse JG and Hoddle MS (2005) Inavasion biology of thrips. *Annual review of Entomology* 51:67-89.
- Mound ,L.A.(1997).Biological diversity.pp197-215.In T. Lewis (ed).Thrips crop pest .CAB International, New Yark.
- Nault LR (1997) Arthropod transmission plant viruses: new synthetic annals of entomological society of America 90:521-541.
- Olivares, B., Lobo, D., Cortez, A., Rodríguez, M.F. & Rey, J.C. Socio-economic characteristics and methods of agricultural production of indigenous community Kashaama, Anzoategui, Venezuela. Rev. Fac. Agron. (LUZ) 2017, 34(2): 187-215. <https://n9.cl/p2gc5>
- Olivares, B., & Hernández, R. Ecoterritorial sectorization for the sustainable agricultural production of potato (*Solanum tuberosum* L.) in Carabobo, Venezuela. Cienc. Tecnol. Agropecuaria 2019a, 20(2): 339-354. https://doi.org/10.21930/rcta.vol20_num2_art:1462
- Olivares, B., Pitti, J., & Montenegro, E. Socioeconomic characterization of Bocas del Toro in Panama: an application of multivariate techniques. Rev. Bras. Gest. Desenvól. Reg, 2020, 16(3):59-71. <https://doi.org/10.54399/rbgdr.v16i3.5871>
- Olivares, B. Descripción del manejo de suelos en sistemas de producción agrícola del sector Hamaca de Anzoátegui, Venezuela. Lgr, 2016, 23(1): 14–24. <https://n9.cl/ycp08>

Formatted: Font: Not Italic

Formatted: Font: Not Italic, Spanish (Spain, International Sort)

Formatted: Font: Not Italic

Formatted: Font: Not Italic, Portuguese (Brazil)

Formatted: Font: Not Italic, Spanish (Spain, International Sort)

Formatted: Font: Not Italic

- Olivares, B., Hernández, R. Regional analysis of homogeneous precipitation areas in Carabobo, Venezuela. RevistaLasallista de Investigación. 2019b, 16(2):90-105. <https://doi.org/10.22507/rli.v16n2a9>
- Olivares B, Rey JC, Lobo D, Navas-Cortés JA, Gómez JA, Landa BB. Fusarium Wilt of Bananas: A Review of Agro-Environmental Factors in the Venezuelan Production System Affecting Its Development. Agronomy, 2021. 11(5):986. <https://doi.org/10.3390/agronomy11050986>
- Pandey Sujay, Singh BK and Gupta RP.(2013).Effect of neem based botanicals, chemical and bio-pesticides for the management of thrips in onion. Indian journal of Agriculture Research 47(6):545-548.
- Parra, R., Olivares, B., Cortez, A. y Rodríguez, M.F. Pluviometric homogeneity patterns in climatic stations of Anzoátegui state, Venezuela. RevistaMulticiencias. 2012, 12 (Extraordinario): 11-17. <https://n9.cl/xbslq>
- Pathak MK, M.K. Pandey, R.C. Gupta and Gupta P.K.(2018).Evaluation of different insecticides against onion thrips in onion seed Production .Int.J.Curr.Microbiol.App.Sci7(07):4204-4207.
- Pitti, J; Olivares, B., & Montenegro, E. The role of agriculture in the Changuinola District: a case of applied economics in Panama. Trop. Subtrop. Agroecosystems 2021, 25 - 1, pp. 1 - 11. <https://n9.cl/quyl2>
- Pozzer, L., T. Nagata, M.I. Lima, E.W. Kitajima, R. de O. Resende, and A.C. de Avila .(1994). Sapeca, an onion disease in the Sub medio Sao Francisco region, Brazil, is caused by tospovirus with a serologically distinct nucleocapsid protein .Fitopatol. Bras. 19-32
- Rueda, A.F.R. Badenes-Perez and A.M. Shelton (2007) Development economic thresholds for onion thrips in Honduras. Crop Prot. 26: 1099-1107.
- Rodríguez-Yzquierdo, G.; Olivares, B.O.; Silva-Escobar, O.; González-Ulloa, A.; Soto-Suarez, M.; Betancourt-Vásquez, M. (2023b). Mapping of the Susceptibility of Colombian Musaceae Lands to a Deadly Disease: Fusarium oxysporum f. sp. cubense Tropical Race 4. Horticulturae 9, 757. <https://doi.org/10.3390/horticulturae9070757>.
- Shitole D.M., Shankar G., Mithyantha M.S. (2002). Evaluation of certain new insecticides against onion thrips .Pestology 26:49-50.
- Shiberu T., Negeri M. (2014). Evaluation of insecticides and botanical against onion thrips thripstabaci (L). Entomology and Applied science 1(2):26-30.

Formatted: Font: Italic

Formatted: Font: Not Italic, Spanish (Spain, International Sort)

Formatted: Left, Space After: 10 pt, Tab stops: Not at 4.31"

Formatted: Font: Not Italic

Formatted: Font: (Default) Times New Roman, 14 pt

- Srinivas, P.S. and Lawande, K.E.(2004). Impact of planting dates on thrips tabaci Lindeman infestation and yield loss in onion (*Allium cepa* L.).*Pest Manag.HorticEcosys*.10:11-18.
- Tirkey S.Kumar A. (2017).Effect of selected insecticides against chilli thrips *Scirtothrips dorsalis*(Hood) on chilli (*Capsicum annum* L.).*Allahabad Journal Pharmacol* .2:41-42.
- WaiganjoM.M.,Mueke J.M.,Gitonega L.M.,(2008).Susceptible onion growth stages for selective and economic protection from onion thrips infestation. *Acta Horticulture* 767:193-200.
- Vega A, Olivares B, Calderón MAR, Rey JC, Lobo D, Gómez JA, Landa BB. Identification of Soil Properties Associated with the Incidence of Banana Wilt Using Supervised Methods. *Plants*, 2022. 11(15):2070. <https://doi.org/10.3390/plants11152070>
- Viloria, J.A.; Olivares, B.O.; García, P.; Paredes-Trejo, F.; Rosales, A. Mapping Projected Variations of Temperature and Precipitation Due to Climate Change in Venezuela. *Hydrology*, 2023, 10, 96. <https://doi.org/10.3390/hydrology10040096>
- Zingaretti, M.L., Olivares, B., Demey Zambrano, J.A., Demey, J.R. Typification of agricultural production systems and the perception of climate variability in Anzoátegui, Venezuela. *Revista FAVE-CienciasAgrarias* 2016. 15 (2): 39-50. <https://doi.org/10.14409/fa.v15i2.6587>
- Zingaretti, ML., Olivares, B. Analysis of the meteorological drought in four agricultural locations of Venezuela by the combination of multivariate methods. *UNED Research Journal* 2018. 10 (1):181-192. <http://dx.doi.org/10.22458/urj.v10i1.2026>

Formatted: Spanish (Spain, International Sort)