

Survey and monitoring of onion thrips(*Thripstabaci*) Lindeman

Abstract

Thrips is a most important global pest of onion crop they are direct impact of onion yield and quality. The nymph and adult of thrips feed on leaves by piercing and rasping the leaf tissues and causes lengthwise, silvery stippling or blotching on the onion leaves, resulting in the loss of chlorophyll and reduced photosynthetic efficiency. Thrips attack onion at all the stages of crop growth stages but their count increases from bulb initiation and remains high up to bulb development and maturity. Onion thrips can also transmit several plant pathogens in onion crop that reduced onion bulb size and quality. The above surveys objective to evaluate the damage percentage of thrips in onion crops. Survey was conducted during Kharif 2022 and Rabi 2022-23 .The maximum thrips damage rating scale of 3(45%) was recorded at 75 DAT in the Kharif season and Rabi season at 75 DAT thrips damage rating scale was recorded at 4 (65%).The survey result show that the thrips population and damage percentage increased due to increased temperature and relative humidity.

Keywords:Survey, Onion thrips, Thripstabaci, Integrated pest management, Crop damage assessment

Introduction

Onion is important commercial crop of India it is mainly used for cuisine and culinary purpose. Onion is cash crop and export oriented most earning of farmers and valuable foreign exchange for the country. Onion use for different medicinal like preparation of Homeopathic, Unani and Ayurvedic medicines. Gupta *et. al.*, 1984 suggested that thrips (*Thripstabaci* Lindeman) is an important pest of onion crop and losses as high as 90% in quality and yield. Srinivas and Lawande (2004) reported that due to crop damage by thrips yield loss recorded in the range of 46-87% in onion crop. According to Waiganjo *et. al.*, (2008) reported that the leaf damage of onion crops by thrips around 40-60% and yield losses 10-20% in the crop. Shibru and Negeri (2014) suggested that due to onion thrips crop damage and yield was reduced 23-85%. Anonymous. 2000, Jaun, 2002 and Kritzman *et. al.*, 2002. reported that onion thrips are a major vector for several plant viruses such as tomato spotted wilt virus failure to control this pest by timely and show marketable yield losses in crops. Maniania *et. al.*, 2003 and Lebedev *et. al.*, 2013 suggested that Insecticides play important role for controlling of thrips population but again and again use of same group insecticides develop resistance power to thrips against insecticides further Shitole *et.*

al.,2002, Alston& Drost 2008 recorded that the farmers are uses some insecticides and repeated again and again develop of resistance to insecticides. Pathak *et. al.*, (2018) reported that the highest seed yield and minimum thrips population were recorded use of fipronil. Tirkey ,Kumar(2017) and Kurbett*et. al.*,(2015) reported that the thiamethoxam were effective for control thrips population in onion. Asgar *et. al.*,(2018)suggested that the insecticides reduced the thrips population compared to the control and the highest yield was recorded by the use of Dimathoate insecticide. Kumar and Singh (2011) and Daset.*al.*, (2017) reported that the spray of Imidacloprid at 15 days interval was recorded the lowest thrips population and highest gross yield. According to Pandey *et al.*, (2013) recorded that the use of fipronil 1.0 ml/L for better controlling of thrips population and increased the yields of onion crop.

Origin of onion thrips

Onion thrips reported first time in 1889 by Russian Entomologist *Karl Eduard Lindeman* based on specimens collected in Bessarabia, Russia, that caused severe damage to tobacco plants

Geographic distribution of onion thrips

Thrips tabaci are a global pest of onion grown between sea level and 2000 m (Lewis 1973) According to Mound 1977 thrips is a native of the Mediterranean region but has become a major pest of agricultural crops throughout most of the world

Biology of onion thrips

Onion thrips lifecycle deepened environmental conditions, like temperature, humidity and nutrient quality of their food sources .The Stages in the developmental life cycle are the egg, first larval stage, second larval stage pre-pupa, pupa, and adult. Because of their small size, this pest species like other thrips can readily be identified as a species even with a hand lens. Morse *et. al.*, 2005 suggested that the adult specimens are usually needed to make species identification under high microscope magnification. Brain A.N.2006 recorded that the biology of onion thrips was as follows the entire life cycle (egg to adult) required about 19 days large populations can develop quickly under dry weather conditions where there are many overlapping generations throughout year. Females have a, saw -like structure that help to make an incision in plant tissue for egg laying. Eggs are placed singly just under the epidermis of succulent leaf, flower, stem, or bulb tissue. Eggs are elliptical, approximately 0.2 mm in length. They are whitish at deposition and change to an orange tint as development conditions. Hatching occurs in 4-5 days. Larvae are whitish to yellowish. There are two larval stages and besides the adults, they are the only damaging stages. Larval development is completed in about 9-10 days.

Thrips tabaci are two non-feeding stages called the pre-pupa and pupa does not feed and occurs primarily in the soil. Combined pre-pupal and pupal development is completed in 4-6 days. Thrips adults about 1 mm in length their body color ranges from pale yellow to dark brown, wings are unbanded and dirty gray. The males are wingless and exceedingly

rare while the female have long, narrow fringed wings. Onion thrips Female live for 12-30 days and lay 50-60 kidney- shaped eggs singly inside leaf tissue with a sharpe ovipositor.

Economic importance

Thrips is a serious pest a wide range of fruit, vegetable, flower and Agronomics crops. Thrips are members of the order Thysanoptera, which contains several genera and species. Among species of thrips that attack onion are onion thrips (*Thripstabaci*). According to (Nault LR 1997) onion thrips incidence is a major problem due to the damage caused by feeding on vegetative parts of crops which caused discoloration, deformities and reduced marketable yields of the crops.

Damage symptoms

According to Bailey 1938 the onion thrips feeding in onion leaves cause silvery leaf spots that turn into white blotches along the leaves due to removal of cellular content followed by the development of silvery patches and curling of leaves. Lewis 1997b suggested that the thripstabaci causes significant yield loss despite decades of research on control strategies worldwide. *T. tabaci* feeding can reduce onion bulb weight (Kendall and Capinera 1987, Fournier *et.al.*, 1995, Rueda *et al.*, 2007, Diaz-Monato *et. al.*, 2010, Waiganjo *et. al.*, 2008 reported that yield loss up to 60% .According to Pozzer *et. al.*, 1994, Kritzman *et. al.*, that the addition injury by feeding of *T. tabaci* transmits IYSV and is the only confirmed vector of this pathogen. IYSV was first identified on onion in southern Brazil in 1981 (Pozzer *et. al.*, 1994) and was confirmed in the United states in 1989 and worldwide (Gent *et. al.*, 2006). (Gent *et.al.*, 2004) reported that IYSV symptoms on leaves appear as lesions (i.e., straw color to white, dry, and sometimes elongate along the edges indicated that IYSV infection can reduced bulb size. According to (Diaz-Montano *et. al.*, 2010) the IYSV infects onion crops early transplanted and onion yield reduce.

Weather factors on outbreak of onion Thrips

Hamdy, M.K. *et. al.*, 1994 reported thrips population depending to relatively high temperatures and lack of rainfall have been increase in onion thrips population, while high relative humidity and rainfall reduce thrips population. Murai T. 2000 suggested that the thrips population and activity depended of temperature and relative humidity further influence the intrinsic rate of natural increase of the thrips. Hamdy, M.K. and Salem M. *et al.*, 1994 reported that the rate of development of thripstabaci is positively affected by increased temperature and decreased by increased relative humidity. According to Bailey 1934 and Rueda *et al.*, 2007 the thrips intensity increased due to climatic condition hot, dry weather and Lewis 1973 suggested the damage severity of thrips injury increased due to water stress. Environmental factors play important role for increased or decreased of thrips population in onion crops.

Material and methods

The survey of onion thrips was conducted during *Kharif* 2022 and Rabi 2022-23 in Nashik district taluka Niphad, Maharashtra. During the survey five villages were selected for survey in and five onion fields in each village for observation of thrips damage scale. The survey was conducted twice a season at 45 days after transplanting and 75 days after transplanting. The thrips damage rating scales at least 10 plants randomly in each field should be scored using rating scale given below.

Thrips damage Rating

List: 1

Scale	Foliage damage rating
1	1-20%
2	21-40%
3	41-60%
4	61-80%
5	81-100%

Results & Discussions

Kharif 2022

The twenty five different onion fields were surveyed in 5 villages during September and October 2022. The survey for thrips infestation in onion crop was done in all the 25 fields in month of September on 29.09.2022 at different Villages like – Chitegaon, Chandori, Saikheda, Chatori and Berewadi of Taluka Niphad District Nashik, Maharashtra. The thrips damage rating scale ranged from 1 to 2 scale were recorded during survey. Based on the survey onion thrips infested fields were recorded 25% at 45 DAT in Saikheda village farmers fields. The mostly farmers transplanting was done of kharif onion in 15th August. During the survey period crop is under vegetative stage.

At 75 DAT further the survey of thrips infestation was done in same villages and same 25 onion fields at Taluka Niphad District Nashik in month of October on 29.10.2022. The thrips damage rating ranged from 1 to 3 scale were recorded during survey. Based on the survey onion thrips infested fields were recorded 45% in Saikheda village farmers fields. During the survey period crop is under bulb maturity stage.

Table 1- Thrips damage rating in villages at 45 DAT

Sr.no	Villages	Damage rating
1	Chitegaon	1
2	Chandori	1
3	Saikheda	2
4	Chatori	1

5	Berewadi	1
---	----------	---

Table 2- Thrips damage rating in villages at 75 DAT

Sr.no	Villages	Damage rating
1	Chitegaon	1
2	Chandori	2
3	Saikheda	3
4	Chatori	2
5	Berewadi	1

The result show that overall highest thrips damage rating was recorded 2 (25% damage) at 45 DAT in farmers village Saikheda Taluka Niphad. Further the same places survey result show that at 75 DAT the overall highest thrips damage rating was recorded 3 (45% damage) at same village Saikheda Taluka Niphad due to higher temperature of this week. During this period temperature range from 20.24 to 32.42 °C and relative humidity range from 77.90 to 86.87% in village Saikheda.

Rabi 2022-23

The survey of onion thrips Nashik district taluka Niphadwas conducted during *Rabi 2022* on 45 DAT and 75 DAT. During the survey five villages were selected for survey in each taluka and five onion fields in each village for observation of thripsdamage scale .The twenty five different onion fields were surveyed in 5 villages during January and February 2023.

The survey for thrips infestation in onion crop was done in all the 25 fields in month of January at different Villages likes –Chitegaon,Chandori,Saikheda ,Chatori and Berewadi of Taluka Niphad District Nashik,Maharashtra.Thethrips damage rating scale ranged 1-2 scale were recorded during survey. Based on the survey onion thrips infested fields were recorded 30% at 45DAT.

At 75 DAT further the survey of thrips infestation was done in same villages and same 25 onion fields at Taluka Niphad District Nashik .The thrips damage rating ranged from 1-4 scale were recorded during survey. Based on the survey onion thrips infested fields were recorded 65%.

Table 1- Thrips damage rating in villages at 45 DAT

Sr.no	Villages	Damage rating
1	Chitegaon	1
2	Chandori	1
3	Saikheda	2
4	Chatori	1

5	Berewadi	2
---	----------	---

Table 2- Thrips damage rating in villages at 75 DAT

Sr.no	Villages	Damage rating
1	Chitegaon	2
2	Chandori	2
3	Saikheda	4
4	Chatori	2
5	Berewadi	3

The result show that overall highest thrips damage rating was recorded 2 (30% damage) at 45 DAT in village Saikheda and Berewadi and minimum thrips damage scale 1 (20%) were recorded in farmers field Chitegaon village Taluka Niphad. Further the same places survey result show that at 75 DAT the overall highest thrips damage rating was recorded 4 (65% damage) at village Saikheda Taluka Niphad due to higher temperature of this week and minimum thrips damage scale 2 recorded in farmersfields villages Chitegaon, Chandori and Chatori. During this period temperature range from 12.05 to 26.28 °C and relative humidity range from 62.26 to 72.83% in village Saikheda.

The survey and monitoring study of onion thrips (*Thrips tabaci*) in Maharashtra reveals that environmental conditions, particularly temperature and relative humidity, play a crucial role in pest population dynamics and the extent of crop damage. Thrips population and damage levels were observed to be highest during the bulb maturity stage, with a strong positive correlation with temperature increases. These findings align with studies in Colombia and Venezuela, where the incidence of pests like thrips in onions, tomatoes, and other vegetables is exacerbated by warm, dry conditions (Campos, 2023; Campos et al. 2023). For instance, thrips thrive in Colombia's warmer regions with prolonged dry spells, indicating a similar response to environmental conditions across different geographic locations (Zingaretti and Olivares, 2018; Rodriguez et al. 2023). Likewise, in Venezuela, studies have documented that high temperatures coupled with low rainfall contribute to thrips outbreaks, highlighting the pest's dependency on climatic factors common to the region.

The influence of humidity on thrips populations in the Maharashtra study also parallels observations in Latin America. Thrips populations in Venezuelan onion fields are notably lower in regions with high rainfall and humidity, as humidity disrupts thrips' ability to feed and reproduce effectively (Viloria et al. 2023). In Colombia, thrips management strategies for onions and other bulb crops often rely on irrigation schedules that enhance moisture levels during critical pest development phases, which has shown to reduce pest

prevalence). This tactic highlights the importance of integrated pest management (IPM) strategies that incorporate environmental control measures to manage pest pressure (Olivares, 2016; Olivares et al. 2021). Both studies demonstrate the need to adapt pest management practices to local climate conditions, emphasizing that temperature and humidity are paramount to thrips population dynamics across these different agricultural systems (Vega et al. 2022; Olivares and Hernandez, 2019a).

Agronomic practices also influence pest dynamics, as evidenced by the current study's recommendations for intercropping, bio-pesticides, and selective insecticides. Colombian and Venezuelan farmers commonly employ intercropping with crops like carrots and tomatoes to manage pest populations, as intercropping reduces pest prevalence by altering microclimates within crop fields (Zingaretti et al. 2016; Vilorio et al. 2023). Additionally, the use of organic solutions such as neem oil and botanical-based pesticides, like in the Maharashtra study, is increasingly popular in Latin America, where small-scale farmers seek cost-effective and ecologically sustainable pest control measures. This approach reduces thrips populations by creating less favorable environments for pests while enhancing plant resilience against pest attacks (Olivares et al. 2017; Olivares et al. 2020). The cumulative findings from India, Colombia, and Venezuela highlight that pest management in bulb crops like onions benefits greatly from holistic approaches that integrate environmental (Hernandez et al. 2020), agronomic (Hernandez and Olivares, 2020) and chemical controls (Hernandez et al. 2018). Future research should focus on developing pest-resistant onion varieties suited to warm, humid environments. Studies could further assess the efficacy of entomopathogenic fungi and natural predators in both regions to offer alternative pest control solutions, reducing reliance on chemical insecticides and mitigating resistance issues (Montenegro et al. 2021). This comparison underscores the interconnectedness of environmental conditions and agronomic practices in shaping effective, sustainable pest management strategies across diverse agricultural landscapes.

Conclusion

The present study concluded that the highest thrips damage rating 2 recorded in kharif season at 45 DAT and rating 3 were recorded at 75 DAT same village Saikheda further highest damage rating 2 in Rabi season at 45 DAT and 4 rating was recorded in Saikheda village at 75 DAT. Above survey show that the thrips population and damage rating increased due to increased temperature and relative humidity.

Disclaimer (Artificial intelligence)

Authors hereby declare that NO generative AI technologies such as large language models and text to image generators have been used during the writing or editing of this manuscript.

Acknowledgement

The authors are thankful to the farmer's taluka Niphad of district Nashik, Maharashtra for helping to us during the survey and provide the details of crops.

Authors, contributions

This work's collaboration between both authors read and approved the final manuscript

References

- Anonymous (2000). Annual Report: National Research Centre for onion and Garlic. pp.63-64.
- Alston D.G., Drost D. (2008). Utah pests fact sheet. Onionthrips (*Thripstabaci*). Internet <http://utahpests.edu/IPM/files/uplod/pdfdocs/factsheet-pdf/ENT-117>
- Asghar Muhammad, Mirza Muhammad Qaddeer Baig, Muhammad Afjal, Naeem Faisal (2018) Evaluation of different insecticides for the management of onion thrips (*Thripstabaci* Lindeman) on onion crop. *Polish journal of entomology* Vol.87:165-176.
- Brain AN (2006) Biology and Ecology of onion thrips on onion fields *PP2. Cornell University*.
- Bailey, S.F. (1938) Thrips economic importance in California, *Univ. Calif. Coll. Agric. Agric. Exp. Stn. Cir. 346*.
- Bertorelli, M., and Olivares, B.O. Population fluctuation of *Spodoptera frugiperda* (J.E. Smith) (Lepidoptera: Noctuidae) in sorghum cultivation in Southern Anzoategui, Venezuela. *J. Agric. Univ. P. R.*, 2020, 104(1):1-16. <https://doi.org/10.46429/jaupr.v104i1.18283>

- Campos, B. O. (2023). Banana Production in Venezuela: Novel Solutions to Productivity and Plant Health. Springer Nature. <https://doi.org/10.1007/978-3-031-34475-6>
- Campos, B. O. O., Araya-Alman, M., & Marys, E. E. (2023). Sustainable Crop Plants Protection: Implications for Pest and Disease Control (p. 200). MDPI-Multidisciplinary Digital Publishing Institute. <https://doi.org/10.3390/books978-3-0365-9150-6>
- Das Ajay Kumar, Wajid Hasan and Sushil Kumar Singh (2017). Management of onion thrips, *Thrips tabaci* Using Chemical and Bio-pesticides for Quality onion production. *Trends in Biosciences* 10(22)4384-4388.
- Diaz-Montano J, Fuchs M, Nault BA, Fail J, Shelton AM (2010). Evaluation of onion cultivars for resistance to onion thrips (Thysanoptera: Thripidae) and Iris yellow spot virus. *J. Econ. Entomol.* 103, 925-937.
- Fournier F., Guy B., Robin S. (1995). Effect of thrips *tabaci* (Thysanoptera: Thripidae) on yellow onion yields and economic thresholds for its management. *Entomological Society of America*. 88(5):1401-1407.
- Gupta, R.P., Srivastva, V.K., Bhardwaj, B.S. and Pandey U.B. (1984). Chemical control of *Thrips tabaci* L. infesting onion crop. *J. Ent. Res.* 8(2)196-198.
- Gent, D.H., HR. Schwartz, and R. Khosla. (2004). Distribution and incidence of Iris yellow spot virus in Colorado and its relation to onion plant population and yield. *Plant Dis.* 88:446-452.
- Gent, D.H., L.J. du Toit, S.F., Fichtner, S.K., Mohan, H.R., Pappu and H.F. Schwartz (2006) Iris yellow spot virus an emerging threat to onion bulb and seed production. *Plant Dis.* 90:1468-1480.
- Hamdy MK and Salem M (1994) .The effect of plantation dates of onion, temperature and relative humidity on the population density of the onion thrips. *Thrips tabaci* Lind in Egypt. *Ann Agric Sci Univ Ain Shams (Egypt)* 39:417-424.
- Hernández, R. & Olivares, B., Arias, A; Molina, J.C., & Pereira, Y. Agroclimatic zoning of corn cultivation for the sustainability of agricultural production in Carabobo, Venezuela *Rev. Univ. de Geo.* 201827(2)139-159 <https://n9.cl/ioupn>.
- Hernandez, R.; Olivares, B.; Arias, A; Molina, J.C., Pereira, Y. Eco-territorial adaptability of tomato crops for sustainable agricultural production in Carabobo, Venezuela. *Idesia*, 2020, 38(2):95-102. <http://dx.doi.org/10.4067/S0718-34292020000200095>.

- Hernández, R. & Olivares, B., Application of multivariate techniques in the agricultural land's aptitude in Carabobo, Venezuela. *Trop. Subtrop. Agroecosystems*, 2020, 23(2):1-12. <https://n9.cl/zeedh>.
- Juan,Anciso(2002).*Onion world,18(3):10*.
- Kurbett A(2015). Studies on elite genotype of chilli against pest complex and their management.Master of sciences (Hort.).*Thesis,University of Horticultural sciences ,Bagalkot,India*.
- Kumar,U.andSingh,S.K.(2011).Evaluation of some chemical, botanical and bio-pesticides against onion thrips under north Bihar condition. *Bihar journal of horticulture*.1:34-35.
- Kritzman,Gera,A.,Racaah,B.,VanLent,J.W.M and Peters,D.(2002).The route of tomato spotted wilt virus inside the thrips body in relation to transmission efficiency. *Arch.Virol.,147:2143-2156*.
- Kendall,D.M. and Capinera .JL.(1987).Susceptibility of onion growth stages to onion thrips(Thysanoptera:Thripidae) damage and machanical defoliation. *Environmental Entomology Journal 16:859-863*.
- Kritzman,A.,M. Lampel,B.Racah and A Gera.(2001) Distribution and transmission of Iris yellow spot virus. *Plant Dis*.85:838-842.
- Lebedev G.,Abo-Moch F.,Gafni G.,Ben-Yakir D.,Ghanim M.(2013).High level of resistance to spinosad, emamectin benzoate and carbosulfan in populations of thripstabaci collected in Isrel. *Pest management science* 69(2):274-277.
- Lewis (1997).Pest thrips in perspective-In: Lewis, T. (ed): Thrips as crop pest *CAB international, Wallingford, UK: 1-14*.
- Lewis,T.(1973).Chemical control pp567-593.In T.Lewis(ed).Thrips as crop pests.*CAB International New Yark*.
- Lindeman, K.(1889) Die schadlichsteninsekten des tabak in Bessarabein .*Bull.Soc.Imp.Nat.Moscou2:10-77*.
- Maniania,N.K.,Sithanantham,S.,Ekesi,S.,AmpongNyarko,K.Baumgartner,J,Lohr,B. and Matoka,C.M. (2003). Afield trial of the entomopathogenous fungus *Metarhiziumanisopliae* for control of onion thrips*Thripstabaci*. *Crop Protect*.22:553-559.
- Murai, T.(2000).Effect of temperature on development reproduction on onion thrips,Thripstabaci Lindeman (Thysanoptera:Thripidae) ,on pollen and honey solution .*Appl.Entomol.Zool*.35:499-504.

- Montenegro, E; Pitti, J; Olivares, B. Identification of the main subsistence crops of the Teribe: a case study based on multivariate techniques. *Idesia*, 2021, 39(3): 83-94. <http://dx.doi.org/10.4067/S0718-34292021000300083>.
- Morse JG and Hoddle MS (2005) Invasion biology of thrips. *Annual review of Entomology* 51:67-89.
- Mound ,L.A.(1997).Biological diversity.pp197-215.In T. Lewis (ed).Thrips crop pest .*CAB International, New York*.
- Nault LR (1997) Arthropod transmission plant viruses: *new synthetic annals of entomological society of America* 90:521-541.
- Olivares, B., Lobo, D., Cortez, A., Rodríguez, M.F. & Rey, J.C. Socio-economic characteristics and methods of agricultural production of indigenous community Kashaama, Anzoátegui, Venezuela. *Rev. Fac. Agron. (LUZ)* 2017, 34(2): 187-215.<https://n9.cl/p2gc5>
- Olivares, B., & Hernández, R. Ecoterritorialsectorization for the sustainable agricultural production of potato (*Solanum tuberosum* L.) in Carabobo, Venezuela. *Cienc. Tecnol. Agropecuaria* 2019a, 20(2): 339-354. https://doi.org/10.21930/rcta.vol20_num2_art:1462
- Olivares, B., Pitti, J., & Montenegro, E. Socioeconomic characterization of Bocas del Toro in Panama: an application of multivariate techniques. *Rev. Bras. Gest.Desenvolv. Reg*, 2020, 16(3):59-71. <https://doi.org/10.54399/rbgdr.v16i3.5871>
- Olivares, B. Descripción del manejo de suelos sistemas de producción agrícola del sector Hamaca de Anzoátegui, Venezuela. *Lgr*, 2016, 23(1): 14–24. <https://n9.cl/yyp08>
- Olivares, B., Hernández, R. Regional analysis of homogeneous precipitation areas in Carabobo, Venezuela. *Revista Lasallista de Investigación*. 2019b, 16(2):90-105.<https://doi.org/10.22507/rli.v16n2a9>.
- Olivares B, Rey JC, Lobo D, Navas-Cortés JA, Gómez JA, Landa BB. Fusarium Wilt of Bananas: A Review of Agro-Environmental Factors in the Venezuelan Production System Affecting Its Development. *Agronomy*, 2021. 11(5):986.<https://doi.org/10.3390/agronomy11050986>
- Pandey Sujay, Singh BK and Gupta RP.(2013).Effect of neem based botanicals, chemical and bio-pesticides for the management of thrips in onion. *Indian journal of Agriculture Research* 47(6):545-548.
- Parra, R., Olivares, B., Cortez, A. y Rodríguez, M.F. Pluviometric homogeneity patterns in climatic stations of Anzoátegui state, Venezuela. *Revista Multiciencias*. 2012, 12 (Extraordinario): 11-17. <https://n9.cl/xbslq>

- Pathak MK, M.K. Pandey, R.C. Gupta and Gupta P.K. (2018). Evaluation of different insecticides against onion thrips in onion seed Production. *Int. J. Curr. Microbiol. App. Sci* 7(07):4204-4207.
- Pitti, J; Olivares, B., & Montenegro, E. The role of agriculture in the Changuinola District: a case of applied economics in Panama. *Trop. Subtrop. Agroecosystems* 2021, 25 - 1, pp. 1 - 11. <https://n9.cl/quyl2>
- Pozzer, L., T. Nagata, M. I. Lima, E. W. Kitajima, R. de O. Resende, and A. C. de Avila. (1994). Sapeca, an onion disease in the Sub medio Sao Francisco region, Brazil, is caused by tospovirus with a serologically distinct nucleocapsid protein. *Fitopatol. Bras.* 19-32
- Rueda, A, F.R. Badenes-Perez and A.M. Shelton (2007) Development economic thresholds for onion thrips in Honduras. *Crop Prot.* 26:1099-1107.
- Rodríguez-Yzquierdo, G.; Olivares, B.O.; Silva-Escobar, O.; González-Ulloa, A.; Soto-Suarez, M.; Betancourt-Vásquez, M. (2023b). Mapping of the Susceptibility of Colombian Musaceae Lands to a Deadly Disease: *Fusarium oxysporum* f. sp. cubense Tropical Race 4. *Horticulturae* 9, 757. <https://doi.org/10.3390/horticulturae9070757>
- Shitole D.M., Shankar G., Mithyantha M.S. (2002). Evaluation of certain new insecticides against onion thrips. *Pestology* 26:49-50.
- Shiberu T., Negeri M. (2014). Evaluation of insecticides and botanical against onion thrips *thrips tabaci* (L.). *Entomology and Applied science* 1(2):26-30.
- Srinivas, P.S. and Lawande, K.E. (2004). Impact of planting dates on thrips *tabaci* Lindeman infestation and yield loss in onion (*Allium cepa* L.). *Pest Manag. Hort. Ecosys.* 10:11-18.
- Tirkey S. Kumar A. (2017). Effect of selected insecticides against chilli thrips *Scirtothrips dorsalis* (Hood) on chilli (*Capsicum annum* L.). *Allahabad Journal Pharmacol.* 2:41-42.
- Waiganjo M.M., Mueke J.M., Gitonega L.M., (2008). Susceptible onion growth stages for selective and economic protection from onion thrips infestation. *Acta Horticulture* 767:193-200.
- Vega A, Olivares B, Calderón MAR, Rey JC, Lobo D, Gómez JA, Landa BB. Identification of Soil Properties Associated with the Incidence of Banana Wilt Using Supervised Methods. *Plants*, 2022. 11(15):2070. <https://doi.org/10.3390/plants11152070>

- Viloría, J.A.; Olivares, B.O.; García, P.; Paredes-Trejo, F.; Rosales, A. Mapping Projected Variations of Temperature and Precipitation Due to Climate Change in Venezuela. *Hydrology*, 2023, 10, 96. <https://doi.org/10.3390/hydrology10040096>
- Zingaretti, M.L., Olivares, B., Demey Zambrano, J.A., Demey, J.R. Typification of agricultural production systems and the perception of climate variability in Anzoátegui, Venezuela. *Revista FAVE-Ciencias Agrarias* 2016. 15 (2): 39-50. <https://doi.org/10.14409/fa.v15i2.6587>.
- Zingaretti, ML., Olivares, B. Analysis of the meteorological drought in four agricultural locations of Venezuela by the combination of multivariate methods. *UNED Research Journal* 2018. 10 (1):181-192. <http://dx.doi.org/10.22458/urj.v10i1.2026>
- MK Pathak, PK Gupta, S Pandey, MK Pandey, S Purshottaman and RC Gupta. A review of onion thrips (*Thrips tabaci* Lindeman) management options and importance. *The Pharma Innovation Journal* 2023; 12(4): 1271-1277