

## Original Research Article

### Diversity of family Scytonemataceae ([Cyanobacteria](#)) from the Wayanad district, Western Ghats regions of Kerala, India.

#### ABSTRACT

The Scytonemataceae is the only traditional cyanobacterial family that exhibits the false branching phenotype and includes ecologically significant species, making it particularly interesting from an ecologic as well as economic perspective. This family is characterized by its taxonomic diversity and widespread distribution. The current study focuses on the diversity of the Scytonemataceae family in the Wayanad district, located in the Western Ghats of Kerala. The Western Ghats is notable for its exceptionally high level of biological diversity and endemism. Additionally, this area is ecologically sensitive and serves as a site for the systematic elucidation of many species. The research found that Scytonemataceae are commonly found on rock surfaces in the Wayanad district. These species display a color range that varies from pale or olive green to light brownish-black. The species were characterized and identified based on various morphological features, including cell dimensions, color, shape, type of branching, sheath characteristics, and cell contents. A total of 19 species were recorded, with the dominant genus being Scytonema with 15 species. Brasilonema follows with 3 species, and Petalonema contributes to one species, all belonging to the order Nostocales. Notably, among the 19 species identified, 11 are new records for the cyanobacterial flora of the Western Ghats in India.

**Keywords:** *Cyanobacteria*, *Diversity*, *Scytonemataceae*, *Nostocales*, *Brasilonema*, *Scytonema*, *Petalonema*.

#### INTRODUCTION

The Cyanobacteria are the largest, most diverse, and widely distributed group of photosynthetic prokaryotes. They are recognized as the first microorganisms on Earth capable of producing oxygen through photosynthesis. These organisms possess the unique ability to perform mutually compatible functions such as nitrogen fixation and photosynthesis. They can thrive in various environments, including extreme climatic conditions and in both oxic and anoxic environments (Whitton and Potts, 2000; Thajuddin and Subramanian, 2005). Among them, Scytonemataceae populations are particularly prominent and are commonly found on rock surfaces. Their occurrence, growth pattern, and distribution vary significantly with changing environmental conditions. This diverse group has various applications in biotechnology, pharmacology, and agriculture.

**Commented [WU1]:** I am confused by counting new record in your taxonomic description. New record 11 or 14? How many new for India, and how many for Kerala? Write clearly in abstract and Result also.

**Commented [WU2]:** Do not use title words

**Commented [WU3]:** Citation required

Cyanobacteria have several advantages as hosts for biotechnological applications owing to minimal growth requirements, easy genetic manipulation, and appealing platforms that enable carbon-neutral production processes (Lau et al. 2015; Kumar et al. 2016). The Scytonemataceae family has significant potential applications in agriculture due to its capacity for nitrogen fixation, making it a promising contributor to this field. Specifically, *Scytonema varium* has demonstrated significant antiviral activity against the human immunodeficiency virus (HIV), supporting its potential use in AIDS therapy (Boyd et al. 1997; Loya et al. 1998; Bokesch et al. 2003). Scytonemin, primarily found in *Scytonema javanicum*, is a UV-A protective pigment known for absorbing harmful ultraviolet radiation (UV-C, UV-B, and UV-A), making it a suitable ingredient for sunscreens. Additionally, scytonemin inhibits the proliferation of human fibroblasts, endothelial cells, and tumor cells, indicating its potential as an anti-inflammatory and antiproliferative agent. Furthermore, Scytonema species have demonstrated antibacterial properties against *Pseudomonas striata*, *Bacillus subtilis*, and *Escherichia coli* (Tyagi et al. 2014). *Scytonema bilaspurense* has demonstrated cytotoxic effects against human cancer cell lines (Ngo et al. 2022). Certain Scytonema species, including *S. scytonema foetidum*, are recognized for producing geosmin, the compound responsible for the earthy aroma in soil (Tawong et al. 2022). Petalonema species play a vital role in enhancing soil fertility through nitrogen fixation. These microorganisms can be used as biofertilizers in agricultural practices to improve soil quality and increase crop yields (Abed et al. 2010). Additionally, Brasilonema species are valuable as a source of high-quality protein and essential nutrients. The biomass of Brasilonema species can serve as a raw material in various industrial sectors, including food production (Menescal et al. 2020).

Commented [WU4]: it

Commented [WU5]: Use abbreviation, S.

The genus Scytonema is an important member of the Cyanophyceae family, characterized by false branching and isopolar filaments. These filaments often appear entangled or irregularly coiled and are typically covered with colored or colorless sheaths. The trichomes are mainly olive green, blue-green, brownish, or yellowish (Komarek, 2013). In contrast, the genus Brasilonema features isopolar filaments that are densely packed and exhibit rare instances of false branching (Fiore et al. 2007). Brasilonema is characterized by its macroscopic filaments, heterocysts, and cell division occurring in a single plane. Additionally, Brasilonema can be distinguished from other taxa by its erect, fasciculate thalli and the presence of vacuole-like spaces in the center of the cytoplasm. While both genera share similar morphological traits, the primary distinction lies in the shape of their young trichomes, which are 'J' or 'C' shaped in Brasilonema, unlike those in Scytonema (Fiore et al. 2007; Sant' Anna et al. 2011). However, the diversity and taxonomic classification of these genera

remain poorly understood (Komarek et al. 2013). The genus *Petalonema* is a filamentous cyanobacterium characterized by false branching and the presence of heterocytes. *Petalonema* species primarily thrive in terrestrial environments, commonly found on calcareous wet rocks, stones, and dripping walls. Morphologically, *Petalonema* resembles the genus *Scytonema*, but it has distinct characteristics that set it apart. These include irregular false branching, very broad and thick lamellated sheaths surrounding the trichomes, and sheaths that are several times broader than the trichomes themselves, with funnel-like divergent ends (Komarek et al. 2013).

Referring to the literature on Cyanobacterial taxa concerning Kerala, research on the diversity in the Scytonemataceae family is scarce. Also, the Western Ghats region of Kerala, which contributes to a spectrum of microhabitats, has not been extensively studied for Cyanobacterial diversity, especially for the family Scytonemataceae. Initial investigations on Kerala's Western Ghats produced astounding findings, indicating that the cyanobacteria in these regions are extremely diversified. These microorganisms show great promise, and gaining a comprehensive understanding of the distribution of the Scytonemataceae family in the Western Ghats region requires a detailed ecological study. Hence the present study has been undertaken on the diversity of family Scytonemataceae in the Wayanad district of Kerala, which geographically falls fully in the Western Ghats mountain chain.

Commented [WU6]: Citation required

## MATERIALS AND METHODS:

### Study area

The Wayanad district is located in the northeastern state of Kerala, India. It is situated within the Western Ghats and lies at the southernmost extremity of the Deccan Plateau. The district shares its western border with Kozhikode and is predominantly covered by dense forests typical of the Western Ghats. The Western Ghats are a mountain range that runs parallel to the western coast of India, extending from Gujarat to Tamil Nadu. This range spans the states of Gujarat, Maharashtra, Goa, Karnataka, Kerala, and Tamilnadu, covering an area of approximately 160,000 square kilometers over a distance of 1,600 kilometers. The Western Ghats are renowned for their exceptional biological diversity and endemism, and they are recognized as one of the world's eight "hottest hotspots" of biodiversity. Wayanad is unique as it is the only plateau in Kerala, situated at an altitude between 700 and 2,100 meters above sea level. The district enjoys a pleasant climate throughout the year, with an average annual rainfall of 2,322 millimeters. Areas such as Lakkidi, Vythiri, and Meppadi experience the highest amounts of rainfall, ranging from 3,000 to 4,000 millimeters each year. During March and April, dry winds blow through the region, while high-velocity

Commented [WU7]: Citation required

Commented [WU8]: Citation required

winds are common during the southwest monsoon season. The high-altitude areas can experience extreme cold, and relative humidity can reach up to 95% during the monsoon.

### Cyanobacterial Sampling

Specimens were collected seasonally (pre-monsoon, monsoon, and post-monsoon) from 13 sites of the Western Ghats, specifically falling in the Wayanad district. The coordinates of sampling sites were recorded using an Etrex 20X Garmin GPS. Cyanobacterial patches were collected from various rock surfaces at different sampling sites by scraping them with sterile scalpels and placing them in collection bottles. Microclimatic parameters associated with the habitat, such as atmospheric temperature (Divinext digital Thermometer), surface temperature (Metravi AVM-08 IR thermometer), humidity (Divinext digital hygrometer), and wind speed (Prova AVM -01 anemometer) were recorded. Data concerning diurnal temperature and relative humidity were acquired from Google Earth Pro (2022). Other habit and habitat characteristics of patches were identified and recorded. Each specimen was assigned a unique voucher number for identification purposes.

**Commented [WU9]:** In map, only 9 sites are seen

**Commented [WU10]:** Better if you give brief description of all 13 sites with latitude, longitude, elevation etc in Study area section.

**Commented [WU11]:** Number of total samples?

**Commented [WU12]:** Did you collect all samples from moist rocks only? Not from water bodies?

**Commented [WU13]:** Is it rock surface? Make it clear.


### Culturing and Identification of Family Scytonemataceae

Two sets of cyanobacterial specimens were collected and brought to the laboratory for further study. To remove any debris and impurities, the samples were first rinsed with distilled water. They were then cultured in a BG-11 medium (Rippka et al. 1979), which is a nitrogen-free medium prepared by omitting sodium nitrate. The specimens were maintained at a temperature of  $25 \pm 2^\circ\text{C}$  and illuminated with 3000 lx, following a light/dark cycle of 16:8 hours. The other set of samples was preserved in 4% (v/v) formaldehyde and deposited in the germplasm collection of cyanobacteria at the Division of Environmental Science, Department of Botany, University of Calicut.

For the identification of cyanobacteria, microscopic analysis was conducted on live specimens. The collected samples were examined, and photomicrographs were taken using a Leica DM6B compound microscope. Morphometric analyses were performed based on key morphological features, including cell dimensions, colour, shape, type of branching, sheath characteristics, and cell contents. These features were essential for the identification and classification of cyanobacteria, following the methods outlined by Komarek and Anagnostidis (1998, 2005) and Komarek (2013). Standard taxonomic manuals by Desikachary (1959) and Komarek (2013) were used for identifying cyanobacterial species. Furthermore, the identity

**Commented [WU14]:** How you measure dimensions?

of the taxonomically accepted species was verified as per Algaebase (<https://www.algaebase.org/>) (Guiry and Guiry 2024).



**Commented [WU15]:** Count the sites in this map, can you find 13 sites here? You have written 13 sites in the text.

**Fig.1.**Map showing the study areas

**Commented [WU16]:** Cite this Fig. 1 at appropriate place in the text

## RESULTS AND DISCUSSION

Kerala is one of the least explored areas in the field of cyanobacteria. The Western Ghats region falling in the eastern side of the state is renowned for its rich ecology and biodiversity, especially due to its diverse habitats. The Western Ghats is characterized by high biological diversity and endemism. Also, the area is ecologically sensitive and constitutes sites for the systematic elucidation of many species. Nevertheless, little research has been attempted on cyanobacteria, particularly the family Scytonemataceae in this region. In this perspective, the study has been attempted with the concept that it will provide comprehensive information on cyanobacterial diversity in this region, that too for the first time. The study revealed a total of ~~nineteen-19~~ species of cyanobacteria belonging to the order Nostocales. Among these, the genus *Scytonema*, represented by ~~fifteen-15~~ species, showed the highest dominance.

### Key to the species

***Scytonema drilosiphon*** [Kutzing] Elenkin et Poljanskij 1922

Komarek, 2013, vol.19, pp. 73, fig.33.

**Description:** Thallus woolly, up to 1-3 mm thick, grey or pale greyish blue-green, sometimes slightly up to densely covered by CaCO<sub>3</sub>- precipitates. Filaments intertwined ± in bundles, 7-13µm wide, with sparse geminate or single false branching. Sheaths are firm, thin, not lamellated, colourless, or slightly yellowish. Trichomes are blue-green, cylindrical, at the ends sometimes slightly widened, not constricted at cross-walls, 5.5-10 µm wide. Cells cylindrical, in the middle of trichomes isodiametric, 2.5-12 µm long; terminal cells rounded. Heterocytes elliptical to quadratic, 8-12 (14) × 7-9µm.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Thirunelli, N11°30'41.79"-E076°14'30.21", Environmental Science Division, University of Calicut, CU No:163831. (Fig. 3, B).

**Comments:** New record to the Western Ghats Cyanobacterial flora of Kerala.

***Scytonema bivaginatatum*** Welsh 1965

Komarek, 2013, vol.19, pp. 85, fig.53.

**Description:** Thallus woolly, dark brown. The main filaments are up to 30 µm wide, rarely geminate, and false branches are about 12 µm wide. Sheaths are firm, smooth, and usually hyaline with a thin, about 2-3 µm thick, yellow layer next to the trichome and a wider, colorless outerpart, about 10 µm thick. In young branches, sheaths are only homogeneous and hyaline. Trichomes in main filaments 6-7 µm wide, in branches 3.5-5 µm wide, cylindrical,

**Commented [WU17]:** Move to introduction at appropriate place.

**Commented [WU18]:** Italicise

**Commented [WU19]:** Describe *Brasilonema* first then *Petalonema* and then *Scytonema* at last (in alphabetic order)

**Commented [WU20]:** Add here, how many new record for India, and how many new for Kerala?

**Commented [WU21]:** Essential. Write the key to the species before taxonomic description.

**Commented [WU22]:** Describe it after *Petalonema*. Arrange *Scytonema* species in alphabetical order.

**Commented [WU23]:** It is not necessary to write "INDIA: Kerala; Wayanad district" for each algae. It has been mentioned in above under Cyanobacterial Sampling

not constricted at cross-walls, sometimes with granular content. Cells 3-8×6-7 µm in old parts, 10-15 ×± 3.5µm in branches. Heterocytes intercalary, sub-rectangular, ±12×8.5 µm.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Edakkal, N11°37'18.97"-E076°13'44.92", Environmental Science Division, University of Calicut, CU No: 163818. (Fig. 2, C).

**Comments:** New record to the Western Ghats Cyanobacterial flora of India.

***Scytonema malaviyanense*** Bharadwaja 1930

Desikachary, 1959, pp. 483; Komarek, 2013, vol.19, pp. 94, fig.86.

**Description:** Thallus thick, spongy, bluish-green. Filaments flexuous, interwoven, young 8-10 µm, old 16.8-20.2 µm wide, up to 4mm long. False branches only geminate, 6.5-8.5 µm wide, up to 14-16 µm wide in older parts. Sheaths firm, at first thin, hyaline or slightly yellow, later thick, lamellated and yellowish-brown, brown to dark brown, 1.4-4 µm thick. Trichomes bluish-green, with older trichomes slightly constricted at cross-walls. Cells isodiametric. Heterocytes are absent in young filaments, but present in old ones, intercalary, quadratic, 11.2-15.4 × 9.8-15.4 µm.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Kanthanpara, N11°31'41.35"-E076°09'25.75", Environmental Science Division, University of Calicut, CU No:173138. (Fig.3, A).

**Comments:** New record to the Western Ghats Cyanobacterial flora of Kerala.

***Scytonema masonianum*** Welsh 1963

Komarek, 2013, vol.19, pp. 94-95, fig.67.

**Description:** Thallus floccose, dark green to brownish. Old filaments intricately intertwined, about 20 µm wide, younger parts of filaments and branches about 13 µm wide; filaments false branched, branches infrequent, V-formed or lateral, commonly single, rarely geminate. Sheaths are well developed, straight, firm, parallelly and faintly, but distinctly lamellated or diffuent, especially in older parts, 2-4 µm thick, greenish-yellow towards ends hyaline. Trichomes brownish-yellow, not constricted at cross-walls. Cells 4-12 µm long. Heterocytes are not numerous, intercalary, or rectangular.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Phantom rock, N11°38'09.96"-E076°12'14.27", Environmental Science Division, University of Calicut, CU No: 173142. (Fig. 3, F).

**Comments:** New record to the Western Ghats Cyanobacterial flora of India.

***Scytonema tenellum*** Gardner [1927](#)

Komarek, 2013, vol.19, pp. 101-102, fig.82.

Filaments flexuous or densely intertwined, erect, 12-16µm wide. Branches geminate, branches perpendicular to gently curved. Sheaths are thin, smooth, lamellated. Trichomes cylindrical, 10-14 µm wide, pale olive or blue-green. Heterocytes are numerous, and golden yellow.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Thirunelli, N11°54'49.33"-E075°59'39.29", Environmental Science Division, University of Calicut, CU No: 163832. (Fig.2, I).

**Comments:** New record to the Western Ghats Cyanobacterial flora of India.

***Brasilonema bromeliae*** Fiore et al. 2007

Komarek, 2013, vol.19, pp. 107-108, fig.88.

**Description:** Thallus is irregularly fasciculated. Filaments joined to flat macroscopic, velvet, blackish green to blackish violet mats, densely fasciculate, more or less creeping on the substrate, slightly irregularly coiled, 10-21µm wide, later joined in dense, erect fascicles. Branches rare. Sheaths firm, thin, colorless to yellow-brown. Trichomes are cylindrical, 8-18 µm wide, and not or very slightly constricted at cross-walls. Cells 1.8-16 µm long greyish-green, olive-green, brownish violet, Heterocyst solitary, intercalary, discoid or cylindrical 4-19x15-16.8 µm.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Edakkal, N11°37'20.12"-E076°13'42.71", Environmental Science Division, University of Calicut, CU No:173146. (Fig. 2, E).

**Comments:** New record to the Western Ghats Cyanobacterial flora of India.

***Brasilonema terrestre*** Sant' Anna ~~et al.~~ Komarek 2011

Komarek, 2013, vol.19, pp. 113, fig.94.

**Description:** Thallus macroscopic, composed of very irregularly fasciculated, partly erect filaments. Filaments cylindrical, 12-17µm wide. Sheaths are thin, firm, colourless to yellow-brown. Trichomes are cylindrical, 9-15µm wide, not constricted at cross walls. Cells ± isodiametric, shorter towards the ends, greyish-green or blue-green. Heterocytes are short barrel-shaped to cylindrical, 6-17x13-14 µm.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Edakkal, N11°37'47.52"-E076°13'44.12", Environmental Science Division, University of Calicut, CU No: 173144. (Fig. 3, I)

**Comments:** New record to the Western Ghats Cyanobacterial flora of India.

Commented [WU24]: Describe it first

***Brasilonema epidendron*** Sant'Anna et al. 2011

Komarek, 2013, vol.19, pp. 108, fig.89.

**Description:** Thallus macroscopic, composed from parallelly fasciculated, erect filaments, irregularly arranged, dark green to blackish. Filaments 7-10.9×12 (14) µm wide, very rarely branched. Sheaths are thin, and firm. Trichomes 5.5-8.2-10 (11) µm wide, cylindrical not constricted at cross-walls, not attenuated towards ends. Cells are cylindrical or shortened. Heterocytes barrel-shaped to cylindrical, (7)8-10 (11.5) ×7-9µm.

**Habitat:** Epilithic.

**Comments:** New record to the Western Ghats Cyanobacterial flora of India.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Phantom rock, N11°38'09.96"-E076°12'14.27", Environmental Science Division, University of Calicut, CU No: 163859. (Fig. 2, F).

***Scytonema spirulinoides*** Gardner 1927

Komarek 2013, vol.19, pp. 133, fig.125.

**Description:** Thallus flocculent. Filaments curved and spirally coiled, 15-20 µm wide, usually geminate branching, Sheaths thick, homogeneous, without lamellation, dark brown. Trichomes are cylindrical, not constricted at cross-walls, in old parts distinctly narrower than towards ends, 10-12 µm wide. Cells cylindrical, isodiametric towards ends shortened. Heterocytes are cylindrical, of the same shape and size as the cells.

**Habitat:** Epilithic.

**Comments:** New record to the Western Ghats Cyanobacterial flora of India.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Govindamoolachira, N11°37'41.96"-E076°14'30.07", Environmental Science Division, University of Calicut, CU (Fig. 2, J).

***Scytonema caldarium*** Setchell 1899

Komarek, 2013, vol.19, pp. 125, fig.107.

**Description:** Thallus expanded, flocculent, Filaments 12-16 µm in diameter, with paired or single false branches. Sheaths firm, Parallelly lamellated, in young parts colourless, in old filaments golden-brown. Trichomes are cylindrical and narrow with long cells, not constricted cross-walls, olive-green or yellowish. Cells are usually longer than wide, 6.9-18.4 µm long, 4-9 µm wide, in meristematic regions shorter; terminal cells rounded. Heterocytes are cylindrical, longer than wide.

**Habitat:** Epilithic.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Phantom rock, N11°38'08.55"-E076°12'11.61", Environmental Science Division, University of Calicut, CU No: 173139. (Fig. 3, G).

***Scytonema variable*** Gardner 1927

Komarek, 2013, vol.19, pp. 133, fig.128.

**Description:** Thallus in the form of a thick layer. Filaments are mostly straight, sparsely branched with single or geminate branches; main filaments are 34-42 µm wide, branches thinner, and all filaments very variable in width in different parts. Sheaths hyaline to yellowish-brown, 6-12 µm thick, homogeneous or with a few divergent layers, subgelatinous. Trichomes are very variable in width, distinctly narrower in older parts and slightly constricted at cross walls, widened toward ends up to 25 µm and constricted distinctly. Cells cylindrical, bright blue-green, in older segments of trichomes long cylindrical and up to 2x longer than wide, towards ends widened, short barrel-shaped; terminal cells widely rounded. Heterocytes are cylindrical, of the same size and shape as the vegetative cells.

**Habitat:** Epilithic.

**Comments:** New record to the Western Ghats Cyanobacterial flora of India.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Edakkal, N11°37'18.97"-E076°13'44.92", Environmental Science Division, University of Calicut, CU No: 173147. (Fig. 2, A).

***Scytonema tolypothrichoides*** Kutzing [sensu Frey 1927 ex Bornet et Flahault 1886](#)

Desikachary, 1959, pp. 479, pl. 91, fig. 1; Komarek, 2013, vol.19, pp. 75, fig.36.

**Description:** Thallus caespitose, floating, flake or tuft-like, green or brownish, up to 3cm in diameter, with radially arranged, coiled filaments. Filaments 10-17 µm wide, 5-6mm long. The sheath is initially colorless, later yellow or orange-brown, sometimes slightly parallelly striated with divergent layers towards ends; sometimes with slight development of ferric precipitates. Trichomes are cylindrical, not constricted at the cross walls, and sometimes narrowed or widened towards ends in narrowed or widened sheaths. Heterocytes are barrel-shaped, almost spherical up to elongated cylindrical, 6-16 µm long.

**Habitat:** Epilithic.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Edakkal, N11°37'47.79"-E076°13'40.58", Environmental Science Division, University of Calicut, CU No: 173145. (Fig. 2, H).

***Scytonema crispum*** [Agardh] ~~Bornet 1889~~ Bornet ex Toni 1907

Komarek, 2013, vol.19, pp. 82-83, fig.48.

**Description:** Thallus caespitose, fasciculated, woolly, green, olive green, blue-green to brownish. Filaments entangled, coiled, (14) 18-30 (36)  $\mu\text{m}$  wide. Branches of the same width as the main filaments. Sheaths firm, colorless, or slightly yellowish brown. Trichomes constricted at cross-walls. Cells 4.6-11.5 $\times$  12-32  $\mu\text{m}$ , in old parts of trichomes up to isodiametric, olive-green, blue-green, or yellowish brown, usually with granular content. Heterocytes ellipsoidal, rounded, solitary.

**Habitat:** Epilithic.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Govindamoolachira, N11°37'41.96"-E076°14'30.07", Environmental Science Division, University of Calicut, CU No: 163820. (Fig. 3, C).

***Petalonema crustaceum*** Agardh ex Kirchner 1900

Komarek, 2013, vol.19, pp. 150, fig.144.

**Description:** Thallus crusty, blackish, or dark brown. Filaments (13) 18-22  $\mu\text{m}$  wide, partly up to 30  $\mu\text{m}$  wide, sometimes densely entangled, richly and repeatedly branched. Branches common, usually in pairs near the base of the thallus, in erect parts mainly solitary: paired branches usually grow in parallel at the point of branching, later divergent. Sheaths yellow-brown to dark brown, with fine or oblique lamellation. Trichomes are cylindrical, not constricted at the cross-walls. Cells are mostly shorter than wide, 4.5-7 $\times$  (6-8)-12 $\mu\text{m}$ , rarely up to quadratic cylindrical. Heterocytes flattened spherical to barrel-shaped, 5.7-8  $\mu\text{m}$  in diameter, rarely elliptical - cylindrical.

**Habitat:** Epilithic.

**Comments:** New record to the Western Ghats Cyanobacterial flora of Kerala.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Thirunelli, N11°30'41.79"-E076°14'30.21", Environmental Science Division, University of Calicut, CU No: 163852. (Fig.3, E).

***Scytonema pseudoguyanense*** Bharadwaja 1934

Desikachary, 1959, pp. 472, pl. 89, fig.2; Komarek, 2013, vol.19, pp. 132, fig.121.

**Description:** Thallus in the form of a thick cushion, with long erect threads, brownish green. Filaments are irregularly curved and densely entangled, 15.7-18.9 (23.1)  $\mu\text{m}$  wide, with nu-

**Commented [WU25]:** Recent name of this algae is: Heteroscytonema crispum (Bornet ex De Toni) G.B.McGregor & Sendall 2018

**Commented [WU26]:** Describe after *Brasilonema*

**Commented [WU27]:** Check the author name. I think the authors are: (Thuret ex Bornet & Flahault) Kirchner 1900

merous false branches which are invariably slightly narrower than the main filaments, forming secondary branches at an early stage; branching usually geminate and sometimes (young) arising in characteristic loops. Sheaths firm, pigmented with parallel stratification, 3.1 (4.5)  $\mu\text{m}$  wide, in old filaments up to 7.3  $\mu\text{m}$  wide, in young branches only narrowed to 2.1  $\mu\text{m}$ . Trichomes are cylindrical, not constricted at cross-walls, 8.4-9.5 (10.5)  $\mu\text{m}$ , in old parts only 5.2  $\mu\text{m}$  wide. Cells almost quadratic, in old parts of filaments cylindrical, up to 4 $\times$  longer than broad, at the ends shortened and flattened. Heterocytes are cylindrical, quadratic, or ellipsoidal, rarely longer or shorter than wide, 7.3-18.6 (21)  $\times$  10.5-12.6  $\mu\text{m}$ .

**Distribution:** India (Gupta 2012), Jharkhand (Gupta 2021), China (Chu et al. 2007).

**Habitat:** Epilithic.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Meppadi, N 11° 30' 41.93" - E076° 0'56.8", Environmental Science Division, University of Calicut, CU No: 163857. (Fig. 2, G).

**Scytonema myochrous** ~~[Dillwyn] C.A.~~ Agardh ex Bornet et Flahault 1886

Desikachary, 1959, pp. 487, pl. 90, fig.3; Komarek, 2013, vol.19, pp. 119, figs.99,100.

**Description:** Thallus flat-hemispherical to expanded, leathery, crusty, brown-black or blackish-green. Filaments entangled, (10-14) 18-36 (40)  $\mu\text{m}$  wide, with usually common false branching; branches usually in pairs, rarely solitary, long. Sheaths are dark yellow, yellow-brown, rarely pale yellow, with distinctly divergent layers. Trichomes are cylindrical, sometimes slightly wider at the ends, not constricted at cross-walls, only slightly constricted at the ends of trichomes in branches. Cells cylindrical, blue-green or olive-green, in older parts of trichomes quadratic or slightly longer or shorter than wide (4-14  $\mu\text{m}$  long), rarely up to 2 $\times$  longer than wide, 6-12  $\mu\text{m}$  (rarely up to 23  $\mu\text{m}$ ) wide, towards ends sometimes shortened and up to slightly short barrel-shaped (as short as 3.5  $\mu\text{m}$  long); end cells rounded, 12-13  $\mu\text{m}$  wide. Heterocytes are cylindrical, rarely spherical or shortly obliquely elliptic, isodiametric-rounded or elongated elliptical, usually 8-10  $\mu\text{m}$  wide.

**Habitat:** Epilithic.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Ambukuthi 19, N11°37'01.80" - E076°14'06.19", Environmental Science Division, University of Calicut, CU No: 163855. (Fig. 23, D).

Commented [WU28]: Fig. 2 or 3? Check your algae plate.

**Scytonema arcangelii** Bornet et Flahault 1887

Komarek, 2013, vol.19, pp. 83, fig.49.

Commented [WU29]: Check this year. I think 1886

**Description:** Thallus pillow-like, olive green. Filaments 12-17 $\mu\text{m}$  wide. False branching usually geminate, branches long and bent of the same morphology as the main filaments.

Sheaths are distinct, thin, firm, colourless. Trichomes green, cylindrical along the whole length, not attenuated or widened at the ends, 9.5-14.5 µm wide, not constricted at cross walls. Cells in the middle of trichomes are cylindrical, quadratic, and rarely indistinctly barrel-shaped. Heterocytes are quadratic or slightly longer or shorter than wide, 6-14.5 µm long.

**Habitat:** Epilithic.

**Comments:** New record to the Western Ghats Cyanobacterial flora of India.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Ambukuthi 19, N11°37'01.80"-E076°14'06.19", Environmental Science Division, University of Calicut, CU No: 163819, Meppadi, N11°33'28.3" E076°07'55.23", Environmental Science Division, University of Calicut, CU No: 163810. (Fig.2, D).

***Scytonema brunneum*** Schmidle ~~in Simmer~~ 1901

Komarek, 2013, vol.19, pp. 116, fig.95.

**Description:** Thallus flat, brownish. Filaments straight, 14-24 µm wide, in lower parts creeping on the substrate. False branching is rare, usually solitary, and rarely branches in pairs. Sheaths wide, obliquely lamellated, yellow-brown. Trichomes brown, at the ends rarely blue-green, cylindrical, in middle parts without constrictions, towards ends widened and constricted at cross-walls. Cells in the middle of trichomes are longer than wide, cylindrical, near the ends barrel-shaped, isodiametric up to distinctly shorter than wide. Heterocytes are usually cylindrical, elliptical, and solitary.

**Habitat:** Epilithic.

**Comments:** New record to the Western Ghats Cyanobacterial flora of India.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Rippon, N11°32'25.02"-E076°08'10.94", Environmental Science Division, University of Calicut, CU No: 163854. (Fig. 2, B).

***Scytonema polycystum*** Bornet et Flahault 1887

Komarek 2013, vol.19, pp. 96, fig.70.

**Description:** Thallus cluster-like, greyish-green. Filaments entangled, 14-24 µm wide, scarcely falsely branched. Branches of the same morphology as the main filaments, geminate or single. Sheaths thin up to 3 µm thick, not lamellated. Trichomes are cylindrical, 10-15 (20) µm wide, not attenuated towards ends, not constricted at cross-walls. Cells cylindrical, 3-6 µm long; terminal cells rounded. Heterocytes 7-21 µm long, cylindrical.

**Habitat:** Epilithic.

**Commented [WU30]:** Check this year. I think it is 1886

**Comments:** New record to the Western Ghats Cyanobacterial flora of India.

**Specimen Examined:** INDIA: Kerala; Wayanad district, Phantom rock, N11°38'09.96"-E076°12'14.27", Environmental Science Division, University of Calicut, CU No: 163813. (Fig. [23](#), H).

**Commented [WU31]:** Fig. 2 or 3? Check your algae plate.

UNDER PEER REVIEW

**Commented [WU32]:** Scales are different in thickness, some are large, some are very small not visible. Better to keep distinctly visible scales with similar thicknesses.




Fig. 2. A. *Scytonema variabile*, B. *S.eytonoma brunneum*, C. *S.eytonoma bivaginatam*, D. *S.eytonoma arcangelii*, E. *Brasilonema bromeliae*, F. *B.rasilonema epidendron*, G. *Scytonema pseudoguyanense*, H. *S.eytonoma tolypothrichoides*, I. *S.eytonoma tenellum*, J. *S.eytonoma spirulinoides*.

Commented [WU33]: Same comment for scale as above plate.




Fig. 3. A. *Scytonema malaviyanense*, B. *S.cytonema drilosiphon*, C. *S.cytonema crispum*, D. *S.cytonema myochrous*, E. *Petalonema crustaceum*, F. *Scytonema masonianum*, G. *S.cytonema caldarium*, H. *S.cytonema polycystum*, I. *Brasilonema terrestre*.

The Present study observed that the members of this family are abundantly seen on rock surfaces. The rock surfaces provide a suitable substrate for the significant growth of Scytonema, Brasilonema, and Petalonema species, often forming mats. Cyanobacteria adhere to rock surfaces by producing and secreting mucilaginous substances known as extracellular polymeric substances. The protective sheath layers of cyanobacteria are made up of polysaccharides that retain moisture, enabling them to survive severe drought conditions (Gloaguen et al. 1995; Adhikary 1998; Bertocchi et al. 1990; Cappitelli et al. 2012; Rossi et al. 2012). Additionally, rocky environments shield cyanobacterial species from various environmental stresses, such as high UV radiation, extreme temperatures, desiccation, and physical removal by wind (Hughes and Lawley 2003).

The study focused on the diversity and species-level identification of members of the Scytonemataceae family. A total of nineteen species of cyanobacteria belonging to the order Nostocales were recorded. Among these, the genus Scytonema, represented by fifteen species, displayed the highest dominance. Additionally, three species from the genus Brasilonema and a single species of Petalonema were identified. The identified species include *Scytonema drilosiphon*, *S.eytonema bivaginatam*, *S.eytonema malaviyanense*, *S.eytonema masonianum*, *S.eytonema tenellum*, *S.eytonema spirulinoides*, *S.eytonema caldarium*, *S.eytonema variabile*, *S.eytonema tolypothrichoides*, *S.eytonema crispum*, *S.eytonema pseudoguyanense*, *S.eytonema myochrous*, *S.eytonema arcangelii*, *S.eytonema brunneum*, *S.eytonema polycystum*, *Brasilonema terrestre*, *B.fasilonema epidendron*, *B.fasilonema bromeliae*, and *Petalonema crustaceum*.

The species *Scytonema pseudoguyanense* Bharadwaja (1934) has been reported in various locations. Adhikary et al. (2015) observed them in stone temples of Bhubaneswar, India. Samad and Adhikary (2008) confirmed their presence on the walls of Parasurameswar temple, Bhubaneswar, Orissa. Mahanandia and Singh (2023) documented their existence on monuments and exterior surfaces of building facades in Odisha state, India. Dasgupta et al. (2017) reported them from suburban waterbodies in Lucknow, Uttar Pradesh. Philip (2021) recorded from Karapuzha, Wayanad, Kerala. In the current study, the same species were noted on the moist rock in Meppadi of Wayanad district.

Fiore et al. (2007) isolated *Brasilonema bromeliae* from sub-aerophytic environments in tropical and subtropical Brazil. Sant' Anna et al. (2011) reported the presence of *Brasilonema epidendron* and *B.fasilonema terrestre* also from Brazil. The current investigation reported *B.fasilonema terrestre* in Edakkal, Wayanad district of the Western Ghats region of Kerala. Additionally, *B.fasilonema epidendron* was recorded at Phantom Rock, Wayanad. This study demonstrates the occurrence of *B.fasilonema bromeliae* from Edakkal, Wayanad district, within the Western Ghats region of Kerala. Biswas (1934) described *Petalonema*

**Commented [WU34]:** Two recent citations will be sufficient, not necessary to write many old citations

**Commented [WU35]:** Copied from <https://doi.org/10.1080/08927014.2013.794224>

**Commented [WU36]:** Repeated. Delete it

**Commented [WU37]:** Italicise

*crustaceum* from Cherrapunji, Meghalaya. Singh and Singh (2019) also described *Petalonema crustaceum* from Sirsi taluk, Uttara Kannada, Karnataka. In the current investigation, the occurrence of this species was observed in Thirunelli, Wayanad district in the Western Ghats region of Kerala.

The occurrence of *Scytonema myochrous* was documented on rock surfaces in Ambukuthi 19, Wayanad districts. This observation is consistent with findings by various researchers. Suresh et al. (2012) recorded *Scytonema myochrous* from the Western Ghats and Eastern Ghats of India. In tropical environments, Budel (1999) observed on moist rock, and Popovic et al. (2018) confirmed its existence on selected monuments in Serbia. Philip (2021) recorded its presence in Eravikulam National Park, Idukki, Kerala. Sabre et al. (2021) reported *Scytonema myochrous*, a cyanobacterium, is a new record for the Egyptian algal inventory.

Philip (2021) recorded *Scytonema caldarium* from Karappuzha, Wayanad, Kerala. The current research recorded its presence in Phantom rock, Wayanad district in the Western Ghats region of Kerala. *Scytonema malaviyanense* (Bharadwaja, 1930) was observed on the bark of trees. Banerjee et al. (2020) reported it from West Bengal, India. The current investigation reported its presence from Kanthanpara, Wayanad district in the Western Ghats region of Kerala. *Scytonema tolypothrichoides* was previously reported by Nikam et al. (2010) from the Western Ghats region of Maharashtra state. Deb et al. (2015) isolated from acidic rice field soil from the Dholai region of the Cachar district in Assam, North-East India. Kumar et al. (2016) recorded from biological soil crust in the eastern region of India. Ram (2022) observed from Sakthikulangara, Kollam. The present study confirmed its presence in Edakkal, Wayanad district of the Western Ghats region of Kerala.

Krawczyk et al. (2014) reported *Scytonema drilosiphon* from historical buildings in the former Auschwitz II-Birkenau concentration camp, in Poland. Banerjee et al. (2020) reported *Scytonema drilosiphon* from West Bengal, India. The current study demonstrates the presence of this species from Thirunelli, Wayanad district in the Western Ghats region of Kerala. Komarek et al. (2013) reported *Scytonema crispum* from Atlantic Rainforest in Southeast Brazil. Philip (2021) recorded from Pazhassi Dam, Kannur, Kerala. In the current investigation, this species was observed in Govindamoolachira, Wayanad, in the Western Ghats region of Kerala.

Sarma et al. (2023) reported *Scytonema bivaginatum* from Tripura, India, and the species were new additions to the flora of Tripura, India. The present work confirmed their presence in Edakkal, Wayanad district. Komarek et al. (2013) reported *Scytonema arcangelii* from Atlantic Rainforest in Southeast Brazil. Caires et al. (2019), recorded from tropical marine environments of Bahia State, Northeastern Brazil. The present study documented its

Commented [WU38]: Sabere

Commented [WU39]: Nowicka-Krawczyk

Commented [WU40]: Aguiar Caires

existence on rocks in Ambukuthi 19, Wayanad district, in the Western Ghats region of Kerala.

Cyanobacteria have evolved several adaptations to survive, compete, and eventually dominate in a wide array of ecosystems. (Rastogi and Madamwar 2015; Menamo and Wolde 2015). Due to their adaptive capacities, cyanobacterial diversity occupies a huge spectrum of habitats on the Earth. Their diversity is due to the cellular mechanisms that allow them to adapt in response to environmental changes and grow easily as a dense population with remarkable speed. Nevertheless, their rapid growth rate varies depending on biotic factors, nutrient levels, climate change, or the recent threats due to global warming (Sukenik et al. 2012; Paerl and Paul 2012).

Commented [WU41]: Delete full stop

## CONCLUSION

The present study investigates the diversity of the family Scytonemataceae in Wayanad district, Western Ghats region of Kerala. This research provides comprehensive information on the cyanobacterial diversity in this area, marking it as the first time study. Species within the Scytonemataceae family are typically abundant in wet environments and often attach to rock surfaces. The present study recorded 19 species from the Western Ghats region of Kerala and among these species, *Scytonema malaviyaensis*, *Scytonema drilosiphon* and *Petalonema crustaceum* are new reports to Kerala and 11 species such as *Scytonema spirulinoides*, *Scytonema masonianum*, *Scytonema variabile*, *Scytonema bivaginatum*, *Scytonema arcangelii*, *Scytonema tenellum*, *Scytonema brunneum*, *Scytonema polycystum*, *Brasilonema bromeliae*, *Brasilonema terrestre*, *Brasilonema epidendron* are new to India.

Wayanad, situated in the Western Ghats, is a biodiversity hotspot known for its unique ecosystems. The area's rocky outcrops and shaded, humid environments provide an ideal habitat for *Scytonema* species belonging to the family Scytonemataceae. However, these populations face several threats. Habitat loss due to construction, quarrying, and deforestation disrupts the rocky habitats vital for Scytonemataceae family. Additionally, pollution and climate change, characterized by altered rainfall patterns, increased temperatures, and changes in water availability further threaten these habitats. Recognizing the ecological and economic value of cyanobacteria is essential for their conservation. By protecting their natural habitats and promoting sustainable practices, we can ensure that these microorganisms continue contributing to Earth's ecosystems and driving future innovations. The biotechnological potential of *Scytonema* species in Wayanad District emphasizes their importance for sustainable development and ecological health. Their ability to produce valuable compounds, such as scytonemin and geosmin, and their roles in nitrogen fixation and soil

stabilization, underscores the need for effective conservation strategies. Protecting these microorganisms will preserve biodiversity and strengthen the resilience of the local ecosystems against environmental threats.

## REFERENCES

1. Whitton B.A & Potts M (2000). "The Ecology of Cyanobacteria: Their Diversity in Time and Space. *Kluwer Academic Publishers*, Dordrecht, Springer Netherlands, 1-704.
2. Thajuddin, N., & Subramanian, G. (2005). Cyanobacterial biodiversity and potential applications in biotechnology. *Current science*, 47-57.
3. Komarek, J., Sant'Anna, C. L., Bohunicka, M., Mareš, J., Hentschke, G. S., Rigonato, J., & Fiore, M. F. (2013). Phenotype diversity and phylogeny of selected Scytonema-species (Cyanoprokaryota) from SE Brazil. *Fottea*, 13(2), 173-200. <http://dx.doi.org/10.5507/ftot.2013.015>
4. Fiore, M. F., Sant'Anna, C. L., Azevedo, M. T. D. P., Komarek, J., Kastovsky, J., Sulek, J., & Lorenzi, A. A. S. (2007). The cyanobacterial genus Brasilonema, gen. nov., a molecular and phenotypic evaluation 1. *Journal of Phycology*, 43(4), 789-798. <http://dx.doi.org/10.1111/j.1529-8817.2007.00376.x>
5. Sant'Anna, C. L., Azevedo, M. P., Fiore, M. D. F., Lorenzi, A. S., Kastovsky, J., & Komárek, J. (2011). Subgeneric diversity of Brasilonema (cyanobacteria, Scytonemataceae). *Brazilian Journal of Botany*, 34, 51-62. <https://doi.org/10.1590/S0100-84042011000100006>
6. Lau, N. S., Matsui, M., & Abdullah, A. A. A. (2015). Cyanobacteria: photoautotrophic microbial factories for the sustainable synthesis of industrial products. *BioMed research international*, 2015(1), 754934. <https://doi.org/10.1155/2015/754934>
7. Kumar, B. N. P., Mahaboobi, S., & Satyam, S. (2016). Cyanobacteria: a potential natural source for drug discovery and bioremediation. *Journal of Industrial Pollution Control*, 32, 508-517.
8. Boyd, M.R., Gustafson, K.R., McMahon, J.B., Shoemaker, R.H., O'Keefe, B.R., Mori, T., Gulakowski, R.J., Wu, L., Rivera, M.I., Laurencot, C.M. and Currens, M.J. (1997). Discovery of cyanovirin-N, a novel human immunodeficiency virus-inactivating protein that binds viral surface envelope glycoprotein gp120: potential applications to microbicide development. *Antimicrobial agents and chemotherapy*, 41(7), pp.1521-1530.
9. Loya, S., Reshef, V., Mizrahi, E., Silberstein, C., Rachamim, Y., Carmeli, S., & Hizi, A. (1998). The inhibition of the reverse transcriptase of HIV-1 by the natural sulfoglycolipids

**Commented [WU42]:** 1.Arrange all references in alphabetic order, instead of number system.  
2.Keep DOI if available (I have added DOI for few references)

from cyanobacteria: contribution of different moieties to their high potency. *Journal of natural products*, 61(7), 891-895.

10. Bokesch, H.R., O'Keefe, B.R., McKee, T.C., Pannell, L.K., Patterson, G.M., Gardella, R.S., Sowder, R.C., Turpin, J., Watson, K., Buckheit, R.W. and Boyd, M.R. (2003). A potent novel anti-HIV protein from the cultured cyanobacterium *Scytonema varium*. *Biochemistry*, 42(9), 2578-2584.
11. Tyagi, R., Kaushik, B. D., & Kumar, J. (2014). Antimicrobial activity of some cyanobacteria. *Microbial diversity and biotechnology in food security*, 463-470.
12. Ngo, T.T., Nguyen, B.L.T., Duong, T.A., Nguyen, T.H.T., Nguyen, T.L., Kieu, K.T., Do, M.H.T., Nguyen, S.V., Thang, N.D. and Pham, H.T. (2022). Polyphasic evaluation and cytotoxic investigation of isolated cyanobacteria with an emphasis on potent activities of a *Scytonema* strain. *Frontiers in Microbiology*, 13, 1025755.
13. Tawong, W., Pongcharoen, P., & Nishimura, T. (2022). Geosmin-producing *Scytonema foetidum* sp. nov.(Scytonemataceae, Cyanobacteria): morphology and molecular phylogeny. *Fottea*, 22(1), 78-92.
14. Abed, R. M., Al Kharusi, S., Schramm, A., & Robinson, M. D. (2010). Bacterial diversity, pigments, and nitrogen fixation of biological desert crusts from the Sultanate of Oman. *FEMS microbiology ecology*, 72(3), 418-428.
15. Menescal, M. T. A., dos Santos Almeida, E., & Sales, E. A. (2020). Identification and Study of a Promising Cyanobacteria Species for Biotechnological Applications. *Journal of bioengineering, technologies, and health*, 3(4), 328-333.
16. GoogleEarthPro.WesternGhats, Kerala. (2022). Retrieved from <https://earth.google.com/>
17. Rippka R., Deruelles, J., Waterbury, J., Herdman, M., & Stanier, R.Y. (1979). Generic assignments, strain histories, and properties of pure cultures of cyanobacteria. *Microbiology* 111, 1–61.
18. Komarek, J. Subwasserflora Von Mitteleuropa, Bd. 19/3: Cyanopokaryota. 3. Teil/3rd Part: Heterocytous Genera. *Springer, Spektrum*. 2013.
19. Desikachary, T. V. *Cyanophyta*. ICAR Monograph on Algae, New Delhi.1959. 1-686.
20. Komarek, J., & Anagnostidis, K. (1998). Cyanopokaryota I: Chroococcales, Süßwasserflora von Mitteleuropa, 19/1. *G. Fischer, Stuttgart*.
21. Komarek, J., & Anagnostidis, K. (2005). Cyanopokaryota -2. Teil/ 2nd Part: Oscillatoriales. In: Budel B., Krienitz L., Gartner G. and Schagerl M. (ed.), Süßwasserflora von Mitteleuropa 19/2. *Elsevier/Spektrum*, Heidelberg.
22. Guiry, M. D., Guiry, G. M. *AlgaeBase*. World-Wide Electronic Publication, National University of Ireland, Galway, 2015, <http://www.algaebase.org> (accessed July 2024)

23. Gloaguen, V., Morvan, H., & Hoffmann, L. (1995). Released and capsular polysaccharides of Oscillatoriaceae (Cyanophyceae, Cyanobacteria). *Algological Studies/Archiv für Hydrobiologie, Supplement Volumes*, 53-69.
24. Adhikary, S. P. (1998). Polysaccharides from mucilaginous envelope layers of cyanobacteria and their ecological significance. *Journal of scientific & industrial research*, 57(8), 454-466.
25. Bertocchi, C., Navarini, L., Cesaro, A., & Anastasio, M. (1990). Polysaccharides from cyanobacteria. *Carbohydrate polymers*, 12(2), 127-153.
26. Cappitelli, F., Salvadori, O., Albanese, D., Villa, F., & Sorlini, C. (2012). Cyanobacteria cause black staining of the National Museum of the American Indian Building, Washington, DC, USA. *Biofouling*, 28(3), 257-266.
27. Rossi, F., Micheletti, E., Bruno, L., Adhikary, S. P., Albertano, P., & De Philippis, R. (2012). Characteristics and role of the exocellular polysaccharides produced by five cyanobacteria isolated from phototrophic biofilms growing on stone monuments. *Biofouling*, 28(2), 215-224.
28. Hughes, K. A., & Lawley, B. (2003). A novel Antarctic microbial endolithic community within gypsum crusts. *Environmental Microbiology*, 5(7), 555-565.
29. Bharadwaja, Y. (1934). *The Taxonomy of Scytonema and Tolypothrix: including some new records and new species from India and Ceylon*. Laboratoire.
30. Adhikary, S. P., Keshari, N., Urzi, C., & De Philippis, R. (2015). Cyanobacteria in biofilms on stone temples of Bhubaneswar, Eastern India. *Algological Studies*, 147, 67-93.
31. Samad, L. K., & Adhikary, S. P. (2008). Diversity of micro-algae and cyanobacteria on building facades and monuments in India. *Algae*, 23(2), 91-114.
32. Mahanandia, S., & Singh, L. (2023). Subaerial cyanobacteria on the monuments and exterior surface of building facades of Odisha state, India. *Taiwania*, 68(2).
33. Dasgupta, C. N., Toppo, K., Nayaka, S., & Singh, A. K. (2017). Algal Diversity and Water Quality Assessment During Summer in Some Suburban Water Bodies of Lucknow, Uttar Pradesh. *Crypt Biodivers Assess*, 2(2), 1-10.
34. Philip, M. (2021). *Cyanobacterial diversity of Western Ghat forests of Kerala, India*. (PhD Thesis). University of Calicut, Malappuram, Kerala.
35. Biswas, K. (1934). *Observations on algal collections from Khasia and Jaintia Hills, Assam, India*. Heinrich.
36. Singh, P. & Singh D.M. (2019). Subaerial non-heterocytous and heterocytous cyanobacteria from Sirsi taluk, Uttara Kannada, Karnataka, India. *International journal of scientific research and reviews*, 8(1): 1248-1274.

37. Suresh, A., Kumar, R. P., Dhanasekaran, D., & Thajuddin, N. (2012). Biodiversity of microalgae in Western and Eastern Ghats, India. *Pakistan Journal of Biological Sciences: PJBS*, 15(19), 919-928.
38. Budel, B. (1999). Ecology and diversity of rock-inhabiting cyanobacteria in tropical regions. *European Journal of Phycology*, 34(4), 361-370.
39. Popovic, S., Stupar, M., Unkovic, N., Subakov Simic, G., & Ljaljevic Grbic, M. (2018). Diversity of terrestrial cyanobacteria colonizing selected stone monuments in Serbia. *Studies in Conservation*, 63(5), 292-302.
40. Saber, A. A., El-Sheekh, M., Nikulin, A. Y., Cantonati, M., & Saber, H. (2021). Taxonomic and ecological observations on some algal and cyanobacterial morphospecies new for or rarely recorded in either Egypt or Africa. *Egyptian Journal of Botany*, 61(1), 283-301. <https://dx.doi.org/10.21608/ejbo.2020.50683.1587>
41. Bharadwaja, Y. (1930). *Scytonema malaviyaensis* sp. nov. *Revue Algologique*, 5, 223-227.
42. Banerjee, S., Singh, A., & Pal, R. (2020). A taxonomic investigation on heterocytous Cyanobacteria of West Bengal, India.
43. Nikam, T.D., Nehul, J.N., Gahile, Y.R., Auti, B.K., Ahire, M.L., Nitnaware, K.M., Joshi, B.N., & Jawali, N. (2010). Cyanobacterial diversity in Western Ghats region of Maharashtra, India. *Methods*, 2008(2009).
44. Deb, S., Rout, J., Sengupta, M., & Chakraborty, B. (2015). Biochemical profile and antimicrobial activity of a cyanobacterium, *Scytonema tolypothrichoides* isolated from acidic rice field soil of Cachar district (Assam), India. *International Journal of Life Science and Pharma Research*, 5(2), L21-L31.
45. Kumar, D., Kastanek, P., & Adhikary, S. P. (2016). Diversity of cyanobacteria in biological crusts on arid soils in the Eastern region of India and their molecular phylogeny. *Current Science*, 1999-2004.
46. Ram, A. T. (2022). *Ecology and diversity of mangrove-associated cyanobacteria of Southern Kerala, India*. (PhD Thesis). University of Calicut, Malappuram, Kerala.
47. Nowicka-Krawczyk, P., Zelazna-Wieczorek, J., Otlewska, A., Kozirog, A., Rajkowska, K., Piotrowska, M., Gutarowska, B. and Zyzik-Bialek, A. (2014). Diversity of an aerial phototrophic coating of historic buildings in the former Auschwitz II-Birkenau concentration camp. *Science of the total environment*, 493, pp.116-123. <https://doi.org/10.1016/j.scitotenv.2014.05.113>
48. Sarma, K., Doli, Gupta, D., Gauri, Kour, N., Singh, J., Saini, A., Kumar, S., Sharma, M., Kumar, N., Das, D., Das, S., & Kant, R. (2023). Diversity and Morpho-Taxonomy of the Genus *Scytonema*: A Heterocystous Cyanoprokaryote from Tripura, India. *Indian Hydrobiology*, 22(1): 163–170

Commented [WU43]: delete

49. Aguiar Caires, T., Leite Sant'Anna, C., & de Castro Nunes, J. M. (2019). Biodiversity of benthic filamentous cyanobacteria in tropical marine environments of Bahia State, Northeastern Brazil. *Brazilian Journal of Botany*, 42, 149-170. <http://dx.doi.org/10.1007/s40415-019-00517-2>
50. Rastogi, P.R., & Madamwar, D. (2015). UV-Induced Oxidative Stress in Cyanobacteria: How Life is able to Survive? *Biochemistry & Analytical Biochemistry*, 4: 173.
51. Menamo, M., & Wolde, Z. (2015). The potential of cyanobacteria growth in different water sources. *Biochemistry and Physiology*, 4, 1-3.
52. Sukenik, A., Hadas, O., Kaplan, A., & Quesada, A. (2012). Invasion of Nostocales (cyanobacteria) to subtropical and temperate freshwater lakes—physiological, regional, and global driving forces. *Frontiers in microbiology*, 3, 86.
53. Paerl, H. W., & Paul, V. J. (2012). Climate change: links to the global expansion of harmful cyanobacteria. *Water Research*, 46 (5), 1349-1363.

UNDER PEER REVIEW