

## Original Research Article

### Influence of secondary and micro nutrients status of soils on grape yield of vineyards in Northern Dry Zone of Karnataka, India

#### Abstract:

A study was conducted to study the influence of secondary and micro nutrients status of soils on grape yield of vineyards during the year 2019-20 in Yelburga taluk of Koppal district. Based on the previous year yield data, the thirty vineyards were classified into three groups namely, low yielding vineyards (LYV), medium yielding vineyards (MYV) and high yielding vineyards (HYV). The soil samples collected after harvest of crop recorded slight increase in the exchangeable Ca and Mg content compared to the soil samples collected before October pruning. The HYV [ $11.16 \pm 2.61$  and  $12.08 \pm 2.76$  c mol (p+)  $\text{kg}^{-1}$ ] soils recorded higher calcium content compared to other groups in both the sampling seasons respectively. HYV [ $6.81 \pm 1.26$  and  $6.88 \pm 1.19$  c mol (p+)  $\text{kg}^{-1}$ ] recorded highest exchangeable Mg content compared to other two groups in both the sampling seasons respectively. The sulphur content in group HYV was significantly higher ( $51.67 \pm 16.46$  and  $64.09 \pm 18.59$  mg  $\text{kg}^{-1}$ ) as compared to the other groups at before October pruning and at after harvest, respectively. Distribution of micronutrients in soil samples collected before October pruning and after harvest were found in the order Mn > Fe > Zn > Cu. Vineyard soil sample showed medium to high content of micronutrients. The concentration of DTPA extractable micronutrients among three groups of vineyards differed significantly. The differences in yields of vineyards is depends on soil fertility status. Hence, it is important to adopt good nutrient management practices to ensure sustainable grape yields in vineyards.

**Keywords:** Vineyards, Fertility, Micro nutrients, Calcium, Magnesium Sulphur etc.

#### Introduction:

Grape (*Vitis vinifera* L.), belonging to the family *vitaceae* is one of the most important commercial fruit crop in India. Though it is temperate fruit crop, it is well adopted to sub-tropical and tropical agro climatic conditions prevailing in the Indian sub-continent. Grapes are being cultivated for table purpose and wine industry. In India, grapes are primarily grown for table use, with 90% of the grapes being consumed fresh. India has some of the highest fresh grape yield levels in the world. Globally, grapes are primarily grown for wine making, with 71% of the world's grape production

used for wine. Other processed grape products include raisins, juice, vinegar, and grape seed extracts and oil. Due to its high commercial value, the crop is intensively managed in terms of soil and crop management practices. The life of vineyards to be sustained for longer periods, the nutrition of vineyards plays a major role. Therefore, grape growing is primarily achieved through drip irrigation in these dry tracts. Most of the essential nutrients (macro, secondary and micronutrients) are added to soil in the form of basal application (through organic manures and fertilizers) and fertigation (through water soluble fertilizers). The majority of grape growers also practice foliar nutrition, in particular for secondary and micronutrients. In enhancing the grapes quality and productivity, the use of growth regulators plays an important role.

Nutritional management is considered as one of the important aspects in successful grape cultivation. The nutrients present in soil and grapevine must be observed regularly and maintained for optimal efficiency. Therefore, it has to constantly changing; the best set up is soil and petiole sampling for regular analysis to overcome nutrient deficiency and reduce excess use of nutrient inputs. Based on the available nutrient content in soil and plant, the fertilizer recommendations are required in vineyards. Indiscriminate use of chemical fertilizers to get better yield can alter the fertility status of the soil and leads to nutrients toxicity. The sustainable soil fertility maintenance is essential for grape production. The available nutrient concentration is often analyzed by properties of soil and nutrient interactions.

The soils of the study area were red in color and texture varies from sand to clay, and having porous and friable structure. The soils have neutral to alkaline pH and red soils generally designated under Alfisols. The secondary and micronutrients are also applied along with nitrogen, phosphorous and potassium fertilizers. Deficiency of any secondary and micronutrients not only reduce the yield but also the fruit quality. Nutrient management is one of the major cost- economic aspects in grape cultivation.

For the optimum use of soil for increased crop production, its fertility status is of prime importance. In the semi-arid region of Karnataka, grape cultivation has claimed great significance. Now the area under grape cultivation in the Koppal district is increasing. Grape production has also been decreasing. In this region, studies have shown that the issues are mainly related to nutrient imbalance. Thus, Assessment of nutritional status and nutritional requirements of vineyards assumes a greater significance for successful viticulture, so the present investigation was undertaken to assess secondary and micronutrients status in vineyards of Koppal district, Karnataka.

## **Material and Methods**

## **Survey of Vineyards**

In this study, thirty vineyards from Yelburga taluk were selected as contact farmers for interview and collection of soil and petiole samples (Fig. 1). These farmers were interviewed initially to collect the information on their vineyards in terms of crop details– variety, age of the orchard, soil type, nutrient management practices, grape yield etc. Classification of vineyards was done based on previous year yield data of grapes. The thirty vineyards were classified into three groups namely, LYV- Low yielding vineyards, MYV- Medium yielding vineyards and HYV- High yielding vineyards presented in Appendix III. Each group contains 10 vineyards.

## **Soil sample collection**

Soil sampling was carried out at two seasons, one before the October pruning i.e. during growth stage and the other one after the harvest of the crop (Feb-March). In each vineyard, soil samples were collected from 6-8 spots at 0-20 cm depth and pooled to get one composite sample of about half kilo gram by quartering technique. Then, the sample was sieved (2 mm) and stored in air tight containers for further analysis. These soil samples were analyzed for secondary nutrients and DTPA extractable micronutrients (metal ions).

## **Analysis of Secondary nutrient**

The exchangeable Ca and Mg (both exchangeable and water soluble) were extracted with neutral normal ammonium acetate and estimated by Versenate titration method using EBT for Ca+Mg and P&R indicator for Ca alone (Jackson, 1973). Available sulphur content was determined by turbidometric method after extraction with 0.15% CaCl<sub>2</sub> using spectrophotometer at 420 nm (Black, 1965).

## **Analysis of DTPA-Extractable micronutrients**

The DTPA extractable micronutrients (metal ions namely Fe, Mn, Zn and Cu) were extracted with DTPA buffer at 1:2 soil to extractant ratio (Lindsay and Norwell, 1978). The concentration of DTPA-Fe, Mn, Zn and Cu in the extract were determined by atomic absorption spectrophotometer (Thermofisher model).

## **Statistical analysis**

The data obtained were subjected to statistical tests using normal one way ANOVA technique, and Descriptive statistical analysis. Simple correlation studies were also made to understand their interaction effects. **One-way ANOVA is a type of ANOVA that uses one**

independent variable. It's used to test for differences among at least three groups. (<https://www.xlstat.com/en/solutions/features/anova-analysis-of-variance>)

## Results and Discussion

### Availability of Secondary nutrients

Extent of distribution of secondary nutrients namely, calcium, magnesium and sulphur are presented in Table 1 and respective mean values of different vineyards are depicted in Fig. 1.

The available exchangeable calcium was recorded low in medium yielding vineyards with  $9.55 \pm 2.87$  c mol (p+)  $\text{kg}^{-1}$  as compared to the low yielding vineyards [ $10.17 \pm 3.04$  c mol (p+)  $\text{kg}^{-1}$ ] and high yielding vineyards [ $11.16 \pm 2.61$  c mol (p+)  $\text{kg}^{-1}$ ]. The soils after harvest of crop recorded slight increase in the exchangeable Ca content compare to the soils before October pruning (Table 1). High yielding vineyards recorded higher [ $6.81 \pm 1.26$  c mol (p+)  $\text{kg}^{-1}$ ] exchangeable Mg content followed by medium yielding vineyards [ $5.64 \pm 1.93$  c mol (p+)  $\text{kg}^{-1}$ ] and low yielding vineyards [ $5.59 \pm 1.83$  c mol (p+)  $\text{kg}^{-1}$ ].

There was an increase in the exchangeable Mg content were observed in the soils after harvest as compared to the soils before October pruning. Exchangeable Ca and Mg content in vineyards soils were found low because red soils generally contain low Ca and Mg content compared to black soils as the cation exchange capacity (CEC) of the red soil is low compared to black soil (Ananthanarayan, 1978). The variation might be due to differences in additions of lime and Mg. They are also known to be influenced by management practices, nutrient inputs, amendments used (Nagaraja 1997). After the harvest, exchangeable Ca and Mg content increased due to high temperature, Soil water evaporation and capillary rise of salts along with water might be attributed. Similar levels of Ca and Mg contents were reported by Bhat *et al.* (2017) in grape soils and Ganai *et al.* (2018) in apple orchards.

Available sulphur content in vineyards ranged between medium to high. Among different groups, high yielding vineyards (HYV) recorded significantly higher available sulphur content in both the seasons ( $51.67 \pm 16.46$  and  $64.09 \pm 18.59$  mg  $\text{kg}^{-1}$  in soils collected before October pruning and after harvest respectively). Least S content was recorded in fields of low yielding vineyards ( $26.45 \pm 4.90$  and  $29.00 \pm 5.33$  mg  $\text{kg}^{-1}$  in soils collected before October pruning and after harvest respectively). High sulphur content in high yielding vineyards (HYV) may be attributed due to application of sulphate form of micronutrient ( $\text{ZnSO}_4$  and  $\text{FeSO}_4$ ). Also, high application of organic matter resulted in higher availability of sulphur in vineyards as reported by Basumurthy and Talukdar (1999). Similar levels of sulphur content in vineyard soils were also reported by Yogeeshappa (2007), Dar *et al.* (2015) and Shivannanavar (2016).

## Availability of DTPA extractable micronutrients

The iron content in soils after harvest was found to be higher as compared to the iron content of soils before October pruning. The mean values of iron were found significantly high in HYV ( $4.73 \pm 0.96 \text{ mg kg}^{-1}$ ). However, the other two groups namely LYV ( $3.70 \pm 0.52 \text{ mg kg}^{-1}$ ) and MYV ( $3.96 \pm 0.82 \text{ mg kg}^{-1}$ ) were on par with each other (Table 2 and Fig. 2).

The LYV recorded significantly low Mn content ( $3.28 \pm 0.44 \text{ mg kg}^{-1}$ ) and manganese content in MYV ( $3.79 \pm 0.68 \text{ mg kg}^{-1}$ ) and HYV ( $4.21 \pm 0.38 \text{ mg kg}^{-1}$ ) were found on par with each other.

The Zn content in LYV recorded significantly low ( $0.92 \pm 0.11 \text{ mg kg}^{-1}$ ), and Zn content of MYV ( $1.11 \pm 0.23 \text{ mg kg}^{-1}$ ) and HYV ( $1.12 \pm 0.08 \text{ mg kg}^{-1}$ ) were found on par with each other. However, the soils after harvest of crop recorded slight increase in the Zn content compared to the soils collected before October pruning.

The mean values of copper were found significantly higher in HYV ( $0.80 \pm 0.30 \text{ mg kg}^{-1}$ ). However, the Cu content in other two groups namely LYV ( $0.33 \pm 0.10 \text{ mg kg}^{-1}$ ) and MYV ( $0.55 \pm 0.34 \text{ mg kg}^{-1}$ ) remained on par with each other.

Distribution of micronutrients in grape soils considered for the study were observed in the order of  $\text{Mn} > \text{Fe} > \text{Zn} > \text{Cu}$  in soils collected before October pruning and after harvest, the values were recorded low to moderate in range. However, the micronutrients (DTPA - Fe, Mn, Zn and Cu) in soils collected after the harvest of the crop recorded higher values.

This variation in concentrations might be due to the nature and behavior of the element in soil as determined by pH and soil organic matter content. Soil micronutrients are enhanced by release and chelation of metal ions (Lindsay and Norwell, 1978) and also direct addition of micronutrients to the crop might have enhanced their availability among soils. Intensive nutrient management in white grape type orchards might have induced higher availability.

Application of Zn-fertilizers under Zn deficiency conditions has been associated with improved yield and quality of grapes (Malakouti, 2007). Samiullah *et al.* (2013) while studying the micronutrient status of peach orchard soils of swat valley, they reported the concentration of Fe in the 0-45 cm soil ranged from 5.73 ug to 116.14 ug  $\text{g}^{-1}$  soil with a mean value of  $27 \pm 18.97$ .

These results indicate that use of organic manures and application of secondary and micronutrients helped to maintain higher availability. Soil micronutrients are enhanced by release and chelation of metal ions (Lindsay and Norwell, 1978) and also direct addition of

micronutrients to the crop might have enhanced their availability among soils. Similar observation recorded by Patiram et al. (2000), Yogeeshappa (2007), Thakreet *et al.* (2013).

Bhargava and Raghupathi (1994) investigated the optimum range of DTPA extractable soil Zn for Anab-e-Shahi was 8 to 24.9 ppm and for Thompson Seedless it was 4.5 to 8.2 ppm. The optimum range of Zn in petioles at bud differentiation stage was 25 to 94 ppm for Anab-e-Shahi and 47 to 88 ppm for Thompson Seedless.

### **Grape yields**

The fruit yields in LYV, MYV and HYV groups were found to be  $19.70 \pm 1.58$ ,  $26.25 \pm 1.02$  and  $33.15 \pm 2.69$  t ha<sup>-1</sup> respectively and HYV recorded significantly higher grape yields as compared to MYV. Thus, significant differences were observed and the yields recorded followed the order of HYV > MYV > LYV (Table 3 and Fig. 3). The differences in yields could be attributed to variations in nutrient status of soil as determined by application of nutrients (Bhargava and Sumner, 1987). Here the recorded yield was found to be directly influenced by the soil fertility status, petiole nutrient content and climatic conditions. The results are in accordance with the Yogeeshappa (2007), Naraboli, 2016 and Kondiet *et al.* (2018).

### **CONCLUSION**

The HYV [ $11.16 \pm 2.61$  and  $12.08 \pm 2.76$  c mol (p+) kg<sup>-1</sup>] soils recorded higher calcium content compared to other groups in both the sampling seasons respectively. HYV [ $6.81 \pm 1.26$  and  $6.88 \pm 1.19$  c mol (p+) kg<sup>-1</sup>] recorded highest exchangeable Mg content compared to other two groups in both the sampling seasons respectively. Distribution of micronutrients in soil samples collected before October pruning and after harvest were found in the order Mn > Fe > Zn > Cu. Vineyard soil sample showed medium to high content of micronutrients. The concentration of DTPA extractable micronutrients among three groups of vineyards differed significantly. It was concluded from the study that the yield of vineyards is largely dependent on nutrient inputs applied, soil fertility status. Thus, it is important to adopt better nutrient management practices by the farmers to ensure higher yields.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

## Reference:

- Ananthanarayana (1978) Dynamics of Mg in relation to crops in soils of Karnataka. *Ph. D. Thesis*, Uni. Agric. Sci. Bangalore (India).
- Basumurthy, A. and Talukdar, M. C. (1999) Long-term effect of integrated nutrients supplies system on secondary nutrient of soil. *Oryza.*, 36: 266-267.
- Bhargava, B. S. and Raghupathi, H. B., 1994, Zinc status of soils and petioles of vineyards of Peninsular India. *Agropedology.*, 4: 113-120.
- Bhargava, B. S. and Sumner, M. E., 1987, Proposal for sampling grape (*Vitis vinifera* L.) petioles for nutritional diagnosis. *Comm. Soil Sci. Plant Anal.*, 18(5): 581-591.
- Bhat, Z. A., Akther, F. A., Padder, S. A., Rehman, H. U. and Ganaie, A. Q., 2017, Nutrient status of grape orchard soils of Jammu and Kashmir, India. *International J. Agric. Sci. Res.*, 7:363-372.
- Black, C. A. (1965) Methods of soil analysis. Part 2, Agronomy monograph No.9, *Am. Soc. Agron.*, Madison, Wisconsin, USA.
- Ganai, A. Q., Dar, M. A., Chesti, M. H., Bhat, Z. A., Khanday, M. and Dar, N. A., 2018, Macronutrients status of apple (cv. Red Delicious) orchard soils of Jammu and Kashmir India. *Int. J. Chem. Stud.*, 6(2): 3314-3319.
- Jackson, M. L. (1973) *Soil Chemical Analysis*, Prentice Hall of India Pvt. Ltd., New Delhi, pp. 205.
- Kondi, A. E., 2016. Nutrient management practices in grape orchards of Bagalkot and Jamakhandi talukas: effects on soil fertility, petiole nutrient contents and yields. *MSc. Thesis.*, Univ. Hort. Sci, Bagalkot (India).
- Lindsay, W. L. and Norvell, W. A. (1978) Development of DTPA soil test for Zn, Fe, Mn and Cu. *J. Am. Soc. Soil Sci.*, 42: 421-428.
- Malakouti, M. J., 2007, Zinc is a neglected element in the life cycle of plants: A review. *Middle E Russ. J. Plant Sci. Biotechnol.*, 1:1-12.
- Nagaraja, M. S. (1997) Biomass turnover, nutrient status and biological processes in different land use systems *Ph.D. Thesis*, Univ. Agric. Sci., Dharwad.

Naraboli, V. C., 2016, Effect of nutrient management practices in grape orchards of Jamakhandi on soil properties, petiole nutrient content and grape yields. *M.Sc Thesis*, Univ. Hort. Sci, Bagalkot (India).

Patiram, R. C., Upadhyaya, C. S., Singh and Munnaram, 2000, Micronutrient cation status of Mandarin orchards of Sikkim. *J. Ind. Soc. Soil Sci.*, 48(2): 246-249.

Samiullah, Z. S., Tariq, M., Shah, T., Latif, A. and Shah, A., 2013, Micronutrients status of peach orchards in Swat valley. *Sarhad J. Agric.*, 29(4): 485-493.

Shivannavar, S. (2016) Effect of nutrient management practices in grape orchards of vijayapura on soil properties, petiole nutrient contents and grape yields, *M.Sc Thesis*, Univ. Hort. Sci., Bagalkot (India).

Thakre, Y. G., Choudhary, M. D. and Raut, R. D., 2013, Available micronutrient cation status of red soils in Wardha region, India. *J. Chem. Pharm. Res.*, 5(8): 64-68.

Yogeeshappa, H. (2007) Yield and quality of grapes (cv. Thompson Seedless) in relation to soil fertility status of vineyards in Bijapur taluk of Karnataka. *M.Sc Thesis*. Univ. Agric. Sci., Dharwad (India).

**Table 1: EDTA exchangeable calcium (Ca), magnesium (Mg) and available sulphur (S) status in soils of different vineyards before October pruning and after harvest**

Vineyard groups	Exchangeable calcium [cmol (p+) kg <sup>-1</sup> ] Mean ± SD		Exchangeable magnesium [cmol (p+) kg <sup>-1</sup> ] Mean ± SD		Available sulphur (mg kg <sup>-1</sup> ) Mean ± SD	
	Before October pruning	After harvest	Before October pruning	After harvest	Before October pruning	After harvest
Low yielding vineyards	10.17 ± 3.04 <sup>a</sup>	10.48 ± 2.46 <sup>a</sup>	5.59 ± 1.83 <sup>a</sup>	5.84 ± 1.54 <sup>a</sup>	26.45 ± 4.90 <sup>b</sup>	29.00 ± 5.33 <sup>c</sup>
Medium yielding vineyards	9.55 ± 2.87 <sup>a</sup>	10.12 ± 2.04 <sup>a</sup>	5.64 ± 1.93 <sup>a</sup>	5.94 ± 1.99 <sup>a</sup>	35.41 ± 5.58 <sup>b</sup>	40.34 ± 6.78 <sup>b</sup>

High yielding vineyards	11.16 ± 2.61 <sup>a</sup>	12.08 ± 2.76 <sup>a</sup>	6.81 ± 1.26 <sup>a</sup>	6.88 ± 1.19 <sup>a</sup>	51.67 ± 16.46 <sup>a</sup>	64.09 ± 18.59 <sup>a</sup>
<b>S.Em ±</b>	<b>0.90</b>	<b>0.77</b>	<b>0.55</b>	<b>0.51</b>	<b>3.30</b>	<b>3.74</b>
<b>C.D. at 5%</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>9.57</b>	<b>10.86</b>

Note: 1. Different letters in mean column imply significant difference at  $P \leq 0.05$

UNDER PEER REVIEW

**Table 2: DTPA extractable micronutrient status in soils of different vineyards before October pruning and after harvest**


Vineyard groups	Fe (mg kg <sup>-1</sup> ) Mean ± SD		Mn (mg kg <sup>-1</sup> ) Mean ± SD	
	Before October pruning	After harvest	Before October pruning	After harvest
Low yielding vineyards	2.63 ± 0.37 <sup>b</sup>	3.70 ± 0.52 <sup>b</sup>	3.28 ± 0.44 <sup>b</sup>	3.97 ± 0.33 <sup>c</sup>
Medium yielding vineyards	3.37 ± 0.72 <sup>a</sup>	3.96 ± 0.82 <sup>b</sup>	3.79 ± 0.68 <sup>a</sup>	4.41 ± 0.39 <sup>b</sup>
High yielding vineyards	3.85 ± 1.11 <sup>a</sup>	4.73 ± 0.96 <sup>a</sup>	4.21 ± 0.38 <sup>a</sup>	4.85 ± 0.45 <sup>a</sup>
<b>S.Em ±</b>	<b>0.25</b>	<b>0.25</b>	<b>0.16</b>	<b>0.12</b>
<b>C.D. at 5%</b>	<b>0.73</b>	<b>0.72</b>	<b>0.48</b>	<b>0.36</b>
Vineyard groups	Zn (mg kg <sup>-1</sup> ) Mean ± SD		Cu (mg kg <sup>-1</sup> ) Mean ± SD	
	Before October pruning	After harvest	Before October pruning	After harvest
Low yielding vineyards	0.67 ± 0.14 <sup>b</sup>	0.92 ± 0.11 <sup>b</sup>	0.33 ± 0.10 <sup>b</sup>	0.42 ± 0.11 <sup>b</sup>
Medium yielding vineyards	0.92 ± 0.27 <sup>a</sup>	1.11 ± 0.23 <sup>a</sup>	0.55 ± 0.34 <sup>b</sup>	0.64 ± 0.33 <sup>ab</sup>
High yielding vineyards	0.91 ± 0.15 <sup>a</sup>	1.12 ± 0.08 <sup>a</sup>	0.80 ± 0.30 <sup>a</sup>	0.87 ± 0.30 <sup>a</sup>
<b>S.Em ±</b>	<b>0.06</b>	<b>0.05</b>	<b>0.08</b>	<b>0.08</b>
<b>C.D. at 5%</b>	<b>0.18</b>	<b>0.14</b>	<b>0.24</b>	<b>0.24</b>

Note: 1. Different letters in mean column imply significant difference at  $P \leq 0.05$


**Table 3: Yields obtained in different vineyard groups**

<b>Vineyard Groups</b>	<b>Yield (t ha<sup>-1</sup>) Mean ± SD</b>
<b>LYV</b> : Low yielding vineyards	19.70 ± 1.58 <sup>c</sup>
<b>MYV</b> : Medium yielding vineyards	26.25 ± 1.02 <sup>b</sup>
<b>HYV</b> : High yielding vineyards	33.15 ± 2.69 <sup>a</sup>
<b>S.Em ±</b>	<b>0.60</b>
<b>C.D. at 5%</b>	<b>1.74</b>


Note: 1. Different letters in mean column imply significant difference at  $P \leq 0.05$



**Fig. 1: DTPA Exchangeable (a) calcium, (b) magnesium and (c) available Sulphur status in soils of different vineyard groups before October pruning and after harvest**



**Fig. 2: DTPA extractable (a) iron, (b) manganese, (c) zinc and (d) copper status in soils of different vineyards groups before October pruning and after harvest**



**Fig. 3: Yield of different vineyard groups.**