

# Post-harvest application and physicochemical properties of jamun (*Syzygium cuminii* L.)

1

## ABSTRACT

Jamun (*Syzygium cuminii* L.) is a subtropical fruit that comes under myrtaceae family originated from the Indian subcontinent. India is the second largest producer of jamun with an annual production of 2.1 million ton which contributes 15.4% of world production. Jamun is a highly perishable seasonal fruit (May–June). Hence, processing and preservation are essential to ensure the availability of its products throughout the year. Engineering and physicochemical properties of the fruit play a crucial role in the design and development of processing equipment, handling of food products and processing. The engineering properties of jamun fruit, including physical, mechanical, and frictional characteristics, were investigated in this study. Additionally, the physicochemical properties of jamun juice, such as pH, total soluble solids (TSS), acidity, total phenolic content, total anthocyanin content, antioxidant activity and vitamin C, were measured as per standard procedures. Jamun contains 80 percent pulp and nearly 20 percent seed. The physical properties of jamun fruit, including length, width, thickness, sphericity, bulk density, and porosity, were determined to be 22.22 mm, 21.93 mm, 20.99 mm, 0.947, 0.67g/cm<sup>3</sup>, and 32.48%, respectively. The firmness of the fruit was measured as 1.09 N and coefficient of friction of the fruit for different materials such as stainless steel, plywood, cardboard and glass were determined to be 1.25, 1.28, 1.56, 1.98, respectively. The physicochemical properties like moisture, pH, TSS, acidity, specific gravity, total phenolic content, antioxidant activity, total anthocyanin content and vitamin c were found to be 85.33 % (wb), 3.12, 13.14%, 0.81, 0.9, 256.76 mg/100ml, 86%, 1281 mg/100ml and 16 mg/100ml, respectively. These properties highlight its significant potential as a functional food and raw material for various food processing applications.

**Keywords:** Postharvest application, Jamun fruit, Jamun juice, Physicochemical properties

## 1. INTRODUCTION

Jamun (*Syzygium cuminii*) is an underutilized fruit tree native to the Indian subcontinent. It belongs to the family of myrtaceae. Jambul, black palm, java palm, and Indian blackberry are the common names of jamun. The promising values of phytochemicals, minerals and vitamins give a great therapeutic value for jamun. It is also referred as “Fruit for the Future” (Madani et al., 2021). Jamun has proven pharmacological properties on the animal system, including hypoglycemic, anti-inflammatory, antimicrobial, antioxidant, antidiarrheal, analgesic, astringent, and gastroprotective effects. Studies on

---

the use of jamun for diabetes have demonstrated positive results. (Jagetia, 2024; Kaur, 2024; Kumar et al., 2022).

India ranks second in jamun production, with an annual output of 13.5 million tonnes in 2013, contributing 15.4% to the world's total estimated production (Pharate et al., 2018). The top producer of jamun is Maharashtra, with Tamil Nadu, Gujarat, Assam, Uttar Pradesh, and other states following closely behind. The jamun tree is frequently grown on homesteads. Ram jamun is a common variety in north India. Other Jamun varieties such as Konkan bhadoli, madhura, Badama, Bhado, Narendra Jamun 6 Narendra, Goma Priyanka, CISH J-42 (Seedless type), CISH J-37 are some of the well-known varieties of jamun (Patil et al., 2011; Singh et al., 2019). From Tamil Nadu in the south to the Indo-Gangetic plains in the north, jamun is grown extensively throughout India's largest regions. Jamun can grow in the saline area also.

Anthocyanin is the bioactive compound that is rich in jamun (Tavares et al., 2016). Jamun is used as the raw material for the production of various value-added food products in many Asian countries like China, India etc due to its high nutritional values (Santhalakshmy et al., 2015). The processed food such as jam, jelly, chips, squash, wine, vinegar, beverages and pickles are made by utilizing the jamun fruit (Lago-Vanzela et al., 2011). Incorporating jamun fruit, juice, seed and powder, several value-added products were made. It provides a good source of nutrition for food security. (Rasheed et al., 2024).



**Fig. 1. Post-harvest processing of jamun fruit**

Jamun is a rich nutritious fruit but it is a seasonal one. The fruits are available during the months of May to June which differ with the genotype. In order to make the availability of jamun through the

year, processing and preservation are important (George et al., 2023). Understanding the raw material is very important for developing good processing techniques, methods and equipment. The physical characteristics of fruits are crucial for machinery design and food processing. They also play a key role in post-harvest handling, such as sorting, grading, packaging, storage, and transportation. (Athmaselvi et al., 2014). The biochemical properties are useful in the design and formulation of innovative food products (Sood et al., 2018).

The objective of the study is to examine the engineering properties of jamun fruit and the physicochemical properties of both the fruit and juice, with the aim of improving quality and enhancing post-harvest handling of jamun fruit.

## 2. MATERIAL AND METHODS

### 2.1 Sample collection

Jamun fruits were procured from local market of Malappuram of Kerala. The following flow chart shows the preparation of jamun juice from jamun fruit.



**Fig. 2. Processing of jamun juice**

### 2.2 Engineering properties

Engineering properties are important in the area of design and development of equipment of post-harvest unit operations of agricultural processing (Sahay & Singh, 2020). Engineering properties of biological materials are classified into physical properties, mechanical properties, frictional properties, physicochemical properties and nutritional properties etc.

#### 2.2.1 Physical properties

Physical properties of agriculture produce include shape, size, density, color, volume, surface area, porosity etc. were determined as per standard procedure.

### 2.2.1.1 Weight

Fruit weight was measured by an electronic balance with the accuracy 0.01 gram.

### 2.2.1.2 Axial dimensions

The axial dimensions are length, width and thickness. These dimensions of the fruits were measured by vernier caliper with least count of 0.02 mm.

### 2.2.1.3 Shape index (SI)

The shape index of the fruit was calculated by the following formula (Mohsenin, 1980):

$$SI = \frac{W+T}{2L} \quad \dots (1)$$

W- width (mm)

T- thickness (mm)

L- length (mm)

### 2.2.1.4 Geometric mean diameter (Dg)

Geometric mean diameter was calculated using the following formula mentioned by (Shousha et al., 2024):

$$Dg = (L * W * T)^{\frac{1}{3}} \quad \dots (2)$$

L - largest intercept

W - largest intercept perpendicular to l

T - largest intercept perpendicular each other

### 2.2.1.5 Arithmetic mean diameter (AMD)

The formula for calculation of AMD is as follows (Sasikumar et al., 2024):

$$Da = \frac{(L*W*T)}{3} \quad \dots (3)$$

### 2.2.1.6 Aspect ratio (AR)

The formula for calculation of AR is as follows (Dash et al., 2008):

$$AR = \left(\frac{W}{L}\right) \times 100 \quad \dots (4)$$

### 2.2.1.7 Elongation ratio

The formula for elongation ratio is as follows:

$$ER = \frac{T}{W} \quad \dots (5)$$

### 2.2.1.8 Sphericity

Sphericity is normally used to express the shape and size of the material. It is calculated using the following formula (Shousha et al., 2024):

$$\emptyset = \frac{(L*W*T)^{\frac{1}{3}}}{L} \quad \dots (6)$$

### 2.2.1.9 Surface area

The fruit surface area was measured by the following formula (Sasikumar et al., 2024);

$$S = \pi * Dg^2 \quad \dots (7)$$

### 2.2.1.10 Fruit volume

Platform scale method was used for measuring the volume (Sahay & Singh, 2020):

### 2.2.1.11 Density

Density of any material can be expressed as ratio of weight of the material to volume of the material.

#### 2.2.1.11.1 Bulk density

A 100 ml cylindrical glass container was filled entirely with jamun fruits in order to calculate the bulk density. Without crushing the fruits, the surplus jamun fruits were removed by repeatedly tapping the container. Using an electronic balance, weight was measured. The formula for bulk density follows (Işikli & Yılmaz, 2014).

$$\text{Bulk density (g/ cm}^3\text{)} = \frac{\text{Bulk mass (g)}}{\text{Total volume (cm}^3\text{)}} \quad \dots (8)$$

#### 2.2.1.11.2 True density

True density was determined using the toluene (C7H8) displacement method (Mohsenin, 1980; Shousha et al., 2024). The formula for true density is

$$\text{True density (g/ cm}^3\text{)} = \frac{\text{Mass of the individual fruit (g)}}{\text{Volume of individual fruit (cm}^3\text{)}} \quad \dots (9)$$

#### 2.2.1.11.3 Density ratio

Using the bulk density and true density measurements, the fruits' density ratio was determined (Işikli & Yılmaz, 2014).

$$D_r(\%) = \left( \frac{\rho_t}{\rho_b} \right) \times 100 \quad \dots (10)$$

$\rho_t$  = true density

$\rho_b$  = bulk density

#### **2.2.1.12 Porosity**

Porosity was calculated using the values of bulk density and true density. The formula for porosity is given by (Bakane et al., 2016):

$$\epsilon = \left( \frac{\rho_t - \rho_b}{\rho_t} \right) \times 100 \quad \dots (11)$$

#### **2.2.1.13 Colour**

The color was determined by colorimeter. It was expressed as L\*, a\*, b\* value (Sasikumar et al., 2024)

#### **2.2.1.14 Yield calculation**

##### **2.2.1.14.1 Pulp content**

The term "pulp content" describes the fruit's edible component. After the seeds were removed from the fruits, it was measured using the formula below (Ghosh et al., 2016).

$$\text{Pulp content (\%)} = \frac{\text{weight of the pulp (g)}}{\text{weight of the fruit (g)}} \times 100 \quad \dots (12)$$

##### **2.2.1.14.2 Juice content**

The quantity of juice from fruit is expressed in terms of juice content. The juice of fruit was extracted using a fruit pulper and juice content was calculated as per the formula mentioned below (Zeng et al. 2019).

$$\text{Juice content (\%)} = \frac{\text{weight of the juice (ml)}}{\text{weight of the fruit (g)}} \times 10 \quad \dots (13)$$

##### **2.2.1.14.3 Seed content**

The inedible portion of the fruit which was separated to make pulp was taken for the calculation (Babu et al., 2019).

$$\text{Seed content (\%)} = \frac{\text{weight of the seed (g)}}{\text{weight of the fruit (g)}} \times 100 \quad \dots (14)$$

#### **2.2.2 Mechanical property**

Using an electronic penetrometer, the firmness of the fruit was tested (Jantra et al., 2018). The meter's sharp edge was positioned on the fruit's surface. To create a rupture on the fruit's surface, it was gradually compressed. It was noted how much force was needed to cause a rupture.

#### **2.2.3 Frictional property**

The frictional properties are important for designing of conveyer and moving parts in processing machineries. The coefficient of friction of fruits were determined by the following formula (Ghosh et al., 2016):

$$\text{Coefficient of friction } (\mu) = \frac{F}{W} \dots (15)$$

F – frictional force required

W – weight of the fruits

### **2.3 Physicochemical properties**

The chemical properties like moisture content, pH, TSS, titratable acidity and nutritional content were determined for pulp and juice.

#### **2.3.1 Moisture content**

Moisture content was determined using the method given by (AOAC, 2012). 10g of jamun pulp was taken in a petri dish. The initial weight was noted. It was then placed in a hot air oven at the temperature of 105°C for 3 hrs. Weight of the samples were checked at frequent interval till the constant weight was reached. The moisture content was calculated by the following formula:

$$\text{Moisture content (wet basis \%)} = \frac{w_i - w_f}{w_i} \times 100 \dots (16)$$

$w_i$  – initial moisture

$w_f$  – final moisture

#### **2.3.2 pH**

The pH of the fruit sample was determined using a digital pH meter (AOAC, 2000). One gram of fruit juice was diluted to 10ml with water respectively. The pH was analysed with buffer (pH 4.7). Triplicate was taken from each sample.

#### **2.3.3 Acidity**

Acidity was determined by titrating sample with 0.1 N alkali (NaOH). One ml of fruit pulp, juice, and one gram of seed were diluted to 10 ml with water respectively and approximately 0.1 ml of phenolphthalein was added. It was titrated against 0.1 N NaOH solution until the pink colour appears. Three replications per each sample were taken. Results were expressed in terms of acid as gram of citric acid per gram (AOAC, 2000).

$$\text{Percentage acidity of pulp} = \frac{\text{ml of 0.1 N NaOH} \times 0.0064}{\text{volume of the sample} \times 100} \dots (17)$$

#### **2.3.4 Total soluble solids**

Total soluble solids were measured at 20°C with a hand refractometer and the results were expressed as °Brix (AOAC, 2005).

### **2.3.5 Specific gravity**

Specific gravity refers to the density difference of the substance. The formula for specific gravity follows:

$$\text{Specific gravity of juice} = \frac{\text{density of juice}}{\text{density of water}} \dots (18)$$

### **2.3.6 Bioactive compounds**

#### ***2.3.6.1 Total phenolic content***

The concentration of total phenols in the extracted samples were determined by using Folin-Ciocalteu Reagent (FCR). To an aliquot of 0.2ml of sample extracts, 2.5 ml of 1:1 FCR reagent was added and left for 5 mins. To that 2ml of unsaturated sodium carbonate solution (7.5% w/v) was added and incubated for 2 hrs at room temperature. Using a spectrophotometer, the absorbance was measured at 760 nm. The amount of total phenolics was determined and reported as mg GAE/100g FW. Calibration curve was made by using gallic acid (Wang et al., 2011).

#### ***2.3.6.2 Total anthocyanin content***

0.5 mL of extract was combined with 10 mL of two buffers: 0.025 M potassium chloride (pH 1.0) and 0.4 M sodium acetate (pH 4.5) and allowed to sit at room temperature for 15 minutes prior to taking absorbance measurements at wavelength of 520 and 700nm (Nguyen & Nguyen, 2018).

$$\text{Total anthocyanin content (mg/100 mL)} = \frac{A \times MW \times DF \times 1000}{\epsilon \times l} \dots (19)$$

where:

$$A = (A_{520\text{nm}} - A_{700\text{nm}})_{\text{pH } 1.0} - (A_{520\text{nm}} - A_{700\text{nm}})_{\text{pH } 4.5}$$

MW (molecular weight) = 449.2 g/mol for cyanidin-3-glucoside

DF = dilution factor established in D

$$l = 1 \text{ cm}$$

$\epsilon$  = 26,900 molar extinction coefficients, in L/mol cm, for cyanidin-3-glucoside

#### ***2.3.6.3 Antioxidant Capacity (DPPH)***

The antioxidant capacity was measured for juice. It was measured using 2,2-Diphenyl-1-picrylhydrazyl (DPPH) assay. 1 mL of the sample extract was mixed with 3 mL of 0.1 mM methanolic DPPH solution then it was vortexed and place at dark for 30 min at room temperature. The absorbance reading was 517 nm. UV-Visible spectrophotometer was used to measure the absorbance reading (Nguyen & Nguyen, 2018). Triplicate reading was taken for the determination. It was calculated as follows:

$$\text{DPPH scavenging effect (\%)} = \left(1 - \frac{A}{A_0}\right) * 100 \dots (20)$$

where

A0 - control solution absorbance

A - DPPH solution absorbance

#### 2.3.6.4 Ascorbic acid

Ascorbic acid present in fresh fruits can be calculated either volumetric method or colorimetric method. Volumetric method was adopted to estimate vitamin C content present in jamun fruit juice. 200 ml of dye solution was prepared in distilled water using 42 mg of sodium bicarbonate and 52 mg of 2, 6, dichloro indophenols. The dye is reduced to a colourless leucobase by ascorbic acid. The titrating medium, 4% oxalic acid, aids in identifying the titration's end point by giving it a pale pink hue. In an acidic media, the dye has the colour of pink. Stock standard solution was prepared by dissolving 100 mg of ascorbic acid in 100 ml of 4% oxalic acid in a standard flask (concentration 1 mg/ml). The working standard solution was obtained by diluting 10 ml stock solution to 100 ml with 4% oxalic acid (concentration 100 µg/ml) (Joy et al., 2015). Sample was prepared by 4% oxalic acid and titrated against the dye.

$$\text{Ascorbic acid mg/100g} = \frac{0.5}{V_1} \times \frac{V_2}{5} \times \frac{100}{V_s} \times 100 \quad \dots (21)$$

V1- titration value of blank

V2 – titration value of sample

Vs – weight of the sample

### 3. Results and discussion

The physicochemical characteristics of a fruit dictate its intended use, including its suitability for direct consumption, technological compatibility and processing potential all of which influence the quality of the final product (Najman et al., 2023). The weight of the fruit was in the range of 2.03g to 15.36g. The weight values indicated that Jamun had an uneven distribution of weight. According to the findings, the fruit's length was greater than its width and thickness. The measurements for the minimum and maximum length, width and thickness were 14.93 mm, 14.95 mm, and 14.12 mm, and 29.11 mm, 28.94 mm, and 28.72 mm, respectively. Varietal changes, species, rootstock utilised, edapho-climatic growing circumstances, and agro-inputs provided to the fruit-bearing trees may all contribute to variations in the physical characteristics of jamun fruit. Fruit weight and colour have an impact on the fruit's market value. The fruit's dimensions were used to determine its maturity (Panday et al., 2021). The geometric mean diameter of the fruit ranges from 14.27 mm at the minimum to 27.93 mm at the maximum. One of the fruit's key physical characteristics is its sphericity, which was measured to range from a minimum of 0.863 to a maximum of 1.0. The surface area was measured using the formula mentioned in the materials and methods section (section 2.2.1.9), with the lowest value being 639.6 mm<sup>2</sup> and the largest area measuring 2455.7 mm<sup>2</sup>. The fruit's volume was measured using the platform

scale method, which is used to calculate the volume of irregularly shaped materials. It was found that the lowest and greatest volumes of jamun fruits were 2.03 mm<sup>3</sup> and 15.06 mm<sup>3</sup>, respectively.

Bulk density was determined by weighing and measuring the volume, with the lowest and highest values being 0.56 g/cm<sup>3</sup> and 0.69 g/cm<sup>3</sup>, respectively. Porosity was measured using the formula mentioned in section 2.2.1.12, with the lowest and highest values being 30.31 and 33.52, respectively. The pulp percentage of the fruit ranged from 74.56% to 82.62%, while the seed percentage varied between 12.78% and 19.05%. The highest juice yield recorded was 50%. Firmness, a mechanical property, was measured using a penetrometer, with a recorded value of 1.09 N. The coefficient of friction was assessed for various materials, including glass, cardboard, stainless steel, and wood, with glass showing the highest value of 1.98. The fruit's moisture content was determined as wet basis, following standard procedures. The physicochemical properties and bioactive components of jamun juice, such as pH, total soluble solids (TSS), acidity, specific gravity, total phenolic content (TPC), total anthocyanin content (TAC), antioxidant activity (DPPH), vitamin C, and colour, were analysed according to standard protocols. The pH ranged from 3.12 to 3.29, TSS from 10 to 13, and acidity from 0.81 to 1.71. Akhila and Umadevi (2018) and Lavanya et al. (2018) were mentioned the similar findings. Table 4. revealed that jamun juice was high in anthocyanin and other bioactive compounds.

Consumer preferences for fruits and vegetables with consistent sizes and shapes are strongly influenced by dimensions, weight, and volume, thus these characteristics must be considered when designing effective grading systems (Afshari et al., 2023). From the table 1 jamun weight and axial dimensions like length, width and thickness were found to be 6.679 g and 22.22, 21.93 and 20.99, respectively. The fruit with dark pulp and an oblong shape was also reported in Kshirsagar et al. (2019). The findings were similar to (Suradkar et al., 2017; Ghosh et al., 2017; Shahnawaz and Sheikh, 2011). From the results of different studies, the weight and axial dimensions are varied according to genotypes. These measures offer important information about the fruit's physical attributes, which helps with the classification and comprehension of its applicability for different botanical or culinary uses. Additionally, this will provide inspiration for postharvest machinery design. For scholars and learners researching its morphology and possible uses, it also provides accurate data. The geometric and arithmetic mean diameters were also computed (Table 1). The surface area affects how gases, such as carbon dioxide and oxygen, are exchanged between the fruit and its surroundings, which is crucial for ripening and preserving freshness (Sasikumar et al., 2024).

Yield calculations of jamun fruit such as pulp content, seed content and juice content values are more or less similar to the findings of (Singh and Swamy, 2021; Gosh et al., 2017; Babu et al., 2019). Ningot et al. (2017) reported the pulp content of 53 to 87 %, respectively. The total phenolic content and antioxidant values were 256.76 mg/100 ml and 86%, respectively, whereas the anthocyanin content in jamun juice was 1824 mg/100 ml. Anthocyanin is the main compound for the fruit color. Ascorbic acid content was in the range of 16 to 18 mg/100ml, respectively. The colour values of juice such as L\*, a\* and b\* values were measured as 17.43, 25.57 and 8.57, respectively. The positive values of the colour indicated that jamun juice having lightness value. The values of the results are similar with

the previous studies. Suradkar et al. (2017) reported that jamun fruit contains ascorbic acid 21.48 mg/100 ml and antioxidant activity of 95.81%, respectively.

**Table 1. Physical properties of jamun fruit**

Properties	values
Weight (g)	6.679±0.92
Axial dimensions	
L(mm)	22.22±3.65
W(mm)	21.93±3.90
T(mm)	20.99±3.78
Shape index	0.96±0.05
Geometric mean diameter (mm)	21.04±3.55
Arithmetic mean diameter (mm)	21.72±3.70
Aspect ratio	0.98±0.06
Elongation ratio	0.95±0.04
Sphericity	0.947±0.03
Surface area ( $mm^2$ )	1427±487
Volume ( $mm^3$ )	7.28 ± 4.1
Bulk density ( $g/cm^3$ )	0.67±0.01
True density ( $g/cm^3$ )	1±0.002
Density ratio	1.48±0.04
Porosity (%)	32.48±1.41
Colour (Pulp)	
L*	13.13±0.75
a*	3.833±0.35
b*	-3.6±0.36
Pulp content (%)	79.20±2.1
Juice content (%)	50.20±5.1
Seed content (%)	17.23±1.7

**Table 2. Mechanical property of jamun fruit**

Property	Value
Firmness (N)	1.09±0.56

**Table 3. Frictional property of jamun fruit**

Materials	Co efficient of friction ( $\mu$ )
Stainless steel	1.25
Plywood	1.28
Cardboard	1.56
Glass	1.98

**Table 4. Physicochemical properties of jamun juice**

Properties	Juice
Moisture content (%)	85.33
pH	3.12 $\pm$ 0.01
TSS (%)	13.14 $\pm$ 0.2
Acidity	0.81 $\pm$ 0.07
Specific gravity	0.9 $\pm$ 0.0
Total phenolic content (mg/100ml)	256.76 $\pm$ 0.52
Total anthocyanin content (mg/100 ml)	1284.14 $\pm$ 1.66
Antioxidant activity (% DPPH)	86 $\pm$ 0.04
Ascorbic acid (mg/100 ml)	16 $\pm$ 1.25
Colour	
L*	17.43 $\pm$ 0.65
a*	25.57 $\pm$ 4.07
b*	8.57 $\pm$ 2.24

\* Moisture content on wet basis

#### 4. CONCLUSION

The study of the engineering and physicochemical properties of jamun fruit provides essential insights for the design and development of efficient food processing machines well as innovative food products. Understanding these properties, such as sphericity, bulk density, porosity, firmness, moisture content and bioactive components, facilitates the optimization of handling, processing and storage techniques. This knowledge supports the production of high-quality jamun-based products while ensuring minimal waste and improved processing efficiency. Consequently, these findings serve as a foundation for advancing value-added product development and promoting the utilization of jamun in

the food industry. These characteristics not only influence consumer preferences but also play a significant role in the processing, storage, and preservation of fruits. By optimizing factors such as ripening conditions and handling procedures, fruit quality can be improved, and post-harvest losses can be minimized.

## ACKNOWLEDGEMENTS

Author acknowledge the work which mentioned in this manuscript was done under the Kerala Agricultural University and All India Coordinated Research Project (AICRP).

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## AUTHORS' CONTRIBUTIONS

Sankar M<sup>1\*</sup> designed the study, wrote the protocol, and wrote the first draft of the manuscript. Rajesh G K<sup>2</sup> managed the analyses of the study. All authors read and approved the final manuscript.

## Disclaimer (Artificial intelligence)

### Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

## REFERENCES

1. Madani, B., Mirshekari, A., Yahia, E.M., Golding, J.B., Hajivand, S. & Dastjerdy, A.M. (2021). Jamun (*Syzygium cumini* L. Skeels): A promising fruit for the future. *Horticultural Reviews*, 48, 275-306.
2. Jagetia, G. C. (2024). Bioactive Phytoconstituents and Medicinal Properties of Jamun (*Syzygium cumini*). *Journal of Exploratory Research in Pharmacology*.
3. Kaur D, Yousuf B., & Qadri OS. (2024). *Syzygium cumini* anthocyanins: recent advances in biological activities, extraction, stability, characterisation and utilisation in food systems. *Food Production, Processing and Nutrition*, 6(1).
4. Kumar, M., Hasan, M., Lorenzo, J.M., Dhupal, S., Nishad, J., Rais, N., Verma, A., Changan, S., Barbhai, M.D., Chandran, D. and Pandiselvam, R. (2022). Jamun (*Syzygium cumini* L.) Skeels) seed bioactives and its biological activities: A review. *Food Bioscience*, 50, 102109.
5. Pharate, K., B. Shubham, K. Holkar, T. Katariya & M. Pharata. (2018). A Review on *Syzygium cumini* - A Valuable Fruit. *Journal of Ready to eat food*, 7, 527–534.

6. Patil, S. S., Thorat, R. M., & Rajasekaran, P. (2011). *Utilization of Jamun Fruit (Syzygium cumini) for Production of Red Wine*. [www.sospublication.co.in](http://www.sospublication.co.in)
7. Singh, S., Singh, A.K., Saroj, P.L. & Mishra, S. (2019). Research status for technological development of jamun (*Syzygium cumini*) in India: A review. *The Indian Journal of Agricultural Sciences*, 89(12), 1991-8.
8. Tavares, I. M. de C., Lago-Vanzela, E. S., Rebello, L. P. G., Ramos, A. M., Gómez-Alonso, S., García-Romero, E., Da-Silva, R., & Hermosín-Gutiérrez, I. (2016). Comprehensive study of the phenolic composition of the edible parts of jambolan fruit (*Syzygium cumini* (L.) Skeels). *Food Research International*, 82, 1–13
9. Santhalakshmy, S., Bosco, S.J.D., Francis, S. & Sabeena, M. (2015). Effect of inlet temperature on physicochemical properties of spray-dried jamun fruit juice powder. *Powder Technology*, 274, 37-43.
10. Lago-Vanzela, Ellen Silva, Ginaldo Vieira dos Santos, Fernanda Arcaro de Lima, Eleni Gomes, & Roberto da Silva. (2011). Physical-chemical, caloric and sensory characterization of light jambolan (*Syzygium cumini* Lamarck) jelly. *Food Science and Technology*, 31, 666-673.
11. Rasheed, H. U., Shoukat, U., Arshad, U., Afzal, S., Khan, M. S., Aleem, S., Fatima, S., Noman, M., Hasan, R. W., & Arshad, M. A. (2024). Adaptation and Agricultural Significance of *Syzygium cumini* L. in Saline Environments: A Global Perspective on Jamun Cultivation and Salt Stress Resilience. *Haya: The Saudi Journal of Life Science*, 9(05), 172–187.
12. George, S., Thomas, A., Kumar, M.V.P., Kamdod, A.S., Rajput, A. & Abdullah, S. (2023). Impact of processing parameters on the quality attributes of spray-dried powders: a review. *European Food Research and Technology*, 249(2), 241-257.
13. Athmaselvi, K.A., Jenney, P., Pavithra, C., & Roy, I. (2014). Physical and biochemical properties of selected tropical fruits. *International Agrophysics*, 28(3).
14. Sood, M., Julie Bandral, I. D., Manmeet Kaur, I., Correspondence Monika Sood, I., Bandral, J. D., & Kaur, M. (2018). Development and quality evaluation of jamun seed powder supplemented noodles. *Journal of Pharmacognosy and Phytochemistry*, 7(3).
15. Sahay, K.M. & Singh, K.K. (2020). *Unit operations of agricultural processing*, Vikas publishing house.
16. Mohsenin, N.N. (1980). *Physical properties of plant and animal materials*. Gordon and Breach Science Publishers, New York.
17. Shousha, E.R., Tayel, S.A. & Ghanem, T.H. (2024). Pomegranate fruit mass modeling with some physical properties. *Al-Azhar Journal of Agricultural Engineering*, 6(1).
18. Sasikumar, R., Vivek, K., Kadirvel, G. & Jaiswal, A.K. (2024). Analysis of physicochemical and phytonutrients properties of bastard oleaster fruits and its mass prediction using artificial neural network model. *Journal of Agriculture and Food Research*, 101224.
19. Dash, A., Pradhan, R., Das, L., & Naik, S. (2008). Some physical properties of simarouba fruit and kernel. *International Agrophysics*.
20. Işıklı ND. & Yılmaz I. (2014). Some physical properties of sun-dried Berberis fruit (*Berberis crataegina*). *Journal of Food Science and Technology*. 51(1), 104-10.

21. Bakane PH, Gajabe M, Khedkar MB & Khakare M. (2016). Physical properties of Jamun Fruit (*Syzyium cumini* L.), *Journal of Food Research and Technology*, 4(2),52-56.
22. Ghosh, P., PRadhaN, R.C., Mishra, S., Patel, A.S. & Kar, A. (2017). Physicochemical and nutritional characterization of jamun (*Syzygium cuminii*). *Current Research in Nutrition and Food Science Journal*, 5(1),25-35.
23. Zeng, D., Xiao, G., Xu, Y., Zou, B., Wu, J. & Yu, Y., (2019). Protein and polyphenols involved in sediment formation in cloudy litchi juice. *Food Science and Biotechnology*, 28(4),945-953.
24. Babu, P., Raghavendra, S. & Tamadaddi, B.L. (2019). Physico-chemical characterization of Jamun (*Syzygium cumini* Skeels) seedling genotypes. *Journal of Pharmacognosy and Phytochemistry*, 8(4),300-303.
25. Jantra, C., Slaughter, D.C., Roach, J. & Pathaveerat, S. (2018). Development of a handheld precision penetrometer system for fruit firmness measurement. *Postharvest Biology and Technology*, 144,1-8.
26. AOAC. (2012). Official method of analysis. (19th edn.) The Association for Official Analysis in Chemistry, Rockville, USA.
27. AOAC. (2000). Official Methods of Analysis. Association of Official Analytical Chemists. Washington, D.C. 16th edition.
28. AOAC. (2005). Official Methods of Analysis. Association of Official Analytical Chemists. Washington, DC.
29. Wang, Zhenbin, Pan, Zhongli, Ma, Haile, Atungulu, & Griffiths. (2011). Extract of Phenolics From Pomegranate Peels. *The Open Food Science Journal*,5,17-25.
30. Nguyen CL, & Nguyen HVH. (2018). Ultrasonic effects on the quality of mulberry juice. *Beverages*,4(3).
31. Joy, P. P., Surya, S., & Aswathy, C. (2015). Laboratory manual of biochemistry. Pineapple research station, Kerala Agricultural University, Vazhakkulam,91-133.
32. Najman, K., Adrian, S., Sadowska, A., Świąder, K., Hallmann, E., Buczak, K., Waszkiewicz-Robak, B. & Szterk, A. (2023). Changes in physicochemical and bioactive properties of quince (*Cydonia oblonga* Mill.) and its products. *Molecules*, 28(7),3066.
33. Pandey, L., Mogra, R. & Goel, A. (2021). Assessment of physico-chemical properties of antioxidant rich fruits and sensory attributes of jamun based RTS beverages. *International journal of chemical studies*, 9(1), 404-409.
34. Akhila H. & Umadevi SH. (2016). Storage effects on vitamin c, anthocyanins, total phenolic and antioxidant capacity of processed jamun fruit products. *International Journal of Pure and Applied Bioscience*,6(6),1335-1347.
35. Lavanya T, Raj D. & Vaghashiya JM. (2018). Standardization of formulation for preparation of health drink by blending Aloe vera, Guava and Jamun. *International Journal of Chemical Studies*, 6(4), 1715-1721.
36. Afshari, K., Javanmard Dakheli, M., Ramezan, Y., Bassiri, A., & AhmadiChenarbon, H. (2023). Physicochemical and control releasing prop -erties of date pit (*Phoenix dactylifera* L.) phenolic

- compounds microencapsulated through fluidized-bed method. *Food Science and Nutrition*, 11(3), 1367–1382
37. Kshirsagar, R.B., Desai, G.B., Sawate, A.R. & Deshmukh, N.M. (2019). Physico-chemical and nutritional properties of jamun (*Syzygium cumini*) seed. *Journal of Pharmacognosy and Phytochemistry*, 8(2), 211-213.
38. Suradkar, N.G., Pawar, V.S. & Shere, D.M. (2017). Physicochemical, proximate and bioactive composition of jamun (*Syzygium cumini* L.) fruit. *International Journal of Chemical Studies*, 5(3), 470-472.
39. Shahnawaz, M. & Sheikh, S.A. (2011). Physicochemical characteristics of Jamun fruit. *Journal of Horticulture and Forestry*, 3(10), 301-306.
40. Singh, Y.S. & Swamy, G.S.K. (2021). Genetic variation for morphological and physico-chemical traits in Jamun (*Syzygium cumini* Skeels). *International Journal of Minor Fruits, Medicinal and Aromatic Plants*, 7 (2), 73-78
41. Ningot EP, Dahale MH, Bharad SG, Nagre PK. (2017). Studies on variability in physico-chemical characteristics of jamun (*Syzygium cumini* Skeels) genotypes from eastern Maharashtra. *Life Science International Research Journal*, 4(1):215-217.