

# Accessibility on utilization of contraceptive methods among youths: an analytical cross-sectional study in Westlands Sub County, Nairobi County.

## Abstract

**Background:** A youth is a person aged between 15 to 24 years. Contraception methods are methods used to prevent unwanted pregnancies and spacing of pregnancies. Information on contraception is abundant among the population but the uptake of contraception is still low, especially among the youth. The study sought to examine how accessibility affects the utilization of contraceptive methods among the youths in the Westlands Sub-County, Nairobi County, Kenya.

**Methods:** The study employed a cross-sectional design and generated both qualitative and quantitative data. 13 health centers in Westland's Sub County were randomly selected and youths visiting the facilities for services were sampled using both systematic and simple random sampling. For this study, 398 participants made up the sample size.

**Results:** The research found a significant correlation ( $p=0.00$ ) between youths' use of modern contraceptive methods and the quality of services they received. From this study, the number of children ( $p=0.503$ ) did not significantly add to the model, but the academic level ( $p=0.003$ ), marital status ( $p=0.021$ ), and gender ( $p=0.001$ ) did.

**Conclusions:** In conclusion utilization of any method of contraceptives translated to a high contraceptive utilization prevalence in youths, the quality of services that health care providers offer to youths, education level, and demographic factors such as gender, and marital status, have continued to greatly affect and lower overall contraceptive utilization prevalence among the youths in developing nations.

**Keywords:** Contraceptives, Reproductive Health, Unwanted Pregnancies, STDs.

## 1.Introduction

A youth is the time between childhood and adulthood and the person is considered young. [1] Globally, the prevalence of contraceptive uptake has been on the increase. The increase in contraceptive uptake is attributed to the development of modern contraceptives and increased coverage and development of family planning and reproductive health programs. [2] Contraceptive use has enabled women especially the youth to postpone, space, limit, and prevent unwanted pregnancies. Contraceptive services are important not only to maternal and child health but also to meet the targets set by the Sustainable Development Goals (SDG). [3] Achieving health and promoting the well-being of people at all ages is the third Sustainable Development Goal (SDG). Young people's contraceptive behavior has positively changed in Africa over the past thirty years. Among Africans, youths under 25 make up 64% of the population, and adolescents make up one in every ten Kenyans.[4] The youths normally experience gradual attraction towards heterosexual relationships which eventually leads to sexual activity.

In Kenya, for instance, 52% of the youths aged between 15 to 25 years are sexually active. In addition, 39% and 65% of unmarried males and females respectively are sexually active in Kenya.[5] Statistics indicate that the utilization of contraceptives is low in the Westlands, even with the growing adoption of this method of birth control.[6] This could be directly associated with the clients' contraceptive utilization. It is against this background that the area was selected for study.

The biggest challenge that youths face is the lack of health services that focus on the needs and desires of young people as a priority. The experiences of adults and their points of view are different from those of youths. To effectively ensure that health services reach the young people youth youth-friendly services should be established to encourage the youth to be advocates and champions of their own social and health welfare.[7] Unwanted pregnancies, unsafe abortions, and sexually transmitted diseases among young people have been reported to be the contributing factors to the increase in morbidity and mortality in

developing countries.[8] Annually an estimated 16 million adolescents become pregnant with about three million of them undergoing unsafe abortion. In addition, adolescents are at risk of pregnancy-related complications as compared to other reproductive-age women.[10]

Youths' access to family planning services and information on reproductive and sexual health issues is restricted in many African and Kenyan communities that specifically devalue youth sexual activity. Persons, communities, and the local and global health systems are impacted by the restricted access to reproductive wellness services and information.[11] Therefore, the differences in access to reproductive health services between people of varied socio-economic status, and those living in urban and rural areas are a global equity issue of importance.[12] Regardless of the extensive promotion of availability and access to health services especially reproductive health globally the disparities among communities especially among young people still remain more so in those living in resource-limited areas.[13] The objective of the study was to assess the accessibility on utilization of contraceptive methods among youths in Westlands Sub County, Nairobi County, Kenya.

## **2. Materials and Methods**

### **2.1 Study design**

A cross-sectional descriptive design was used for this research, which produced quantitative data on the factors influencing youths in Nairobi County's Westlands Sub-County's use of contraceptive services.

### **2.2 Sampling technique**

Health facilities were sampled randomly therefore Westlands Health Center, Gichagi Dispensary, Lower Kabete Dispensary, Mji wa Huruma Dispensary, Karura Health Center, Kangemi Health Centers, MSK Kangemi, Amurt Health Center, Eagle Health Center, Chiromo Medical Center, Medanta Africare Medical center and Avenue Hospital were considered in the research. In each health center, young people between the ages of 15 and 24 were chosen through systematic random sampling. The possibility of bias from human selection was minimized by using systematic random sampling. The participants were selected as they attended the selected health centers for the services until the desired number of 398 was reached.

### **2.3 Study population**

The study targeted youths aged between 15 to 24 years who were residents of Westlands and sought services at health centers in Westland Sub-County when the study was conducted.

### **2.4 Sample size determination**

The size of the sample for the present investigation was determined using Sloven's formula, which led to the recruitment of 398 study participants.

### **2.5 Data collection**

Structured questionnaires were used to gather primary data. Data from youths enrolled in particular healthcare facilities was gathered using the questionnaires. There were five sections on the questionnaire, some of which asked for demographic information and others which focused on the objectives of the research.

### **2.6 Data analysis**

In the present investigation, descriptive statistics were used to analyze the data. Data analysis was done using the Statistical Packages for Social Science (SPSS) version 23. When coding with the designated coding scale, statistical values were produced. A chi-square test for independence was used to assess the relationship between the independent and dependent variables. Percentages and frequencies were used in descriptive statistics.

Tables were utilized to help create the discussions and provide a clearer explanation of the findings.

### **2.7 Ethical consideration**

The Institutional Research Ethics and Review Committee (IREC) at Mount Kenya University (MKU) granted ethical approval for the research to be conducted. The National Commission for Science, Technology, and Innovation (NACOSTI) was consulted in order to obtain permission to conduct the research. Permissions from the Westlands sub-county health department were also acquired. The respondent's informational discretion was maintained. The researcher asked study participants and emancipated minors to sign informed written consent forms, and respondents' involvement in the investigation was entirely voluntary. Psychosocial support was provided to reduce the possibility of stigmatization by directing the impacted party to a reputable hospital psychologist for suitable care and assistance. Because those who participated were assigned identification numbers rather than names, their anonymity was strictly maintained.

### 3 Results

#### 3.1 Response rate

The study targeted 431 youths living in Westlands Sub County visiting various sampled health facilities for contraceptive services. Out of these 398 questionnaires were filled and considered for analysis, representing a response rate of 92.3% and a non-response rate of 7.7% (33 respondents) occasioned by refusals and withdrawal from the study.

#### 3.2 Level of awareness on the use of contraceptive methods among youths

To assess the level of awareness of the use of contraceptive methods among youths, the variable measuring academic level was used. The total percentages of the academic level categories were used to indicate the shared proportion of the usage of contraceptive methods within each of the combinations as shown in table 1 below. The study established that 41% of the participants had attained primary-level education. In addition, 11% of youths with primary-level education are currently using modern contraceptives. In addition, among the sampled youths to participate in the study 30% who indicated to have primary level education did not use any modern contraceptive method. 28% of the youths who participated in the study had secondary-level education. Among the youth with secondary education, only 19% were using modern contraceptives compared to 9.40% who were not using any modern method of contraception. 31% of the sampled youths reported to have attained tertiary level of education. Among those with tertiary education, 25% were using modern contraceptive methods as compared to 6% who were not using any method.

**Table 1. Level of Education and use of modern contraceptives**

			Yes	No	Total
Level of Education	Primary	Expected Count	43	118	162
		% of Total	11%	30%	41%
	Secondary	Expected Count	75	37	112
		% of Total	19%	9%	28%
	Tertiary	Expected Count	100	25	125
		% of Total	25%	6%	31%
Total	Expected Count	218	180	398	
	% of Total	55%	45%	100%	

#### 3.3 Accessibility and utilization of contraceptive methods among youth

As indicated in Table 2, the test is statistically not significant,  $p\text{-value} = 0.609 > 0.05$  (ns). This implies that the utilization of the methods of contraceptives is not associated with the accessibility of the services the youths seek from the health facilities they visit.

**Table 2. Relationship between health facility accessibility and contraceptive use**

	Value	df	Asymptotic Significance (2-sided)	Exact Sig. (2-sided)	Exact Sig. (1-sided)
Pearson Chi-Square	.262 <sup>a</sup>	1	0.609		
Continuity Correction <sup>b</sup>	0.15	1	0.698		
Likelihood Ratio	0.262	1	0.609		
Fisher's Exact Test				0.622	0.35
Linear-by-Linear Association	0.261	1	0.61		
N of Valid Cases	398				

### 3.4 Quality of service and utilization of contraceptives among youths

The study evaluated whether there existed any relationship between the utilization of modern contraceptives and the quality of services that youths receive when seeking services for their health concerns, the Chi-square test was used having its procedure fulfilled. Based on the Chi-square Tests in Table 3 results, the test is Statistically noteworthy,  $p < 0.05$ . This implies that the quality of services youths receive is correlated significantly to the utilization of modern methods of contraceptives by youths.

**Table 3. Relationship between quality of service and use of contraceptives**

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	83.740 <sup>a</sup>	3	0.000
Likelihood Ratio	89.875	3	0.000
Linear-by-Linear Association	81.753	1	0.000
N of Valid Cases	398		

### 3.5 Influence of demographic factors on utilization of modern contraceptives among youths

Youths' use of contraceptives is thought to be impacted by a complex interplay of numerous demographic factors, such as gender, educational attainment, marital status, and number of children. The relationship or

combined impact of these four variables on the use of contemporary contraceptive methods was examined. To find out how these demographic characteristics affected a participant's likelihood of currently using any form of contraception, a logistic regression analysis was carried out. The results of the study from the Hosmer and Lemeshoe Test indicate that the logistics regression model is statistically significant,  $\chi^2(7) = 4.673$ ,  $p\text{-value} = 0.700$ .

From the findings of this study, As indicated in Table 4 below, Academic level ( $p=0.003$ ), Status of marriage ( $p=0.021$ ), and gender ( $p=0.001$ ) were added significantly to the model, but the number of children ( $p=0.503$ ) did not add significantly to the model. Higher levels of education were associated with an increased likelihood of a youth utilizing any method of contraceptives. As the youths move from one aspect of marital status, that is from being single to married, divorced, or widowed, in this order the likelihood of utilization of any methods of contraceptives decreases. Youths of female gender were 3.47 times more likely to be utilizing any method of contraceptives.

**Table 4: Influence of demographic factors on utilization of modern contraceptives among youths**

	<b>B</b>	<b>S.E.</b>	<b>Wald</b>	<b>df</b>	<b>Sig.</b>	<b>Exp(B)</b>
Academic level (1)	0.766	0.342	65.253	1	0.003	2.063
No of children (1)	0.002	0.261	0.449	1	0.503	1.785
Step 1 <sup>a</sup>						
Marital status (1)	-1.106	0.558	14.225	1	0.021	0.122
Gender (1)	1.907	0.566	2.571	1	0.001	3.477
<b>Constant</b>	<b>20.38</b>	<b>602.711</b>	<b>0.253</b>	<b>1</b>	<b>0.619</b>	<b>52964.05</b>

#### 4. Discussion

This research was designed to study the aspects influencing the use of contraceptive methods among the youths in Westlands, Nairobi County, Kenya. Factors that were investigated in this study included accessibility of the health facilities and services that youths seek, the quality of services youth receive when seeking services for their health concerns, education status which may be limiting their knowledge on utilization of methods of contraceptives, and demographic factors such as gender, marital status and number of children one has or desires to have. The sample of this study comprised 64.1% female participants and 35.9% male participants. A higher proportion of female participants was found to be utilizing any method of contraceptives compared to males at the time of this study.

The accessibility factor was not a significant factor affecting the utilization of the method of contraceptives. 57.1% of the youths who participated in this study accessed the health facilities they visited by walking, out of which 54.1% took less than one hour to arrive. 22.9% of those walking to the health facilities were currently using any method of contraceptives. The increased likelihood of a youth utilizing any method of contraceptives. As the youths move from one aspect of marital status, that is from being single to married, divorced, or widowed, in this order the likelihood of utilization of any methods of contraceptives decreases. Youths of female gender were 3.47 times more likely to be utilizing any method of contraceptives.

The proportion of those using cars to get to the health facilities was 42.9% with 39.0% taking less than one hour to arrive and 31.8% currently using any method of contraceptives. The research findings deviate from those of an Ethiopian study that found women who lived in remote areas far from medical facilities had lower access to and usage of contraceptives.[14] The two studies' different settings may have contributed to the disparity in the results.

The education status of the youths was a significant factor indicating lower levels of education are associated with low utilization of methods of contraceptives. The study established that of the youths who participated in the study, 25.0% had a tertiary level of education and were currently using any method of

contraceptives while only 6.3% at this level were not using any method of contraceptives. Only 10.9% of those with a primary level of academic were utilizing any method of contraceptives and 29.7% were not using any method of contraceptives. Additionally, out of the 45.3% of the entire sample not utilizing any method of contraceptive, 29.7% of them had a primary or lower level of academic. This associated youths with lower levels of education with low contraceptive prevalence in their lives.

This study's results are comparable to those of a study done in South Kivu, the Democratic Republic of the Congo, which found that while some or completed primary school was not significantly associated with contraception use, having some secondary education was significantly associated with contraception use (AOR 1.77 [95% CI 1.18–2.67]). According to the results of another Ethiopian study, women who were between the ages of 24 and 49 and who had completed at least their primary education were more inclined to utilize contraceptives. In addition, the study concluded that having a husband of secondary education and above also increased the chances of a woman using contraceptive services.[15]

The quality of services offered in health facilities by the health care providers formed a very crucial aspect of studying how the utilization of methods of contraceptives could be affected if this factor is considered. Based on the results of this study, substantial evidence was obtained associating the good quality of services that generally youths receive in any health center they visit with higher percentages of youths utilizing methods of contraceptives. The results of this study sample, 25.0% of youths using any method of contraceptives rated the services they receive as very good and only 6.3% were not using and rated the services as very good. Out of the 20.3% of the participants who rated the services they receive as very poor, 17.2% were not currently using any method of contraceptives and 3.1% were found to be using any method of contraceptives. This shows that more youths receiving quality services translated to high contraceptive prevalence.

The results of the research were similar to those of an investigation done in Senegal, which found that the quality of services had a highly significant impact on youths' use of contraceptives ( $p=0.000$ ).[16] In addition, research in Nigeria indicated that the majority of the participants who rated the quality of contraceptive services as good were using contraceptives and were adherent to the contraceptive re-visits. [17] The study differed from a study conducted in Uganda that indicated that most youths using contraceptives did not care much about the quality of services. [18] The study concluded that the majority of the participating adolescent girls focused more on the benefits of contraceptive use such as prevention of pregnancy and STIs.

## **5. Conclusion**

Generally, for utilization of any method of contraceptives to translate to high contraceptive prevalence in youths, the quality of services that health care providers offer to youths, education level, and demographic factors such as gender, and marital status, have continued to greatly affect and lower overall contraceptive prevalence among the youths. In this study, some factors offered substantial evidence and could be associated with a high likelihood that youths were not currently using any modern method of contraceptives.

## **6. Recommendations**

There is a need for healthcare providers to improve the quality of services they offer while focusing on areas of confidentiality and privacy, adequate information, and counseling regarding contraceptive methods in addition a more targeted campaign should be started to target different segments of the youthful population than has low access and utilization of contraceptives. Youth-friendly services and the quality of services in contraceptive clinics should be improved and made more accessible to improve the utilization of contraceptive services among youths.

## References

1. Aemro E, Abdo M, Deksisa A, Alemayehu A, Mulatu T, Ahmed Hassen T, et al. Immediate postpartum intrauterine contraceptive device utilization and associated factors among women who gave birth in public health facilities of Adama town, Ethiopia. *SAGE Open Med* [Internet]. 2022 Jan 1 [cited 2024 Feb 13];10. Available from: <https://journals.sagepub.com/doi/10.1177/20503121221142412?icid=int-sj-abstract.similar-articles.6>
2. Ahmed NMAEH, Mostafa RH, Abuelghar W, Elbishry G. Discontinuation Rates among Copper Intrauterine Device Users in Primary Healthcare Unit and University Clinic. Is There a Difference? *Egypt J Hosp Med*. 2018 Jul 1;72(11):5658–65.
3. Anguzu R, Sempeera H, Sekandi JN. High parity predicts use of long-acting reversible contraceptives in the extended postpartum period among women in rural Uganda. *Contracept Reprod Med*. 2018 Dec;3(1).
4. Adedini SA, Babalola S, Ibeawuchi C, Omotoso O, Akiode A, Odeku M. Role of Religious Leaders in Promoting Contraceptive Use in Nigeria: Evidence From the Nigerian Urban Reproductive Health Initiative. *Glob Heal Sci Pract* [Internet]. 2018 Oct 10 [cited 2024 Feb 12];6(3):500. Available from: [/pmc/articles/PMC6172128/](https://pubmed.ncbi.nlm.nih.gov/31111111/)
5. Kamuyango A, Hou WH, Li CY. Trends and contributing factors to contraceptive use in Kenya: A large population-based survey 1989 to 2014. *Int J Environ Res Public Health*. 2020 Oct 1;17(19):1–12.
6. KDHS. Kenya Demographic and Health Survey 2022. 2022; Available from: <https://dhsprogram.com/pubs/pdf/PR143/PR143.pdf>
7. Animen S, Lake S, Mekuriaw E. Utilization of intra uterine contraceptive device and associated factors among reproductive age group of family planning users in Han Health Center, Bahir Dar, North West Amhara, Ethiopia, 2018. *BMC Res Notes*. 2018 Dec 22;11(1).
8. Tefera LB, Abera M, Fikru C, Tesfaye DJ. Utilization of Immediate Post-Partum Intra Uterine Contraceptive Device and Associated Factors: A Facility based Cross Sectional Study among Mothers Delivered at Public Health Facilities of Sidama Zone, South Ethiopia. *J Pregnancy Child Heal*. 2017;04(03).
9. Khatri B, Khadka A, Amatya A, Shrestha SM, Paudel R. Perception And Use Of Intrauterine Contraceptive Devices (IUCD) Among Married Women Of Reproductive Age In Bhaktapur, Nepal. *Open Access J Contracept* [Internet]. 2019 Nov [cited 2024 Feb 12];10:69. Available from: [/pmc/articles/PMC6886550/](https://pubmed.ncbi.nlm.nih.gov/31111111/)
10. Franklin SG, O'Neal M, Arneus A, Colvin C, Aung M, Jolly PE. Effectiveness of an intrauterine device informative intervention among post-natal women in Western Jamaica. *Reprod Health*. 2021 Dec 1;18(1).
11. Wolde TF, AYana K, Bekele F. Determinants of Intrauterine Contraceptive Device Discontinuation Among Women Using Family Planning, in Southwest Ethiopia: Unmatched Case–Control Study. *Open Access J Contracept*. 2022 Apr;Volume 13:39–47.
12. Ajayi AI, Nwokocha EE, Adeniyi OV, Ter Goon D, Akpan W. Unplanned pregnancy-risks and use of emergency contraception: a survey of two Nigerian universities. *BMC Heal Serv Res*. 2017 Jun 2;17(1):382.
13. Kabagenyi A, Reid A, Ntozi J, Atuyambe L. Socio-cultural inhibitors to use of modern contraceptive techniques in rural Uganda: a qualitative study. *Pan Afr Med J*. 2016;25.
14. Ontiri S, Ndirangu G, Kabue M, Biesma R, Stekelenburg J, Ouma C. Long-Acting Reversible Contraception Uptake and Associated Factors among Women of Reproductive Age in Rural Kenya. *Int J Environ Res Public Health* [Internet]. 2019 May 1 [cited 2023 Oct 4];16(9). Available from: [/pmc/articles/PMC6539670/](https://pubmed.ncbi.nlm.nih.gov/31111111/)
15. Safari W, Urassa M, Mtenga B, Changalucha J, Beard J, Church K, et al. Contraceptive use and discontinuation among women in rural North-West Tanzania. *Contracept Reprod*

- Med. 2019 Dec;4(1).
16. Haakenstad A, Angelino O, Irvine CMS, Bhutta ZA, Bienhoff K, Bintz C, et al. Measuring contraceptive method mix, prevalence, and demand satisfied by age and marital status in 204 countries and territories, 1970–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet*. 2022 Jul 23;400(10348):295–327.
  17. Kungu W, Agwanda A, Khasakhala A. Prevalence of and factors associated with contraceptive discontinuation in Kenya. *African J Prim Heal Care Fam Med* [Internet]. 2022 [cited 2023 Oct 4];14(1). Available from: </pmc/articles/PMC9210174/>

UNDER PEER REVIEW