

Evaluation of red wine grape varieties for growth and yield parameters in semi-arid condition of India

Abstract

The study was carried out at the ICAR-National Research Centre for Grapes, Pune, during the 2022–2023 and 2023–2024 fruiting seasons. Among the red wine grape varieties, **Cabernet Sauvignon** demonstrated superior growth parameters, including pruned biomass (1.080 kg/vine). The variety **Syrah** exhibited the highest shoot length (120.44 cm), shoot diameter (9.75 mm), number of leaves per fruiting shoot (25.61), and number of bunches per vine (60.50). **Grenache Noir** showed the shortest time to bud sprout (8.93 days) and the highest number of berries per bunch (177.47), while **Petit Syrah** recorded the maximum internodal length (6.94 cm). The **Caladoc** variety displayed a higher number of leaves (24.42) and the largest leaf area per fruiting shoot (2931.25 cm²), whereas **Nielluccio** had the highest individual leaf area (141.25 cm²). **Cabernet Franc clone** had the highest chlorophyll content (13.52 mg/ml). Yield-related parameters highlighted **Aliquate Bauschet** for its highest bunch weight (264.80 g) and yield (8.84 kg/vine and 10.70 MT/acre on a pooled mean basis), while **Red Muscat** had the highest 100-berry weight (221.94 g).

Introduction

Grapes (*Vitis* spp. L.) The *Vitis* genus are believed to have originated around 54 million years ago, but humans only began domesticating and cultivating ~~them~~ grapes approximately 6,000 years ago (McGovern, 2003). Grapes hold a significant position in the fruit industry both in terms of cultivation area and economic value, and they are extensively grown in subtropical and temperate climates. Although grapes are fundamentally a temperate-origin crop, they are primarily cultivated in subtropical and tropical agro-climatic conditions (Ghule *et al.*, 2021; Somkuwar *et al.*, 2021). In India, roughly 98% of grape cultivation is dedicated to table grapes or raisin production, while only about 2% of the total grape yield is utilized for juice and winemaking (Ausari *et al.*, 2024).

According to the International Organization of Vine and Wine (OIV), wine is a food product produced exclusively by the complete or partial alcoholic fermentation of fresh grapes or their must, either pressed or unpressed. Chemically, wine is a complex beverage consisting of water, ethanol, sugars, amino acids, polyphenolic compounds, anthocyanins, and a range of organic and inorganic substances (Karataş *et al.*, 2015; Bora *et al.*, 2016). During 2023, global wine production, excluding juices and musts, was estimated at 237 mhl

Commented [RA1]: The correct name of the variety is ALICANTE HENRI BOUSCHET (see VIVC database: <https://www.vivc.de/index.php?r=cultivarnames%2Findex&ultivarnamesearch%5Bcultivarnames%5D=&Cultivarnamesearch%5Bcultivarnames%5D=cultivarn&Cultivarnamesearch%5Btext%5D=Alicante>)

Commented [RA2]: The abbreviation should be disclosed, or use the full term: million hectoliters.

(OIV, 2024). Wine is a widely popular beverage made from grapes through fermentation conducted under controlled conditions (Somkuwar *et al.* 2019). The wine market has experienced rapid growth in recent years due to increasing global demand and consumption, with production rising by over 40% in the past decade (OIV, 2024). Berries with high acidity, low sugar content, and low pH are beneficial for winemaking, as the quality of wine largely depends on its acid content (Jones *et al.*, 2014). Grape composition is influenced not only by climate, which can cause disruptions, but also by vineyard management practices that can enhance the desired levels of acidity, sugar, and pH (Lavras, 2017).

Commented [RA3]: As the topic is general, there should be added some other authors.

Commented [RA4]: See the previous comment.

The quality of grape wine is directly influenced by various factors, including the grape variety, environmental and climatic conditions, physical and chemical soil characteristics, the winemaking process, transportation and storage methods, the degree of agro-technical practices employed, and the vineyard's geographical location (Bora *et al.*, 2015a; Condruso *et al.*, 2015). A suitable climate is crucial for maintaining stable productivity, while year-to-year variations in atmospheric conditions can significantly impact grape quality (Jones and Goodrich, 2008) and, in turn, the quality of the wine.

Commented [RA5]: The same.

Wineries produce wine using more than 30 different types of white grapes, with approximately 90% of the world's wine grapes originating from the *Vitis vinifera* species. Key white grape varieties used in winemaking include Cabernet Sauvignon, Syrah, Cabernet Franc, Grenache, Pinot Noir, Merlot, and Petit Verdot, among others. The quality of wine is primarily influenced by the grape variety cultivated, the prevailing climate during growth, vineyard management practices, and the fermentation techniques employed (Karibasappa, 2013). The micro climates of grape growing regions of Maharashtra specifically Nashik and Pune are suitable to grow wine varieties and produce acceptable quality wine.

Commented [RA6]: ??? More attention, please.

Materials and Methods

Experimental site: In the current study, growth, yield and berry quality parameters of 15 red grape wine varieties (*V. vinifera* L.) grafted on Dogridge rootstock planted at National Research Centre for Grapes, Pune was undertaken during two seasons (2022-23 and 2023-24). The age of the vineyard was seven years old with good health and regular crop. The vines were trained to a mini-Y trellis system with single cordons trained in the horizontal direction while shoots were placed in a vertical position. The soil in the region is heavy black with pH 7.75 and EC 0.46 dS m⁻¹.

All the varieties were planted at a spacing of 3 m between the rows and 1.5 m between the vines in a completely randomized setup with three replications and five vines per replication.

Procedure to record observations

Fifteen red wine grape varieties (Aliquate Bauschet, Cabernet Sauvignon, Cabernet Franc, Cabernet Franc clone, Syrah, Petit Syrah, Tempranillo, Grenache, Grenache Noir, Cinsaut, Caladoc, Nielluccio, Merlot, Petit Verdot and Red Muscat) were used as per the treatments for production and for further use in the study.

Five vines were selected and marked within each replication and means of five vines was calculated for each parameter. After pruning, pruned material from each vine was collected immediately and weighed using a weighing balance (Param weighing scale). The mean weight of biomass was calculated and expressed in kg/vine.

Days to bud sprout were calculated after each pruning (foundation and fruit pruning). The first sprouted bud with fully expanded leaf was taken as an indicator to count the days to bud sprout (Satisha *et al.*, 2010)

Five canes were selected randomly and tagged to measure cane length using measuring tape at 90 days after fruit pruning (DAP) and was expressed in cm.

Cane diameter was measured between fifth and sixth node of cane from five different vines and the mean was expressed in millimeters (mm) at 90 days after foundation pruning.

Internodal length was measured between fifth and sixth nodes of canes using a measuring scale at 90 DAP during the fruit pruning. The mean was calculated and reported in centimeters (cm).

Number of leaves per fruiting shoot was recorded by counting the number of leaves from selected canes at 90th day of fruit pruning. The mean number of shoots was worked out.

Five shoot were selected from each vines and it was measured using BIOVIS, leaf area meter at days after foundation pruning and their mean was expressed in cm².

The chlorophyll content in the various parts of vine was estimated using the method of Witham *et al.* (1971).

After harvesting bunches, five healthy bunches/vine were selected for recording the observations for yield and quality parameters.

Number of bunches per vine was recorded by counting number of bunches from three different vines and the mean of the three vines was calculated.

Number of berries was recorded by counting number of berries from five different bunches and mean of the five bunches was calculated.

Five healthy bunches per replication were selected randomly at the time of harvesting and their mean weight was recorded using weighing balance. The mean average bunch weight was expressed in grams.

A hundred berries from five bunches were selected under each replication and their mean weight was recorded using weighing balance. The mean 100 berry weight was expressed in grams.

At the time of harvest, five vines were selected and tagged. The harvested grapes from these vines under each treatment were weighed using weighing balance. The mean yield of each vine calculated and was expressed in Kg.

Results and Discussion:

1. Growth parameters

1.1 Pruned biomass

Cabernet Sauvignon exhibited the highest pruned biomass weight during both years. This suggests that Cabernet Sauvignon vines are more vigorous. Cabernet Franc clone recorded the lowest pruned biomass weight, indicating less vigorous growth compared to other varieties. In the year 2022-23, Cabernet Sauvignon showed highest weight of pruned biomass. However, lowest weight of pruned biomass was recorded in Cabernet Franc Clone. During fruit pruning season of 2023-24, higher pruned biomass was recorded in Cabernet Sauvignon variety which was statistically higher than all other varieties while, lower pruned biomass was reported in Merlot variety. Vine vigor is measured by pruning weight, an essential growth characteristic that helps differentiate grape varieties into vigorous and non-vigorous categories based on their growth rates (Benz *et al.*, 2006 and Somkuwar *et al.*, 2024a). Similar significant variation for pruning weight was found by Jayalakshmi *et al.*, (2019); Nidhi *et al.*, (2023) and Somkuwar *et al.*, (2024a).

1.2 Days taken to bud sprout

During 1st year of trial, Grenache Noir variety took minimum days to sprout which was followed by Aliquant Bauschet, Cabernet Franc Clone, Petit Syrah, Cabernet Franc,

Merlot and Red Muscat. While, maximum days to bud sprout were taken by Grenache Noir. During season 2023-24, minimum days for bud sprout were taken by Aliquant Bauschet, Cabernet Franc, Cabernet Franc Clone, Petit Syrah, Grenache Noir (9.00 days after fruit pruning) while, maximum days to bud sprout were taken by Cinsaut. According to pooled mean data of two years, Grenache Noir showed early sprouting which was followed by Aliquant Bauschet, Cabernet Franc Clone, Petit Syrah, Cabernet Franc and Merlot while Cinsaut was late to sprout. Bud burst is a varietal character as it marks the beginning of seasonal growth and is strongly influenced by temperature. The data on the growth parameter clearly indicated that prevailing temperature after pruning affects the time required for bud break in the same variety and the influence of temperature is more than that of variety (Somkuwar *et al.*, 2024a). Similar studies were reported by Gupta *et al.*, (2015).

1.3 Shoot length

Shoot length recorded during both season was maximum in Syrah variety whereas the minimum shoot length in Merlot. Highly vigorous varieties generally produce longer shoots. In contrast, Veena *et al.*, (2015) found that vigorous varieties have produced comparatively shorter shoot than less vigorous varieties. This may be due to number of buds retained on the cane after pruning.

1.4 Shoot diameter

Syrah recorded maximum shoot diameter during season 2022-23 and pooled mean of experiment, respectively which was followed by Red Muscat and minimum in Petit Verdot. During season 2023-24, highest shoot diameter was observed in Syrah which was followed by Red Muscat while Nielluccio exhibited lowest shoot diameter. The production of canes and their dimension depends upon vigour of the vine, which in turn depends upon the extent of stored food material in the vine. Vigorous varieties having greater stored food material produce higher number of canes. Several workers recorded similar observations *i.e.*, Pina and Bautists (2006), Havinal *et al.*, (2008), Soni *et al.*, (2019) and Nidhi *et al.*, (2023).

1.5 Internodal length

Petit Syrah recorded the highest pooled mean internodal length. Syrah, Nielluccio and Red Muscat closely followed to Petit Syrah however, Merlot had the lowest pooled mean internodal length. Syrah showed highest internodal length during 2022-23 season which was closely followed by Petit Syrah, Red Muscat, Nielluccio and Tempranillo and minimum internodal length was noted in Cabernet Sauvignon. During 2023-24 season, Petit Syrah

exhibited long internodal which was followed by Syrah, Nielluccio, Caladoc and Red Muscat while short internodal recorded in Merlot, Petit Verdot, Grenache Noir, Aliquant Bauschet and Cabernet Sauvignon. Higher internodal length in Syrah may be due to more shoot length which increases internodal length in between cane. Mostly shorter internodes accumulate higher carbohydrates food reserves which are pre-requisite for flower bud initiation (Jayalakshmi *et al.*, 2019; Somkuwar and Ramteke, 2008)

1.6 Number of leaves per fruiting shoot

During both season, Caladoc recorded the maximum number of leaves which was followed by Cinsaut while, Syrah had the fewest leaves per fruiting shoot. Such variation among the colored and white varieties for leaf number may be attributed to difference in number of canes and vigour of the vine and the inherent varietal character (Veena *et al.*, 2015). Similar results were also reported by Anupama *et al.*, (2016).

1.7 Leaf area

During both season, Nielluccio continued to show the highest leaf area and followed by Caladoc however, Merlot and Cabernet Sauvignon had the minimum leaf area. Such, high variation among the red wine varieties for leaf area may be attributed to varietal character (Veena *et al.*, 2015). The increased leaf area results in highest active photosynthesis rate which helps to store more carbohydrate in the sink, bunch (Somkuwar *et al.* 2013). Somkuwar *et al.*, (2024c, 2024d, 2024e) revealed that leaf area was decreased with increased number of leaves per shoot.

1.8 Leaf area per fruiting shoot

During 2022-23 season and pooled mean, Caladoc had the highest leaf area per fruiting shoot which was followed by Cinsaut. On the lower end, Merlot observed the lowest leaf area per fruiting shoot. During 2023-24 season, Caladoc recorded significantly higher leaf area per fruiting shoot among all varieties studied. However, Cabernet Sauvignon reported minimum leaf area per fruiting shoot. The increase in leaf area per shoot and vine with more leaves is due to the direct correlation between the number of leaves and the overall vegetative growth of the vine. Leaf area is strongly correlated with the annual shoots and cane growth; the most vigorous varieties usually have the highest annual growth (Borca *et al.*, 2020). Similar results were reported by Georgeta *et al.*, (2021) in red wine varieties. The optimum leaf number enhanced the overall leaf area, potentially contributing to higher

photosynthetic capacity (source) and resource distribution (sink) for grape development (Somkuwar *et al.*, 2024e and Thoke *et al.*, 2024).

1.9 Chlorophyll content in the leaves

When considering the pooled mean across both seasons, Cabernet Franc clone maintained the highest chlorophyll content, closely followed by Cabernet Sauvignon. However, Caladoc consistently exhibited the lowest pooled chlorophyll content. The chlorophyll content in leaf gives an indication of the efficiency of leaf to prepare food through photosynthesis (Somkuwar *et al.*, 2024d). More leaves can increase overall photosynthetic capacity. There may be an optimum leaf number beyond which chlorophyll content per leaf might begin to decrease. Maintaining an appropriate number of leaves can produce maximum chlorophyll content and photosynthetic efficiency without any negative effect (Somkuwar *et al.*, 2024d). Petrie *et al.*, (2000) and Somkuwar *et al.*, (2014b) observed that leaf removal led to an increase of chlorophyll content. Somkuwar *et al.*, (2024c) reported that the results illustrated variations in chlorophyll a, chlorophyll b and total chlorophyll concentrations between different treatments. Chlorophyll content in leaf ranged from 29.15 to 25.30 mg/ml and 21.35 mg/ml to 19.14 mg/ml among the treatments.

2 Yield Parameters

2.1 Number of bunches per vine

Pooled mean values across both seasons was significant with Syrah had the highest number of bunches per vine. On the lower end, Petit Verdot recorded the lowest pooled mean. During 2022-23 season, Syrah exhibited the highest number of bunches per vine which was followed by Grenache Noir, while lowest in Petit Verdot. During 2023-24 season, Syrah recorded a high number of bunches per vine while lowest in Cabernet Franc and its clone, along with Petit Verdot. The increase in number of bunches per vine in Syrah might be due to highest fruitfulness of spur. Similar line of work in grapes was reported by Chalak *et al.*, (2011), Anupama *et al.*, (2016), Leao *et al.*, (2017), Anjanawe *et al.*, (2020), Tecchio *et al.*, (2022), Nidhi *et al.*, (2023) and Somkuwar *et al.*, (2024a).

2.2 Number of berries per bunch

During 2022-23, Grenache Noir had the highest average number of berries per bunch and Merlot recorded the fewest berries per bunch. During the 2023-24 season, Grenache Noir again led with higher berries per bunch while Red Muscat had fewest berries per bunch. The pooled mean over both seasons showed that Grenache Noir exhibited the highest average

number of berries per bunch (177.47). However, minimum number of berries per bunch observed in Red Muscat (79.67) and Merlot (80.00). Somkuwar *et al.*, (2024a) reported that the maximum number of berries/bunch were recorded in Cabernet Sauvignon (111.00) during the year 2014-15 whereas, in 2016-17 and pooled mean it was maximum in Cinsaut (227.33 and 134.55) and in 2015-16 was recorded maximum in Nielluccio (145.00) variety. While minimum number of berries per bunch was recorded in Merlot (69.00 and 79.00 respectively) during both the year 2015-16 and 2016-17 while, in 2014-15 and pooled data was recorded in Nielluccio and Cabernet Franc (62.33 and 81.55). Jayalakshmi *et al.*, (2019) found maximum number of berries per bunch (278.54) in Manjari Medika, 43.65 numbers of berries per bunch in Zinfandel. Similar line of findings was earlier reported by Havinal *et al.*, (2008); Tecchio *et al.*, (2022).

2.3 Average bunch weight

When analyzing the pooled mean along with both seasons, it was found that Aliquate Bauschet consistently emerged as the variety with the highest bunch weight. In contrast, Merlot and Petit Verdot had the lowest pooled mean bunch weight. The variation in the bunch weight among different varieties may be attributed to inherent genetic character of the variety, number of berries per bunch, difference in number of canes and berry size and also the size of vine canopy where varieties with larger canopy sizes were noted to have higher bunch weights (Somkuwar *et al.*, 2024a; Nidhi *et al.*, 2023; Anjanawe *et al.*, 2020; Havinal *et al.*, 2008;). Tecchio *et al.*, (2022) found that Cabernet Sauvignon was one of the cultivars that had the lowest bunch mass (85.5 g). Moreover, this value was 76% lower than those detected by Rizzon and Miele (2002) under temperate conditions. Similar line of work was done earlier by Leao *et al.*, (2017); Ingole *et al.*, (2018).

2.4 100 berries weight

The analysis of 100 berry weight for various red wine grape varieties over two seasons (2022-23 and 2023-24) demonstrated noticeable differences among the varieties. Red Muscat variety had the maximum 100 berry weight followed by Aliquate Bauschet. The Petit Verdot variety showed the minimum 100 berry weight, making it the variety with the lightest berries overall. The variation in berry weight may arise from differences in both the diameter of the berries and the number of berries/bunch (Thakur *et al.*, 2008). The present studies are in confirmation with the results of Ratnacharyulu (2010); Ghosh *et al.* (2012); Anupama *et*

al., (2016); Pal *et al.*, (2018); Soni *et al.*, (2019); Nidhi *et al.*, (2023); Somkuwar *et al.*, (2024a, 2024b).

2.5 Yield

Aliquate Bauschet recorded the highest yield and lowest yield was observed in Petit Verdot and Merlot varieties. Disparities in yield per vine among various grape cultivars may be attributed to variations in bunch weight, number of bunches, and berry weight besides their successful adoption to the varying agro-climatic conditions under which they are cultivated (Havinal *et al.*, 2008). The positive correlation of yield per vine with average bunch weight and berry weight was recorded. Crop yield was found to increase in proportion to the number of clusters/vine, a trend similarly observed in the findings of Myers *et al.* (2008) in Sangiovese grape wines. Orlando *et al.* (2008) assessed the productive performance of Syrah and Cabernet Sauvignon under similar climatic conditions and they observed that Syrah was also more productive than Cabernet Sauvignon regardless of the rootstock. Similar line of work is reported by Somkuwar *et al.* (2024a, 2024b), Tecchio *et al.*, (2022); Anjanawe *et al.*, (2020); Ingole *et al.* (2018); Vijaya *et al.*, (2018); Leao *et al.*, (2017); Veena *et al.* (2015); Chalak *et al.*, (2011).

Conclusion

The evaluation of growth and yield parameters of different red wine grape varieties over the 2022-23 and 2023-24 seasons revealed significant varietal differences. The variety Cabernet Sauvignon showing superior growth parameters like pruned biomass while higher shoot length, shoot diameter, number of leaves and yield parameter like number of bunches was recorded in Syrah. Minimum days taken to bud sprout and number of berries was recorded in Grenache Noir. Maximum internodal length was recorded in Petit Syrah. Higher number of leaves in Caladoc, leaf area was recorded in Caladoc and Nielluccio varieties. Higher chlorophyll content was recorded in Cabernet Franc clone. Yield-related metrics like bunch weight and yield was recorded higher in Aliquate Bauschet, 100 berry weight (221.94 g) in Red Muscat for pooled mean basis.

References

- Anjanawe, S. R.; Naruka, I. S.; Sharma, A. and Mishra, P. (2020). Evaluation of Wine Purpose Varieties of Grapes under the Environmental Condition of Malwa Plateau. *Current Journal of Applied Science and Technology*, **39**(43): 98-107.
- Anupama, H.; Hipparagi, K.; Rani, S.; Ravindranath and Balesh G. (2016). Evaluation of wine grape varieties for growth and yield under northern dry zone of Karnataka. *International Journal of Scientific Research*, **5**(2): 409-411.
- Ausari, P.K.; Gurjar, P.K.S.; Somkuwar, R.G.; Naruka, I.S.; Sharma, A.K. and Gharate, P.S. (2024). Effect of rootstocks on yield and wine quality of Sauvignon Blanc variety. *Plant Archives*, **24**(1): 1477-1482.
- Benz, M.J.; Anderson, M.M.; Williams, M.A.; Barnhisel, K. and Wolpert, J.A. (2006). Viticultural performance of five Merlot clones in Oakville, Napa Valley. *American Journal of Enology and Viticulture*, **57**: 233-237.
- Bora, F.D.; Bunea, C.I.; Rusu, T. and Pop, N. (2015). Vertical distribution and analysis of micro, macroelements and heavy metals in the system soil-grapevine-wine in vineyard from North-West Romania. *Chemistry Central Journal*, **9**(1): 19.
- Bora, F.D.; Donici, A.; Oslobanu, A.; Fitiu, A.; Babes, A.C. and Bunea, C.I. (2016). Qualitative assessment of the white wine varieties grown in Dealu Bujorului vineyards, Romania. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, **44**(2): 593-602.
- Borca, F.; Roxana, N.; Alina, D.; Eleonora, N. and Dobrei, A. (2020). The anthropogenic influence of soil management on grape yield and economic parameters in grapevine growing. *Journal of Horticulture, Forestry and Biotechnology*, **24**(4): 58-65.
- Burg, P.; Burgova, J.; Masan, V. and Vachun, M. (2017). Leaf surface area estimation in different grapes varieties using a AM-300 leaf area meter. In International scientific conference Rural Development 2017: 24-30.
- Chalak S.U.; Kulkarni, S.S.; Kolse, S.V. and Garad, B.V. (2011). Effect of pruning levels in some red wine grape varieties for yield and growth parameters under Western Maharashtra conditions. *Eco. Env. & Cons.* **17**(1): 87-93.
- Chalak, S.U.; Kulkarni, S.S.; Kishrsagar, A.V. and Nimbalkar, C.A. (2012). Pruning studies in some white wine grape varieties for yield and yield contributing parameters under Western Maharashtra conditions. *Asian Journal of Horticulture*, **7**(2): 1468-1472.

- Concurso, C.; Cioncatta, F.; Tripodi, G.; Sparacio, A.; Giglio, D.M.F.; Sparla, S. and Verzera, A. (2015). Effects of cluster thinning on wine quality of Syrah cultivar (*Vitis vinifera* L.). *European Food Research and Technology*, **246**(10): 1719-1726.
- Georgeta, D.A.; Eleonora, N.; Daniela, D.; Petcov, D.Z. and Dorei, A.I. (2021). Evaluation of grape yield and quality of some red wines varieties from Minis-Maderat Vineyard. *Journal of Horticulture, Forestry and Biotechnology*, **25**(2): 74-79.
- Ghosh, S.N.; Bera, B.; Roy, S. and Kundu, A. (2012). Adaptation and commercialization of viticulture in West Bengal - a new area in India. *Acta Horticulturae*, **931**: 389-399.
- Ghule, V.S.; Ranpise, S.A.; Somkuwar, R.G.; Kulkarni, S.S.; Wagh, R.S.; Naik, R.M. and Nimbalkar, C.A. (2021). Effect of rootstocks on growth parameters of red globe grapevines (*Vitis vinifera* L.). *International Journal of Chemical Studies*, **9**(1): 3483-87.
- Gupta, N.; Gill, K.K.; Babuta, R.; Gill, M. and Arora, N.K. (2015). Thermal requirement and phenological development of different grape varieties under South Western Punjab. *Annals of Agricultural Research*, **36**(4): 377-383.
- Havinal MN, Tambe TB and Patil SP (2008) Comparative studies on vine vigour and fruitfulness of grape wine varieties. *Asian J Hort* 3(1):180-182.
- Huang, H. and Lu, J. (2000). Variation and correlation of bud breaking, flower opening and fruit ripening in Muscadine grape cultivars. *Proceedings of the Florida State Horticultural Society*, **113**: 46-47.
- Ingole, R.H.; Tambe, T.B. and Bobade, D.H. (2018). Influences of various rootstocks on yield and quality of wine grape varieties (*Vitis vinifera* L). *Journal of Pharmacognosy and Phytochemistry*, **7**(2): 2466-2468.
- International Organisation of Vine and Wine -OIV, 2023. Standards and technical documents. 2023. Available at: <https://www.oiv.int/what-we-do/standards>. Access in: 20 April 2024.
- Jayalakshmi, C.; Saraswathy, S.; Subbiah, A.; Ilamurugu, K. and Balachandar, D. (2019). Evaluation of wine varieties of grapes (*Vitis vinifera* L.) during winter pruning under Cumbum valley condition of Tamil Nadu. *Journal of Pharmacognosy and Phytochemistry*, **8**(3): 3770-3773.
- Jones, G.V. and Goodrich, G.B. (2008). Influence of climate variability on wine regions in the western USA and on wine quality in the Napa Valley. *Climate Research*, **35**(3): 241-254.

- Jones, J.E.; Kerslake, F.L.; Close, D.C. and Dambergs, R.G. (2014). Viticulture for sparkling wine production: a review. *American Journal of Enology and Viticulture*, **65**: 407-416.
- Karatas, D.; Aydin, F.; Aydin, I. and Karatas, H. (2015). Elemental composition of red wines in Southern Turkey. *Czech Journal of Food Sciences*, **33**(3): 228-236.
- Karibasappa G.S. (2013). Manual and good agricultural practices for quality wine production. *Indian Council of Agricultural Research* 2013, 5-16.
- Lavras, M.G. (2017). Planting density for Chardonnay Grapevines in The South of Minas Gerais- M.Sc. Thesis. Luciana Wilhelm De Almeida.
- Leao, P.C. de Souza; Nunes, B.TG.; Carvalho de Souza, E.M.; Rego, J.I. de Souza and Bernardino do Nascimento, J.H. (2017). Evaluation of Some New Wine Grape Cultivars under Sao Francisco Valley Conditions. *Journal of Agricultural Science and Technology*, **B**(7): 320-325.
- McGovern, P.E. (2003). *Ancient Wine: The Search for the Origins of Viniculture* (PDF). Princeton University Press, Princeton.
- Myers, J.K.; Wolpert, J.A. and Howell, G.S. (2008). Effect of shoot number on the leaf area and crop weight relationship of young Sangiovese grapevines. *American Journal of Enology and Viticulture*, **59**(4): 422-424. DOI: 10.5344/ajev.2008.59.4.422
- Orlando, T.G.S, Pedro Junior, M.J.; Santos, A.O. and Hernandez, J.L. (2008). Comportamento das cultivares de Cabernet Sauvignon e Syrah em diferentes porta-enxertos. *Ciência e Agrotecnologia (Science and Agrotechnology)*, **32**: 749-755. <https://doi.org/10.1590/S1413-70542008000300007>
- Pal, R.; Deb, P. and Ghosh, S.N. (2018). Performance of Some Grape Cultivars for Commercial Cultivation in West Bengal. *International Journal of Bio-resource and Stress Management*, **9**(5): 568-572.
- Patel, Nidhi; Soni, N.; Ausari, P.K.; Meena, K.C.; Patidar, B.K.; Patidar, D.; Haldar, A. and Thakur, R. (2023). Evaluation of Different Wine varieties of Grapes Grafted on Dogridge Rootstocks. *Biological Forum- An International Journal*, **15**(8a): 521-525.
- Petrie, P.R.; Trought, M.T. and Howell, G.S. (2000). Influence of leaf ageing, leaf area and crop load on photosynthesis, stomatal conductance and senescence of grapevine (*Vitis vinifera* L. cv. Pinot noir) leaves. *Vitis*, **39**(1): 31-36.
- Pina, S. and Bautista, D. (2006). Evaluation of vegetative growth on several table grape cultivars under semiarid tropic conditions in Venezuela. *Magazine of the Faculty of Agronomy of the University of Zulia*, **23**: 402-413.

- Ratnacharyulu SV (2010). Evaluation of coloured grapes varieties for yield, juice recovery and quality. M.Sc., Thesis. Grapes Research Station, Andhra Pradesh Horticultural University, Rajendranagar, Hyderabad, A. P.
- Rizzon, L.A. and Miele, A. (2002). Avaliação da cv. Cabernet bSauvignon para elaboração de vinho tinto. *Food Science and Technology*, **22**: 192-198. <https://doi.org/10.1590/S0101-20612002000200015>
- Satisha, J.; Somkuwar, R.G.; Sharma, J.; Upadhyay, A.K. and Adsule, P.G. (2010). Influence of rootstocks on growth yield and fruit composition of Thompson Seedless grapes grown in the Pune region of India. *South African Journal of Enology and Viticulture*, **31**: 1-8.
- Somkuwar, R.G. and Ramteke, S.D. (2008). Effect of shoot orientation on growth, biochemical changes and fruitfulness in grape (*Vitis vinifera* L.) variety Thompson Seedless. *Journal of Plant Sciences*, **3**(2): 182-187.
- Somkuwar, R.G.; Bhor, V.A.; Ghule, V.S.; Hakale, D.P.; Shabeer, A. and Sharma, A.K. (2021). Rootstock affects stress relieving enzymatic activity during bud break in 'Red Globe' grapevine under semi-arid condition. *Vitis*, **60**: 93-99.
- Somkuwar, R.G.; Hakale, D.P. and Sharma, A.K. (2019). Studies on biochemical composition of different parts of berries and wine quality of wine grape varieties (*Vitis vinifera* L.). *International Journal of Current Microbiology and Applied Sciences*. **8**(03): 155-164. doi: <https://doi.org/10.20546/ijcmas.2019.803.022>
- Somkuwar, R.G.; Kakade, P.B.; Dhemre, J.K.; Gharate, P.S.; Deshmukh, N.A. and Nikumbhe, P.H. (2024c). Leaf area influences photosynthetic activities, raisin yield and quality in Manjari Kishmish grape variety. *Archives of Current Research International*, **24**(6): 613-622.
- Somkuwar, R.G.; Kakade, P.B.; Dhemre, J.K.; Thutte, A.S.; Nikumbhe, P.H. and Deshmukh, N.A. (2024d). Leaf retention affects photosynthetic activity, leaf area index, yield and quality of Crimson Seedless grapes. *Journal of Advances in Biology & Biotechnology*, **27**(9): 123-130.
- Somkuwar, R.G.; Kakade, P.B.; Jadhav, A.S.; Ausari, P.K.; Nikumbhe, P.H. and Deshmukh, N.A. (2024e). Leaf area index, photosynthesis and chlorophyll content influences yield and quality of Nanasaheb Purple Seedless grapes under semi-arid condition. *Journal of Scientific Research and Reports*, **30**(9): 750-758.

- Somkuwar, R.G.; Ramteke, S.D. and Satisha, J. (2010). Effect of cluster clipping and berry thinning on yield and quality of Thompson Seedless grapes. *Acta Horticulturae*, **758**: 229-231.
- Somkuwar, R.G.; Sharma, A.K.; Nilima, G. and Ausari, P.K. (2024a). Performance of red wine varieties under Pune region of Maharashtra. *International Journal of Advanced Biochemistry Research*, **8**(5): 981-988.
- Somkuwar, R.G.; Sharma, A.K.; Nilima, G. and Ausari, P.K. (2024b). Evaluation of White Wine Varieties for Growth, Yield, Berry and Wine Quality under Pune Region of Maharashtra India. *Pant Archives*, **24**(2): XX-XX.
- Soni, N.; Patil, P.; Meena, K.C.; Haldar, A.; Patidar, D.K. and Tiwari, R. (2019). Evaluation of Different Colored Varieties of Grapes under Nontraditional Area of Malwa Plateau: A Thin Line Tool for Doubling the Farmer Income. *International Journal of Current Microbiology and Applied Sciences*, **8**(03): 1968-1976. doi: <https://doi.org/10.20546/ijcmas.2019.803.234>
- Stanley, H.G. (2001). Sustainable Grape Productivity and the Growth-Yield Relationship: A Review. *American Journal of Enology and Viticulture*, **52**(3): 165-174.
- Tecchio, M A.; Silva, M.J.R.; Sanchez, C.A.P.C.; Callili, D.; Vedoato, B.T.F.; Hernandes, J.L. and Moura, M.F. (2022). Yield performance and quality of wine grapes (*Vitis vinifera*) grafted onto different rootstocks under subtropical conditions. *Bragantia*. **81**: e1622. <https://doi.org/10.1590/1678-4499.20210214>
- Tecchio, M.A.; Hernandes, J. L.; Pires, E.J.P.; Terra, M.M. and Moura, M.F. (2018). Cultivo da videira para mesa, vinho e suco. In: R. Pio (Ed.). Cultivo de fruteiras clima temperado em regiões subtropicais e tropicais (2. ed., p. 512-584). Lavras: Editora UFLA.
- Thakur, A.; Arora, N.K. and Singh, S.P. (2008) Evaluation of some grape varieties in the Arid Irrigated region of North West India. *Acta Horticulturae*, **785**: 79-83.
- Thoke, S.; Patil, S.N.; Gollagi, S.G.; Hipparagi, K.; Hiremath, V.; Peerajade, D. and Kore, D. (2024). Evaluation of Different Table Grape Varieties in Response to Growth and Physiological Traits during Foundation Bud Pruning. *Journal Agriculture of Experimental International*, **46**(2): 126-140.
- Veena, J.; Kumar, V.; Debnath, M.; Pattanashetti, S.; Variath, M.T.; Khadakabhavi, S. (2015). Multivariate analysis of colored and white grape grown under semi-arid tropical conditions of Peninsular India. *International Journal of Agriculture and Crop Sciences*. **8**(3): 350-365.

Vijaya, D.; Reddy, G.R.; Joshi, V.; Mamatha and Kumari, D.A. (2018). Evaluation of juice and wine varieties of grapes (*Vitis spp*) for petiole nutrient content, bud break, yield and yield components. *International Journal of Chemical Studies*, **6**(6): 2739-2745.

Witham, F.H.; Blaydes, D.F. and Devlin, R.M. (1971). *Experiments in plant physiology*. Litton Educational Publishing Inc. D. Van Nostrand Company. New York.

UNDER PEER REVIEW

Table 1: Pruned biomass, days taken to bud sprout and shoot length data of red grape wine varieties during two growing seasons (2022-23 and 2023-24) and pooled mean data

Varieties	Pruned biomass (Kg)			Days taken to bud sprout			Cane length (cm)		
	2022-23	2023-24	Pooled Mean	2022-23	2023-24	Pooled Mean	2022-23	2023-24	Pooled Mean
Aliquate Bauschet	0.499	0.492	0.496	8.97	9.00	8.98	53.89	54.50	54.20
Cabernet Sauvignon	1.053	1.107	1.080	11.83	10.33	11.08	62.44	61.20	61.82
Cabernet Franc	0.400	0.397	0.399	9.17	9.00	9.08	77.00	77.10	77.05
Cabernet Franc clone	0.367	0.387	0.377	9.00	9.00	9.00	62.02	62.69	62.35
Syrah	0.583	0.680	0.631	10.73	10.03	10.38	115.44	125.43	120.44
Petit Syrah	0.575	0.616	0.595	9.13	9.00	9.07	91.20	88.67	89.93
Tempranillo	0.674	0.661	0.668	12.17	12.00	12.08	99.33	103.33	101.33
Grenache	0.645	0.627	0.637	11.33	11.67	11.50	80.33	87.03	83.68
Grenache Noir	0.521	0.527	0.524	8.87	9.00	8.93	57.16	56.84	57.00
Cinsaut	0.649	0.612	0.631	13.00	13.00	13.00	87.22	93.90	90.56
Caladoc	0.841	0.836	0.839	12.00	11.33	11.67	91.11	90.33	90.72
Nielluccio	0.720	0.875	0.798	11.00	10.33	10.67	73.67	91.43	82.55
Merlot	0.383	0.376	0.380	9.33	9.67	9.50	49.66	53.00	51.33
Petit Verdot	0.450	0.384	0.418	12.67	12.33	12.50	59.44	61.10	60.27
Red Muscat	0.823	0.839	0.831	10.07	10.00	10.03	71.07	75.00	73.04
S.Em. (±)	0.011	0.009	0.007	0.510	0.456	0.325	3.563	3.393	2.170
C.D. (0.05)	0.031	0.027	0.021	1.486	1.329	0.947	10.263	9.775	6.251

Table 2: Shoot diameter, internodal length and number of leaves per shoot data of red grape wine varieties during two growing seasons (2022-23 and 2023-24) and pooled mean data

Varieties	Cane diameter (mm)			Internodal length (cm)			Number of leaves/shoot		
	2022-23	2023-24	Pooled Mean	2022-23	2023-24	Pooled Mean	2022-23	2023-24	Pooled Mean
Aliquate Bauschet	6.02	6.55	6.28	5.04	5.01	5.02	15.67	16.25	15.96
Cabernet Sauvignon	5.22	4.90	5.06	4.01	5.03	4.53	16.44	17.60	17.02
Cabernet Franc	6.67	6.20	6.43	5.18	5.67	5.43	14.67	16.17	15.42
Cabernet Franc clone	6.57	6.93	6.75	5.50	5.97	5.73	15.30	15.65	15.47
Syrah	9.89	9.60	9.75	6.93	6.90	6.92	13.55	14.73	14.14
Petit Syrah	8.52	7.71	8.12	6.84	7.04	6.94	17.93	17.52	17.73
Tempranillo	6.22	5.93	6.08	6.28	6.23	6.26	15.78	16.50	16.14
Grenache	6.89	4.90	5.90	6.23	5.90	6.07	18.34	18.17	18.25
Grenache Noir	8.94	9.09	9.02	4.81	4.79	4.80	14.58	15.31	14.95
Cinsaut	6.11	5.70	5.91	6.01	6.13	6.08	22.89	24.67	23.78
Caladoc	7.86	4.67	6.27	5.98	6.53	6.26	23.67	25.17	24.42
Nielluccio	8.56	4.63	6.60	6.30	6.90	6.60	15.44	16.50	15.97
Merlot	5.22	4.83	5.03	4.59	4.40	4.50	16.50	16.87	16.68
Petit Verdot	5.00	4.80	4.90	4.52	4.60	4.56	17.17	17.17	17.17
Red Muscat	9.27	9.37	9.32	6.33	6.47	6.40	19.26	19.32	19.29
S.Em. (±)	0.281	0.171	0.157	0.255	0.222	0.189	0.314	0.371	0.280
C.D. (0.05)	0.809	0.493	0.453	0.736	0.639	0.545	0.906	1.069	0.806

Table 3: Leaf area per leaf, leaf area per fruiting shoot and cane chlorophyll content of red grape wine varieties during two growing seasons (2022-23 and 2023-24) and pooled mean data

Varieties	Leaf area/leaf			Leaf area/fruiting shoot			Chlorophyll content (mg/ml)		
	2022-23	2023-24	Pooled Mean	2022-23	2023-24	Pooled Mean	2022-23	2023-24	Pooled Mean
Aliquate Bauschet	90.03	94.48	92.25	1409.44	1533.08	1471.26	9.78	10.39	10.09
Cabernet Sauvignon	72.52	46.45	59.49	1193.90	816.92	1005.41	12.38	13.68	13.03
Cabernet Franc	73.60	83.56	78.58	1073.68	1353.81	1213.74	10.85	12.48	11.67
Cabernet Franc clone	92.62	84.38	88.50	1416.24	1317.22	1366.73	13.11	13.92	13.52
Syrah	102.96	116.78	109.87	1392.52	1719.76	1556.14	12.29	13.58	12.94
Petit Syrah	103.84	112.88	108.36	1858.19	1968.76	1913.47	11.34	12.78	12.06
Tempranillo	111.37	97.33	104.35	1761.16	1609.35	1685.26	8.10	8.77	8.43
Grenache	99.52	113.48	106.50	1822.23	2059.65	1940.94	12.22	12.98	12.60
Grenache Noir	107.20	103.67	105.43	1562.63	1590.86	1576.74	7.11	8.19	7.65
Cinsaut	104.52	100.46	102.49	2385.10	2467.03	2426.06	9.78	10.59	10.19
Caladoc	107.90	131.20	119.55	2553.85	3308.65	2931.25	5.91	6.79	6.35
Nielluccio	129.77	152.73	141.25	2004.07	2513.12	2258.60	11.73	12.96	12.34
Merlot	55.46	53.48	54.47	912.18	900.07	906.13	8.60	9.32	8.96
Petit Verdot	78.66	76.65	77.66	1349.22	1312.01	1330.61	10.64	12.24	11.44
Red Muscat	112.17	89.96	101.07	2162.22	1738.04	1950.13	9.22	9.79	9.51
S.Em. (±)	6.396	9.496	5.636	106.827	177.142	103.640	0.284	0.303	0.184
C.D. (0.05)	18.426	27.356	16.236	307.731	510.286	298.553	0.818	0.872	0.529

Table 4: Number of bunches per vine, number of berries per bunch and Average bunch weight of red grape wine varieties during two growing seasons (2022-23 and 2023-24) and pooled mean data

Varieties	Number of bunches/vine			Number of berries/bunch			Average bunch weight (g)		
	2022-23	2023-24	Pooled Mean	2022-23	2023-24	Pooled Mean	2022-23	2023-24	Pooled Mean
Aliquate Bauschet	32.66	34.04	33.36	140.33	134.50	137.42	261.17	267.83	264.50
Cabernet Sauvignon	37.17	38.37	37.77	96.83	97.33	97.08	105.92	102.57	104.25
Cabernet Franc	29.50	24.33	26.92	93.83	94.50	94.17	116.34	113.27	114.81
Cabernet Franc clone	28.28	24.58	26.43	103.83	104.50	104.17	122.33	118.90	120.62
Syrah	43.17	43.27	43.22	111.44	107.03	109.24	137.89	144.27	141.08
Petit Syrah	35.80	36.03	35.92	131.00	130.67	130.83	165.30	163.70	164.50
Tempranillo	35.00	35.00	35.00	114.00	115.30	114.65	201.00	197.67	199.34
Grenache	35.50	35.67	35.58	142.50	140.33	141.42	157.30	149.27	153.28
Grenache Noir	39.03	39.00	39.02	175.60	179.33	177.47	179.23	185.57	182.40
Cinsaut	30.50	29.00	29.75	116.83	118.63	117.73	151.57	149.80	150.69
Caladoc	36.00	36.67	36.33	89.00	85.33	87.17	147.08	145.50	146.29
Nielluccio	25.67	26.33	26.00	128.00	117.70	122.85	183.45	181.60	182.53
Merlot	27.17	27.00	27.08	81.67	78.33	80.00	92.57	92.60	92.59
Petit Verdot	24.00	25.00	24.50	97.67	101.00	99.33	94.53	93.40	93.97
Red Muscat	30.08	30.29	30.19	85.33	74.00	79.67	179.33	171.50	175.42
S.Em. (±)	1.827	1.635	1.316	3.540	3.724	2.447	6.423	4.548	3.292
C.D. (0.05)	5.262	4.710	3.790	10.198	10.727	7.048	18.502	13.101	9.483

Table 5: 100 berry weight, yield/vine and yield/acre of red grape wine varieties during two growing seasons (2022-23 and 2023-24) and pooled mean data

Varieties	100 berry weight (g)			Yield/vine (Kg)			Yield/acre (MT)		
	2022-23	2023-24	Pooled Mean	2022-23	2023-24	Pooled Mean	2022-23	2023-24	Pooled Mean
Aliquate Bauschet	186.52	199.15	192.83	8.57	9.12	8.84	10.37	11.03	10.70
Cabernet Sauvignon	109.70	105.49	107.59	3.92	3.94	3.93	4.75	4.77	4.76
Cabernet Franc	123.87	120.01	121.94	3.44	2.76	3.10	4.16	3.34	3.75
Cabernet Franc clone	117.61	113.94	115.77	3.48	2.91	3.20	4.21	3.52	3.87
Syrah	124.42	134.89	129.65	5.93	6.25	6.09	7.18	7.56	7.37
Petit Syrah	126.36	125.28	125.82	5.92	5.91	5.91	7.16	7.15	7.15
Tempranillo	176.31	171.63	173.97	7.06	6.92	6.99	8.55	8.38	8.46
Grenache	110.64	106.45	108.54	5.60	5.33	5.46	6.77	6.44	6.61
Grenache Noir	102.24	103.89	103.07	7.00	7.24	7.12	8.47	8.76	8.61
Cinsaut	130.89	126.53	128.71	4.63	4.34	4.48	5.60	5.25	5.43
Caladoc	165.53	170.51	168.02	5.30	5.34	5.32	6.40	6.46	6.43
Nielluccio	143.37	154.89	149.13	4.75	4.78	4.76	5.75	5.78	5.77
Merlot	113.71	119.56	116.63	2.51	2.50	2.51	3.04	3.03	3.03
Petit Verdot	96.96	92.55	94.75	2.27	2.34	2.30	2.74	2.83	2.79
Red Muscat	212.15	231.72	221.94	5.39	5.20	5.30	6.52	6.29	6.41
S.Em. (±)	7.438	5.147	4.057	0.389	0.270	0.217	0.470	0.327	0.263
C.D. (0.05)	21.428	14.826	11.687	1.119	0.777	0.626	1.354	0.941	0.758