

Varietal Evaluation of Tuberose (*Agave amica* (Medik.) For Smart Farming Applications

Commented [k1]: Rewrite the title. Since none of the smart farming technique is involved in the methodology the title is not suitable. Title can be rewited as **Varietal Evaluation of Tuberose for commercial traits**

ABSTRACT

Aims: To evaluate the tuberose varieties for growth, yield and quality
Study design: The experiment was laid out in Randomized block design with three replications.

Place and Duration of Study: Department of Floriculture and Landscaping, College of Agriculture, Padannakkad, Kasargod during 2019-2020.

Methodology: Ten genotypes of tuberose viz. A. Shringar, A. Niranthara, A. Prajwal, A. Sugandhi, A. Suvasini, A. Vaibhav, Culcutta Single, Culcutta Double, Bidhan Ujwal and Phule Rajani were assessed for further evaluation in smart farming. Treatments were replicated thrice. The accessions were screened for vegetative, floral, quality and post-harvest parameters.

Results: The result of the study showed that out of the ten genotypes of tuberose *A. Prajwal* has been found suitable for commercial farming with minimum days to 50% flowering, longest floret, highest diameter of floret, longest spike, maximum yield per unit area and good aroma. Tuberose needs an improved package of practices for better yield and quality of flowers.

Conclusion:

In view of these, further evaluation of *A. Prajwal* for standardization of manuring, drip irrigation, fertilization, spacing and mulching for smart farming is undergoing at Department of Floriculture and Landscaping at College of Agriculture, Vellayani. A semi- automatic controller prototype will be developed based on nutrient utilization model derived in terms of growth parameters of the tuberose as well as the nutrient content of soil and plant.

Commented [k2]: In smart farming system

Commented [k3]: Mention the numerical values of each parameters

Commented [k4]: In smart farming systems.

Commented [k5]: It is the future line of work, please give a specific conclusion based on the results.

Keywords: Tuberose, Accessions, A. Prajwal, Precision Farming

1. INTRODUCTION

Floriculture is a fast-emerging major venture in the world. Now a days, floriculture is lucrative profession with more returns per unit area than other agricultural or horticultural crops. In India, commercial floriculture has increased its credibility by diversification in agriculture and also evolved as a foreign exchange earner. India is bestowed with varied agro-climatic zones conducive for production of sensitive and delicate floriculture products.

Tuberose (*Agave amica* (Medik.) L. Family: Amaryllidaceae) is one of the most popular tropical ornamental bulbous flowering plants grown on a commercial scale throughout different states of India. They are much adored by the aesthetic world for their colour, elegance and fragrance. Tuberose occupies an important position among the commercially cultivated flowers in India due to its popularity as loose flower as well as cut flower. Loose flowers have great demand for making garland, veni, worshipping and offerings in religious functions and auspicious days. The long-lasting flower spikes are largely used for the vase decoration and preparation of bouquets. The aromatic oil extracted from the fragrant white flowers have great demand for essential oil industry, which fetches a very good price in the international market.

Climatic factor plays a critical role for successful production of tuberose. Tropical to subtropical and temperate climates are best suited for its cultivation. Tuberose prefer to grow in open sunny location. It grows in mild climate without extremes of high or low temperature. It can be successfully grown under warm humid areas having temperature around 30°C and approximate day length of 16 hours which make it suitable for cultivation under Kerala condition.

The quality and production of any crop or variety largely influenced by its genetic constitution and climatic condition under which they are grown. Hence, the cultivar which performs well in one region may not perform well in other regions of varied agro-climatic conditions. Therefore, in order to select suitable and high yielding cultivar for a particular region it is very much necessary to collect and evaluate all the available genotypes.

Smart farming is a comprehensive system designed to optimize production. Using the key elements of information, technology, and management, smart farming can be used to increase production efficiency, improve product quality, enhance the efficiency of crop chemical use, conserve energy and protect the environment. This experiment is carried out as a pre-evaluation study to further evaluate smart farming techniques in tuberose for its commercial cultivation.

2. MATERIAL AND METHODS

A genetic screening of different varieties of tuberose (*Agave amica* (Medik.)) were carried out at College of Agriculture, Padannakkad, Kasargod during 2019-2020. The experiment was laid out in Randomized block design with ten accessions (Plate 1.) viz. A. Shringar, A. Niranthara, A. Prajwal, A. Sugandhi, Culcutta Single, Bidhan Ujwal Phule Rajani (single type), A. Vaibhav (semi-double), A. Suvasini, Culcutta Double (double type). Treatments were replicated thrice. Bulbs of uniform size were planted at a depth of 5 cm at a spacing of 20cm x 25 cm. Uniform cultural operations and crop management practices were followed for all varieties. Fertilizers, farm yard manure and lime were given as per the package of practices recommendations for crops (KAU, 2016). Five plants were randomly selected and tagged from each treatment and replication for the purpose of recording observation.




Plate 1. Tuberose varieties used for the experiment

3. RESULTS AND DISCUSSION

The study indicated that genotypic differences among varieties and variation in environmental factors significantly influence vegetative and floral characters of tuberose. Varietal influence on vegetative and floral characters are listed in Table 1 and Fig. 1.

The maximum plant height was observed in variety Arka Prajwal (54.27 cm) and minimum in Bidhan Ujwal (31.60 cm). The variation among the plant height might be due to the differences in the genetic makeup and the diversified origin of the cultivar, which results into changes in the expression of phenotype under specific geographical location. The results of variation in plant height due to genotypic differences and environmental variation are in agreement with the findings of Sateesha *et al.* (2011), Ranchana *et al.* (2015), Dimri (2017) in tuberose.

Duration from planting to flowering indicates the early or delayed flowering habitat of a cultivar, which is an important parameter for selecting a variety for a particular area. Early flower emergence is favourable for tuberose growers as it leads to early availability of flowers in market. Minimum number of days for 50 per cent flowering was observed in variety Arka Prajwal (108.38) followed by variety Bidhan Ujwal (108.62). The variation among blooming

period is also reported by Ramachandrudu and Thangam (2009), Singh *et al.* (2018) in tuberose and Simmy (2012) in gladiolus.

Flower related parameters like length of floret, diameter of floret and weight florets plays an important role when the quality of flower is concerned. For the better appearance of spike, it should have floret with bigger diameter. As the diameter of the floret increases tendency of crowding by overlapping will increase this will further enhance the beauty of spike. For the preparation of garlands, veni etc. longest florets are more attractive. In the present experiment maximum length of floret was observed in variety Culcutta Single (7.03 cm) which was statistically on par with Arka Prajwal and Arka Niranthara (6.97 and 6.69 cm, respectively). Maximum diameter of floret was observed in variety Arka Prajwal (3.53 cm) followed by Culcutta Double and Arka Suvasini (3.14 and 2.79 cm, respectively). Similar finding in tuberose was observed by Bindiya *et al.* (2018).

Longer spike length is a desirable character for cut flower. In the present study maximum length of spike was observed in variety Arka Prajwal (111.80 cm) followed by Arka Suvasini (109.07 cm) which was on par. The longest spike in Arka Prajwal may be due to more internodal distance. The variation in spike length is also reported by Susila (2013), Krishnamoorthy (2014), Prakash *et al.* (2015) and Bindiya *et al.* (2018) in tuberose. This variation among spike length might be due to difference in genetic makeup.

Maximum loose flower yield per unit area were recorded by the variety Arka Suvasini followed by Arka Prajwal which was on par (2626.26 and 2125.50 gm⁻², respectively). Increased floret size and weight of individual florets leads to increased yield of loose flower. This is supported with the findings of Mahoviya (2003) in tuberose.



Fig. 1 Tuberose varietal influence on plant height, spike length, floret length and diameter of floret

Table 1. Tuberose varietal influence on vegetative and floral characters

Treatments	Plant height (cm)	Days to 50 % flowering	Length of floret (cm)	Diameter of floret (cm)	Length of spike (cm)	Loose flower yield per unit area (g/m ²)
T1	47.35	110.09	5.68	2.43	91.17	1545.73
T2	54.27	108.38	6.97	3.53	111.80	2125.50
T3	38.60	124.61	6.69	2.60	92.53	1024.67
T4	35.39	108.83	6.07	2.51	75.20	1029.33
T5	48.33	111.23	5.28	2.79	109.07	2626.26
T6	32.00	109.03	5.59	2.21	65.87	1049.80
T7	31.60	108.62	5.65	2.67	65.73	1026.50
T8	45.07	126.15	5.78	2.65	76.60	986.00
T9	46.48	122.93	7.03	2.38	106.00	849.47
T10	40.54	126.03	5.78	3.14	101.07	2086.58
C.D (0.05)	3.75	2.83	0.41	0.44	13.73	564.41
SEm (±)	1.25	0.95	0.14	0.15	4.59	188.50

Commented [k6]: Rewrite as Varietal influence on vegetative and Floral traits of tuberose

4. CONCLUSION

Out of the ten genotypes of tuberose *A. Prajwal* has been found suitable for commercial farming with highest plant height, minimum days to 50% flowering, longest floret, highest diameter of floret, longest spike, maximum loose flower yield per unit area and good aroma. Tuberose needs an improved package of practices for better yield and quality of flowers. In view of these, further evaluation of *A. Prajwal* for standardization of manuring, drip irrigation, fertilization, spacing and mulching for precision farming is undergoing at Department of Floriculture and Landscaping at College of Agriculture, Vellayani, Thiruvananthapuram.

Commented [k7]: In smart farming systems

REFERENCES

- Kerala Agricultural University (KAU). (2016). *Package of practices recommendations: Crops* (14th ed.). Kerala Agricultural University, Thrissur, India.
- Sateesha, G. R., Kumar, A., & Biradar, M. S. (2011). Performance of different tuberose (*Polianthes tuberosa* L.) varieties under field conditions. *Journal of Plant Archives*, 11(1), 359–360.
- Ranchana, P., Kannan, M., & Jawaharlal, M. (2015). Chemical composition of concrete content of tuberose (*Polianthes tuberosa* L.) growing in Tamil Nadu. *Progressive Horticulture*, 47(2), 267–270.
- Dimri, S. (2017). *Screening of suitable germplasm of tuberose (Polianthes tuberosa L.) for mid hill conditions of Garhwal Himalayas* (Master's thesis). VCSG Uttarakhand University of Horticulture and Forestry, Uttarakhand, India.
- Ramachandrudu, K., & Thangam, M. (2009). Performance of tuberose (*Polianthes tuberosa* L.) cultivars in Goa. *Journal of Horticultural Science*, 4(1), 76–77.
- Singh, A., Singh, A. K., Sisodia, A., Padhi, M., & Pal, A. K. (2018). Performance of tuberose varieties for flowering and flower yield parameters under Indo-Gangetic plains of eastern Uttar

Pradesh, India. *International Journal of Current Microbiology and Applied Sciences*, 7(8), 1129–1133.

Simmy, A. M. (2004). *Enhancement of spike qualities of gladiolus* (Master's thesis). Kerala Agricultural University, Thrissur, India

Bindiya, C., Naik, B., Kamble, S., Tirakannavar, S., & Parit, S. (2018). Evaluation of different genotypes of tuberose (*Polianthes tuberosa* L.) for growth, flowering, and yield characters. *International Journal of Current Microbiology and Applied Sciences*, 7(7), 4135–4141.

Susila, T. (2013). Performance of tuberose cultivars under north coastal Andhra Pradesh. *Indian Journal*, 33(2), 161–162.

Krishnamoorthy, V. (2014). Assessment of tuberose (*Polianthes tuberosa* L.) varieties for growth and yield characters. *Asian Journal of Horticulture*, 9(2), 515–517.

Prakash, S., Arya, J. K., Singh, R. K., & Singh, K. P. (2015). Varietal performance of tuberose in Muzaffarnagar under western plain zone conditions. *Asian Journal of Horticulture*, 10(1), 149–152.

Mahoviya, R. (2003). *Studies on performance of different tuberose (Polianthes tuberosa Linn) cultivars and their vase life* (Master's thesis). Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, Madhya Pradesh, India.

Commented [k8]: Follow the APA style of reference and it should be in Alphabetical order