

## EFFECT OF GA<sub>3</sub> AND TIME OF SEED SOAKING ON GERMINATION AND SEEDLING VIGOUR OF SANDALWOOD

### ABSTRACT

The experiment entitled, "effect of GA<sub>3</sub> and time of seed soaking on germination and seedling vigour of sandalwood" was carried out under greenhouse condition with two factors *i.e.*, different GA<sub>3</sub> levels viz., Control (g<sub>0</sub>), seed soaking in GA<sub>3</sub> 100 ppm (g<sub>1</sub>), seed soaking in GA<sub>3</sub> 200 ppm (g<sub>2</sub>), seed soaking levels viz., in GA<sub>3</sub> 300 ppm (g<sub>3</sub>), seed soaking in GA<sub>3</sub> 400 ppm (g<sub>4</sub>), seed soaking in GA<sub>3</sub> 500 ppm (g<sub>5</sub>) and seed soaking for 12 hours (t<sub>1</sub>), seed soaking for 18 hours (t<sub>2</sub>) and seed soaking for 24 hours (t<sub>3</sub>). The results of the study revealed that, significantly minimum days for germination (21.44), higher survival percentage at 60 days after sowing (65.35), number of roots per seedling at the time of final survival (4.40), maximum length of roots per seedling at 60 days after sowing (7.58 cm) and height of plant at 90 days after sowing (12.27 cm), maximum survival percentage (100) at 15 days after bagging and stem diameter at 90 days after sowing (0.24 cm) was recorded with treatment g<sub>5</sub> (Seed soaking in GA<sub>3</sub> 500 ppm). From the pooled data, significantly early germination (22.00 days), maximum survival percentage (59.87) at 60 days after sowing, number of roots (4.00) per seedling at the time of final survival, length of roots per seedling (7.76 cm) at 60 days after sowing, height of plant at 90 days after sowing (12.03 cm), maximum survival percentage (99.69) at 15 days after bagging and stem diameter at 90 days after sowing (0.23 cm) was recorded with treatment t<sub>3</sub> (Seed soaking for 24 hours). From the present investigation, sandalwood seed soaking treatment with GA<sub>3</sub> 500 ppm for 24 hours recorded early germination, higher survival and superior in seedling vigour.

**Key word:** GA<sub>3</sub>, Green house, Seed, Sandalwood and Time of soaking.

### INTRODUCTION

Sandalwood (*Santalum album* L.) is the fragrant heartwood of some species of genus *Santalum*. The widely distributed and economically important *Santalum* genus belongs to the family *Santalaceae*. Sandal is one of the most primitive precious useful plants since ancient times. Ecologically sandal has adapted to various agroclimatic and soil conditions for in situ regeneration with an exception of waterlogged areas and very cold places. Sapwood is white to pale yellow, sharply demarcated from the yellowish brown to dark brown heartwood, hard, heavy, lustrous

**Commented [P1]:** Soaking in water/cold water/warm water/normal water, specify?

**Commented [P2]:** Several passive sentences are used in the abstract section; explanations of outcomes are unclear and can be rewritten for better clarity.

**Commented [P3]:** Reference?

and straight-grained to slightly wavy, fine textured with pleasant characteristic smell (Rao et al., 2007). Sandalwood trees are famous and very costly because of its fragrant heartwood and oil.

The various PGR's treatment for specific period of time also helpful for early growth phases. GA triggers the respiratory and metabolic activities and activates certain genes resulting in the production of various components required for the active growth and development of embryo. Thus GA overcomes the dormancy and induces germination. In case of sandal, the basic problem is seed germination artificially. The seeds were treated with hot and cold water, GA<sub>3</sub> in different concentrations for its artificial germination.

After a certain period, seedlings were started to germinate. By observing all these peculiarities the germinated seeds were transferred to polypots and hycopots. Now, it is being realized that the only way to counter smuggling activities and to meet demand for sandalwood is to encourage sandal domestication, which may be the only option. But, an easy technology is to be explored for the plantation to be taken up by the general people. Keeping all these views in mind, the authors undertook the venture for the same since over a couple of years for its propagation and survivality to explore the areas of suitable as well as congenial edaphic factors. Day to day growth and development were observed and recorded, all these data were properly which have been exhibited in this context.

The main aims and objectives are to study the integrated scientific approach of seed germination, its growth behaviour in nursery for production of quality and healthy planting stock and raising successful plantations subsequently. There has been no any work related to the seed germination of sandalwood was carried out under Gujarat state, so, keeping above things in mind, the attempt was being taken to study the "effect of GA<sub>3</sub> and time of seed soaking on germination and seedling vigour of sandalwood" was carried out under greenhouse conditions.

## MATERIAL AND METHODS

An experiment entitled, effect of GA<sub>3</sub> and time of soaking on germination of sandalwood seed, was carried out under greenhouse condition at Department of Horticulture, C. P. College of Agriculture, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar. Present experiment were evaluated with two factors viz., different GA<sub>3</sub> levels viz., Control (g<sub>0</sub>), seed soaking in GA<sub>3</sub> 100 ppm (g<sub>1</sub>), seed soaking in GA<sub>3</sub> 200 ppm (g<sub>2</sub>), seed soaking in GA<sub>3</sub> 300 ppm (g<sub>3</sub>), seed

**Commented [P4]:** Add full form of PGRs. Add some relevant studies on PGRs importance as seed treatment and seedling growth, with references.

**Commented [WU5]:** Reference?

**Commented [P6]:** Discuss the gap of previous studies in Santalum album and highlight the importance of present study in the Introduction part

soaking in GA<sub>3</sub> 400 ppm (g<sub>4</sub>), seed soaking in GA<sub>3</sub> 500 ppm (g<sub>5</sub>) and seed soaking for 12 hours (t<sub>1</sub>), seed soaking for 18 hours (t<sub>2</sub>) and seed soaking for 24 hours (t<sub>3</sub>).

**Commented [WU7]:** Clarify the seed soaking method. Water soaking/cold water/hot water/normal water, etc.

Seeds of White Sandal (*Santalum album* L.) were tried in three trials viz., First trial: 17<sup>th</sup> May to 17<sup>th</sup> August, Second trial: 22<sup>nd</sup> July to 22<sup>nd</sup> October, Third trial: 27<sup>th</sup> September to 27<sup>th</sup> December were carried out in a plug tray under greenhouse conditions. Seed treatment of GA<sub>3</sub> was given as per treatment and standard media i.e. Vermiculite: Poultry Manure: Vermicompost (2:1:1) were used on volume basis in plug tray. The experiment was laid out in Factorial Completely Randomized Design as described by Nigam and Gupta (1979) with three repetitions. The treatments were evaluated and observations were recorded periodically in relation to germination, survival and seedling vigour.

**Commented [WU8]:** Replications or repetitions?. If replications, how many replications were made for each treatment?

**Commented [P9]:** Add standard methodologies, formulas, statistical reference in the methodology part.

## RESULTS AND DISCUSSION

### Effect of time of seed soaking (T):

The pooled data of three trial revealed that the significantly early germination (22.00 days), maximum survival percentage (59.87) at 60 days after sowing, number of roots (4.00) per seedling at the time of final survival, length of roots per seedling (7.76 cm) at 60 days after sowing and height of plant at 90 days after sowing (12.03 cm) were recorded with treatment t<sub>3</sub> (Seed soaking for 24 hours) and which was significantly superior over all other treatments.

From the pooled data, it was noted that not significant the maximum survival percentage (99.69) at 15 days after bagging and stem diameter at 90 days after sowing (0.23 cm) was recorded with treatment t<sub>3</sub> (Seed soaking for 24 hours).

It was observed that soaking of seeds in gibberellic acid is one of the most successful methods for accelerating the germination of the seeds of sandalwood (Rai, 1990, and Tennakoon et al., 2000). Similar results have been reported by Raviraja Shetty et al. (2016) in *Celastrus paniculatus* a wild threatened medicinal plant, where the seeds treated with lower concentration of GA<sub>3</sub> i.e. 400ppm.

**Commented [P10]:** Delete

**Commented [P11]:** Delete, Modify the reference part also.

**Formatted:** Subscript

### Effect of level of GA<sub>3</sub> (G):

The results of three trial pooled data found that significantly minimum days for germination (21.44), higher survival percentage at 60 days after sowing (65.35) and number of roots per seedling

at the time of final survival (4.40) was recorded with treatment g<sub>5</sub> (Seed soaking in GA<sub>3</sub> 500 ppm), which was significantly superior over other treatments. The significantly maximum length of roots per seedling at 60 days after sowing (7.58 cm) and height of plant at 90 days after sowing (12.27 cm) were recorded with treatment g<sub>5</sub> (Seed soaking in GA<sub>3</sub> 500 ppm) and it was statistically at par with treatment g<sub>4</sub> (Seed soaking in GA<sub>3</sub> 400 ppm). The cytological basis of gibberellic acid involves the promotion of cell elongation and cell division ultimately increasing the length of root and shoot and also the plant height (Achard et al., 2009).

From the pooled data, it was noted that not significant survival percentage (100) at 15 days after bagging and stem diameter at 90 days after sowing (0.24 cm) was recorded with treatment g<sub>5</sub> (Seed soaking in GA<sub>3</sub> 500 ppm).

Gibberellic acid is known to induce the expression of genes encoding enzymes that mobilize the reserved food material stored in the endosperm which includes starch, protein and lipids. This helps in increasing the biological yield of the plant. The overall increase in the growth of the plant is due to gibberellic acid which leads to the increase in rate of photosynthesis (Patel et al., 2018).

## Conclusion

From the present investigation, sandalwood seed soaking treatment with GA<sub>3</sub> 500 ppm for 24 hours recorded early germination, higher survival and superior in seedling vigour.

## REFERENCES

- Achard P, Gusti A, Cheminant S, Alioua M, Dhondt S, Coppens F. (2009). Gibberellin signaling controls cell proliferation rate in *Arabidopsis*. *Current Biology*. 19:1188–1193.
- Nigam A.K. and Gupta V.K. (1979). Handbook on Analysis of Agriculture Experiments. *Indian Agricultural Statistics Research Institute*. New Delhi. pp 39-54.
- Patel HS, Tandel MB, Prajapati VM, Amlani MH, Prajapati DH. (2018). Effect of different pre-sowing treatments on germination of Red sanders (*Pterocarpus santalinus* L. f.) in net house condition. *International Journal of Chemical Studies*. 6(2):876-879.
- Rai SN. (1990). Status and cultivation of sandalwood in India. Proceedings of the symposium on sandalwood in the Pacific, April 9-11, Honolulu-Hawaii, 1990, 66-71.

**Commented [WU12]:** Expand the discussion section with related study work on seed pre treatment by additional species of the same genus, family, or others with similar recommendations.

**Formatted:** Font: Italic

**Formatted:** Font: Italic

Rao RV, Hemavathi TR, Sujatha M, Shashikala S (2007). Status on Wood structure, heartwood formation, age, gaps and research needs in *Santalum album* L. Proceedings of National Seminar held in Institute of Wood Science and Technology (ICFRE), 12-13 December, Bangalore, India pp 165-171.

Raviraja Shetty G, Poojitha KG, Souravi K, Rajasekharan PE. (2016). Seed germination studies in *Celastrus paniculatus* Willd: A threatened medicinal a plant. *International Journal of Applied and Agricultural Research*. 11(1):51-56.

Tennekoon KU, Ekanayake SP, Etampawala ERLB. (2000). An overview of *Santalum album* research in SriLanka. *Sandalwood Research Newsletter.*; 11:1-4.

**Formatted:** Font: Italic

**Commented [P13]:** Correct the spelling of author after comparing from the text.

UNDER PEER REVIEW

**Table: 1 Effect of GA<sub>3</sub> and time of seed soaking on germination and survival of sandalwood (*Santalum album* L.)**

Treatments	Days taken for germination				Survival percentage at 60 days after sowing				Survival percentage at 15 days after bagging			
	Trial I	Trial II	Trial III	Pooled	Trial I	Trial II	Trial III	Pooled	Trial I	Trial II	Trial III	Pooled
<b>Time of seed soaking (T):</b>												
<b>t<sub>1</sub> (Seed soaking for 12 hours)</b>	22.72	23.16	23.33	23.07	50.16	53.40	51.97	51.84	99.53	99.53	94.44	97.83
<b>t<sub>2</sub> (Seed soaking for 18 hours)</b>	22.44	23.33	22.77	22.85	52.21	54.86	52.96	53.34	98.61	99.53	100.0	99.38
<b>t<sub>3</sub> (Seed soaking for 24 hours)</b>	21.61	22.33	22.05	22.00	58.33	60.73	60.55	59.87	99.07	100.0	100.0	99.69
<b>S.Em.±</b>	0.17	0.31	0.24	0.14	0.92	0.79	0.64	0.45	0.75	0.37	3.20	1.10
<b>C.D. at 5 %</b>	0.50	NS	0.70	0.40	2.64	2.28	1.85	1.29	NS	NS	NS	NS
<b>GA<sub>3</sub> levels (G):</b>												
<b>g<sub>0</sub> (Control)</b>	24.55	24.77	25.33	24.88	46.29	47.40	48.14	47.28	100.0	100.0	88.88	96.29
<b>g<sub>1</sub> (Seed soaking in GA<sub>3</sub> 100 ppm)</b>	22.22	23.11	22.88	22.74	51.47	52.18	51.84	51.83	97.22	99.07	100.0	98.76
<b>g<sub>2</sub> (Seed soaking in GA<sub>3</sub> 200 ppm)</b>	21.88	22.44	22.33	22.22	50.73	54.70	53.70	53.04	98.14	100.0	100.0	99.38
<b>g<sub>3</sub> (Seed soaking in GA<sub>3</sub> 300 ppm)</b>	22.00	22.88	22.66	22.51	51.47	56.92	55.18	54.52	99.07	100.0	100.0	99.69
<b>g<sub>4</sub> (Seed soaking in GA<sub>3</sub> 400 ppm)</b>	21.66	22.66	21.77	22.03	55.55	61.36	57.36	58.09	100.0	99.07	100.0	99.69
<b>g<sub>5</sub> (Seed soaking in GA<sub>3</sub> 500 ppm)</b>	21.22	21.77	21.33	21.44	65.88	65.44	64.73	65.35	100.0	100.0	100.0	100.0
<b>S.Em.±</b>	0.24	0.44	0.34	0.20	1.30	1.12	0.91	0.65	1.07	0.53	4.53	1.56
<b>C.D. at 5 %</b>	0.71	1.26	1.00	0.57	3.73	3.23	2.62	1.82	NS	NS	NS	NS
<b>T X G</b>												
<b>S.Em.±</b>	0.43	0.76	0.60	0.35	2.25	1.95	1.58	1.12	1.85	0.92	7.85	2.70
<b>C.D. at 5 %</b>	1.23	NS	NS	NS	6.47	5.60	4.54	3.16	NS	NS	NS	NS
<b>CV %</b>	3.35	5.78	4.60	4.71	7.29	6.00	4.97	6.14	3.24	1.61	13.27	8.21

**Table: 2 Effect of GA<sub>3</sub> and time of seed soaking on number of roots and length of roots per seedling of sandalwood (*Santalum album* L.)**

Treatments	Number of roots per seedling at the time of final survival				Length of roots per seedling (cm) at 60 days after sowing			
	Trial I	Trial II	Trial III	Pooled	Trial I	Trial II	Trial III	Pooled
<b>Time of seed soaking (T):</b>								
t <sub>1</sub> (Seed soaking for 12 hours)	3.55	3.66	3.88	3.70	6.22	5.75	5.77	5.91
t <sub>2</sub> (Seed soaking for 18 hours)	3.83	3.61	3.66	3.70	6.18	6.25	6.24	6.22
t <sub>3</sub> (Seed soaking for 24 hours)	3.94	3.88	4.16	4.00	8.25	7.41	7.63	7.76
S.Em.±	0.10	0.11	0.10	0.06	0.23	0.22	0.15	0.12
C.D. at 5 %	0.29	NS	0.29	0.17	0.67	0.65	0.44	0.33
<b>GA<sub>3</sub> levels (G):</b>								
g <sub>0</sub> (Control)	3.22	3.11	3.44	3.25	5.55	4.98	4.98	5.17
g <sub>1</sub> (Seed soaking in GA <sub>3</sub> 100 ppm)	3.44	3.55	3.66	3.55	6.04	5.96	6.08	6.03
g <sub>2</sub> (Seed soaking in GA <sub>3</sub> 200 ppm)	4.00	3.66	3.88	3.85	7.13	6.68	6.55	6.79
g <sub>3</sub> (Seed soaking in GA <sub>3</sub> 300 ppm)	3.66	3.66	3.88	3.74	7.00	6.77	6.60	6.79
g <sub>4</sub> (Seed soaking in GA <sub>3</sub> 400 ppm)	3.88	4.00	4.11	4.00	7.80	7.10	7.46	7.45
g <sub>5</sub> (Seed soaking in GA <sub>3</sub> 500 ppm)	4.44	4.33	4.44	4.40	7.78	7.33	7.62	7.58
S.Em.±	0.14	0.16	0.14	0.08	0.33	0.32	0.21	0.17
C.D. at 5 %	0.41	0.47	0.41	0.24	0.95	0.91	0.62	0.47
<b>T X G</b>								
S.Em.±	0.24	0.28	0.24	0.15	0.57	0.55	0.37	0.29
C.D. at 5 %	NS	NS	NS	NS	1.64	NS	1.08	0.82
CV %	11.39	13.18	11.01	11.87	14.43	14.83	10.00	13.30

**Table: 3 Effect of GA<sub>3</sub> and time of seed soaking on stem diameter and height of plant at 90 DAS of sandalwood (*Santalum album* L.)**

Treatments	Stem diameter at 90 days after sowing (cm)				Height of plant at 90 days after sowing (cm)			
	Trial I	Trial II	Trial III	Pooled	Trial I	Trial II	Trial III	Pooled
<b>Time of seed soaking (T):</b>								
<b>t<sub>1</sub> (Seed soaking for 12 hours)</b>	0.21	0.20	0.21	0.21	11.00	10.23	10.30	10.51
<b>t<sub>2</sub> (Seed soaking for 18 hours)</b>	0.21	0.21	0.21	0.21	12.24	10.57	10.73	11.18
<b>t<sub>3</sub> (Seed soaking for 24 hours)</b>	0.22	0.22	0.23	0.23	13.02	10.40	11.68	12.03
<b>S.Em.±</b>	0.01	0.009	0.01	0.006	0.16	0.23	0.15	0.10
<b>C.D. at 5 %</b>	NS	NS	NS	NS	0.46	0.68	0.44	0.30
<b>GA<sub>3</sub> levels (G):</b>								
<b>g<sub>0</sub> (Control)</b>	0.20	0.20	0.20	0.20	10.54	9.11	9.15	9.60
<b>g<sub>1</sub> (Seed soaking in GA<sub>3</sub> 100 ppm)</b>	0.22	0.20	0.20	0.20	11.12	10.18	10.50	10.60
<b>g<sub>2</sub> (Seed soaking in GA<sub>3</sub> 200 ppm)</b>	0.20	0.21	0.21	0.20	12.34	10.94	10.90	11.39
<b>g<sub>3</sub> (Seed soaking in GA<sub>3</sub> 300 ppm)</b>	0.22	0.22	0.24	0.22	12.67	11.12	11.38	11.72
<b>g<sub>4</sub> (Seed soaking in GA<sub>3</sub> 400 ppm)</b>	0.23	0.22	0.23	0.22	10.74	11.20	11.62	11.85
<b>g<sub>5</sub> (Seed soaking in GA<sub>3</sub> 500 ppm)</b>	0.24	0.24	0.24	0.24	13.11	11.85	11.86	12.27
<b>S.Em.±</b>	0.01	0.01	0.01	0.008	0.23	0.33	0.21	0.15
<b>C.D. at 5 %</b>	NS	NS	NS	NS	0.66	0.97	0.62	0.43
<b>T X G</b>								
<b>S.Em.±</b>	0.02	0.02	0.02	0.01	0.39	0.58	0.37	0.26
<b>C.D. at 5 %</b>	NS	NS	NS	NS	NS	NS	NS	NS
<b>CV %</b>	19.53	17.76	18.37	18.58	5.71	9.46	5.99	7.14