

## Comparative Analysis of Phytochemical Composition, Proximate Content, and Heavy Metal Contamination in Naturally and Calcium Carbide-Ripened Bananas."

### ABSTRACT

The use of artificial ripening agents, particularly calcium carbide, in fruits such as bananas has raised concerns about their impact on nutritional quality. This study compared the phytochemical composition, proximate analysis, and heavy metal contamination of naturally ripened (NR) and calcium carbide-ripened (CR) bananas and analysed the adverse risk associated with the consumption of calcium carbide -ripened bananas.

Mature unripe bananas obtained from the Institute of Agricultural Research and Development (IARD) were divided into two groups: one ripened naturally (NR) at room temperature and the other ripened using calcium carbide (CR). Phytochemical analysis detected phenols, alkaloids, flavonoids, tannins, saponins, and terpenoids in both groups, but with significant differences in concentration. Notably, cyanogenic glycoside and oxalate levels were higher in CR bananas ( $7.0 \times 10^{-4}\%$  and 1.82%, respectively) compared to NR bananas ( $6.0 \times 10^{-4}\%$  and 0.50%). Additionally, NR bananas showed higher concentrations of flavonoids (11.64%), alkaloids (4.50%), saponins (9.84%), and tannins ( $7.5 \times 10^{-4}\%$ ), indicative of high antioxidant properties. In contrast, CR bananas had reduced levels of these compounds (10.22%, 4.2%, 6.41%, and  $3.16 \times 10^{-4}\%$  respectively), suggesting decreased antioxidant potential.

Proximate analysis revealed that CR bananas had higher moisture content (82.05%) compared to NR bananas (79.83%). However, NR bananas contained higher levels of protein (1.88%), carbohydrate (19.68%), and fiber (2.40%).

Heavy metal analysis identified elevated phosphorus levels in CR bananas (184.5 ppm), exceeding the FAO's permissible limits (80–120 ppm).

In conclusion, our findings highlight the potential nutritional compromises and safety risks associated with calcium carbide-ripened bananas. The study emphasizes the need for strict regulation of artificial ripening agents to protect consumer health.

**Keywords:** Calcium Carbide, Banana, Phytochemical analysis, Proximate analysis, Heavy Metals, Artificial ripening, Food safety.

## Introduction

Bananas (*Musa spp.*) are one of the most popular and widely consumed fruits globally, recognized for their rich nutritional content and health benefits. They are a significant source of vitamins, minerals, dietary fibre, and bioactive compounds, making them a staple in many diets around the world (Annor et al., 2016). Ripening is a genetically programmed highly coordinated irreversible phenomenon which includes many biochemical changes including tissue softening, pigment changes, aroma and flavour volatile production, reduction in astringency, and many others (Maduwanthi, et al 2019). In many developing countries, bananas play a crucial role in food security and are consumed either raw or as processed products. However, due to their perishable nature, bananas are often subjected to artificial ripening processes to meet consumer demand and extend their market availability (Olufemi, 2024)

Calcium carbide, a common chemical agent used for artificial ripening, has raised significant health and safety concerns. When calcium carbide encounters moisture, it produces acetylene gas, which accelerates the ripening process. Ripening is initiated by allowing the temperature to rise to 15 to 18°C and by introducing ethylene gas (usually 1000 ppm) and closing rooms for 24 hr and then ventilating and lowering the temperature to 15°C. (Marriott, 1980) This method, while efficient, may introduce harmful substances into the fruit, such as phosphine, and negatively impact the fruit's nutritional composition (Islamiyat et al., 2016). Several studies have highlighted the adverse effects of calcium carbide on the quality and safety of ripened fruits, including potential toxicities and nutrient degradation (Asif, 2012; Igbinađuwa et al., 2022). Despite these risks, calcium carbide remains widely used in many countries, including Nigeria, due to a lack of regulatory enforcement and consumer awareness (Ibeachu and Nwidobie)

The nutritional and phytochemical profile of fruits, including bananas, is largely influenced by the ripening process. Naturally ripened bananas undergo a series of biochemical changes that enhance their sweetness, flavor, and nutritional value (Ukwo et al., 2023). Conversely, artificially ripened fruits may experience an altered balance of bioactive compounds, potentially affecting their health benefits. Phytochemicals, such as phenols, flavonoids, and alkaloids, are particularly sensitive to the ripening process and play a crucial role in antioxidant defense and disease prevention (Izundu et al., 2016).

Given the widespread use of calcium carbide as a ripening agent and the growing concerns about its safety and associated health risk, this study aims to compare the phytochemical composition, proximate analysis, and heavy metal content of naturally ripened and calcium carbide-ripened banana fruits. By analyzing the differences in these parameters, the study seeks to provide insight into the potential health risks associated with consuming artificially ripened fruits, as well as highlight the importance of adopting safer, natural ripening methods. The findings of this study will contribute to the growing body of knowledge regarding food safety and quality and inform public health policies on the use of ripening agents in fruit production.

## **2 METHODOLOGY**

### **2.1 SAMPLE COLLECTION AND PREPARATION**

**Banana Acquisition:** Mature unripe bananas were sourced from the Institute of Agricultural Research and Development (IARD). The bananas were then split into two equal groups. One group was left to ripen naturally at room temperature, while the other was artificially ripened with calcium carbide. For artificial ripening, 5 g of calcium carbide was applied per 1 kg of bananas, in accordance with the method outlined by Igbinauwa et al. (2018) and similar techniques used in agricultural ripening studies (Oboh et al., 2016).

**Sample Processing:** Selected ripe banana samples (both naturally and artificially ripened) were washed with distilled water, sliced, sun-dried, ground into powder using a mortar and pestle, and then sieved through a 2 mm sieve. The powdered samples were stored in airtight containers until further analysis.

## 2.2 PHYTOCHEMICAL ANALYSIS

**2.2.1 Terpenoids:** About 5 ml of each extract was mixed with 2 ml of chloroform and 3 ml of concentrated  $H_2SO_4$ . A reddish-brown coloration indicated the presence of terpenoids (Ejikeme et al., 2014).

**2.2.2 Steroids:** A mixture of 2 ml of acetic anhydride and 0.5 g of ethanol extract was treated with 2 ml of  $H_2SO_4$ , changing color from violet to blue/green to indicate the presence of steroids.

**2.2.3 Alkaloids:** Alkaloids were extracted by weighing 5 g of the dried sample into a 250 ml beaker, adding 200 ml of 10% acetic acid in ethanol, and allowing it to stand for 4 hours before filtration and precipitation with concentrated ammonium hydroxide (Harborne, 1973).

**2.2.4 Saponins:** A 20 g sample was extracted with 100 ml of 20% aqueous ethanol, and heated for 4 hours, and the combined extract was purified with n-butanol (Obadoni and Ochuko, 2001).

**2.2.5 Flavonoids:** Extract 10 g of the sample with 100 ml of 80% aqueous methanol, filter, and evaporate to dryness (Sufowara, 1993).

**2.2.6 Tannins:** Extraction involved boiling 0.1 g of the sample in water, followed by color development with Folin-Denis reagent (Ejikeme et al., 2014).

**2.2.7. Phenols:** The sample was boiled in distilled water, treated with NaOH, and titrated with  $Na_2S_2O_2$ .

**2.2.8 Cyanogenic Compounds:** 5 g of sample was allowed to hydrolyze overnight before distillation and titration with silver nitrate (Hikino et al., 1984).

**2.2.9 Oxalates:** 25  $cm^3$  of the solution was titrated with potassium permanganate after heating with dilute sulfuric acid (Ejikeme et al., 2014).

**2.2.10. Phytate Acid:** A sample was extracted with HCl, neutralized, treated with  $FeCl_3$ , and analyzed calorimetrically.

## 2.3 PROXIMATE ANALYSIS

**2.3.1 Carbohydrate:** 0.1 g of the sample was treated with perchloric acid, and the resulting solution was analyzed using an Anthrone reagent.

**2.3.2 Protein:** Digestion involved concentrated sulfuric acid, followed by distillation and titration to determine nitrogen content.

**2.3.4 Moisture:** 1 g of sample was dried at 105°C, and moisture percentage was calculated.

**2.3.5 Lipid:** Lipids were extracted using a Soxhlet extractor with acetone.

**2.3.6 Ash:** The sample was incinerated in a muffle furnace, and ash content was calculated.

**2.3.7 Crude Fiber:** The sample was subjected to a series of extractions and weight measurements to determine fibre content.

## 2.4 HEAVY METAL ANALYSIS

Arsenic and Phosphorus: 5 g of dried sample was ashed at 500°C, dissolved in HNO<sub>3</sub>, and analyzed using Atomic Absorption Spectrophotometry (Miroslav and Vladimir, 1999).

## STATISTICAL ANALYSIS

Data were analyzed using one-way Analysis of Variance (ANOVA) with post-hoc LSD tests, considering results significant at  $p < 0.05$

## RESULTS

Table 1. **Phytochemical Analysis of natural ripened and Calcium Carbide Ripened Banana Fruit**

	%	%	%	%	%	%	%	%	%
	Cyanogenic glycoside	Phenols	Phylate	Oxalate	Alkaloids	Flavonoid	Tannin	Saponin	Terpenoid
<b>CR</b>	7.0 x 10 <sup>-4</sup>	0.094	1.0 x 10 <sup>-4</sup>	1.82	4.2	10.22	7.5 x 10 <sup>-4</sup>	6.41	+++
<b>NR</b>	6.0 x 10 <sup>-4</sup>	0.094	1.0 x 10 <sup>-4</sup>	0.50	4.50	11.64	3.16 x 10 <sup>-4</sup>	9.84	+++

## Phytochemical Analysis

The phytochemical analysis of naturally ripened (NR) and calcium carbide-ripened (CR) banana fruits revealed differences in the concentration of several compounds (Table 1). Cyanogenic glycoside levels were higher in calcium carbide-ripened bananas ( $7.0 \times 10^{-4}$ ) compared to naturally ripened bananas ( $6.0 \times 10^{-4}$ ), though the difference was not statistically significant ( $p > 0.05$ ). Phenols (0.094%) and phytate ( $1.0 \times 10^{-4}$ ) were consistent in both ripening methods.

However, oxalate and tannin levels were significantly elevated in calcium carbide-ripened bananas (1.82) and ( $7.5 \times 10^{-4}$ ) respectively) compared to the naturally ripened ones (0.50) and ( $3.16 \times 10^{-4}$ ). Flavonoid, Alkaloid, and saponin concentrations were higher in naturally ripened bananas (11.64%, 4.50%, and 9.84%, than in calcium carbide-ripened fruits, where alkaloids and saponin levels dropped to 4.2% and 6.41%, respectively. Terpenoid levels remained unchanged between the two groups.

**Table 2. Proximate Analysis of Natural Ripened and Calcium Carbide Ripened Banana Fruit**

	%	%	%	%	%	%	%
	Moisture	M/crushed	Ash	Lipid	Protein	Carbohydrates	Fibre
CR	82.05	83.03	0.21	1.23	1.61	14.16	0.49
NR	79.83	81.11	0.52	1.65	1.88	19.68	2.40

The proximate analysis showed a notable difference in moisture content, which was slightly higher in calcium carbide-ripened bananas (82.05%) compared to naturally ripened bananas (79.83%) (Table 2). The moisture content of calcium carbide-ripened bananas also translated to a crushed moisture content of 83.03%, compared to 81.11% in naturally ripened bananas.

Ash content was lower in calcium carbide-ripened bananas (0.21%) than in naturally ripened bananas (0.52%). The lipid content also dropped in calcium carbide-ripened bananas (1.23%) compared to naturally ripened bananas (1.65%). Similarly, protein levels in calcium carbide-ripened bananas (1.61%) were slightly reduced compared to naturally ripened ones (1.88%). Carbohydrate content was lower in calcium carbide-ripened bananas (14.16%) versus naturally ripened bananas (19.68%). Fiber content showed a sharp decline in calcium carbide-ripened bananas (0.49%) compared to naturally ripened fruits (2.40%).

**Table 3. Heavy Metal Analysis of Natural Ripened and Calcium Carbide Ripened Banana Fruit**

<b>Heavy metals</b>	<b>Naturally ripened (mg/kg)</b>	<b>Calcium carbide ripened (mg/kg)</b>	<b>FAO STANDARD (ppm)</b>
<b>Arsenic</b>	<b>Not detectable</b>	<b>Not detectable</b>	0.5 -2.0
<b>Phosphorous</b>	77.19	184.5	80 - 120

The heavy metal analysis indicated that arsenic was undetectable in both naturally ripened and calcium carbide-ripened bananas (Table3). However, there was a significant increase in phosphorus content in calcium carbide-ripened bananas (184.5 mg/kg) compared to naturally ripened bananas (77.19 mg/kg). The phosphorus content in calcium carbide-ripened bananas also exceeded the FAO standard of 80–120 ppm, highlighting the potential risk of heavy metal contamination from artificial ripening methods.

## **Discussion**

### **Phytochemical Analysis**

The phytochemical composition of both naturally ripened and calcium carbide-ripened banana fruits indicates the presence of several bioactive compounds such as phenols, tannins, alkaloids, saponins, flavonoids, and glycosides. These compounds are known for their medicinal

properties, which makes bananas a valuable source of health-promoting nutrients (Kasolo et al., 2010). However, significant differences were observed between the two ripening methods.

For instance, the cyanogenic glycoside content was found to be higher in calcium carbide-ripened bananas. This glycoside is a natural plant toxin, which can release cyanide, potentially causing serious health issues like growth retardation and neurological disorders such as neuropathy, seizures, encephalopathy, Amyotrophic lateral sclerosis (ALS) and dementia (Islamiyat et al., 2016). A similar trend was observed in other studies, which reported an increase in cyanogenic glycosides with fruit ripening (Alagbaoso et al., 2017). While the phenol content remained consistent between the two types of ripened bananas, phenolic compounds are known for their antioxidant properties and their ability to protect against diseases like cancer and cardiovascular conditions (Hollman, 2001).

The oxalate content was notably higher in calcium carbide-ripened bananas, which raises concerns about potential kidney issues due to the formation of calcium oxalate stones Gao et al., 2012). This is significant, as bananas are commonly consumed in large quantities, and the accumulation of oxalates over time could pose a health risk (Gbakon et al., 2018).

Interestingly, the alkaloid and flavonoid contents were higher in naturally ripened bananas. Alkaloids are valuable for their antimicrobial, analgesic, and antiplasmodic properties (Iheagwam et al., 2019), while flavonoids contribute to antioxidant defense mechanisms, helping prevent oxidative stress (Izundu et al., 2016). On the other hand, tannin levels were higher in calcium carbide-ripened bananas. Tannins have astringent properties and are beneficial for wound healing, but in higher doses, they can negatively affect nutrient absorption (Ugwu et al., 2003; Offor et al., 2015).

The study also found a higher concentration of saponins and terpenoids in naturally ripened bananas, which are known for their cleansing, pesticidal, and anticancer properties (Gurfinkel

& Rao, 2003). These findings suggest that naturally ripened bananas may offer superior health benefits compared to those ripened with calcium carbide.

### **Proximate Analysis**

The proximate composition of calcium carbide-ripened bananas revealed an increased moisture content (82.05%) compared to naturally ripened bananas (79.83%). This observation aligns with findings from Nahu et al. (2020), where the moisture content increased with the use of calcium carbide. The elevated moisture content is a result of weakened banana peel fiber, which allows for higher water absorption during ripening. Ash content, which reflects the inorganic mineral content, was found to be lower in calcium carbide-ripened bananas (0.21%) compared to naturally ripened bananas (0.52%), but both fell within the FAO's permissible limits (2.9%). Similar trends were reported by Nahu et al. (2020), where calcium carbide affected mineral absorption in banana peels, although the mineral content remained within safe levels.

The percentage of lipid content was also found to be lower in calcium carbide-ripened bananas, differing from reports by Adeyemi et al. (2018), where calcium carbide increased lipid levels. Protein content was significantly reduced in calcium carbide-ripened bananas, suggesting that calcium carbide may impair the nutritional quality of bananas, particularly their ability to provide essential amino acids and energy (Nura et al., 2018). Carbohydrate content followed a similar pattern, with higher levels in naturally ripened bananas (19.68%) compared to those ripened with calcium carbide (14.16%). This could be attributed to the increased respiration rate in calcium carbide-treated bananas, which accelerates the conversion of carbohydrates into carbon dioxide (Adeyemi et al., 2018). Fiber content was higher in naturally ripened bananas, which is important for digestive health. Calcium carbide-ripened bananas, on the other hand, showed a decrease in fiber content, which contrasts with other studies, such as Adeyemi et al. (2018), where calcium carbide was associated with increased fibre.

## **Heavy Metal Analysis**

The heavy metal analysis revealed no detectable levels of arsenic in either naturally ripened or calcium carbide-ripened bananas. This aligns with the findings of Igbinađuwa et al. (2018), who reported that laboratory-ripened bananas did not contain arsenic, although market-sourced bananas did. The absence of arsenic in this study suggests that proper handling and control of calcium carbide during ripening may minimize arsenic contamination.

However, phosphorus levels were significantly higher in calcium carbide-ripened bananas (184.5 mg/kg) compared to the naturally ripened ones (77.19 mg/kg), exceeding the FAO standard of 80-120 mg/kg (FAO and WHO, 1998). Prolonged consumption of phosphorus in such high quantities may lead to health issues such as memory loss, mental confusion, and seizures (Fattah & Ali, 2010). This finding is consistent with previous studies that have reported high phosphorus levels in calcium carbide-ripened bananas (Nuhu et al., 2020; Igbinađuwa et al., 2018).

## **Conclusion**

The study suggests that calcium carbide ripening significantly impacts the phytochemical and proximate composition of bananas, potentially reducing their nutritional quality and increasing the risk of exposure to harmful substances like cyanogenic glycosides and excessive phosphorus. While it accelerates ripening, calcium carbide poses nutritional and health risks, highlighting the need for stricter regulations and increased public awareness about artificial ripening agents. Given the global trend of banning calcium carbide in countries like India, Sri Lanka, and Bangladesh, it is crucial for Nigeria to implement similar measures to protect public health..

## Disclaimer (Artificial intelligence)

### Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

### Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

1. Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

2.

3.

## REFERENCES

- Adeyemi, O. S., Fashina, A. Y., Oguntibeju, O. O., & Oguntoyinbo, A. (2018). Effects of artificially ripened banana fruits on oxidative stress parameters in male Wistar rats. *Journal of Food Biochemistry*, 42(2), e12497.
- Alagbaoso, S. O., Nwachukwu, J., & Ibeanu, V. N. (2017). Comparative study of phytochemical and antioxidant properties of avocado pear with ripening. *International Journal of Biochemistry Research & Review*, 17(1), 1-10.
- Amic, D., Davidovic-Amic, D., Bešlo, D., & Trinajstić, N. (2003). Structure-radical scavenging activity relationships of flavonoids. *Croatica Chemica Acta*, 76(1), 55-61.
- Annor, G. A., Asamoah-Bonti, P., & Sakyi-Dawson, E. (2016). Fruit physical characteristics, proximate, mineral and starch characterization of FHIA 19 and FHIA 20 plantain and FHIA 03 cooking banana hybrids. *SpringerPlus*, 5, 1-12.

- Asif, M. (2012). Physico-chemical properties and toxic effect of fruit-ripening agent calcium carbide. *Annals of Tropical Medicine & Public Health*, 5(3).
- Ejikeme, P. M., Okwu, D. E., & Uche, A. (2014). Phytochemical screening and toxicity assessment of some medicinal plants in Nigeria. *African Journal of Plant Science*, 8(2), 92-98.
- Eleazu, C. O., Eleazu, K. C., & Awa, E. (2010). Phenolic compounds and antioxidant activity of palm fruit. *Journal of Medicinal Food*, 13(4), 1035-1040.
- F.A. O and W.H.O. (1998). Carbohydrates in Human nutrition. Reports of a Joint FAO/WHO expert consultation, Rome. FAO and Nutrition paper 66.
- FAO (2018). Banana Market Review. Preliminary Results for 2018. Food and Agriculture Organization of the United Nations, Rome.
- Fattah, S. A., & Ali, M. E. (2010). Phosphorus toxicity: Clinical effects and toxicology. *Journal of Environmental Science and Health, Part C*, 28(4), 352-375.
- Gao, Z., Zhang, J., Wang, H., & Wang, Y. (2012). Oxalate accumulation and regulation is closely related to the reproductive strategy in higher plants. *PLOS One*, 7(5), e36787.
- Gbakon, I. E., Awolola, G. V., & James, I. O. (2018). Oxalate content and health implications of commonly consumed vegetables in Northern Nigeria. *Nigerian Journal of Nutritional Sciences*, 39(1), 37-45.
- Gurfinkel, D. M., & Rao, A. V. (2003). Saponins as anticarcinogens. *Journal of Nutritional Biochemistry*, 14(6), 224-231.
- Harborne, J. B., & Harborne, J. B. (1973). Phenolic compounds. *Phytochemical methods: A guide to modern techniques of plant analysis*, 33-88.
- Herrmann, K. (1991). Flavonoids and fruit ripening. *Zeitschrift für Lebensmittel-Untersuchung und -Forschung*, 192(4), 426-429.
- Hikino, H., Kato, Y., & Mizuno, K. (1984). Studies on the mechanism of cyanogenic glycoside sformation in the seeds of *Sorghum bicolor* (L.) Moench. *Phytochemistry*, 23(9), 1871-1875.
- Hollman, P. C. (2001). Evidence for health benefits of plant phenols: Local or systemic effects? *Journal of the Science of Food and Agriculture*, 81(9), 842-852.
- Igbinaduwa, P. O., Aigbogun, C. O., & Okoh, C. I. (2018). Assessment of heavy metal contamination in banana fruits ripened with calcium carbide and naturally. *Nigerian Journal of Basic and Applied Sciences*, 26(1), 49-54.
- Igbinaduwa, P. O., Okeke, U. B., Dike, K. C., & Olohigbe, A. E. (2022). Effect of calcium carbide induced ripening on the vitamin C and mineral composition of banana (*Musa acuminata*) and papaya (*Carica papaya*) fruits sourced from Benin city, Nigeria. *West African Journal of Pharmacy*, 33(1), 92-98.
- Iheagwam, P. N., Onyeike, E. N., & Amadi, B. A. (2019). Phytochemical and vitamin contents of *Mangifera indica* (Mango) fruits subjected to ripening by artificial methods. *Int. J. Environ. Agric. Biotech*, 4(3), 677-684.

- Imam-Hossain, M., & Abu-Zubair, M. (2015). Phosphorus toxicity and its effects on health. *Journal of Environmental Health*, 78(1), 101-110.
- Islamiyat, FB; Moruf, OO; Sulaiman, AO; Adeladun, S.A (2016). Toxicology - New Aspects to this Scientific Conundrum. Intech Open Publishers, United Kingdom. Pp. 218.
- Izundu, A. I., & Ogbu, A. I. (2016). Flavonoid content and antioxidant activities in different plant species. *International Journal of Research in Pharmacy and Biosciences*, 3(8), 47-53.
- Joon, K., Kim, S., & Lee, Y. (2001). Respiration, sugar content, and ripening of artificially ripened banana fruits. *Postharvest Biology and Technology*, 24(3), 301-308.
- Kasolo, J. N., Bimenya, G. S., Ojok, L., Ochieng, J., & Ogwal-Okeng, J. W. (2010). Phytochemicals and uses of *Moringa oleifera* leaves in Ugandan traditional medicine. *Journal of Medicinal Plants Research*, 4(9), 753-757.
- Kim, K., Kim, J. H., & Cho, S. K. (2003). Anti-cancer properties of saponins in various vegetables. *Journal of the Korean Society of Food Science and Nutrition*, 32(7), 1235-1242.
- Materska, M., & Peruka, I. (2005). Flavonoid content and antioxidant activity of fruits and vegetables. *Polish Journal of Food and Nutrition Sciences*, 55(1), 15-20.
- Miroslav, P., & Vladimir, V. (1999). Method for determination of arsenic and phosphorus in food products. *Journal of Food Science*, 64(3), 439-442.
- Nuhu, H. S., Lamido, A., & Ahmad, B. (2020). Effects of calcium carbide on ripening, chemical composition, and shelf-life of banana fruits. *Journal of Food Science and Nutrition*, 58(1), 45-53.
- Obadoni, B. O., and Ochuko, P. E. (2001). Phytochemical studies and comparative efficacy of the crude extracts of some homeostatic plants in Edo and Delta States of Nigeria. *Global Journal of Pure and Applied Sciences*, 7(3), 385-388.
- Offor, I. F., and Akwagiobe, U. R. (2015). Tannin content and its health implications. *Journal of Medicinal Plants Research*, 9(30), 789-794.
- Price, K. R., and Fenwick, G. R. (1987). The chemistry and biological significance of saponins in foods and feedingstuffs. *Critical Reviews in Food Science and Nutrition*, 26(1), 27-135.
- Robinson, T. (1996). *The Organic Constituents of Higher Plants: Their Chemistry and Interrelationships*. Cordus Press.
- Rodríguez, K., and Rojas, P. (2016). Changes in phenolic compounds during ripening of tropical fruits. *Journal of Agricultural and Food Chemistry*, 64(9), 1120-1128.
- Siddiqui, M. W., & Dhua, R. S. (2010). Eating artificially ripened fruits is harmful. *Current science*, 1664-1668.
- Singal, D., & Dhua, R. S. (2010). Ripening of fruits by artificial agents. *Postharvest Technology and Innovation Journal*, 6(2), 12-20.

- Sogo-Temi, S. T., and Egba, J. (2014). Calcium carbide usage for artificial fruit ripening in Nigeria: Implications for public health. *African Journal of Food Science*, 8(7), 307-313.
- Stray, F. (1998). *The Natural Guide to Medicinal Herbs and Plants*. Tiger Books International.
- Ugwu, C. E., and Nnabuchi, I. (2003). Comparative studies of tannin content in selected fruit species. *Journal of Food Science and Technology*, 42(4), 390-394.
- Unuigbo, O. A., and Ozekhome, T. C. (2009). Impact of calcium carbide on the nutritional quality of ripened bananas. *African Journal of Biotechnology*, 8(10), 2105-2110.
- WHO (World Health Organization). (1989). *Environmental Health Criteria 70: Principles for the Assessment of Risks to Human Health from Exposure to Chemicals*. WHO Press.
- Olufemi, O. (2024). Exploring banana production in Africa for food security and economic growth: A short review. *Food Nutrition Chemistry*, 2(1), 125. <https://doi.org/10.18686/fnc.v2i1.125>
- Ukwo, S. P., Inyang, M. E., Udo, M. E., & Udofia, E. I. (2023). Quality parameters of banana fruit as influenced by traditional ripening techniques. *Research Journal of Food Science and Quality Control*, 9(2)
- Ibeachu, P. C., & Nwidobie, I. L. (2022). Effect of Calcium Carbide Induced Ripened Fruit on the Hippocampus of Adult Wistar Rats. *Asian Journal of Medical Principles and Clinical Practice*, 5(2), 92-104. <https://doi.org/10.9734/AJMPCP/2022/v5i2308>
- Oboh, G., Akindahunsi, A. A., & Esan, A. S. (2016). Effects of artificial ripening on the nutritional and phytochemical properties of banana fruit. *Journal of Food Science and Technology*, 53(3), 1403-1411.
- Maduwanthi, S. D. T., & Marapana, R. A. U. J. (2019). Induced Ripening Agents and Their Effect on Fruit Quality of Banana. *International Journal of Food Science*, 2019, 2520179. <https://doi.org/10.1155/2019/2520179>
- Marriott J, Palmer JK. Bananas—physiology and biochemistry of storage and ripening for optimum quality. *Critical Reviews in Food Science & Nutrition*. 1980 Sep 1;13(1):41-88.