

## Original Research Article

# Association of neutrophil-lymphocyte ratio, platelet-lymphocyte ratio and ABO blood group in early pregnancy loss

**Background:** Intrauterine pregnancy loss referred to natural pregnancy loss before the twentieth weeks of gestation. Women who show signs of stress are three times more likely to lose their pregnancy during the first three weeks. Blood group system has been linked with development of mental stress, anxiety or disease. Medical literature has also increasingly supported neutrophil-lymphocyte ratio (NLR) and platelet-lymphocyte ratio (PLR) as strong indicators of inflammation and oxidative stress.

**Place and Duration of Study:** Department of Haematology, Gynaecology and Obstetrics unit both of Enugu State University of Science and Technology Teaching Hospital, between August and December 2023.

**Methodology:** A total of 100 participants were enrolled comprising individuals who had experienced between 1-2 miscarriages. Fifty healthy normal pregnant women who had no history of miscarriage were used as control. Demographic information such as age and number of miscarriages was obtained through questionnaire. Neutrophil, lymphocyte, monocyte, eosinophil, packed cell volume (PCV) and platelet (PLT) were analyzed by automated haematology analyzer, while ABO and Rhesus blood group were done by tile method. The data were analyzed using SPSS version 25. Analysis of variance (ANOVA), Chi-square test and pie chart were used. P-Value < 0.05 was considered as statistically significant.

**Results:** Out of the 100 women, blood group A was 27, B was 24, AB was 1 and O was 48. Platelet count and NLR values were higher in miscarriage group compared to healthy normal pregnant women ( $p=0.039$  and  $<0.001$  respectively). Lymphocyte count were found to be lower in miscarriage group compared to healthy normal pregnant women ( $p=0.042$ ). There was no significant relationship between NLR, PLR and ABO blood group ( $p=0.199$ ).

**Conclusion:** Increased platelet count and NLR were observed in women who had early pregnancy loss. Measurement of low cost test like platelet count and NLR before twentieth weeks should be recommended as one of the ways to predict pregnancy leading to spontaneous abortion.

## ABSTRACT

Keywords: ABO, NLR, pregnancy, loss, ratio, miscarriage, platelet

### 1. Introduction

Miscarriage is the loss of a pregnancy before twentieth weeks. A spontaneous abortion or miscarriage is a process that can be grouped into 4 stages—threatened, inevitable, incomplete, and complete [1]. Miscarriage is an incident that usually leads to various maternal complications such as anemia, injuries to the uterus and endometrium, and infection. The combination of oxidative stress and defective placentation can cause increased serum ischemia-modified albumin levels, which in turn can contribute to the pathophysiology of early pregnancy loss [2, 3]. In general, miscarriage rate is reported as 15-20%, meaning that 15-20% of recognized pregnancies result in miscarriage. Maternal age increases the

occurrence of spontaneous miscarriage. Around 80% of miscarriages occur within the first trimester. And the rate of miscarriage reduces with increasing gestational age [4]. The immune response remodel leads to abnormal placental function, with ischemia of epithelial covering of the highly vascular embryonic placental villi and the presence of products are the crucial reasons for endothelial integrity damage. The incidence of endothelial dysfunction activates an inflammatory response within blood vessels that requires intravascular leukocytes, as well as the clotting and complement systems [5]. It is assumed that the increase in systemic inflammatory response markers like neutrophil-lymphocyte ratio (NLR), platelet-lymphocyte ratio (PLR), mean platelet volume (MPV), and plateletcrit are associated to the pathology of missed abortion and preeclampsia. It can be assumed that these markers would have notable changes in spontaneous abortion [6]. Several studies have been done on neutrophil-lymphocyte ratio and platelet-lymphocyte ratio focusing on pathologic conditions such as gestational diabetes [7], endometriosis [8] and gynecological cancers [9]. In addition, the ability of prognosticating early pregnancy loss and missed miscarriage of NLR, PLR, and MPV values has been studied [10]. Complete blood count (CBC), ABO and Rhesus blood group are simple, inexpensive, practical, and most sensitive test, are among the normal screening tests during pregnancy. Several studies had reported relationship between ABO blood group and spontaneous abortion [11, 12, and 13]. In 1972, study on ABO incompatibility as a cause of spontaneous abortion was done by Takno and Miller [14]. Neutrophil-lymphocyte ratio (NLR) and platelet-lymphocyte ratio (PLR) can be extracted from the complete blood count test results [15]. Bearing in mind the foremost importance of finding out the diagnostic and prognostic factors connected with miscarriage, the results of this study will be of great help in reducing the incidence of miscarriage. This study aims to determine the NLR and PLR in blood of women presenting miscarriage at first trimester.

## **2. Materials and Methods**

### **2.1.1 Study Area**

A case-control cross sectional study was conducted at the Enugu State University Teaching Hospital (ESUTH), gynaecology and obstetrics unit from August to December, 2023.

### **2.1.2 Methodology**

One hundred women presenting miscarriage in their first trimester at gynaecology and obstetrics units in the age range of 16-45 years were recruited for the study. The approval for this study was given by the Research Ethics Committee of Enugu State University of Science and Technology Teaching Hospital (ESUTH) Enugu. Written informed consent was obtained from each woman. Those who were unable to sign informed consent form were excluded. Fifty ages matched pregnant women with no history of miscarriage were recruited as controls.

### **2.1.3 Sample collection**

Three milliliters of blood samples were collected via venepuncture. Complete blood count (CBC) was analyzed by automated haematology analyzer (Mindray/BC-5150). ABO and Rhesus blood group were done by tile method.

**2.1.4 Statistical analysis:** This was done using the SPSS version 25; means of variables are reported as mean  $\pm$  standard deviation. Test of significant difference between means of variables was determined using the student "T" test, one way ANOVA. Chi-square test of independence was used to determine association between ABO blood group and inflammatory response markers. A value of  $\leq 0.05$  was considered significant.

## **3. Results:**

Table 1 showed mean  $\pm$  SD of some haematological parameters. There were statistically significant increase in the values of platelet ( $249.93 \pm 142.23 \times 10^9/l$ ) and NLR ( $2.11 \pm 1.54$ ) in the subjects that had experienced miscarriage, compared to control subjects (PLT  $212.27 \pm 100.02 \times 10^9/l$ ,  $p = 0.039$ ; MCHC  $1.61 \pm 0.03$ ,  $p = 0.012$ ). The mean values of lymphocytes and PCV were significantly decreased

in the subjects that had experienced miscarriage, (34.95 $\pm$ 16.74 %; 28.30 $\pm$ 5.18 %), compared to control subjects (39.18  $\pm$ 13.68 %,  $p = 0.042$ ; 33.80 $\pm$ 2.24 %,  $p = 0.001$ )

**Table 1: Mean  $\pm$  SD of some haematological parameters**

Parameters	test (N=100)	control (N=50)	p-value
Neutrophils (%)	54.81 $\pm$ 18.11	51.82 $\pm$ 14.49	0.227
Lymphocytes (%)	34.95 $\pm$ 16.74	39.18 $\pm$ 13.68	0.042*
Monocytes (%)	6.86 $\pm$ 2.25	5.82 $\pm$ 1.60	0.368
Eosinophils (%)	2.58 $\pm$ 1.89	3.86 $\pm$ 2.44	0.182
PCV (%)	28.30 $\pm$ 5.18	33.80 $\pm$ 2.24	<0.001*
Platelet count (10 <sup>9</sup> /L)	249.93 $\pm$ 142.23	212.27 $\pm$ 100.02	0.039*
NLR	2.11 $\pm$ 1.54	1.61 $\pm$ 0.03	0.012*
PLR	107.52 $\pm$ 85.39	92.79 $\pm$ 41.39	0.125

Table.2 compared mean  $\pm$  SD of NLR of ABO blood group in women who had experienced loss of 1-2 pregnancies with pregnant women who had not experience pregnancy loss. There was statistically significant increase in the values of NLR of blood group B rhesus D positive subjects (1.96  $\pm$ 1.14), compared to control subjects (0.89  $\pm$  0.35,  $p=0.001$ ).

Table.2: mean  $\pm$  SD of NLR in ABO blood group

	A +ve	B +ve	O +ve
Test (N=94)	1.79 $\pm$ 1.22	1.96 $\pm$ 1.14	2.91 $\pm$ 1.61
control (47)	1.53 $\pm$ 0.82	0.89 $\pm$ 0.35	1.83 $\pm$ 1.10
p value	0.233	<0.001	0.183

Abbreviation: +ve = positive

Table 3: compared mean $\pm$ -SD of PLR of ABO blood group in women who had experienced loss of 1-2 pregnancies with women who had not experience pregnancy loss. The result showed no significant difference between test and control.

Table 3: mean $\pm$ -SD of PLR in ABO blood group

	A +ve	B +ve	O +ve
Test (N=94)	87.16 $\pm$ 45.96	82.82 $\pm$ 47.10	92.82 $\pm$ 47.13
control (47)	86.72 $\pm$ 50.98	92.89 $\pm$ 18..66	99.07 $\pm$ 34.92
p value	0.965	0.171	0.462

Table 4 compared mean $\pm$ -SD of NLR and PLR among A B O Rhesus D positive women that had experienced loss of 1-2 pregnancies. However, the result showed no statistically significant changes.

.Table 4: Mean+/- SD of NLR and PLR among A B O Rhesus D positive blood groups

	NLR	PLR
A +ve	1.795 ± 1.224	87.159 ± 45.939
B +ve	1.939 ± 1.139	82.818 ± 47.103
O +ve	2.190 ± 1.610	92.818 ± 47.127
F (p) value	0.958 (0.386)	0.507 (0.604)
A vs B	0.839	0.900
A vs C	0.410	0.836
B vs C	0.683	0.582

Table 5 showed relationship between NLR, PLR and ABO blood group of women who had experienced loss of 1-2 pregnancies using Chi-square test. The result showed no significant relationship between NLR, PLR and ABO blood groups ( $p = 0.199$ )

Table.5: Relationship between NLR, PLR and ABO blood group using Chi-square test

	Value	df	Asymp. Sig (2-sided)
Pearson Chi-square	6.000 <sup>a</sup>	4	0.199
Likelihood Ratio	6.592	4	0.159
Linear-by-Linear association	0.859	1	0.354
N of valid cases	3		




Fig 1: Blood group distribution of women with miscarriage




Fig 2: Blood group distribution of normal pregnant women

### Discussion

Neutrophil-to-lymphocyte ratio is simply calculated from the differential cell count. It is immediately available upon admission in all the patients. It is also a reflection of physiologic stress [16]. Study has shown that in both early pregnancy failures and preeclampsia, that inflammatory markers and oxidative biomarkers are significantly high in maternal blood. The high level of markers is investigated in preeclampsia, but there are still limited studies investigating them in the early pregnancy loss [17]. In this study, NLR significantly increased in women with miscarriage compared to normal pregnancies. The finding is consistent with the results of another study, which reported increased NLR in miscarriage compared to normal pregnancy [18]. Increased platelet counts were observed in women with miscarriage compared to women with normal pregnancy. The finding is consistent with recent report on risk of early foetal demise [19]. Blood group antigens are markers on surface-uncovered red cell proteins. Blood groups study also showed that an incompatibility of the blood groups can have an effect on reproduction. Miscarriage could occur in couples with incompatible blood groups [20]. In this study blood group 'O' was the dominant group occurring in 48% of the women with miscarriage, followed by groups A (27%) and group B (24%). Only 1% of the women with miscarriage were of blood group 'AB'. Although the distribution of ABO blood groups differ from one population to the other, in most studies blood group 'O' has been reported to be the upper most group and blood group 'AB' as the lowest which is in line with this study and previous studies in Enugu State and Nigeria in general [21, 22]. The ABO prevalence in Nigeria differs slightly in different regions, but blood group 'O' has been reported as the most dominant in all reported areas of studies. In this study, blood group 'O' was found to be the most dominant blood group among the normal pregnant women without history of miscarriage. This observation is consistent with previous studies in Nigeria [23, 24]. In this study NLR was found to be significantly increased in blood group B subjects with miscarriage compared to blood group B women with normal pregnancy. The finding is consistent with previous study on ABO blood groups [25]. Many studies have reported NLR as inflammatory indicator and were closely linked to the prognosis of various diseases [26]. As a marker of inflammation, NLR has been found to be high in various diseases and complications that affect pregnancy outcome [27]. Pregnancy complications, such as gestational diabetes and preeclampsia have been associated with elevated NLR, but little is known on the direct relationship with different blood groups.

### Conclusion

In conclusion, increased platelet count and NLR were observed in women who had early pregnancy loss. Measurement of low cost test including platelet count and NLR before twentieth weeks should be recommended as one of the ways to predict pregnancy leading to spontaneous abortion.

### Conflict of interests

Author declared that no competing interests exist

Guarantor of submission: The corresponding author is the guarantor of submission

## References

1. Yang X, Hu R, Shi M, Wang L, & Yan J, Gong J, et al. (2023). Placental malfunction, fetal survival and development caused by sow metabolic disorder: The impact of maternal oxidative stress. *Antioxidants*.12:360.
2. Cengiz H, Dagdeviren H, & Kanawati A, et al. (2015). Ischemia-modified albumin as an oxidative stress biomarker in early pregnancy loss. *J Matern Fetal Neonatal Med*. Sep 18. 1-4.
3. Calleja-Agius J, Jauniaux E, Pizzey AR, & Muttukrishna S. (2012). Investigation of systemic inflammatory response in first trimester pregnancy failure. *Hum Reprod*.27(2):349-57.
4. Arck PC, Rucke M, & Rose M, et al. (2008). Early risk factors for miscarriage: a prospective cohort study in pregnant women. *Reprod Biomed Online*. 17(1):101-13.
5. Mihiu D, Razvan C, Malutan A, & Mihaela C. (2015). Evaluation of maternal systemic inflammatory response in preeclampsia. *Taiwan J Obstet Gynecol*. 54:160–166.
6. Yücel B, & Ustun B. (2017). Neutrophil to lymphocyte ratio, platelet to lymphocyte ratio, mean platelet volume, red cell distribution width and plateletcrit in preeclampsia. *Pregnancy Hypertens*. 7:29–32.
7. Sargin MA, Yassa M, Taymur BD, Celik A, Ergun E, & Tug N. (2016). Neutrophil-to-lymphocyte and platelet-to-lymphocyte ratios: Are they useful for predicting gestational diabetes mellitus during pregnancy? *Ther Clin Risk Manag*. 12: 657-665.
8. Bas FY, Tola EN, Sak S, & Cankaya BA. (2018). The role of complete blood inflammation markers in the prediction of spontaneous abortion. *Pak J Med Sci*. 34:1381–1385.
9. Oğlak SC, & Aydın MF. (2020). Are neutrophil to lymphocyte ratio and platelet to lymphocyte ratio clinically useful for the prediction of early pregnancy loss? *Ginekol Pol*. 91: 524-527.
10. Biyik I, Albayrak M, & Keskin F. (2020). Platelet to lymphocyte ratio and neutrophil to lymphocyte ratio in missed abortion. *Rev Bras Ginecol Obstet*. 42:235–239
11. Clarke CA. (1972). Practical effects of blood group incompatibility between mother and fetus. *Br Med J*. 2:90-5.
12. Schaap T, Shemer R, & Palti Z et al. (1984). ABO incompatibility and reproductive failure. I. Prenatal selection. *Am J Hum Genet*. 36:143-51.
13. Malekasgar AM. (2004). ABO blood group prevalence in spontaneously repeated abortion. *Turk J Haematol*. 21:181-7.
14. Takano K, & Miller JR. (1972). ABO incompatibility as a cause of spontaneous abortion: evidence from abortuses. *J Med Genet*. 9:144-50.
15. Liu D, Huang X, Xu Z, Chen M, & Wu M. (2022). Predictive value of NLR and PLR in missed miscarriage. *J Clin Lab Anal*. 36:e24250.
16. Depu Zhou, Jie Wang and Xiaokun Li (2021). The platelet-lymphocyte ratio associated with depression in diabetes patients in the US National Health and Nutrition Examination Survey. *Int. J. Gen. Med*. 14:7825-7832.
17. Kim Y. (2020). Retrospective analysis of prognostic value of the neutrophil-to-lymphocyte ratio in early miscarriages: A 8-year survey. *Medicine*. 99:e20888.
18. Biyik I, Albayrak M, & Keskin F. (2020). Platelet to lymphocyte ratio and neutrophil to lymphocyte ratio in missed abortion. *Rev Bras Ginecol Obstet*. 42:235–239..
19. Xiaowen S, Dandan W, Yue X, Ling G, Hui Y, Jieru Z, Jiayi L, Jie Q, Jiajing C, Lihua S, & Yaozu X. (2021). The high platelet counts as predictor for early foetal demise. *Ann Med*. 53(1):1505–1511.
20. Omu AE, Al-Mutawa M, & Al-Qattan F. (1998). ABO blood group and expression of antisperm antibodies in infertile couples in Kuwait. *Gynecol Obstet Invest*. 45:49–53.
21. Ngwu A. M, Obi G. O, Anigolu M. O, & Eluke B. C. (2015). Distribution of ABO and Rhesus blood groups among voluntary blood donors in Enugu. *International Blood Research & Reviews*. 3(3):109-116.

22. Ifeanyichukwu M.O, Amilo G.O, NgwuA.M, Obi G.O,& Okoye A.E. (2014). Ferritin and serum iron levels among the ABO blood groups in Enugu, South Eastern Nigeria. *Journal of Blood Disorders & Transfusion*. 5: 204. Doi: 104172/2155-9864. 1000204.
23. Akpan, U.O., Basse, I.E., Nwatu, N.N., &Offor, J. (2018). A comparative Study on ABO blood group and fertility hormones in infertility women in Calabar, Southern Nigeria. *Annals of Medical Physiology*. 2:19-24.
24. Iyola OA, Igunnugbemi OO, Anifowoshe AT,& Raheem UA. (2011). Gene frequencies of ABO and Rh (D) blood group alleles in Ilorin, North-central Nigeria. *World J Biol Res*. 4(1):6-14
25. Poole EM, Gates MA, &High BA, et al. (2012). ABO blood group and risk of epithelial ovarian cancer within the Ovarian Cancer Association Consortium. *Cancer Causes Control*. 23:1805–1810
26. Zhang A, Ning L., & Han J, et al. (2019). Neutrophil-To-lymphocyte ratio as a potential biomarker of neovascular glaucoma. *Ocul Immunol Inflamm*. 2019;10:1–8
27. Viktoria C., Zafeiris Z., George D., Theodoros K., &Charalampos S. (2020). First trimester neutrophil to lymphocyte ratio (NLR) and pregnancy outcome. *J Obstet Gynaecol*. 40(1):59-64