

GROWTH PERFORMANCE OF *Oreochromis niloticus* FRY FED WITH TWO DIFFERENT STARTER DIETS

Abstract

This research was carried out at Joseph Sarwuan Tarka University, Benue State, Nigeria to evaluate the growth performance of Nile tilapia (*Oreochromis niloticus*) fry fed Coppens and Skretting starter diets separately. 200 fries were stocked at a density of 50 fry per 50 litre bowls. The experimental feeds used were Coppens and Skretting starter diets. The fry was fed 4 times daily to satiation for 8 weeks (56 days) in two replicates for each feed therefore, 4 plastic bowls were used, 50 fry for each bowl. The bowls were aerated throughout the study period. Sampling was done weekly to determine SGR and survival rate of the fry from each bowl. Statistical analysis was done and there was observable difference, although statistically there was no significant difference between the two treatments in weight $P=0.001$ ($p<0.05$) and length $P=0.01$ ($p<0.05$). The average temperature during the experimental period was $27.45^{\circ}\text{C} \pm 0.77$ while the pH was 7.35 ± 0.15 . Results of the studies shows that both feed types compete favorably in terms of growth rate and survival rate. But due to high cost of Coppens feed, Skretting may be recommended as a starter feed.

Keywords: Growth performance, Nile Tilapia, Fry, Coppens, Skretting

1.0 INTRODUCTION

Feeding plays a crucial role in the successful culture of *Oreochromis niloticus* (Nile Tilapia), especially during the fry stage when nutritional needs are particularly high. Proper nutrition at this early stage of development is essential for maximizing growth rate ensuring healthy development and improving survival rates. Fish nutrition plays a significant role in promoting the growth performance as well as sustaining the health of cultured fish. Artificial diets manufactured from various feed ingredients act as the primary source of nutrition in intensive aquaculture systems. Cultured fish are fed with different diets of complete feeds based on species, age, and production purpose in which protein is the limiting factor. Protein, carbohydrates, and lipids are distinct groups of nutrients that the body metabolizes to produce the energy needed for physiological processes and physical activities (Otoh *et al.*, 2022; Otoh *et al.*, 2023; Otoh *et al.*, 2024 a; b; c; d).

Food is an essential component of an organism, as its growth, development, reproduction, and other physiological activities are dependent on the energy generated by the consumed food material. Feeding is an important physiological activity of fishes like other living organisms. Fish require nutrients for growth, reproduction, and other normal physiological functions. In a natural aquatic environment, phytoplankton, zooplankton, plant materials, insects, insects' larvae, worms, and smaller fish are the major food types of fish (George, *et al.*, 2011). Fish tend to show a preference for particular food items within their environment. The availability of food in any aquatic environment determines the well-being and reproductive potential of fish (George, *et al.*, 2013; Keyombe *et al.*, 2015). The weight and size of fish are a reflection of food availability in the aquatic ecosystem (Bolarinwa and Popoola, 2014).

Nile tilapia (*Oreochromis niloticus*) is the most preferred cultured fish species in many tropical and subtropical countries of the world and constitute the third largest farmed finfish with annual production growth rate of about 11.5%. Nile tilapia (*O. niloticus*) is a well-known tropical food fish native to Africa. *O. niloticus* are principally herbivorous, although occasionally omnivorous. The fish is an efficient converter of waste foodstuff and appears to thrive well on artificial supplemental feed (Omoriege, *et al.*, 2009).

Tilapia farming is more acceptable, technically and economically more viable and sustainable. Tilapia culture is a profitable enterprise and even subsistence farmers in Nigeria can afford to culture tilapia to augment their income. Tilapia is consumed by poor people as it is relatively low-priced commodity. They are mainly of commercial importance in aquaculture because they are highly resistant to diseases, exhibit rapid growth, efficient feed conversion, breed in captivity, and have good consumer acceptance (Nyonje, *et al.*, 2011). A variety of factors influencing its growth rate include sex, stocking density, decrease in water temperature, and supplemental feeding (Liti, *et al.*, 2005; Mizanur, *et al.*, 2014).

Many environmental factors, such as water temperature, food availability, stocking density and environmental conditions, influence the food selection behavior of fish. The size of food items and the size and age of fish can also determine their food selection behavior (Otieno *et al.*, 2014). Mainly, fish feed on items that can fit

into their mouth and what their stomach can digest. As fish grow, the stomach becomes longer and their digestive system becomes more developed. However, the feeding rate relative to body weight decreases, whereas the absolute rate of food consumed increases (Wakil, *et al.*, 2014). Therefore, administering quality feeds to fish is of great importance as it plays an important role in its growth as well as behavior of the fish.

It is important to note that the early-stage development of organism (fish) determine the successful growth and development of that organism. The fry stage of Nile Tilapia (*O. niloticus*) is a critical and fundamental stage. They are most fragile and reactive to any sudden change, positive or negative. Good management and feeding during this period are important for sustainable growth.

The selection of an appropriate starter diet is thus a critical decision for aquaculture operations seeking to enhance production efficiency and sustainability. This study focuses on comparing the growth performance of *O. niloticus* fry fed with Coppens and Skretting starter diets, evaluating factors such as growth rates, survival rates and proximate composition. By identifying the more effective feed for this species, the study aims to provide insights that can help improve Tilapia farming practices.

2.0 Materials and Methods

2.1 Study Area

The study was carried out in the departmental hatchery unit at Joseph SarwuanTarka University Makurdi, the state capital of Benue State. The state lies in the middle of the country and share boundaries with five others namely, Nasarawa to the North, Taraba to the East, Cross-River to the South, Enugu to the South- West and Kogi to the West. The State also shares a common boundary with the Republic of Cameroon on the South-East.

2.2 Experimental Fish

A total of 200 Nile tilapia (*Oreochromis niloticus*) fry were used as experimental fish.

2.3 Experimental Diets

The starter diets used to evaluate the growth performance of *Oreochromis niloticus* are Coppens and Skretting starters. The diets were obtained from a fish feed seller in Makurdi.

2.4 Experimental Setup

The experiment was conducted for a period of 8 weeks. 4 (four) plastic bowls (50-litre), each half-filled with water was aerated continuously using air compressor. A total number of 200 fry were obtained from fish seed production unit at the Joseph SarwuanTarka Research Fish Farm and acclimatized to laboratory conditions for 7 days and distributed randomly into the four-plastic bowls (50 fish per tank) representing two starter dietary treatments. The fry was divided into 4 groups (50 each) and placed into 4 plastic bowls. Two replicates of plastic bowls were used for each experimental diet. The fry was fed 4 times daily to satiation; in the morning at 9:00 am, afternoon at 12:00 pm, in the late afternoon at 3:00 pm and evening at 6:00 pm. The water quality parameters such as dissolved oxygen, pH, temperature, Total dissolved solids, electrical conductivity, were monitored using multichannel water quality parameter kit.

2.4.1 Water quality parameters

Water quality parameters that were measured include temperature, electrical conductivity, dissolve oxygen, total dissolved solids and pH. These parameters were measured based on the standard methods describe by (APHA, 2005)

2.4.2 Growth indicators and Feed utilization

Growth indicators and feed conversion, in terms of survival rate (SR), weight gain (WG), percentage of weight gain (%WG), daily growth rate (DGR), specific growth rate (SGR), condition factor (K), feed conversion ratio (FCR), and protein efficiency ratio (PER) were determined using the following equations:

$$SR = (\text{final fish number} / \text{initial fish number}) \times 100,$$

$$WG = \text{final weight (g)} - \text{initial weight (g)}$$

$$WG (\%) = 100 \times [\text{final weight (g)} - \text{initial weight (g)}] / \text{initial weight (g)}$$

DGR = weight gain (g) / experimental period

SGR = $100 \times [\text{Ln}(\text{final weight}) - \text{Ln}(\text{initial weight})] / \text{experimental period}$

K = $[\text{body weight (g)} / \text{body length}^3(\text{cm}^3)] \times 100$

FCR = feed intake (g) / body weight gain (g)

PER = weight gain (g) / protein intake (g)

2.5 Data Analysis

Analysis of variance (ANOVA) was used at 95% significance level to test for significant differences between the various treatment means obtained for the growth and feed utilization (WG, WG%, DGR, SGR, FCR and PER) and body Fisher's Least Significant Differences test (LSD-test) will be applied to compare the average values (means \pm standard deviation) and the significant differences was defined at $p < 0.05$.

3.0 Results

3.1 Growth performance of *Oreochromis niloticus* fry Fed Different Starter Diets

Table 1 below shows the mean growth performance of *Oreochromis niloticus* fry fed two different treatment diets. There is an observable significant difference between treatment 1 and treatment 2 ($P < 0.05$). Coppens diet was higher than Skretting in all the parameters studied, mean weight gain, mean length gain, growth rate, specific growth rate and survival rate, 1.02 ± 0.03 , 3.47 ± 0.07 , 0.018 ± 0.00 , 0.039 ± 0.00 and 75.00 ± 1.00 respectively.

Table 1: Meangrowthperformanceof*Oreochromis niloticus* fryfeddifferentstarterdiets

Growth Parameters	Treatments		P-Value
	Coppens	Skretting	
Mean Weight Gained	1.02 ± 0.03^a	0.63 ± 0.04^b	0.01
Mean Length Gained	3.47 ± 0.07^a	2.75 ± 0.05^b	0.01
Growth Rate	0.018 ± 0.00^a	0.01 ± 0.00^a	0.13
Specific growth Rate	0.039 ± 0.00^a	0.031 ± 0.00^b	0.00
Survival Rate	75.00 ± 1.00^a	41.00 ± 1.00^b	0.00

*Means in the same row with different superscripts differ significantly

3.2 Water Quality Parameters

Table 2 below shows the mean water quality parameters of each of the treatment. Here, the water quality parameters in the two treatments do not differ significantly except for TDS that is slightly higher in Coppens diet.

Table 2: Mean Water Quality Parameters of Experimental Media

*Means in the same row with different superscripts differ significantly

Water Quality Parameters	Treatments		P-Value
	Coppens	Skretting	
Dissolved Oxygen	4.10 ± 0.20^a	3.80 ± 0.20^a	0.40
Temperature	27.50 ± 0.00^a	27.45 ± 0.15^a	0.77
pH	7.35 ± 0.15^a	7.35 ± 0.05^a	1.00
Total Dissolved solids	210.50 ± 0.50^a	230.10 ± 0.00^a	0.02
Electrical Conductivity	610.50 ± 2.50^a	633.00 ± 0.00^a	0.42

*Means in the same row with different superscripts differ significantly

3.3 Proximate Composition of *Oreochromis niloticus* Fry fed different Starter Diets

Table 3 below shows the proximate composition of *Oreochromis niloticus* fry fed two different treatment diets. There is no significant difference between the proximate composition of the two treatments fish, but Coppens fed fry had higher CP (56.96 ± 0.08) and fat (20.68 ± 0.47) than that of Skretting which recorded 56.72 ± 0.16 and 20.00 ± 0.35 for CP and fat but lower in Ash. No significant difference was observed in all the parameters.

Table3: Mean proximate composition of *Oreochromis niloticus* fry fed different starter diets

Proximate Components	Treatments		P-Value
	Coppens	Skretting	
Crude Protein	56.96±0.08 ^a	56.72±0.16 ^a	0.32
Fat	20.68±0.47 ^a	20.00±0.35 ^a	0.37
Ash	12.72±0.25 ^a	13.01±0.12 ^a	0.39
Moisture	4.74±0.60 ^a	5.55±1.09 ^a	0.58
Crude Fibre	0.00±0.00	0.00±0.00	—

*Means in the same row with different superscripts differ significantly




Figure 1: Mean Weekly Weight of *Oreochromis niloticus* fry fed different starter diets




Figure 2: Mean Weekly Length of *Oreochromis niloticus* fry fed different starter diets.

5.0 Discussion, Conclusion and Recommendations

4.1 Discussion

The present study was aimed at comparing the growth performance of *O. niloticus* fry fed with two different Starter Diets (Coppens and Skretting). The water quality parameters showed little variation in electrical conductivity and TDS. However, the range of temperature (27.4°C-27.5°C), pH (7.3-7.3) and dissolved oxygen (3.8-4.1mg/l) obtained were favorable for fish culture and were within the range described as optimal by Jamabo *et al.*, (2015) and Jonah and George (2019). Balogun *et al.*, 2004 reported a higher specific growth rate (SGR) value of 3.19% d⁻¹ using Coppens, Vital and Ajanla feed, which may be due to feeding frequency and the type of feed. With respect to specific growth rates, *O. niloticus* fed Coppens diets showed a higher performance than those fed with Skretting diet. The high growth rate recorded in Coppens fed fry may be attributed to the high nutritive value of the feeds and tolerable physico-chemical parameters.

The problem associated with the low survival rate in Skretting fed fry arose due to the change in the physico-chemical parameters of the water. The ability of an organism to utilize nutrients especially protein will positively influence its growth rate Sogbesan and Ugwumba, (2008). The growth of fish depends upon the ingredients and its percentage in the formulated feed Glencross *et al.*, (2007).

The results of this study shows that mean weight of Coppens fed fry increases up to week 7 and then reduces on week 8 while Skretting fed fry increases up to week 8. However, Coppens fed fry had higher mean weight gain, mean length gain, growth rate, specific growth rate and survival rate than Skretting fed fry. Coppens fed fry also had better results in terms of proximate composition. The better growth rate exhibited by Coppens fed fry can be attributed to good feed quality in terms of its proximate composition (CP, fat and ash), acceptance of feed by fish and tolerance to physio-chemical water parameters, feed utilization, which resulted in better growth performance and high survival rate. Conclusively, among the experimental feeds, Coppens is the best feed because the fish fed in this group showed better results than those fed with Skretting diet.

4.2 Conclusion

The growth performance of *O. niloticus* fry fed with Coppens and Skretting starter diets revealed that fry fed with Coppens showed superior results in key growth parameters such as mean weight gain, length gain, specific growth rate and survival rate. In addition, the proximate composition of fry fed with Coppens was also more favourable. Therefore, it is recommended that Coppens starter diets be used for *O. niloticus* fry in aquaculture, as they promote better overall growth and health compared to Skretting.

5.0. REFERENCES

- APHA (2005). *Standard Methods for the examination of Water and Waste Water*. 21st ed., Washington DC, USA: American Public Health Association, 1498 Pp.
- Balogun, J. K., Abdullahi, S. A., Auta, J. and Ogunlade, O. P. (2004). Feed conversion, protein efficiency, digestibility and growth performance of *Oreochromis niloticus* fed Delonix regia seed meal. Proceedings of National Conference of Fisheries Society of Nigeria, pp823-829.
- Bolarinwa, J.B. and Popoola. (2014). Length-Weight Relationships of Some Economic Fishes of Ibeshe Waterside, Lagos Lagoon, Nigeria *Journal of Aquaculture Research & Development* 05(1):10.4172/2155-9546.1000203
- George, U. U., Ekanem, A. P., and George, E. (2011). Food and Feeding Habits of *Ophiocephalus obscura* (African Snakehead in the Cross-River Estuary, Cross River State, Nigeria. *International Journal of Fisheries & Aquaculture*, 3(13):231-238.
- George, U. U., Idung, J. U., Andem, A. B., Okorafor, K. A. and Mowang, D. (2013). Diet Composition and condition Factor of *Ethmalosa fimbriata* in the Cross-River Estuary. *Greener Journal of Biological Science*, 3(6):244-252.
- Glencross, B. D., Booth, M. and Allan, G. L. (2007). A feed is only as good as its ingredients: A review of ingredient evaluation strategies for aquaculture feeds. *Aquac. Nutr.*; 13: 17-34.
- Jamabo, N. A., Fubara, R. I. and Dienye, H. E. (2015). Feeding frequency on growth and feed conversion of *Clarias gariepinus* (Burchell, 1822) fingerlings. *International journal of fisheries and aquatic studies*;3(1):353-356.
- Jonah, U. E. and George, U. U. (2019). Influence of Water Quality on Zooplankton Community Structure of Etim Ekpo River, Akwa Ibom State, South-South, Nigeria. *World Rural Observation*; 11(3):49-57.
- Keyombe, J.L., Edna Waithaka and Beatrice Obegi. (2015). Length-weight relationship and condition factor of *Clarias gariepinus* in Lake Naivasha, Kenya. *International Journal of Fisheries and Aquatic Studies*; 2 (6), 382-385.
- Liti D, Cherop L, Munguti J, Lim, C. (2005) Growth and economic performance of Nile tilapia (*Oreochromis niloticus* L.) fed on two formulated diets and two locally available feeds in fertilized ponds. *Aquaculture Research*; 36: 746-752.
- Mizanur, R.M., Yun H, Moniruzzaman, M., Ferreira F. and Kang-woong, K. (2014). Effects of feeding rate and water temperature on growth and body composition of juvenile Korean Rockfish, *Sobastesschegeli* (Hilgendorf 1880). *Journal Animal Science*; 27: 690-699
- Omoriegie, E. Igoche, L. Ojobe, T.O., Absalom, K.V., and Onusiriuka, K.V. (2009). Effect of varying levels of sweet potato (*Ipomea batatas*) peels on growth, feed utilization and some biochemical responses of the cichlid (*Oreochromis niloticus*). *African Journal Food Agri Nutr and Dev*;9: 700-712.

- Otieno, N.O., Kitaka, N. and Njiru, J.M. (2014). Length-weight relationship, condition factor, length at first maturity and sex ratio of Nile tilapia, *Oreochromis niloticus* in Lake Nalvasha, Kenya. *International Journal of Fisheries and Aquatic studies*; 2(2):67-72
- Otoh A. J., M. T. Udo and U. U. George. (2022) Comparative Effect of Inducing Broodstock with Natural and Artificial Hormones on Reproductive Performances of *Heterobranchus longifilis*. *Tropical Freshwater Biology*, 31: 95-102.
- Otoh A. J., Udo, M. T., and George, U. U. (2023). Comparative Growth Performance and Sex Ratio of *Heterobranchus longifilis* and its Offspring Induced with Synthetic Hormone and Pituitary Gland of *Heterobranchus longifilis*. *Journal of Wetlands and Waste Management*; 5(1): 106- 111.
- Otoh, A. J., Ekanem, I. E., Okoko, A. C., Asangusung, P. S. and George, U. U (2024 d). Comparative Study of Inducing Broodstock with Natural and Artificial Hormones on Reproductive Performances of *Clarias gariepinus*. *Asian Journal of Fisheries and Aquatic Research*; 26(3):31-38.
- Otoh, A. J., Ekanem, I. E., Okoko, A. C., Asangusung, P. S., George, U. U., Idiong, T. E. (2024 b) Effect of different ages of African catfish (*Clarias gariepinus*) broodstock on reproductive performance and fry production. *Asian Journal of Fisheries and Aquatic Research*; 26(3): 39-47.
- Otoh, A. J., George, U. U., Asangusung, P. S., Okoko, A. C., Ekanem, I. E. and Umaha, M. N. (2024 a). Investigation into the impacts of varied batches of stripped eggs on the reproductive success of *Clarias gariepinus*. *Asian Journal of Research in Zoology*; 7(2):28-35.
- Otoh, A. J., Okoko, A. C., Ekanem, I. E., George, U. U. and Asangusung, P. S. (2024 c). Effects of Multiple Batches of Stripped Eggs on the Reproductive Performances of *Heterobranchus longifilis*. *Asian Journal of Research in Zoology*; 7(2): 9-16.
- Sogbesan, A.O. and Ugwumba, A.A.A. (2008). Nutritional Values of Some Non-Conventional Animal Protein Feedstuffs Used as Fishmeal Supplement in Aquaculture Practices in Nigeria. *Turkish Journal of Fisheries and Aquatic Sciences* 8: 159-164.