

## Shelf stability of tortilla chips enriched with microwave stabilized wheat bran

### ABSTRACT

This study investigates the effects of incorporating microwave-stabilized wheat bran into tortilla chips, focusing on nutritional, sensory, and storage-related characteristics. Tortilla chips were developed with varying wheat bran levels (0–30%) and evaluated for sensory attributes, moisture content, water activity, lipid stability, and microbial load over a 90-day storage period. Results showed that tortilla chips with 20% wheat bran achieved the highest sensory scores for taste, texture, and appearance, with overall acceptability reaching 7.8 on a nine-point hedonic scale. However, higher bran levels (25–30%) led to declines in sensory qualities, likely due to increased moisture content. Moisture and water activity ranged from 2.78% to 4.10% and 0.44 to 1.73, respectively, and both values increased with bran levels and storage time. The addition of bran also affected lipid stability; free fatty acid content rose from 0.44% in the control (T1) to 1.73% in the 30% bran treatment (T7), while peroxide values increased from 3.14 to 7.99 meq O<sub>2</sub>/kg over 90 days. Microbial counts followed similar trends, with the highest mean load of  $2.18 \times 10^4$  cfu/g in the 30% bran formulation, yet all counts remained within acceptable limits. This study highlights the potential for wheat bran-enriched tortilla chips as a fiber-rich snack alternative, contributing valuable data for the formulation of healthier, shelf-stable snacks.

**Keywords:** Wheat bran, Tortilla chips, Nutritional, Shelf life, Storage

### 1. INTRODUCTION

In recent years, snacking has become an integral part of dietary habits globally, shaping food choices and consumption patterns across diverse populations. With changes in lifestyle, such as increased urbanization, busier schedules, and the rise in single-person households, the demand for convenient, ready-to-eat foods has surged, making snacks a vital component of modern diets [1]. Snacks can vary widely in their nutritional composition, from healthy options like fruits, nuts, and yogurt to highly processed, energy-dense products often high in sugar, fat, and salt [2]. The growing market for snacks has encouraged food researchers to explore new product development, focusing on creating healthier snack options that meet consumer preferences for taste, convenience, and nutrition [3].

Health-conscious consumers have driven an increase in snack products that emphasize functionality, such as high-protein, high-fiber, and plant-based ingredients, catering to a broader range of dietary needs and trends [4]. Additionally, food technology advancements have

enabled the development of innovative processing methods, improved preservation techniques, and novel ingredients, all contributing to the enhancement of snack quality and nutritional value [5]. Understanding the balance between consumer demands for taste and convenience and the need for nutritious and sustainable snack options has become crucial for food scientists and product developers.

Among popular snack options, tortilla chips hold a prominent place due to their versatility, taste, and appeal across various demographics. Traditionally made from corn, tortilla chips have seen numerous innovations in recent years as food scientists work to enhance their nutritional profile while preserving desirable sensory qualities. One promising approach involves the incorporation of stabilized wheat bran and nixtamalized corn flour, which introduces fiber and beneficial nutrients into a product typically characterized by refined carbohydrates and lower fiber content. Stabilized wheat bran, known for its high fiber and mineral content, not only enhances the nutritional value of product but also improves texture and adds a mild, nutty flavor [6]. Furthermore, the use of nixtamalized corn flour contributes to superior flavor and digestibility, as well as a rich source of calcium and B vitamins, resulting from the alkaline cooking process [7].

The integration of stabilized wheat bran and nixtamalized corn flour in tortilla chips aligns with consumer demand for functional foods that provide health benefits without compromising convenience [8]. These modified tortilla chips cater to a range of dietary needs, including increased fiber intake and the need for complex carbohydrates. In addition, they provide an alternative to traditional snacks, often criticized for their high fat, salt, and sugar content [9]. Research on optimizing ingredient ratios, processing techniques, and sensory evaluation is essential in the development of these enhanced tortilla chips, ensuring that they retain the taste and texture consumers expect. This approach underscores the role of food technology in creating innovative snacks that bridge the gap between nutrition and indulgence, making tortilla chips a viable option for health-conscious snackers.

**Table 1: Treatment details for preparation of wheat bran enriched tortilla chips**

<b>Treatment</b>	<b>Corn flour (%) (Masa)</b>	<b>Stabilized wheat bran (%)</b>
T <sub>1</sub>	100	0
T <sub>2</sub>	95	5
T <sub>3</sub>	90	10
T <sub>4</sub>	85	15
T <sub>5</sub>	80	20
T <sub>6</sub>	75	25
T <sub>7</sub>	70	30

## 2. MATERIALS AND METHODS

**2.1. Procurement of raw materials:** The study was conducted in Division of Food Science and Technology, Sher-e-Kashmir University of Agricultural Sciences and Technology of Jammu. Wheat bran used for the study was purchased from Amar flour mills, Gangyal, Jammu. The sample was exposed to microwave stabilization, followed by packaging in low-density polyethylene (LDPE) pouches (150 gauge) and stored at ambient conditions for a period of 90 days. White corn (Ganga safed 2 cultivar) was purchased from the local market of Jammu and was cleaned to remove impurities and dirt and stored at room temperature till further use. Refined oil and salt used for the preparation of tortilla chips were purchased from the local market.

To microwave the wheat bran, a microwave-resistant glass plate was preheated at 100% power for 3 minutes. Next, 100 g of wheat bran was spread evenly on the plate to a thickness of 0.5 cm. The plate was then placed in a microwave oven (Samsung Electronics Corp, CE137NEL) and heated at 900 Watts for three minutes at a frequency of 2450 MHz.[10]. The temperature of the sample after heating was  $105 \pm 2^{\circ}\text{C}$ . The treated wheat bran was allowed to cool to room temperature and packed in airtight LDPE bags (150 gauge).

## **2.2. Quality assessment of tortilla chips**

**2.2.1. Water activity:** Water activity was measured using an Aqualab water activity meter (Model series 3TE) and readings were corrected at  $20^{\circ}\text{C}$  [11].

**2.2.2. Moisture content:** Moisture content in the samples was determined by following the oven drying method as the loss in weight due to evaporation from sample at a temperature of  $105 \pm 1^{\circ}\text{C}$ . The weight loss in each case represented the amount of moisture present in the sample [11].

**2.2.3. Free fatty acid:** 1 gram of ground sample was mixed with 50 ml of benzene in a 100 ml volumetric flask and allowed to stand for 30 minutes for extraction of free fatty acids [3]. In a separate flask, benzene (5ml) and alcohol (10ml) was added to 1 ml of extract followed by addition of phenolphthalein as an indicator and solution was titrated using 0.02N KOH until disappearance of pink colour indicating that all the free fatty acids have been neutralized [12].

**2.2.4. Peroxide value:** To determine the peroxide value (PV), 30 ml of acetic acid: chloroform (2:1) was added to the weighed grounded sample. It was kept under a cool and dark place for 30 minutes and 30 ml of distilled water was added. The mixture was then shaken. This was

slowly titrated with 0.1 N sodium thiosulphate with vigorous shaking until the yellow colour disappeared. Then 0.5 ml of 1.0 per cent starch solution was added and titrated continuously with vigorous shaking to release all iodine from chloroform layer until pink colour just disappeared [11].

**2.2.5. Total Microbial count:** Spread plate technique, described by Palczar and Chan [13] was followed. 1 g of ground sample was aseptically transferred into test tube containing 9 ml of sterile water and was mixed vigorously. After mixing, 1 ml of this mixture was transferred to a test tube containing 9 ml sterile water for further dilution. The process was continued until 4<sup>th</sup> diluents ( $10^4$ ). Nutrient agar was inoculated with 0.1 ml of diluted sample ( $10^4$ ), by spread plating technique and incubated at 37°C for 48 hours. Colonies were counted and multiplied by dilution factor.

**2.2.6. Sensory evaluation:** In evaluating the tortilla chips made with stabilized wheat bran and nixtamalized corn flour, a nine-point hedonic scale can be applied to assess the impact of these ingredients on consumer acceptance. Sensory attributes, such as crunchiness, color, and taste are particularly important in snack foods like tortilla chips, where texture and flavor play a significant role in consumer preference [14].

**3. Statistical analysis:** The data, collected in triplicate, were subjected to statistical evaluation using the OPSTAT package program for analysis of variance.

## 4. RESULTS AND DISCUSSION

**4.1 Water activity:** Water activity, which refers to the amount of unbound or free water available in a food product, is a critical factor in determining food stability and shelf life. Water activity values range from zero (indicating no available water) to 1.0 (pure water) [15]. In this study, the water activity of tortilla chips was found to vary significantly with increasing levels of wheat bran from 0 to 30% (Table 2). As wheat bran levels increased, water activity values also increased from 0.237 in T1(100:00::CM:SWB) to 0.322 in T7(70:30::CM:SWB), consistent with findings from previous studies on bran-enriched cereal products [16]. This increase in water activity may be attributed to the higher moisture content absorbed due to the hygroscopic nature of the product [17]. Similar trends were observed in bran-enriched baked meat biscuits, where increasing bran content corresponded to an increase in water activity levels [18].

During a storage period of 90 days, the mean water activity values of the tortilla chips continued to increase from initial value of 0.239 to 0.311 at the end of storage period, likely due to additional moisture absorption during storage, which may be attributed to the hygroscopic properties of bran [19]. Our findings align with other studies showing similar increases in water activity over time in bran-enriched pasta products [20]. A gradual increase in water activity over extended storage periods has also been observed in tortilla chips and other grain-based snack products [21].

**4.2 Moisture content:** The moisture content of tortilla chips decreased significantly from 37.25% in corn masa to 2.78% after processing, a reduction attributed to the high temperatures used during tortilla chip production. Water molecules, being polar, are heated and expelled from the surface through oil during the frying process [22]. This reduction in moisture due to frying is consistent with findings in other fried products, such as surimi sausage incorporated with oat powder, where significant moisture loss occurs during high-temperature cooking [23]. In this study, the moisture content of tortilla chips ranged from 2.78% to 4.10% (Table 2), with a noticeable trend of increasing moisture content as wheat bran levels increased from 0% to 30%. This rise in moisture content can likely be attributed to the water-holding capacity of wheat bran fibers, which contain polysaccharides capable of retaining moisture [24]. Similarly, studies on bran-enriched foods, such as pasta and baked goods, have reported an increase in moisture retention with higher wheat bran incorporation, aligning with the results observed in tortilla chips [25]. This increase in moisture content has also been documented in products like flatbreads, biscuits, and pasta with added wheat bran or similar fibers [26].

Throughout the 90-day storage period, the mean moisture content of the tortilla chips rose from an initial value of 3.11% to 3.61%. This gradual increase in moisture could be due to the hygroscopic nature of bran, which absorbs moisture over time [27]. Similar findings were reported in buckwheat-based snacks, where moisture content increased progressively during storage [28]. Additionally, a significant increase in moisture content was observed in cereal bran-enriched snacks after six months of storage, likely due to bran's moisture-retentive properties [29].

UNDER PEER REVIEW

**Table 2: Effect of treatment and storage period on water activity (aw) and moisture content (%) of tortilla chips**

Treatments	Water activity					Moisture content				
	Storage days					Storage days				
	0	30	60	90	Mean	0	30	60	90	Mean
T1(100:00::CM:SWB)	0.190	0.230	0.256	0.274	<b>0.237±0.034</b>	2.46	2.71	2.89	3.08	<b>2.78±0.23</b>
T2 (95:05::CM:SWB)	0.210	0.245	0.257	0.279	<b>0.248±0.026</b>	2.63	2.86	3.03	3.18	<b>2.92±0.23</b>
T3 (90:10::CM:SWB)	0.215	0.252	0.269	0.286	<b>0.255±0.025</b>	2.87	3.09	3.24	3.35	<b>3.13±0.18</b>
T4 (85:15::CM:SWB)	0.238	0.271	0.297	0.310	<b>0.279±0.030</b>	3.10	3.28	3.40	3.56	<b>3.33±0.19</b>
T5 (80:20::CM:SWB)	0.257	0.282	0.307	0.329	<b>0.294±0.034</b>	3.37	3.61	3.73	3.82	<b>3.63±0.20</b>
T6 (75:25::CM:SWB)	0.276	0.298	0.321	0.343	<b>0.309±0.026</b>	3.53	3.70	3.89	3.94	<b>3.76±0.18</b>
T7 (70:30::CM:SWB)	0.287	0.311	0.334	0.359	<b>0.322±0.025</b>	3.80	4.02	4.22	4.35	<b>4.10±0.21</b>
CD(P=0.05) Treatment (T)	0.005					0.03				
Storage (S)	0.004					0.03				
T x S	NS					0.07				

**Table 3: Effect of treatment and storage period on free fatty acids (%oleic acid) and peroxide value(meqO<sub>2</sub>/ Kg)of tortilla chips**

Treatments	Free fatty acids					Peroxide value				
	Storage days					Storage days				
	0	30	60	90	mean	0	30	60	90	Mean
T1(100:00::CM:SWB)	0.33	0.42	0.49	0.54	<b>0.44±0.08</b>	2.82	4.17	5.48	7.27	<b>4.93±0.74</b>
T2 (95:05::CM:SWB)	0.56	0.67	0.75	0.80	<b>0.69±0.10</b>	2.90	4.40	5.87	7.45	<b>5.15±0.72</b>
T3 (90:10::CM:SWB)	0.75	0.84	0.91	0.96	<b>0.86±0.08</b>	3.03	4.70	6.23	7.67	<b>5.41±0.67</b>
T4 (85:15::CM:SWB)	0.97	1.10	1.19	1.26	<b>1.13±0.11</b>	3.12	4.97	6.52	7.95	<b>5.64±0.69</b>
T5 (80:20::CM:SWB)	1.12	1.21	1.28	1.32	<b>1.23±0.16</b>	3.24	5.20	6.79	8.19	<b>5.85±0.75</b>
T6 (75:25::CM:SWB)	1.30	1.41	1.50	1.58	<b>1.45±0.11</b>	3.36	5.74	7.27	8.47	<b>6.21±0.64</b>
T7 (70:30::CM:SWB)	1.59	1.70	1.79	1.86	<b>1.73±0.11</b>	3.48	6.07	7.54	8.97	<b>6.51±0.69</b>
CD(P=0.05) Treatment (T)	0.03					0.03				
Storage (S)	0.02					0.03				
T x S	0.06					0.07				

**4.3 Free fatty acids:** Free fatty acids (FFAs) are generated through the enzymatic or microbial breakdown of lipids, serving as indicators of fat stability during storage. In this study, the FFA content in treatment T1 (100:00::CM:SWB) was found to be lower than in corn and corn masa. This reduction may be due to the denaturation of lipase enzymes caused by high processing temperatures used in tortilla chip production (baking at 280°C and frying at 190°C). The incorporation of stabilized wheat bran (SWB) led to an increase in the mean FFA value, from 0.44% in T1 (100:00::CM:SWB) to 1.73% in T7 (70:30::CM:SWB) (Table 3). This increase could be attributed to the higher initial FFA content in bran, which, when added, can alter the lipid composition of the product [30]. Similar increases in FFA have been observed in other bran-enriched products, such as pasta, where the addition of cereal bran contributed to elevated FFA content [20]. Research on wheat bran-enriched baked goods has also reported increases in FFA, supporting these findings [31].

Over the 90-day storage period, the mean FFA content of tortilla chips increased significantly from 0.95% to 1.18% as oleic acid. This increase in FFA during storage is likely due to residual lipolytic activity of lipase, which may be enhanced by the rise in moisture content during storage [32]. Similar observations have been made in other studies where hydrolytic rancidity was noted in response to elevated storage temperatures [33]. Progressive increases in FFA content with extended storage have also been documented in bran- and tuber-enriched snack foods, such as breakfast snacks and root-based chips [34].

**4.4 Peroxide value:** Hydroperoxides are the primary products of lipid oxidation and serve as indicators of the early stages of oxidation. The peroxide value (PV) is a widely used to measure the extent of lipid oxidation in foods, including tortilla chips, throughout storage. To maintain product quality, the PV of fried tortilla chips should not exceed 10 meq O<sub>2</sub> per kg [35]. Secondary oxidation products, generated from the breakdown of hydroperoxides, contribute to rancidity by breaking the double bonds in unsaturated fatty acids [36].

In this study, the PV of T1 (100:00::CM:SWB) tortilla chips was significantly lower than that of raw corn and corn masa, as well as other treatments. This decrease in PV may be due to the effects of boiling and frying, which align with findings from studies on fried products, where pre-treatment cooking reduced peroxide levels [37]. In our samples, the mean PV of tortilla chips ranged from 4.93 to 6.51 meq O<sub>2</sub> per kg (Table 3). Replacing corn flour with microwave-stabilized wheat bran resulted in higher peroxide value in the tortilla chips, potentially due to the increased moisture content. Higher moisture levels may activate the lipoxygenase enzyme, facilitating the formation of hydroperoxides and raising peroxide

value [19]. Similar observations were reported in bran-enriched biscuits, where increased peroxide value was linked to bran's influence on moisture and enzyme activity [38].

Over a 90-day storage period, the mean peroxide value of tortilla chips increased significantly from 3.14 to 7.99 meq O<sub>2</sub> per kg. This progressive increase is likely due to residual lipase activity, which may be stimulated by the gradual gain in moisture content during storage [39]. Comparable trends have been reported in bran-enriched snack products, including tortilla chips and fried potato crisps, where storage-induced moisture uptake led to elevated peroxide value [40].

**4.5 Total microbial count:** The microbial count of a food product reflects the sanitary quality of the raw materials, processing conditions, and storage practices [41]. In this study, the microbial load of tortilla chips was significantly influenced by both treatment and storage duration (Table 4). The highest mean microbial count was observed in treatment T7 (70:30::CMSWB) with  $2.18 \times 10^6$  cfu/g, while the lowest count of  $1.12 \times 10^6$  cfu/g was recorded in T1 (100:00::CM:SWB). This increase in microbial load with higher levels of wheat bran supplementation may be due to increased moisture content, which supports microbial growth [42]. The addition of microwave-stabilized wheat bran appears to have contributed to the increased microbial count, aligning with findings in wheat bran-enriched flatbreads [43] and pasta [16].

Over the storage period, the total microbial plate count in tortilla chips increased significantly. This increase in microbial count is likely attributable to gradual moisture gain during storage, which provides a favorable environment for microbial growth [44]. However, all microbial counts remained within acceptable safety limits ( $\leq 10^6$  cfu/g), below the maximum level considered safe for consumption for stored foods [45]. Previous studies have also reported increases in microbial load over extended storage periods, particularly in bran-enriched snacks and high-fiber meat products [46&47].

**Table 4: Effect of treatment and storage period on total microbial count ( $\times 10^4$ cfu/g) of tortilla chips**


Treatments	Water activity				Mean
	Storage days				
	0	30	60	90	
T1(100:00::CM:SWB)	ND	ND	1.02	1.23	<b>1.12±0.11</b>
T2 (95:05::CM:SWB)	ND	ND	1.15	1.32	<b>1.23±0.09</b>
T3 (90:10::CM:SWB)	ND	ND	1.30	1.43	<b>1.36±0.07</b>
T4 (85:15::CM:SWB)	ND	ND	1.43	1.58	<b>1.50±0.08</b>
T5 (80:20::CM:SWB)	ND	ND	1.60	1.78	<b>1.69±0.09</b>
T6 (75:25::CM:SWB)	ND	ND	1.87	2.03	<b>1.95±0.06</b>
T7 (70:30::CM:SWB)	ND	ND	2.10	2.27	<b>2.18±0.09</b>
CD( $P=0.05$ )					
Treatment (T)	0.05				
Storage (S)	0.06				
T x S	0.09				

**4.6 Sensory evaluation:** A hedonic scale is commonly used in sensory evaluation to measure consumer preferences for food products. This scale, typically ranging from "dislike extremely" to "like extremely," provides a quantitative measure of sensory attributes such as taste, texture, aroma, appearance, and overall acceptability [48]. In this study, tortilla chips were evaluated for appearance, texture, taste, and overall acceptability using a nine-point hedonic scale. The overall acceptability scores were calculated based on the sensory quality scores for appearance, texture, and taste [49]. As the level of microwave-stabilized wheat bran increased from 5% to 20%, the scores for appearance, texture, taste, and overall acceptability followed an upward trend. However, at 25% and 30% wheat bran, these scores began to decline (Table5; Figure 1).

The overall acceptability of the wheat bran-enriched tortilla chips exceeded that of the control, which aligns with previous findings by [50] who reported that consumers preferred bran-enriched cookies over refined flour versions. Notably, the 20% wheat bran formulation achieved the highest taste scores, likely due to the distinctive, pleasant bran flavor [51]. The reduced appearance score at higher wheat bran levels may stem from increased oxidation, resulting in a darker hue [52], similar to findings in bran-enriched baked goods. Comparable effects of bran on sensory scores have been documented in biscuits [53], while [54] also observed diminishing sensory appeal in biscuits with 20% bran. [55] highlighted similar trends, noting sensory drawbacks at high bran concentrations. Among the tested formulations, T5 (80:20::CM:SWB) achieved the highest appearance, texture, taste, and

overall acceptability ratings. This finding concurs with [56], who reported that bran-enriched biscuits were well-received. The reduction in sensory scores with higher bran levels is consistent with trends reported in pasta [25].

A significant decline in sensory characteristics was observed over a 90-day storage period. The mean scores for appearance, texture, taste, and overall acceptability decreased, likely due to non-enzymatic browning (Maillard reaction) and fat oxidation [57]. These changes align with results from [58], who documented similar declines in wheat bran-enriched meat products. A reduction in sensory scores with storage was also observed in bran-enriched chevon rolls [59].



UNDER PEER REVIEW

**Table 5: Effect of treatment and storage period on sensory evaluation (hedonic score) of tortilla chips**

Treatments	Taste				Texture				Appearance			
	0	30	60	90	0	30	60	90	0	30	60	90
T1(100:00::CM:SWB)	7.00	6.82	6.70	6.58	7.12	7.01	6.95	6.87	7.50	7.48	7.44	7.39
T2 (95:05::CM:SWB)	7.28	7.11	6.96	6.84	7.25	7.19	6.97	6.81	7.56	7.53	7.49	7.43
T3 (90:10::CM:SWB)	7.50	7.35	7.08	6.87	7.50	7.25	7.05	6.85	7.81	7.76	7.72	7.67

T4 (85:15::CM:SWB)	7.68	7.55	7.39	7.19	7.87	7.53	7.29	7.10	8.00	7.95	7.89	7.84
T5 (80:20::CM:SWB)	8.12	8.07	7.89	7.76	7.98	7.77	7.63	7.32	8.10	8.03	7.97	7.91
T6 (75:25::CM:SWB)	6.96	6.73	6.51	6.33	7.00	6.90	6.82	6.60	6.50	6.44	6.39	6.35
T7 (70:30::CM:SWB)	6.56	6.43	6.20	6.08	6.62	6.47	6.26	6.10	6.00	5.95	5.91	5.86
CD( $P=0.05$ )												
Treatment (T)		0.02				0.02					0.02	
Storage (S)		0.03				0.02					0.01	
T x S		0.06				0.04					0.04	

UNDER PEER REVIEW

## 5. CONCLUSION

Based on the results of this study, it is evident that incorporating microwave-stabilized wheat bran into tortilla chips can enhance certain nutritional and sensory properties, while also impacting shelf stability and microbial safety. The addition of wheat bran at levels up to 20% significantly improved the taste and overall acceptability of tortilla chips, as consumers appreciated the unique, slightly branny flavor. However, beyond 20% bran inclusion, sensory qualities like appearance and texture began to decline, possibly due to oxidative effects and increased moisture retention associated with higher fiber content. Water activity, and moisture content, increased with the inclusion of wheat bran, which is naturally hygroscopic and can potentially reduce shelf life, highlighting the need for optimized packaging to manage moisture uptake and maintain crispness or texture. During storage, the increase in moisture content also appeared to contribute to an elevation in microbial load, though levels remained within acceptable limits. Consequently, additional control measures may be needed for extended storage, especially in high-humidity conditions. Free fatty acids and peroxide values, both critical indicators of lipid stability, increased with the addition of wheat bran and over the storage period. These increases reflect bran's naturally higher free fatty acid content and the greater susceptibility to oxidation due to bran incorporation. Interestingly, the initial processing of tortilla chips through high-temperature baking and frying effectively limited initial peroxide values, suggesting a protective effect from thermal processing. However, during storage, the slight rise in peroxide and free fatty acid values underscores the need for antioxidant strategies in bran-enriched snacks.

The sensory evaluation results support the development of a tortilla chip formulation with 20% stabilized wheat bran, which was preferred for its taste, texture, and appearance. This formulation achieved high consumer acceptance on the hedonic scale, confirming that bran enrichment can positively influence sensory characteristics at moderate levels. The decrease in sensory scores during storage, attributed to Maillard browning and fat oxidation, aligns with common challenges in bran-enriched products. Despite these declines, the product retained acceptable sensory attributes up to 90 days, especially with proper storage measures to limit moisture uptake and oxidation.

Disclaimer (Artificial intelligence)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

## REFERENCES

1. Popkin BM, Reardon T. Obesity and the food system transformation in Latin America. *Obesity Reviews*. 2018; 19(2):102-116.
2. Monteiro CA, Cannon G, Lawrence M, Costa Louzada MD, Pereira Machado P. Ultra-processed foods, diet quality, and health using the NOVA classification system. *Public Health Nutrition*. 2019; 22(1):5-13.
3. Onipe OO, Jideani AIO, Beswa D. Composition and functionality of wheat bran and its application in some cereal food products. *International Journal of Food Science and Technology*. 2015; 50(12) 2509-2518.
4. Ketkaew C, Oktaviani RD, Padthar S, Leraanaksiri R, Meethonglang T, Hanna, A. Enhancing sustainable food innovation through consumer participatory co-creation: A case study on plant based dietary snack bars in Thailand. *Future Foods*. 2021; 10: 2-11.
5. Mahajan R, Panesar PS. Technological advancements in snacks processing: A review. *Journal of Food Processing and Preservation*. 2018; 42(10): 13710.
6. Tyl C, Bresciani A, Marti, A. Recent progress on improving the quality of bran enriched extruded snacks. *Foods*. 2021;10(9):1-20.
7. Serna-Saldivar SO, Gutierrez-Uribe JA, Garcia-Lara S. Phytochemical profiles and nutraceutical properties of corn and wheat tortillas. In *Tortillas*. AACC International Press, Mexico. 2015; 65-96.
8. Gonzalez R, Cockburn M. Consumer perceptions and sensory characteristics of fiber-enriched tortilla chips. *Journal of Sensory Studies*. 2020; 35(4): 125-159.
9. Prajapati PS, Joshi DC. Developing functional snacks using modified ingredients: A review. *Food Science and Nutrition*. 2021; 9(2):1032-1048.
10. Laukova M, Karovicova J, Minarovicova L, Kohajdova Z. Wheat bran stabilization and its effect on cookies quality. *Potravinarstvo Slovak Journal of Food Sciences*. 2019; 13(2):109-115.
11. AOAC. *Official Methods of analysis*. 14<sup>th</sup> Edition. Association of Official Analytical Chemists, Washington, DC; 2005.
12. AACC. *Approved Methods of American Association of Cereal Chemists*, Saint Paul, Minnesota, USA, 2000.
13. Palczar, MJ, Chan ECS. *Laboratory Exercise in Microbiology*. Black Dot Inc., New York. 1991.

14. Kumar R, Chambers E, Chambers DH, Lee J. Generating New Snack Food Texture Ideas Using Sensory and Consumer Research Tools: A Case Study of the Japanese and South Korean Snack Food Markets. *Foods*. 2021;10(2):474-496.
15. Nielsen OF, Bilde M, Frosch M. Water activity. *Journal of Spectroscopy*. 2012; 27(6): 565-569.
16. Dar BN, Sharma S, Nayik GA. Effect of storage period on physicochemical, total phenolic content and antioxidant properties of bran enriched snacks. *Journal of Food Measurement and Characterization*. 2016; 10(4): 755-761.
17. Thompson M, Green L. Moisture Absorption in Hygroscopic Foods: A comparative study of bran enriched and standard products. *Journal of Food Processing and Preservation*. 2019; 43(7): e13921.
18. Kumar P, Chatli MK, Mehta N, Malav OP, Verma AK, Kumar D. Quality attributes and storage stability of chicken meat biscuits incorporated with wheat and oat bran: Fiber enriched meat. *Journal of Food Quality*. 2016; 39(6): 649-657.
19. Ashraf S, Sood M, Bandral JD, Gupta N, Rahman R, Nishu. Effect of stabilization methods on bioactive components and stability indicators of wheat bran. *AMA, Agricultural mechanization in Asia, Africa and Latin America*. 2023; 54(8):309-322.
20. Kaur G, Sharma S, Nagi HPS, Dar BN. Functional properties of pasta enriched with variable cereal brans. *Journal of Food Science and Technology*. 2012;49(4): 467-474.
21. Taylor P, Roberts S. Storage stability of whole grain and bran-enriched snacks: water activity trends over time. *Journal of Food Storage and Preservation*. 2019; 45(6), 161-175.
22. Bordin K, Kunitake MT, Aracava KK, Trindade CSF. Changes in food caused by deep fat frying-A review. *Archivos Latinoamericanos de Nutricion*. 2013; 63(1): 5-13.
23. Muzamil Z, Razali UH, Qhairul N, Zamri NH. Physicochemical and sensory analysis of surimi sausage incorporated with rolled oat powder subjected to frying. *International Food Research Journal*. 2021;28(3):457-466.
24. Lee H, Park Y. Water holding capacity of dietary fibers in baked and extruded foods. *Food Hydrocolloids*. 2019; 91: 134-142.
25. Amel K, Amel M and Samir F. Effect of the incorporation of wheat bran on pasta quality. *Wulfenia*. 2015; 22(4): 225-234.
26. Sobota A, Rzedzicki Z, Zarzycki P, Kuzawinska E. Application of common wheat bran for the industrial production of high fibre pasta. *International Journal of Food Science and Technology*. 2015; 50(1): 111-119.

27. Thomas L, Green M. 2022. Hygroscopicity of bran-enriched snack foods during storage. *Food Research International*. 2022; 153: 110762.
28. Ajala O, Adelusi OA, Kajihansa OE, Onabanjo OO, Oyewole OB, Obadina AO. Effect of storage temperature and time on the microbial quality and sensory properties of extrudes produced from pearl millet and Bambara groundnut flour blends. *Journal of Stored Products Research*. 2024; 109: 102442.
29. Miller K, Tran Q. Effects of bran enrichment on moisture content in long-term storage of snack products. *Journal of Food Quality*. 2021; Article ID 5678923
30. Oghbaei M, Prakash J. Nutrient availability in deep fried flat breads prepared with bran enriched and mineral fortified whole and refined wheat flour. *Food and Nutrition*. 2018; 1(2):1-8.
31. Lee Y, Zhang M. Lipid Oxidation in Wheat Bran-Enriched Baked Products. *Food Chemistry*. 2021; 3(4): 78-99.
32. Sharma S, Kaur S, Dar BN, Singh, B. Storage stability and quality assessment of processed cereal brans. *Journal of Food Science and Technology*. 2014;5(3): 583-588.
33. Akhter S, Ahiduzzaman MD. Effect of storage life of rice bran on the quality of oil. *Journal of Food and Nutrition Science*, 2017; 5(1): 11-15.
34. Zhao Q, Li X. Effect of storage time on free fatty acid content in root vegetable chips. *Food Quality and Preference*. 2022; 9(6): 104-106.
35. Brown L, Lee Y. Standards for oxidation in fried foods: peroxide value guidelines. *Food Chemistry*. 2019; 27: 650-657.
36. Shobha D, Kumar HVD, Sreeramasetty TA, Puttaramanaik GKTP, Shivakumar GB. Storage influence on the functional, sensory and keeping quality of maize flour. *Journal of Food Science and Technology*. 2014; 55(11): 3154-3162.
37. Davis M, Johnson H. Thermal treatment effects on peroxide levels in fried foods. *International Journal of Food Science and Technology*. 2018; (4):1043-1050.
38. Kumar P, Chatli MK, Mehta N, Malav OP, Verma AK, Kumar D. Quality attributes and storage stability of chicken meat biscuits incorporated with wheat and oat bran: fibre-enriched meat snacks. *Journal of food quality*. 2016; 39(6):649-657.
39. Tangariya P, Sahoo A, Awasthi P, Pandey A. Quality analysis of composite flour and its effectiveness for Chapatti formulation. *Journal of Pharmacognosy and Phytochemistry*. 2018;7(4):1013-9.
40. Taylor P, Kim S. Effects of extended storage on lipid stability in tortilla chips. *Food Quality and Safety*. 2019; 3(2):123-130.

41. Zhao M, Liu Q. Sanitary quality and microbial indicators in processed foods. *Journal of Food Quality and Safety*. 2021; 15(1): 32-38.
42. Nagi HPS, Dar BN, Sharma S. Effect of storage period and packaging on the shelf life of cereal bran incorporated biscuits. *American Journal of Food Technology*. 2012; 7(5): 310-310.
43. Kim J, Chen R. Effect of bran enrichment on microbial quality in flatbread products. *Journal of Cereal Science*. 2020; 89: 101-107.
44. Chakroborty T, Chakroborty SC. Effect of storage temperature on the quality and microbial Content of salt-smoke-dried shoal (*Ophiocephalus striatus*). *Journal of Fisheries and Livestock Production*. 2017; 5(1): 1-4.
45. Singh A, Patel N. Microbial safety standards for stored food products. *Food Safety and Health Standards*. 2020; 10: 133-140.
46. Wang X, Yao T. Impact of storage on microbial stability in high-fiber foods. *Food and Bioprocess Technology*. 2018; 11(4): 222-229.
47. Zhao Y, Lin J. Microbial growth patterns in high-fiber and bran-enriched meat products. *Meat Science*. 2020; 1(6): 108-128.
48. Stone H, Bleibaum RN, Thomas HA. *Affective testing in sensory evaluation practices 5<sup>th</sup> edition*, Academic press. 2021; 297-336. <https://doi.org/10.1016/B978-0-12-815334-5.00004-5>.
49. Chaudhry P. Bran-enriched snack sensory evaluation using hedonic scale. *International Journal of Food Quality and Preference*. 2020; 95, 102-118.
50. Latif A. (2019). Consumer preference for bran-enriched cookies over refined flour varieties. *Journal of Cereal Science*, 90, 87-93.
51. Hussein Y, Saleh S. Flavor and texture preferences in wheat bran-enriched snacks. *Food Science and Technology*. 2020; 78: 1105-1114.
52. Mishra N. Utilization of waste defatted rice bran in formulation of functional cookies and its effect on physiochemical characteristics of cookies. *International Journal of Advanced Science and Research*. 2017; 2(5): 64-68.
53. Xhabiri G, Sinani V, Hoxha I. The impact of the wheat bran on the quality of biscuits and their nutritional value. *Anglisticum Journal*. 2016; 3(3): 66-73.
54. Nandeesh K, Jyotsna R, Venkateswara RG. Effect of differently treated wheat bran on rheology, microstructure and quality characteristics of soft dough biscuits. *Journal of Food Processing and Preservation*. 2009; 35 (2): 179-200.

55. Onipe OO, Jideani AIO, Beswa D. Composition and functionality of wheat bran and its application in some cereal food products. *International Journal of Food Science and Technology*. 2021; 50(12): 2509-2518.
56. Silky, Gupta MP, Tiwari A. Development of high fibre biscuits using wheat bran. *International Journal of Engineering and Innovative Technology*. 2014; 4(6): 90-94.
57. Thompson R, Richardson T. Maillard Reaction and fat oxidation in stored foods. *Food Chemistry*. 2021; 358, 129856.
58. Yadav S, Pathera AK, Islam RU, Malik AK, Sharma DP. Effect of wheat bran and dried carrot pomace addition on quality characteristics of chicken sausages. *Asian-Australasian Journal of Animal Science*. 2018; 31(5):729-737.
59. Parkash J, Yadav S, Sharma DP, Pathera AK, Islam RU. Effect of dried apple pomace and corn bran incorporation on quality characteristics of chevon rolls during refrigerated storage. *Journal of Animal Research*. 2016;6(3): 465-470.

UNDER PEER REVIEW