

Influence of nitrogen levels and zinc application on Morphological and Physiological attributes of rice varieties

Abstract

A field experiment was conducted during Kharif and Rabi, 2018 at College of Agriculture, Rajendranagar, Hyderabad to study the effect of different nitrogen levels and zinc application on growth and development in paddy. The experiment was laid out in split plot design with three varieties as main plots, six nutrient levels as sub plots and replicated thrice. Among the varieties Kunaram Sannalu has recorded lowest Plant height and Telangana Sona has recorded Highest Plant height (63.3, 98.8 and 103.3 cm at vegetative, flowering and grain filling stage respectively), Maximum Stem thickness 5.37, 6.09 and 6.17 mm at vegetative, flowering and grain filling stage respectively), Crop growth rate (13.451 25.77 and 8.86 g m⁻² day⁻¹ at vegetative, flowering and grain filling stage respectively), Total dry matter (2959, 7585 and 8909 kg ha⁻¹ at vegetative, flowering and grain filling stage respectively) and had taken more number of days to panicle initiation (67), flowering (86) and maturity (122). Tella Hamsa has recorded less Stem thickness, Crop growth rate, Total dry matter and had taken less number of days to panicle initiation (64), flowering (83) and maturity (118). Application of 25 % higher than RDN + 0.5 % ZnSO₄ foliar spray resulted in maximum Plant height, Stem thickness, Crop growth rate, Total dry matter.

Keywords: Plant height, Stem thickness, Crop growth rate, Panicle Initiation

1. INTRODUCTION

Rice is one of the most important cereal crops of the world which feeds half of the world's population providing 35-60% of the total calorie (Tayefe et al., 2014) [1]. During past few decades, rice production increased mostly due to adoption of high yielding varieties, increase in irrigated area and use of chemical fertilizers. However, the rate of increase in rice yield is static and if the rate is not possible to increase, severe food shortage is likely to occur in near future. To push up the yield ceiling, sustainable technologies are essential, which are economically viable and environmentally friendly. Cost minimization by saving resources and development of low cost technologies must be considered in rice production. Among the crop management practices, judicious application of nitrogenous fertilizer is paramount important for yield enhancement of rice.

Among the major nutrient elements, nitrogen (N) is the most limiting nutrient for rice crop growth and yield which is required in higher amounts compared to other nutrients (Djaman et al., 2018) [2]. N influences rice yield by playing major role in the photosynthesis, biomass accumulation, effective tillering, and spikelets formation (Yoshida et al., 2006) [3]. Therefore, N fertilization is imperative for modern rice varieties in order to exploit their full yield potential (Chamely et al., 2015) [4]. High yielding modern rice varieties show a greater response to applied nitrogen, while they differ in N demand depending on their genotype and agronomic traits under different climatic conditions (Rahman et al., 2007) [5]. On the other hand, excessive N application can lead to ground water pollution, increased production cost, reduced yield and environmental pollution (Djaman et al., 2018)[2].

Therefore, it is essential to achieve efficient use of nitrogen in chemical fertilizers, through cultivation techniques and fertilizer management with high nitrogen use efficiency and reducing nitrogen inputs from farming to the environment. Evaluating the reaction of rice to diverse doses of nitrogen will aid in the development of high nitrogen use efficiency varieties, and the screening of appropriate genotypes for all cultivated condition. Nitrogen use

efficiency is complex because rice yield was influenced by inherent factors such as the number of productive culms, grains per panicle and 1000 grain weight, in addition to plant management conditions. However, measuring genotypic differences in dry matter production and nitrogen use efficiency at the vegetative growth stage eliminates those additional variables affecting yield. Understanding the mechanisms regulating the processes of nitrogen uptake, assimilation, utilization efficiency and remobilization are crucial for the improvement of nitrogen use efficiency in crop plants. One important approach is to develop an understanding of the plant response to different nitrogen regimes and studying plants that show better growth under nitrogen limiting conditions. Studies on impacts of elevated nitrogen on growth dynamics, biomass partitioning, chaffy grain and nitrogen use efficiency are limited.

The higher dose of nitrogen causes excessive vegetative growth that leads to lodging of the crop and a consequent decline in filled grains per panicle (Zhang et al., 2014) [6]. Applied nitrogen has been found to have a synergistic effect with zinc in rice. It has been reported that the uptake and concentration of zinc increases substantially with an increase in the rate of nitrogen application (Jiang et al., 2008) [7]. Hence the present study was conducted to evaluate the effect of different levels of nitrogen and zinc application on morphological and physiological parameters in paddy.

2. MATERIALS AND METHODS

Field experiment was conducted on sandy clay soil in college farm, College of Agriculture, Rajendranagar, Hyderabad during Kharif and Rabi, 2018. The experiment was laid out in a split plot design with three replications. The seedlings of different rice varieties G1- Kunaram Sannalu, G2 - Tella Hamsa and G3 - Telangana Sona were selected as main plots. Fertilizers were given as N1 - RDN (120 Kg N ha⁻¹), N2 - 25 % less than RDN (90 Kg N ha⁻¹), N3 - 25 % higher than RDN (150 Kg N ha⁻¹), N4 - 25 % less than RDN + 0.5 % ZnSO₄ Foliar spray, N5 - 25% higher than RDN + 0.5 % ZnSO₄ Foliar spray, N6 - Control taken as sub plots.

The varieties were sown separately in raised bed nursery and 25 days old seedlings were transplanted into 15 m² (5 m X 3 m) plots by adopting a spacing of 15 cm between rows and 15 cm within a row. Nitrogen applied as per treatment in form of urea in 3 splits as basal, maximum tillering and flowering stage. Similarly, 0.5 % ZnSO₄ foliar spray was applied 3 times at tillering, panicle initiation and flowering stage. Phosphorus was applied as single super phosphate at the rate of 60 kg ha⁻¹ and Potash as muriate of potash at the rate of 40 kg ha⁻¹ as a basal dose at the time of transplanting. Irrigation and weed management was done time to time.

For analysis of physiological characters, in each plot five plants were tagged and observations were recorded at vegetative, flowering and grain filling stages. Plant height, Stem thickness, Crop Growth Rate, Total dry matter at vegetative, flowering and grain filling stages were recorded. Number of days taken in each genotype in each plot was noted in days to panicle initiation. Number of days taken for 50 % of plants to flower in each genotype in each plot was noted in days to 50 % flowering. In rice, as physiological maturity approaches the erect flag leaves starts desiccating. The number of days taken from sowing to physiological maturity was recorded in each plot and each replication was recorded and reported as days to maturity in different treatments.

The plant height was expressed in centimeters (cm) and measured from base of the plant to the tip of the terminal leaf or panicle on main stem. It was measured during the maximum vegetative, flowering and harvest stages. Stem thickness was expressed in centimeters (cm) and measured at the base of the mother stem using digital calipers. It was measured during the maximum vegetative, flowering and maturity stages. Crop growth rate is

the rate of drymatter production per unit land area per unit time. It was measured during the maximum vegetative, flowering and maturity stages. It was calculated by using this formula, $CGR = \frac{W_2 - W_1}{T_2 - T_1}$ and it is expressed as $g\ m^{-2}\ day^{-1}$. Five plants were cut at ground level from each plot of three replications for determining the dry matter accumulation at vegetative, flowering and harvest stage. The plants were then oven dried at 60 °C for 3 days until constant weight was attained and expressed in $kg\ ha^{-1}$. The experimental data recorded on different parameters were analyzed statistically by applying the technique of analysis of variance for split-plot design by using windostat software version 9.2.

3.RESULTS AND DISCUSSION

3.1 Plant height (cm)

Significant differences were observed between the genotypes in plant height at different stages of crop growth significantly influenced by nitrogen and zinc foliar spray (Table 1). Plant height recorded was 59.6 to 63.3 cm at vegetative, 91.4 to 98.8 cm at reproductive and 93.9 to 103.3 cm at harvest. Among the three genotypes studied highest plant height was recorded in Telangana Sona (G3). Plant height significantly varied at different growth stages with application of fertilizer. Mean values of height observed at three growth stages was 61.7, 95.7 and 99.8 cm. At 25 % higher than RDN + 0.5 % $ZnSO_4$ foliar spray (N5) application resulted in maximum growth at vegetative phase of the rice crop (64.8 cm) and at reproductive phase (98.0 cm) and at harvest stage (102.3 cm). Swaroopa and Lakshmi (2015) [8] conducted experiment with four treatments of nitrogen and results revealed that application of 150 % RDN recorded maximum plant height, while shorter plants were observed at 75 % RDN. The superior performance of treatments might be due to quick and better utilization of zinc through foliar feeding at different growth stages of rice which in turn increased the nutrient content and total uptake (Zayed *et al.*, 2011) [9].

Significant interaction was observed between genotype and fertilizer for plant height at various crop growth stages. Treatment N5 and G3 recorded maximum growth at vegetative (66.3 cm), flowering (101.5 cm) and harvest stage (105.7 cm). Genotype Telangana Sona (G3) has obtained taller plants at all three stages of crop growth. While, Kunaram Sannalu (G2) was found to be lowest height at all stages of crop growth. It can be concluded from the above experimental results that plant height was maximum in Telangana Sona (G3) and treatment 25 % higher than RDN + 0.5 % $ZnSO_4$ foliar spray (N5) Genotypes were more responsive to fertilizer treatments as indicated by higher plant height.

Table 1: Plant height (cm) in rice as influenced by different nitrogen levels and zinc foliar spray during *kharif* and *rabi*, 2018

Plant height Pooled (Kharif and Rabi)												
Treatment	Vegetative stage				Flowering stage				Harvest stage			
	G ₁	G ₂	G ₃	Mean	G ₁	G ₂	G ₃	Mean	G ₁	G ₂	G ₃	Mean
N ₁	60.8	62.9	64.0	62.6	91.9	97.0	99.3	96.1	94.7	103.0	104.2	100.6
N ₂	58.0	61.6	62.7	60.8	90.0	95.8	97.3	94.4	93.9	101.2	102.5	99.2
N ₃	62.6	64.3	65.4	64.1	93.3	98.1	100.6	97.3	95.7	104.0	105.1	101.6
N ₄	58.8	62.4	63.8	61.6	91.5	97.1	98.2	95.6	94.4	102.7	103.5	100.2
N ₅	63.4	64.8	66.3	64.8	93.8	98.7	101.5	98.0	96.5	104.6	105.7	102.3
N ₆	54.2	56.0	57.7	56.0	87.8	94.2	95.8	92.6	88.8	98.0	99.7	95.5
Mean	59.6	62.0	63.3	61.7	91.4	96.8	98.8	95.7	93.9	102.1	103.3	99.8
CD (5%)	Genotype(G)		0.30		0.91		0.12					
	Treatment(N)		0.34		0.38		0.33					
Main plots : Genotypes												
G ₁		59.6		91.4		93.9						

G₂	62.0	96.8	102.1
G₃	63.3	98.8	103.3
Mean	61.7	95.7	99.8
SEm_±	0.10	0.30	0.04
CD (5 %)	0.30	0.91	0.12
Subplots : Fertilizer treatments			
N₁	62.6	96.1	100.6
N₂	60.8	94.4	99.2
N₃	64.1	97.3	101.6
N₄	61.6	95.6	100.2
N₅	64.8	98.0	102.2
N₆	56.0	92.6	95.5
Mean	61.7	95.7	99.8
SEm_±	0.10	0.13	0.11
CD (5%)	0.34	0.38	0.33
Interaction			
Rice genotypes at same level of fertilizer treatments			
SEm_±	0.24	0.25	0.11
CD (5%)	0.58	0.74	0.29
Interaction			
Fertilizer treatments at same or different rice genotypes			
SEm_±	0.19	0.37	0.18
CD (5%)	0.57	1.11	0.54

3.2 Stem Thickness

Pooled data indicate that there was an increase in the stem thickness from maximum vegetative stage (4.72 mm) to harvest stage (6.02 mm) (Table 2). At vegetative stage, stem thickness ranged from 4.36 to 5.03 mm, at flowering stage from 5.60 to 6.09 mm and at harvest stage 5.77 to 6.17 mm. Telangana Sona (G3) was found maximum stem thickness at different crop growth stages (5.03, 6.09 and 6.17 mm at vegetative, flowering and harvest stage respectively).

Stem thickness varied significantly due to application of different dose of fertilizer treatments and ranged from maximum vegetative stage 4.12 to 4.96 mm, at flowering stage 4.50 to 6.30 mm and at harvest stage 4.64 to 6.42 mm. Application of fertilizer at 25 % higher than RDN + 0.5 % ZnSO₄ foliar spray recorded maximum thickness at different crop growth stages (4.96, 6.30 and 6.42 mm at vegetative, flowering and harvest stage respectively). Wujun *et al.* (2016) [10] found that stem diameter in rice increased rapidly with increased nitrogen application. Interaction of genotype and fertilizer application showed significant variation in the stem thickness at various crop growth stages. Treatment N5 and G3 recorded maximum growth at vegetative stage (5.22 mm), flowering stage (6.41 mm) and maturity stage (6.46 mm). Such interaction effects are common with application of varied doses of fertilizer application. Genotype Telangana Sona (G3) has recorded significantly higher thickness at all three stages of crop growth. Fertilizer treatments at same or different genotypes revealed maximum thickness with N5 at vegetative, reproductive and maturity stages.

Table 2: Stem thickness (mm) in rice as influenced by different nitrogen levels and zinc foliar spray during *kharif* and *rabi* 2018

Stem thickness Pooled (Kharif and Rabi)			
Treatment	Vegetative stage	Flowering stage	Maturity stage

	G ₁	G ₂	G ₃	Mean	G ₁	G ₂	G ₃	Mean	G ₁	G ₂	G ₃	Mean
N ₁	4.90	4.43	5.17	4.83	6.26	5.83	6.36	6.15	6.40	6.00	6.46	6.29
N ₂	4.77	4.37	5.05	4.73	6.15	5.71	6.28	6.05	6.33	5.86	6.37	6.20
N ₃	4.98	4.49	5.24	4.90	6.34	5.91	6.41	6.22	6.48	6.09	6.48	6.35
N ₄	4.83	4.40	5.11	4.78	6.20	5.77	6.34	6.10	6.38	5.93	6.42	6.24
N ₅	5.02	4.54	5.31	4.96	6.41	6.00	6.48	6.30	6.55	6.17	6.53	6.42
N ₆	4.12	3.94	4.31	4.12	4.51	4.35	4.66	4.50	4.59	4.59	4.73	4.64
Mean	4.77	4.36	5.03	4.72	5.98	5.60	6.09	5.89	6.12	5.77	6.17	6.02
CD (5%)	Genotype(G)	0.020			0.008			0.009				
	Treatment(N)	0.015			0.014			0.013				
Main plots : Genotypes												
G ₁	4.77			5.98			6.12					
G ₂	4.33			5.60			5.77					
G ₃	5.03			6.09			6.17					
Mean	4.72			5.89			6.02					
SEm _±	0.007			0.003			0.003					
CD (5 %)	0.020			0.008			0.009					
Subplots : Fertilizer treatments												
N ₁	4.83			6.15			6.29					
N ₂	4.73			6.05			6.20					
N ₃	4.90			6.22			6.35					
N ₄	4.78			6.10			6.24					
N ₅	4.96			6.30			6.42					
N ₆	4.12			4.50			4.64					
Mean	4.72			5.89			6.02					
SEm _±	0.005			0.004			0.004					
CD (5%)	0.015			0.014			0.013					
Interaction												
Rice genotypes at same level of fertilizer treatments												
SEm _±	0.009			0.008			0.007					
CD (5%)	0.026			0.024			0.022					
Interaction												
Fertilizer treatments at same or different rice genotypes												
SEm _±	0.010			0.008			0.007					
CD (5%)	0.028			0.026			0.023					

3.3 Crop growth rate (CGR)

Pooled results on crop growth rate were significantly affected by genotypes at various nitrogen levels (Table 3). Pooled results on crop growth rate among the genotypes noted a mean to range from vegetative to grain filling was 12.69 to 8.69 g m⁻² day⁻¹. Among the genotypes Telangana Sona recorded maximum CGR at different growth stages (13.45, 25.77 and 9.86 g m⁻² day⁻¹ at vegetative, flowering and grain filling respectively). Mean values among the various treatments ranged from vegetative to grain filling was 12.69 to 8.69. Application of 25 % higher than RDN + 0.5 % ZnSO₄ foliar spray was found to be highest CGR at different growth stages (13.87, 26.12 and 9.70 g m⁻² day⁻¹ at vegetative, flowering and grain filling respectively).

Interaction of genotype and fertilizer application showed significant variation in the CGR at various crop growth stages. Treatment N5 and G3 recorded maximum CGR at vegetative (14.86 g m⁻² day⁻¹), at flowering (28.24 g m⁻² day⁻¹) and at grain filling stage (11.12 g m⁻² day⁻¹). Such interaction effects are common with application of higher doses of

fertilizer application. Genotype Telangana Sona (G3) at vegetative, flowering and grain filling recorded highest CGR, while Tella Hamsa (G1) was observed lowest. Mannan *et al.* (2012) [11] found in rice that the supply of available nitrogen progressively increased the crop growth rate. These findings were in conformity with the results obtained by Roy *et al.* (2004) [12].

3.4 Total dry matter

The dry matter accumulation increased progressively in all the genotypes with the advancement of crop age (Table 4). Significant differences were observed between the genotypes in dry matter at all the stages of crop growth. Mean ranged from 2730 to 2959 kg ha⁻¹ at vegetative stage, 7108 to 7585 kg ha⁻¹ at flowering stage and 8604 to 8909 kg ha⁻¹ at harvest stage. Maximum dry matter was observed in Telangana Sona (G3) at different crop growth stages (2959, 7585 and 8909 kg ha⁻¹ at vegetative, flowering and harvest stage respectively).

Significant increase in dry matter was recorded due to nitrogen application and dry matter was significantly different with varied dose of fertilizer treatments. At maximum vegetative stage ranged from 2070 to 3379 kg ha⁻¹, at flowering stage from 6172 to 7905 kg ha⁻¹ and at harvest stage from 7123 to 9419 kg ha⁻¹. Application of fertilizer treatment N5 recorded highest dry matter at different crop growth stages (3379, 7905 and 9419 kg ha⁻¹ at vegetative, flowering and harvest stage respectively). The increase in plant height and tiller number would have contributed to higher dry matter accumulation which is the outcome of photosynthetic activity of the plant and its capacity to utilize available nutrients. Several research workers also observed significant increase in dry matter accumulation with foliar application of ZnSO₄ (Rao, 2003) [13].

Genotype x fertilizer application showed significant interaction for dry matter production. Treatment N5 and G3 recorded highest in dry matter at 3611 kg ha⁻¹ at vegetative, 8621 kg ha⁻¹ at flowering and 9769 kg ha⁻¹ at harvest stage. Such interaction effects are common with application of higher doses of fertilizer application. Genotype Telangana Sona (G3) and N5 has recorded highest dry matter.

3.5 Days to Panicle initiation, 50 % Anthesis and Maturity

Pooled data on days to PI, 50 % anthesis and maturity as influenced by nitrogen supply in rice genotypes is presented in table 5. Results from the pooled data revealed that number of days taken ranged from 64 to 67 days for to panicle initiation, 83 to 86 days for 50 % anthesis and 118 to 122 days for maturity. Tella Hamsa (G2) was taken less number of days for panicle initiation (64 days), 50 % anthesis (83 days) and days to maturity (118 days), while more number of days to panicle initiation (67 days), 50 % anthesis (86 days) and days to maturity (122 days) was taken by the genotype Telangana Sona (G3).

Panicle initiation, 50 % anthesis and days to maturity recorded was significantly different with the fertilizer treatments and ranged from 63-68 days for panicle initiation, 83 to 87 days for 50 % anthesis and 117 to 123 days for maturity. Application of fertilizer at 25 % higher than RDN (N3) and 25 % higher than RDN + 0.5 % ZnSO₄ foliar spray (N) were at par and were taken more number of days to panicle initiation (68 days), 50 % anthesis (87 days) and maturity (123 days). Variation in phenological characters depends on genotypic constituent, micro and macro environments (Shahidullah *et al.*, 2009) [14]. Interaction effect of genotypes and fertilizer levels for days to PI, 50 % anthesis and maturity was found to be non significant.

Table 3: Crop growth rate (g m⁻² day⁻¹) in rice as influenced by different nitrogen levels and zinc foliar spray during *kharif* and *rabi* 2018

Crop growth rate Pooled (Kharif and Rabi)

Treatment	Vegetative stage				Flowering stage				Grain filling stage			
	G ₁	G ₂	G ₃	Mean	G ₁	G ₂	G ₃	Mean	G ₁	G ₂	G ₃	Mean
N ₁	13.35	11.85	14.05	13.35	24.84	22.39	26.60	24.61	9.26	7.76	10.28	9.10
N ₂	12.36	11.31	13.23	12.30	23.49	21.13	24.73	23.12	8.22	7.16	9.47	8.28
N ₃	14.07	12.25	14.67	13.66	25.93	22.86	27.57	25.45	9.53	7.94	11.00	9.49
N ₄	12.67	11.51	13.38	12.52	24.09	21.6	25.32	23.67	8.33	7.40	9.63	8.45
N ₅	14.33	12.43	14.86	13.87	26.71	23.42	28.24	26.12	9.60	8.37	11.12	9.70
N ₆	10.27	9.80	11.19	10.42	20.79	19.65	22.15	14.86	7.04	6.61	7.67	7.11
Mean	12.84	11.78	13.45	12.69	24.30	21.84	25.77	22.97	8.66	7.54	9.86	8.69
CD (5%)	Genotype(G)	0.031			0.145			0.071				
	Treatment(N)	0.067			0.188			0.092				
Main plots : Genotypes												
G ₁	12.84			24.30			8.66					
G ₂	11.78			21.84			7.54					
G ₃	13.45			25.77			8.86					
Mean	12.69			22.97			8.69					
SEm±	0.011			0.049			0.024					
CD (5%)	0.031			0.145			0.071					
Subplots : Fertilizer treatments												
N ₁	13.35			24.61			9.10					
N ₂	12.30			23.12			8.28					
N ₃	13.65			25.45			9.49					
N ₄	12.52			23.67			8.45					
N ₅	13.87			26.12			9.70					
N ₆	10.42			20.86			7.11					
Mean	12.69			23.96			8.69					
SEm±	0.023			0.065			0.031					
CD (5%)	0.067			0.188			0.092					
Interaction												
Rice genotypes at same level of fertilizer treatments												
SEm±	0.040			0.113			0.055					
CD (5%)	0.117			0.326			0.159					
Interaction												
Fertilizer treatments at same or different rice genotypes												
SEm±	0.038			0.114			0.056					
CD (5%)	0.116			0.313			0.143					

Table 4: Total dry matter (Kg ha⁻¹) in rice as influenced by different nitrogen levels and zinc foliar spray *kharif* and *rabi* 2018

Total dry matter Pooled (Kharif and Rabi)												
Treatment	Vegetative stage				Flowering stage				Harvest stage			
	G ₁	G ₂	G ₃	Mean	G ₁	G ₂	G ₃	Mean	G ₁	G ₂	G ₃	Mean
N ₁	2750	2452	2948	2717	7254	6741	7718	7237	8920	8437	9219	8859
N ₂	2418	2153	2617	2396	6774	6360	7138	6757	8304	8121	8520	8315
N ₃	3081	2833	3379	3098	7568	6972	8050	7530	9153	8770	9419	9114
N ₄	2650	2302	2965	2639	7088	6758	7535	7127	8737	8321	8953	8670
N ₅	3428	3097	3611	3379	7817	7270	8628	7905	9353	9136	9769	9419
N ₆	2054	1921	2236	2070	6144	5929	6443	6172	7156	6640	7572	7123
Mean	2730	2460	2959	2716	7108	6672	7585	7122	8604	8238	8909	8583
CD (5%)	Genotype(G)	64.11			74.45			76.10				
	Treatment(N)	51.75			83.28			86.89				

Main plots : Genotypes			
G ₁	2730	7108	8604
G ₂	2460	6672	8238
G ₃	2959	7585	8909
Mean	2716	7122	8583
SEm _±	22.05	25.17	26.64
CD (5 %)	64.11	74.45	76.10
Subplots : Fertilizer treatments			
N ₁	2716	7237	8859
N ₂	2396	6757	8315
N ₃	3098	7530	9114
N ₄	2639	7127	8670
N ₅	3379	7905	9419
N ₆	2070	6172	7123
Mean	2716	7122	8583
SEm _±	17.83	28.69	29.94
CD (5%)	51.75	83.28	86.89
Interaction			
Rice genotypes at same level of fertilizer treatments			
SEm _±	29.42	39.20	40.71
CD (5%)	84.58	113.66	114.15
Interaction			
Fertilizer treatments at same or different rice genotypes			
SEm _±	31.50	41.41	43.71
CD (5%)	92.50	118.04	119.89

Table 5: Days to panicle initiation, 50 % anthesis and maturity of rice as influenced by different nitrogen levels and zinc foliar spray during *kharif* and *rabi* 2018

Treatment	Panicle initiation Pooled				50 % Anthesis Pooled				Days to maturity Pooled			
	G ₁	G ₂	G ₃	Mean	G ₁	G ₂	G ₃	Mean	G ₁	G ₂	G ₃	Mean
N ₁	67	64	68	66	86	83	86	85	122	118	123	121
N ₂	65	63	66	65	85	82	85	84	121	117	122	119
N ₃	69	66	70	68	88	84	89	87	123	119	125	122
N ₄	65	63	66	65	85	82	85	84	121	117	122	119
N ₅	68	65	70	68	88	84	89	87	123	119	125	122
N ₆	64	61	64	63	82	80	84	82	119	113	120	117
Mean	66	64	67	66	85	83	86	85	121	118	122	120
CD (5%)	Genotype (G)	0.21			0.28				0.35			
	Treatment(N)	0.50			0.45				0.45			
Main plots : Genotypes												
G ₁	66			85				121				
G ₂	64			83				118				
G ₃	67			86				122				
Mean	66			85				120				
SEm _±	0.07			0.10				0.14				
CD (5 %)	0.21			0.28				0.35				
Subplots : Fertilizer treatments												
N ₁	66			85				121				
N ₂	65			84				119				

N₃	68	87	122
N₄	65	84	119
N₅	68	87	122
N₆	63	82	117
Mean	66	85	120
SEm_±	0.17	0.15	0.15
CD (5%)	0.50	0.45	0.45
Interaction			
Rice genotypes at same level of fertilizer treatments			
SEm_±	0.36	0.27	0.30
CD (5%)	NS	NS	NS
Interaction			
Fertilizer treatments at same or different rice genotypes			
SEm_±	0.38	0.27	0.28
CD (5%)	NS	NS	NS

Conclusions and recommendations

REFERENCES

1. Tayefe M, Gerayzade A, Amiri E, Zade AN. Effect of nitrogen on rice yield, yield components and quality parameters. *African Journal of geology, ecology, and landscapes biotechnology*. 2014 Jan 1;13(1):91-105.
2. Djaman K, Mel VC, Ametonou FY, El-Namaky R, Diallo MD, Koudahe K. Effect of nitrogen fertilizer dose and application timing on yield and nitrogen use efficiency of irrigated hybrid rice under semi-arid conditions. *Journal of Agricultural Science and Food Research*. 2018 May 21;9(2):223. DOI: <https://hdl.handle.net/10568/102040>.
3. Yoshida H, Horie T, Shiraiwa T. A model explaining genotypic and environmental variation of rice spikelet number per unit area measured by cross-locational experiments in Asia. *Field Crops Research*. 2006 Jun 1;97(2-3):337-343.
4. Chamely SG, Islam N, Hoshain S, Rabbani MG, Kader MA, Salam MA. Effect of variety and nitrogen rate on the yield performance of boro rice. *Progressive Agriculture*. 2015 Aug 12;26(1):6-14. DOI: <https://doi.org/10.3329/pa.v26i1.24508>.
5. Rahman MH, Ali MH, Ali MM, Khatun MM. Effect of different level of nitrogen on growth and yield of transplant Aman rice CV BRR1 dhan32. *International Journal of Sustainable Crop Production*. 2007 Feb;2(1):28-34.
6. Zhang, W.J., Li, G.H., Yang, Y.M., Li, Q., Zhang, J., Liu, J.Y., Wang, S.H., Tang, S. and Ding, Y.F. Effects of nitrogen application rate and ratio on lodging resistance of super rice with different genotypes. *Journal of Integrative Agriculture*. 2014: 13(1): 63-72.
7. Jiang, W., Struik, P.C., Van Keulen, H., Zhao, M., Jin, L.N and Stomph, T.J. Does increased zinc uptake enhance grain zinc mass concentration in rice?. *Annals of Applied Biology*. 2008:153(1): 135-147.
8. Swaroopa, V. J and Lakshmi, M. B. Effect of nitrogen and foliar fertilization on yield components and quality parameters of machine transplanted rice. *Current Biotica*. 2015:9(3): 230-238.
9. Zayed, B.A., Salem, A.K.M and Sharkawy, H.M. Effect of different micronutrient

- treatments on rice (*Oryza sativa* L.) growth and yield under saline soil conditions. *World Journal of Agricultural Sciences*.2011:7(2):179-184.
10. Wujun, Z., Long-mei, W., Yan-feng, D., Fei, W., Xiao-ran, W., Gang-hua, L., Zheng-hui, L., She, T., Cheng-qiang, D and Shao-hua, W. Top-dressing nitrogen fertilizer rate contributes to decrease culm physical strength by reducing structural carbohydrate content in japonica rice. *Science Direct*. 2016:15(5): 992-1004.
 11. Mannan, M A., Bhuiya, M.S.U., Akhand, M.M and Zaman, M.M. Growth and yield of basmati and traditional aromatic rice as influenced by water stress and nitrogen level. *Journal of Science Foundation*. 2012:10: 52-62.
 12. Roy, B.C., Leihner, D.E., Hilger, T.H and Steinmueller, N. Tillering pattern of local and modern T.aman varieties as influenced by nitrogen rate and management practices. *Journal of Subtropical Agricultural Research Development*. 2004:2(2): 6-14.
 13. Rao, S.N. Rice (*Oryza sativa* L.) response to time and method of zinc application. M.Sc. (Ag) thesis submitted to Acharya N G Ranga Agricultural University, Hyderabad, India. 2003.
 14. Shahidullah, S.M., Hanafi, M.M., Ashrafuzzaman, M., Razi Ismail, M and Salam, M.A. Phenological characters and genetic divergence in aromatic rice. *African Journal of Biotechnology*.2009:8(14): 3199-3207.
 15. [THE EFFECT OF ZN-ENRICHED BIOCHAR ON PEROXIDE ENZYME ACTIVITY, AUXIN CONTENT, CHLOROPHYLL COUNT AND ZN OF RICE IN DIFFERENT FLOODING SYSTEMS](#) CELL BIOTECHNOLOGY AND MOLECULAR BIOLOGY 19 (5-6), 196–204