

Relationship analysis of Phenology, Stress Tolerance, and Mean Productivity in Wilt and Cold Stressed Chickpea (*Cicer arietinum* L.) Following Synthetic PGRs Application

ABSTRACT

To understand the phenotypic response to mitigate stress tolerance for achieving maximum mean productivity, a comparative study of four synthetic PGRs—Abscisic Acid (ABA), Naphthyl Acetic Acid (NAA), Salicylic Acid (SA), and Fusaric Acid (FA)—was conducted as a pot experiment against artificially inoculated *Fusarium oxysporum* (a wilt pathogen) and cold exposure in four chickpea varieties. Additionally, the relationship between traits and PGRS application was looked at to evaluate their role in the stress tolerance mechanism. The results showed that ABA at 5 and 2 ppm was effective in delaying flowering, therefore extending the vegetative development phase in plants. In this way, flowering promotes stress tolerance while evading the damaging impacts of wilting and cold. This resulted in a lower percentage of wilt and a reduced incidence of cold compared to all other treatments, which raised mean productivity. The use of ABA at 5 and 2 ppm has been shown to positively correlate with both the prolongation of vegetative development and the delay of flowering. But fusaric acid (FA), a fungal toxin, is what caused early flowering, which allowed the flower to coincide with the development of wilt and cold. The occurrence of wilt at seedling and cold during flowering accelerated the incidence of wilt and cold, which led to a reduced mean productivity after giving fusaric acid (FA) @ 10 and 20 ppm. According to the investigation, the application of fusaric acid (FA) at 10 and 20 ppm was found to be positively and highly correlated with an increase in the incidence of wilt and cold and consequently negatively correlated with the mechanism possessed by ABA at 5 and 2 ppm.

Key word: Phenology, Abscisic Acid (ABA), Fusaric Acid (FA), Correlation, Stress Tolerance, Mean Productivity

Introduction: After field peas and dry beans, chickpeas (*Cicer arietinum* L.) are the most significant winter-season grain legume crop globally. It is also a crucial part of crop rotation. Currently, chickpeas are grown on around 13.8 million hectares of land globally, yielding 9.83 q ha⁻¹ and producing 13.65 million tons. Grown on around 10.5 million hectares, it yields 11.15 million tons and 10.56 q ha⁻¹ productivity in India. Madhya Pradesh has 3.5 million hectares of total chickpea land, with a productivity of 10.82 q ha⁻¹ and a production of 4.59 million tons (1). Protein (220 g kg⁻¹), total carbohydrates (670 g kg⁻¹), starch (470 g kg⁻¹), and fat (50 g kg⁻¹) are all present in significant amounts in chickpea seeds. They

thus have a major impact on the human dietary system (2). In most chickpea-growing regions of the world, Fusarium wilt, caused by *Fusarium Oxysporum*, significantly reduces chickpea yield. Fusarium wilt is regarded as a significant foreign disease that can cause annual chickpea output losses ranging from 10% to 15%, with the potential to cause the crop to completely collapse under certain circumstances (3). Furthermore, chickpeas exhibit poor pod set, sterile pods, abortion of pods, poor flower set, and abortion of flowers when they are subjected to cold stress during the reproductive phase, particularly during flowering and pod formation (4). The following can be used to categorize the many phenological stages of chickpea growth: emergence, flowering, pod initiation, pod set, seed development, and physiological maturity. Since the duration of the reproductive phase and the environmental circumstances that prevail during flowering impact the percentage of fruit set and the eventual yield, flowering is regarded as the key period (5). Planning, scheduling, and carrying out our farm tasks on time can be made easier with knowledge of the timing and variety of phenological phenomena. With this knowledge, we may also modify our farming practices to maintain our farm's produce in the face of anticipated climate change (6). Acquiring an understanding of the correlation between phenological events and other qualities is also necessary to manipulate or perform agricultural plant behavior in a way that interests us (7). **Plant hormones**—which are primarily organic compounds—modify the developmental pattern and yield response of crops. With the assistance of PGRs, which primarily regulate the defensive responses of plants through antagonistic and synergistic interactions known as signaling crosstalk, plants have evolved sophisticated mechanisms to detect environmental signals and can initiate an ideal reaction to stressful circumstances (8). Therefore, the present investigations are aimed to studying the relationship between phenology, stress tolerance, and mean productivity in wilt and cold-stressed chickpea (*Cicer arietinum* L.) after the application of different synthetic PGRs.

Materials and methods: In the years 2020–2021 and 2021–2022, the research was conducted at the Herbal Garden Department of Plant Physiology, JNKVV, Jabalpur (MP). Four varieties (V1 (JG74, susceptible to wilt), V2 (JG11, susceptible to cold), V3 (RAJAS, resistance to wilt), and V4 (PBG5, tolerant to cold) were used in the factorial completely randomized design of the crop planted in pots. At various concentrations Spraying of PGRs, like Abscisic Acid (ABA), Naphthyl Acetic Acid (NAA), Salicylic Acid (SA), and Fusaric Acid (FA), was performed two times, once during the early seedling stage and again at the flower initiation stage.

Method of wilt and cold application: When crops reach the 20 DAS stage, inoculation of wilt in pots was performed according to the method used by (9), and in accordance with the treatment schedule, the

first foliar spray of various PGRs was applied when plants showed signs of wilting. At the floral initiation stage, cold treatment was also artificially done as per the method used by (9), and after applying wilt and cold treatments, a second spraying of the PGRs was done.

Phenological studies: Throughout all crop seasons, visual observations were made every two days to track the crop's phenological development from the beginning of flowering until maturity.

Wilt incidence (%): After applying wilt inoculum in a pot, the number of plants that displayed noticeable signs such as wilting, chlorosis, and browning of the vascular system was counted, and the Mayee and Datar (1986) (10) methodology was used to determine the percentage wilt incidence.

Cold incidence (%): It was calculated using the method proposed by Mayee and Datar (1986) (10), with appropriate modifications.

Mean productivity - Mean Productivity was calculated by using formula -

$$MP = (Y_{pi} + Y_{si}) / 2$$

Y_{si} and Y_{pi} are the mean grain yields of individual treatments under stressed and non-stressed conditions.

Statistical analysis - The data were statistically analyzed through a completely randomized design given by Fisher (1955), and comparison of means was performed on the basis of the least significant difference test (LSD) according to the method given by Gopinath et al. (2021) (11). The association between the features we took into consideration in our study and the application of PGRs, as well as their respective contributions to the stress tolerance mechanism for achieving maximum mean productivity was subsequently studied using PCA biplot analysis.

Results and Discussion

From the pooled data (figure 1) obtained from years (2020-2021 and 2021-2022), it was found that hormonal treatment differed significantly ($p > 0.05$) for days required to flower initiation, completion of flowering, pod initiation, and seed development. Higher number of days required for flowering (figure 1) (48.39 & 47.20 DAS), complete flower formation (70.40 & 69.95 DAS), pod initiation (79.86 DAS), and seed development (87.35 DAS) were taken under the treatment of ABA @ 5 PPM and ABA @ 2 PPM. The application of ABA delayed flowering, increasing the length of vegetative growth and preventing

flowers from being negatively impacted by cold treatment. This prevented flowers from being susceptible to cold shock and lessened its damaging effects. Increased photo assimilated amounts that the crop plants used for their reproductive growth cycle are also attributed to longer vegetative growth durations (12). A PCA-based biplot analysis (figure 2, 3, 4) was used to attribute the relationship between plant modulators and traits that show stress tolerance and boost crop yields under wilt and cold-induced conditions. The PCA-biplot analysis's trait vector angles show how the variables are correlated. When the angle between two trait vectors is less than 90°, it is positively correlated, when it is more than 90°, it is negatively correlated, and when it is equal to 90°, it is correlated independently. The characteristics linked to stress tolerance were distinguished into four categories using the PCA-biplot analysis: strong positive correlations, positive correlations, independent correlations, and negative correlations. The results showed that both treatments had a favorable correlation with the start and completion of flowering and the number of days required for pod initiation and seed development; as a result, the more these features were valued by the plants after these treatments were sprayed, On the other hand, using PCA-biplot (figure 2, 3, 4) analysis's a negative correlation was found between ABA @ 5 PPM and ABA @ 2 PPM with fusaric acid treatment @ 20 ppm and 10 ppm. This resulted (figure1) in the earliest flowering (40.24 & 41.57 DAS) and complete flower formation (63.28 & 64.83 DAS), as well as a shorter duration of days required for pod initiation and seed development under both of these treatments. One possible explanation is that fusaric acid, a fungal toxin, causes wilt in treated plants, and plants that experience abnormalities from internal or external sources, attempt to complete their life cycle as quickly as possible, which leads to early flowering when fusaric acid is applied (13).




FIGURE1: Effects of different PGRs spray and varietal difference on phenophases, wilt & cold incidence and mean productivity of wilt and cold stressed chickpea.

Variety V4 (PBG5) (figure1) had a greater number of days needed for flower initiation and flowering completion than variety V1 (JG74), which had the earliest flower initiation and completion times, at 41.55 & 63.17 days, respectively. Similarly, V₄ (PBG5) (figure1) required more number of days for pod initiation (79.86 DAS) and seed formations (87.35 DAS) due to its normal growth period, which resulted from its tolerance capacity to nullify adverse impacts imposed by wilt and cold, whereas V1 (JG74) had a requirement of a smaller number of days for pod initiation (69.20 DAS) and seed formations (73.61 DAS), which might be due to changes in the growth period imposed by wilt and cold. Similar findings were also observed by (14), who discovered that the variety with wilt incidence (17.20%) required significantly less time to reach 50% flowering than the variety with incidence (11.50%), which required more time. Here again, a positive correlation (figure 2, 3, 4) is found between phenophase and stress tolerance. The variety that can tolerate stress better, grows normally, or has a longer lifespan. These traits are similar to the trend observed with the hormone.

The number of days needed for physiological maturity (107.78 & 105.53 DAS) and harvestable maturity (119.96 & 116.89 DAS) (figure1). increased as a result of treating the plants with ABA @ 5 ppm and ABA @ 2 ppm. Again, both treatments had a positive correlation with both of these traits and achieved a higher value than other treatments. Whereas, due to the negative correlation (figure 2, 3, 4) of fusaric acid treatment @ 20 ppm and @ 10 ppm with ABA @ 5 ppm and ABA @ 2 ppm, it took a shorter time to acquire physiological (92.90 & 95.54 DAS) and harvestable maturity (101.70 & 104.50 DAS) (figure1). under fusaric acid treatment @ 20 ppm and @ 10 ppm. Our results are consistent with the findings of (15), who observed that crop plants reduce their growth period and attempt to complete their life cycle in a shorter amount of time by making the most use of available resources when they experience abnormalities brought on by any internal or external factors. In comparison (figure1), genotype V1 (JG74) needed the fewest days (93.50 DAS) and (101.28 DAS) for physiological development (104.01 DAS) and harvestable maturity (115.60 DAS), while genotype V4 (PBG5) required the maximum days (figure1). Stress tolerance and susceptibility traits vary among genotypes, and this is reflected in the range of days needed for physiological maturity and harvestable maturity (16). For physiological development and harvestable maturity, genotype V4 (PBG5) (figure1), needs the greatest number of days, while genotype V1 (JG74), which is more sensitive, needs the fewest days.




FIGURE 2: Relationship study between traits under investigation where MP – mean productivity, PI – days required for pod initiation, FI - days required for flower initiation, CF - days required for complete flowering , DSF - days required for seed formation, DPM- days required for physiological maturity and DHM- days required for harvestable maturity




FIGURE 3: Relationship study between different PGRs spray in wilt and cold stressed chickpea.




FIGURE 4: Combined study of relationship between PGRs spray and there effect on traits under investigations in wilt and cold stressed chickpea.

Pooled data from both years (2020-2021 and 2021-2022) (figure1) shows that treatments with FA @ 20 ppm and FA @ 10 ppm exhibited the highest disease incidence of 31.25% and 30.71%, respectively. One well-known fungus toxin that causes chickpea death is fusaric acid (17). FA's synergistic actions with wilt inoculums caused foliar sprays to accelerate wilting in wilt-contaminated pots, which in turn caused the chickpea crop to die soon. Due to alterations in crop plant structure and functional activities, plants became more susceptible to several kinds of stresses when disease incidence increased under FA @ 20 ppm and FA @ 10 ppm (18). Consequently, under FA @ 20 ppm and FA @ 10 ppm, cold incidence (23.96% and 22.29%) (figure1) was likewise demonstrated to be higher. One possible explanation for plants' vulnerability to these treatments could be their inability to activate osmoregulation mechanisms during their active stage of development (19). Again, a favorable link (figure2, 3, 4) has been shown between fusaric acid treatment and the incidence of cold and wilt, and it might be due to the shorter duration of phenophases as depicted in the PCA biplot analysis, which showed a negative association with cold and wilt incidence.

After artificial wilt inoculation, plants receiving ABA @ 5 and ABA @ 2 ppm were the least wilted (18.75 and 20.21%) (figure1). It might be due to the fact that ABA is a major PGR responsible for activating defensive mechanisms in infected plants through upregulation of ABA-dependent transcription pathways (20) or enhancement of tolerance-responsive gene expression, both of which are important for providing crop plant tolerance against potential stress damage (21). Here, we can observe the relationship between ABA treatment and phenophase delay, which allows the plant more time to express the genes, involved in the defensive system or activates its own defense mechanism, ultimately leading to tolerance and a lower wilt incidence in comparison to other PGRs that were used in our study.

It was also discovered (figure1) that with both treatments, the percentages of cold occurrence were reduced, at 16.46% and 17.29%. This result could be explained by the fact that plants biosynthesize various types of small molecules known as stress proteins, or heat-shock proteins (Hsp), in response to wilt stress. These molecules may act as stress-mitigating agents by maintaining the water potential in cell sap and forming a barrier against membrane disruption and solute leakage, giving the plant the ability to cope with stress by protecting intracellular activity (22). Based on earlier findings, it is now evident that delayed phenophases following ABA

application at 5 ppm and 2 ppm are crucial in reducing the percentage of cold and wilt occurrences. The cultivar JG 74 (which is prone to wilt) had the highest incidence of wilt (36.67%), while the lowest (18.72%) was found in PBG 5.

Due to wilt pathogen disruptions in crop plant structure and functional activities, JG 74 also showed an increased incidence of cold (25.83%) (figure1). All other cultivars utilized in the research had a higher cold incidence than PBG 5 (15.52%). Under simulated exposure to wilt and cold, the plant sprayed with ABA at 5 ppm and ABA at 2 ppm was able to sustain its stress tolerance response and displayed its maximum mean productivity (12.73 & 12.33 g plant⁻¹). Similarly, the best mean productivity (12.50 g plant⁻¹) was achieved by variety PBG 5. Since FA is a natural fungal toxin, substantial plant mortality happened when it was sprayed at 20 ppm and 10 ppm, which decreased plant productivity under these treatments. Genotype JG 74's mean production was likewise reduced (10.40 g plant⁻¹) because of the obstacle imposed by stressful circumstances.

Conclusion

Plants sprayed with ABA @ 5 ppm and ABA @ 2 ppm effectively manage the damage caused by the wilt at an early stage and the cold at a later stage, thus providing tolerance to the crop to withstand unfavorable conditions and minimizing the yield loss due to the impact of wilt and cold. We discovered a strong and positive correlation between phenology and hormones by using PCA biplot analysis. Hormones that lengthen the life of crop plants may also help in lowering the probability of cold and wilt, same was found in the cases of ABA @ 5 ppm and ABA @ 2 ppm. Similarly, variety PBG 5 performed well under both (wilt and cold) adverse-induced conditions due to its longer life span.

References

1. Anonymous 2019. ANNUAL REPORT, Department of agriculture cooperation and farmer's welfare Ministry of Agriculture & Farmers Welfare Government of India
2. Kaur R and Prasad K. 2021. Technological, processing and nutritional aspects of chickpea (*Cicer arietinum*). A review. Trends Food Sci. Technol. 109, 448–463.
3. Achari SR, Mann RC, Sharma M and Edwards J. 2023. Diagnosis of *Fusarium oxysporum* f. sp. *ciceris* causing *Fusarium* wilt of chickpea using loop-mediated isothermal amplification (LAMP) and conventional end-point PCR. Sci Rep. 13(1):2640.

4. Rani A, Kiran A, Sharma KD, Prasad PVV, Jha UC, Siddique KHM and Nayyar H. 2021 .Cold Tolerance during the Reproductive Phase in Chickpea (*Cicer arietinum* L.) Is Associated with Superior Cold Acclimation Ability Involving Antioxidants and Cryoprotective Solutes in Anthers and Ovules. *Antioxidants* (Basel).10(11):1693.
5. Asha Kiran, Sanjeev Kumar, Harsh Nayyar and Kamal Dev Sharma. 2019. Low temperature-induced aberrations in male and female reproductive organ development cause flower abortion in chickpea.42 (7):2075-2089.
6. Banday ZZ and Nandi AK. 2015. Interconnection between flowering time control and activation of systemic acquired resistance. *Front. Plant Sci.* 6, 174.
7. Biasi R, Brunori E, Ferrara C and Salvati L. 2019 Assessing Impacts of Climate Change on Phenology and Quality Traits of *Vitis vinifera* L.: The Contribution of Local Knowledge. *Plants*, 8, 121.
8. Zhang Y, Xu J, Li R, Ge Y, Li Y and Li R. 2023 Plants' Response to Abiotic Stress: Mechanisms and Strategies. *Int J Mol Sci.*; 24(13):10915.
9. Thakur S, Tiwari G, Kumawat RK, Kumhare A, Singh S, Singh R. 2023. Synthetic PGR's Modify Phenology, Stress Tolerance and Mean Productivity in Wilt and Cold Stressed Chickpea (*Cicer arietinum* L.). *International Journal of Environment and Climate Change.* 13(11):1055-68.
10. Mayee CD and Datar VV 1986. *Phytopathometry*. Marathwada Agric. Univ. Parbhani Tech. Bull.-1: 66
11. Gopinath Pratheesh P , Parsad Rajender , Joseph Brigit and Adarsh VS. 2021. grapesAgri1: Collection of Shiny Apps for Data Analysis in Agriculture. *Journal of Open Source Software.* 6(63):34-37
12. Santiago Julián Kelly, María Gabriela Cano, Diego Darío Fanello, Eduardo Alberto Tambussi, Juan José Guiamet, 2021, Extended photoperiods after flowering increase the rate of dry matter production and nitrogen assimilation in mid maturing soybean cultivars, *Field Crops Research*, 265: 108104.
13. Singh VK, Singh HB, Upadhyay RS. 2017 Role of fusaric acid in the development of 'Fusarium wilt' symptoms in tomato: Physiological, biochemical and proteomic perspectives. *Plant Physiol Biochem.* 118:320-332.
14. Veeramani P and Sendhilvel V. 2020. Evaluation of Chickpea (*Cicer aritinum* L.) Varieties against Wilt Disease in North Eastern Hilly Zone of Tamil Nadu *J Krishi Vigyan*, 9 (1):114-117
15. Qi F and Zhang F 2020. Cell Cycle Regulation in the Plant Response to Stress. *Front. Plant Sci.* 10:1765. doi: 10.3389/fpls.2019.01765
16. Wabena Darkwa, Daniel Ambachew, Hussein Mohammed, Asrat Asfaw and Matthew W. Blair, 2016. Evaluation of common bean (*Phaseolus vulgaris* L.) genotypes for drought stress adaptation in Ethiopia, *The Crop Journal*, 4(5): 367-376,
17. Perincherry L, Lalak-Kańczugowska J and Stępień Ł. 2019 *Fusarium*-Produced Mycotoxins in Plant-Pathogen Interactions. *Toxins* (Basel).11(11):664.

18. Seleiman MF, Al-Suhaibani N, Ali N, Akmal M, Alotaibi M, Refay Y, Dindaroglu T, Abdul-Wajid HH and Battaglia ML 2021. Drought Stress Impacts on Plants and Different Approaches to Alleviate Its Adverse Effects. *Plants (Basel)*; 10(2):259.
19. Zhao S, Zhang Q, Liu M, Zhou H, Ma C and Wang P. 2021. Regulation of Plant Responses to Salt Stress. *Int J Mol Sci.*22(9):4609.
20. Tischer SV, Wunschel C, Papacek M, Kleigrew K, Hofmann T and Christmann A.2017. Combinatorial interaction network of abscisic acid receptors and coreceptors from *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. U. S. A.* 114 (38):10280–1028
21. Sabagh EL, Hossain A, Islam A, Dinajpur MS, Iqbal B, and Aftab MA, 2021. Prospective role of plant growth regulators for tolerance to abiotic stresses, Springer Nature Switzerland AG 1–38.
22. Sun W, Van Montagu M and Verbruggen N. 2002. Small heat shock proteins and stress tolerance in plants. *Biochimica et Biophysica Acta (BBA)-Gene Structure and Expression*, 1577(1)1-9.