

Evaluation of Capsicum (*Capsicum annuum*) Varieties in Naturally Ventilated Poly House Condition

Comment [D1]: under

ABSTRACT

An experiment entitled, "Evaluation of Capsicum (*Capsicum annuum*) Varieties in Naturally Ventilated Poly House Condition." was conducted at Horticulture Research Farm, Department of Horticulture, during the year 2022-23. Ten Capsicum Varieties, Names Sweet Pepper Nemalite, Sweet Pepper Cebrail, Sweet Pepper Bungl. Capsicum Varun, Sweet Pepper Volante, Shehzadi. Alpine 8510, Alpine 8248, Capsicum Orange and Capsicum IMP were evaluated at SHUATS, Prayagraj in randomized block design with three replications during winter season 2022-23 for find out best performing variety in terms of growth, yield and quality. The Sweet Pepper Volante was found maximum performance with the maximum fruit weight (154.20g), Fruit length (8.83cm), Fruit yield per plant (1.47 kg), Fruit yield (17.72 q/200 sq. m.). Sweet Pepper Volante was found superior based on overall performance in term of growth, yield and quality and highest cost benefit ratio (2.79) was found for capsicum in Sweet Pepper Volante variety under naturally ventilated polyhouse condition.

Comment [D2]: under

Comment [D3]: location of research

Comment [D4]: viz.,

Comment [D5]: to evaluate

Comment [D6]: remove

Comment [D7]: the performance of capsicum

Comment [D8]: to be promising

Comment [D9]: lower case

Comment [D10]: lower case

Comment [D11]: and

Comment [D12]: lower case

Comment [D13]: express as per plant or per ha

Comment [D14]: . The highest cost benefit ratio (2.79) was recorded in Sweet Pepper Volante variety under naturally ventilated polyhouse condition.

Comment [D15]: remove

Comment [D16]: add more key words

Comment [D17]: Italicize

Comment [D18]: Italicize

Keywords: Capsicum, Performance, Varieties.

INTRODUCTION

Capsicum (*Capsicum annuum* L. var. *grossum*) belonging to Solanaceae family, is native to Mexico (Bukasov, 1930; CABI, 2019) with Guatemala as secondary centre of origin (Bukasov, 1930). In 16th century, Portuguese and Spanish explorers introduced bell pepper from South America to Asia through trade routes (CABI, 2019). It is highly variable herb, or sub-shrub type annual crop attending 0.5 to 1 m in height with highly branching characters (CABI, 2019), and is available in diverse colors, such as green, red, black, orange, purple etc (Kumar *et al.*, 2015). It is grown worldwide for cooked vegetable, salad, pickle and processing purpose. The reason behind its worldwide admiration is its delicate taste, pleasant flavor, and varied colors with nutritious qualities. Its beautifully shaped form and vivid diverse color entitled it as "The Christmas ornaments of the vegetable world" (Sharma *et*

al., 2019). Furthermore, presence of vitamin C and zinc makes it better for strong and healthy immune system. Fatty acids, flavonoids, volatile oil, carotene pigment, beta carotene, iron, potassium, calcium, vitamin A and rutin (a bioflavonoid) are some compounds which are vastly available in bell pepper (*Agarwal et al., 2007*).

The traditionally grown green capsicum, depending upon variety and season, usually yields 20-40 tons per hectare in about 4-5 months. In greenhouse, the crop duration of green and coloured capsicums is about 7 -10 months and yields about 80-100 t ha⁻¹. Despite its economic importance, growers are not in a position to produce good quality capsicum with high productivity due to various biotic (pest and diseases), abiotic (rainfall, temperature, relative humidity and light intensity) and crop factor (flower and fruit drop), which ultimately affects the crop productivity adversely (*Ochigbo and Harris, 1989*). Growing of capsicum under polyhouse has been reported to give high productivity of good quality produce in developed countries and several hybrids have been developed recently. Hence, there is a need to evaluate their performance under naturally ventilated polyhouse conditions for getting higher productivity of excellent quality under Indian conditions. Capsicum is one such vegetable which befits into cultivation under protection to increase the production and yield. There are several capsicum varieties and hybrids available and being cultivated by farmers on commercial scale under open field condition. Hence, there is a need to evaluate various commercial capsicum hybrids for recommending its cultivation under greenhouse condition.

Comment [D19]: per ha/ha⁻¹

Different types of structures are being used for improving the productivity and profitability of horticultural crops as well as growing of the crops and planting material throughout the year like green house, shade house, lath house, mist house etc. In modern research, green house technology can be utilized for control of environmental parameters such as temperature, relative humidity, light intensity, light duration, CO₂ level, irrigation, nutrient supply and uptake, spacing, growing medium and root development (*Baghel et al., 2003*).

Materials and Methods

This experiment was carried out during 2023 at Crop Research Farm, Department of Horticulture, SHUATS, Prayagraj, UP, which is located at 25.28°N latitude, 81.54°E longitude and 98 m altitude above the mean sea level. This area situated on the right side of the river Yamuna by the side of PrayagrajRewa Road about 5 km away from Prayagraj, city. The climate of this region is typically sub-tropical and semi-arid with monsoon commencing

by the third week of June and with drawing by end of September. The temperature reached up to 48°C and in winter it goes down to as low as 2-3°C. During the summer hot scorching winds known as “Loo” and frost during winter months are common features. Experimental mechanical analysis of the soil was sand 59.50 (%), silt 24.10 (%), clay 16.40 (%) and textural class silt loam, while chemical analysis of soil was available nitrogen (242 kg/ ha), available phosphorus(24.50 kg/ ha), available potassium(95.00 kg/ ha), organic carbon(0.40 %), pH(7.50) and EC(0.19 dS/ m). The material for the study was comprised of 10Capsicumvariety. The variety were raised in field experiment in randomized block design with three replications. Growth, yield and yield attributing parameters were studied and data were collected from the five randomly selected plants from each plot. The data collected on different parameters was subjected to statistical analysis as per method of analysis prescribed by Panse and Sukhatme (1995).

Comment [D20]: taken

Results and Discussion

Vegetative parameters

Data presented in Table 1 showed that plant height and Number of branches of Capsicum varied significantly under the influenced by different variety at harvest stage. Plant height increased continuously with advancement of crop age and attained its maximum value at the maturity stage irrespective of variety. Maximum plant height and Number of branches (77.47 and 4.60), at harvest was recorded V₄ (Capsicum Varun), followed by V₅ (Sweet Pepper Volante) and minimum plant height and Number of branches (56.40 and 3.47) was recorded under V₉ (Capsicum Orange). Environment is the aggregate of all external conditions which influence the growth and development of crop, which play dominant role in crop production. Each crop has its own set of environmental conditions under which it grows best. Generally, crops are not profitable unless they are adapted to the region in which they are produced. Raising a crop successfully means the crop must be productive and economical to grow under prevailing conditions (Reddy, 1999). Temperature is the major regulator of development processes. It influences the flower and fruit development. The effect of temperature on net photosynthesis is a vital concern in crop production. The higher temperatures have more adverse influence on net photosynthesis than lower temperatures leading to decreased production of photosynthates above a certain temperature (Reddy, 1999). The temperature can be controlled and regulated under protected condition, therefore healthy and better growth of plants can be expected under protected condition. The

Comment [D21]: unit

Comment [D22]: in V₄

Comment [D23]: unit

differential response of vegetative growth of the different variety may be due to differences in genetic constituents of the varieties and microclimate condition (Bergefurdet *et al.*, 2011).

Table 1. Vegetative parameters of Different Varieties in Capsicum Varieties in naturally ventilated polyhouse.

Comment [D24]: under

Varieties	NAME OF VARIETIES	Height of Plant (cm)	Number of Branches/plant
V1	Sweet Pepper Nemalite	68.20	4.13
V2	Sweet Pepper Cebrail	70.27	4.33
V3	Sweet Pepper Bungi	67.27	4.27
V4	Capsicum Varun	77.47	3.53
V5	Sweet Pepper Volante	72.87	3.47
V6	Shehzadi	58.20	4.53
V7	Alpine 8510	61.00	4.60
V8	Alpine 8248	60.67	4.27
V9	Capsicum Orange	56.40	4.20
V10	Capsicum IMP	57.40	4.07
F - TEST		S	S
S.E.(d)		2.70	0.32
C.D at 5%		5.66	0.66
C.V.		5.08	9.36

Reproductive parameters

Data presented in Table 2 showed that Days to First Flower Initiation of Capsicum varied significantly under the influenced by different variety at harvest stage. Days to First Flower Initiation increased continuously with advancement of crop age and attained its maximum value at the maturity stage irrespective of variety. Maximum Days to First Flower Initiation

Comment [D25]: lower case

Comment [D26]: lower case

Comment [D27]: lower case

Comment [D28]: lower case. Follow similar pattern thereafter

(20.07), at harvest was recorded V₉ (Capsicum Orange), followed by V₆ (Shehzadi) and minimum Days to First Flower Initiation (9.13) was recorded under V₂ (Sweet Pepper Cebrail). This might be due to genetic constitution of the different hybrids that leads to differ in flower initiation. Similar findings were noticed by Rajender and Shukla (2013) and Ahmed *et al.* (2015) in capsicum. In case of Fruit Length increased continuously with advancement of crop age and attained its maximum value at the maturity stage irrespective of variety. Maximum Fruit Length (8.83), at harvest was recorded V₅ (Sweet Pepper Volante), followed by V₂(Sweet Pepper Cebrail) and minimum Fruit Length (4.90) was recorded under V₉ (Capsicum Orange). This could be due to higher uptake of nutrients and higher leaf area which enables higher build up of sufficient photosynthates enabling the increase in length of fruit in that hybrid. This is in concordance to the results of Biwalkeret *al.* (2015), Yellava and Patil (2009) in capsicum. Similarly in Fruit Diameter increased continuously with advancement of crop age and attained its maximum value at the maturity stage irrespective of variety. Maximum Fruit Diameter (6.40), at harvest was recorded V₅ (Sweet Pepper Volante), followed by V₃ (Sweet Pepper Bungi) and minimum Fruit Diameter (4.50) was recorded under V₁₀ (Capsicum IMP). The least Fruit Diameter (6.40) were recorded by V₅ (Sweet Pepper Volante). This could be attributed to the increased Fruit Diameter. These results are in conformity with Bhatt and Rao (1993) and Kurubetta (2008) in capsicum. However in Number of fruits per plant increased continuously with advancement of crop age and attained its maximum value at the maturity stage irrespective of variety. Maximum Number of fruits per plant (9.40), at harvest was recorded V₅ (Sweet Pepper Volante), followed by V₄ (Capsicum Varun) and minimum Number of fruits per plant (3.67) was recorded under V₁₀ (Capsicum IMP). Higher fruit yield plant-1 in V₅ (Sweet Pepper Volante) was due to more number of flowers and fruits plant-1, high average fruit weight and maximum sized fruits with higher values of fruit length and fruit width. This is in consonance with the findings of Zende (2008), Kurubetta and Patil (2009) and Aruna and Sudagar (2010) in capsicum. Also in Average fruit weight (g) increased continuously with advancement of crop age and attained its maximum value at the maturity stage irrespective of variety. Maximum Average fruit weight (g) (154.20), at harvest was recorded V₅ (Capsicum Varun), followed by V₁ (Sweet Pepper Nemalite) and minimum Average fruit weight (g) (138.27) was recorded under V₉ (Capsicum Orange). V₅ (Sweet Pepper Volante) recorded maximum Maximum Average fruit weight (g) (154.20) which was significantly superior over all the variety. The minimum Average fruit weight (g) (138.27) was recorded by V₉ (Capsicum Orange). This might be due to good vegetative growth besides effective pollination and fertilization and

Comment [D29]: in

Comment [D30]: in

Comment [D31]: in

Comment [D32]: remove

Comment [D33]: in

Comment [D34]: per plant

Comment [D35]: in

Comment [D36]: lower case

lower abscission rate of flowers. Similar results were recorded by Backer (1989) in sweet pepper. Among the Fruit yield per plant (kg) increased continuously with advancement of crop age and attained its maximum value at the maturity stage irrespective of variety. Maximum Fruit yield per plant (kg) (1.47), at harvest was recorded V₅ (Sweet Pepper Volante), followed by V₁ (Sweet Pepper Nermalite) and minimum Fruit yield per plant (kg) (0.29) was recorded under V₁₀ (Capsicum IMP). Similarly V₅ (Sweet Pepper Volante) recorded maximum number of fruits per plant (1.47) which was significantly superior over the other variety. The least number of fruits per plant (0.29) were recorded by V₁₀ (Capsicum IMP), might be due to more number of flowers and high value of per cent fruit set because of vigorous and healthy plants. These results are line with Backer (1989) in sweet pepper. Number of fruits per plant second harvest increased continuously with advancement of crop age and attained its maximum value at the maturity stage irrespective of variety. Maximum Number of fruits per plant second harvest (3.27), at harvest was recorded V₂ (Sweet Pepper Cebrail), followed by V₅ (Sweet Pepper Volante) and minimum Number of fruits per plant second harvest (0.47) was recorded under V₉ (Capsicum Orange). And Total yield increased continuously with advancement of crop age and attained its maximum value at the maturity stage irrespective of variety. Maximum Total yield (17.72), at harvest was recorded V₅ (Sweet Pepper Volante), followed by V₁ (Sweet Pepper Nermalite) and minimum Total yield (9.27) was recorded under V₁₀ (Capsicum IMP). Atmospheric moisture also plays a significant role in crop growth and development. The relative humidity increases the availability of net energy for crop growth and prolong the survival of crops under moisture stress conditions. Relative humidity reduces the evaporation losses from the plants which lead to optimum utilization of nutrients. It also maintains the turgidity of the cells, which is useful in enzyme activity leading to a higher yield (Reddy, 1999). Higher values of fruit length, fruit diameter and no. of fruit/plant contributed for maximum average fruit weight in the variety, V₅ (Sweet Pepper Volante). Similar observations were recorded by Valsikova and Belko (2004), Zende (2008) and Kurubetta and Patil (2009) in capsicum.

Comment [D37]: remove

Comment [D38]: in

Comment [D39]: in

Comment [D40]: which

Comment [D41]: in

Comment [D42]: lower case

Comment [D43]: in

Table 2. Reproductive parameters of Different Varieties in Capsicum varieties in naturally ventilated polyhouse.

Name of Varieties	Days to First Flower Initiation	Fruit Length in (cm)	Fruit Diameter (cm)	Number of fruits /plant	Average fruit weight (g)	Fruit yield /plant (kg)	No. of fruits /plant 2 nd harvest	Total yield (q)/200sq. m
V1	14.27	8.19	5.78	8.07	138.27	1.11	2.33	15.15

Comment [D44]: express in per hectare

V2	9.13	8.79	5.88	8.00	130.13	1.05	3.27	15
V3	15.87	6.14	6.35	7.87	96.27	0.74	2.53	12.75
V4	11.4	7.71	4.67	8.6	114.2	0.97	2.6	14.7
V5	9.67	8.83	6.4	9.4	154.2	1.47	3.13	17.72
V6	18.33	5.5	6.36	4.4	91	0.37	2	9.69
V7	11.8	5.65	4.59	4.6	88.47	0.39	1.33	9.91
V8	15.07	5.67	5.8	4.6	92.53	0.43	1.67	10.06
V9	20.07	4.9	4.56	4.53	87.13	0.4	0.47	10.09
V10	13.73	5.97	4.5	3.67	87.53	0.29	0.87	9.27
F - TEST	S	S	NS	S	S	S	S	S
S.E.(d)	0.57	0.29	0.85	0.45	6.27	0.09	0.77	1.052
C.D at 5%	1.19	0.61	1.79	0.95	13.16	0.18	1.63	2.172
C.V.	4.98	5.3	2.24	8.68	7.11	14.67	46.9	18.227

Quality parameters

Data presented in Table 3 showed that TSS ($^{\circ}$ brix) and Vitamin C (mg/100g) of Capsicum varied significantly under the influenced by different variety at harvest stage. TSS ($^{\circ}$ brix) and Vitamin C (mg/100g) increased continuously with advancement of crop age and attained its maximum value at the maturity stage irrespective of variety. Maximum TSS and Vitamin C (mg/100g) (obrix) (9.03 and 154.66), at harvest was recorded V₁ (Sweet Pepper Nemalite), followed by V₃ (Sweet Pepper Bungl) and minimum TSS (obrix) and Vitamin C (mg/100g) (7.23 and 56.40) was recorded under V₄ (Capsicum Varun).

Higher Vitamin-C content was recorded in V₄ (Capsicum Varun) (154.66 mg/100 g) variety which was superior over all variety evaluated. Generally, the higher ascorbic acid content would increase the nutritive value of capsicum, which would help better retention of colour and flavour. Capsicum varieties and hybrids possessing high ascorbic acid content are of great demand in export markets as opined by Sweta Rani (2003), Banaras *et al.* (2005) and Choudhary *et al.* (2011) in capsicum. Pevicharova *et al.* (2007) also recorded lower amount of vitamin-C content under higher temperature and stated that in bell pepper fruits high day temperature decreased the total vitamin-C of fruits. Higher TSS content was recorded in Inspiration (9.03 $^{\circ}$ Brix) which was found to be superior over all other treatments. Since capsicum is used for salad making, fruits with high TSS are highly preferred. TSS is an

important quality attribute of capsicum fruit which increases the palatability and also it determines the varieties for processing industries. Similar findings were recorded by Hajarica and Phookan (2005) in tomato and Sood *et al.* (2007) in capsicum.

Table 3 Quality parameters of Different Varieties of Capsicum in naturally ventilated polyhouse condition.

Varieties	NAME OF VARIETIES	TSS (°brix)	Vitamin C (mg/100g)
V1	Sweet Pepper Nemalite	9.03	154.10
V2	Sweet Pepper Cebrail	8.63	154.48
V3	Sweet Pepper Bungi	8.87	152.56
V4	Capsicum Varun	7.23	154.66
V5	Sweet Pepper Volante	8.00	154.05
V6	Shehzadi	8.60	152.00
V7	Alpine 8510	8.70	153.22
V8	Alpine 8248	8.43	152.03
V9	Capsicum Orange	8.43	152.15
V10	Capsicum IMP	7.90	152.56
F - TEST		S	NS
S.E.(d)		0.38	1.19
C.D at 5%		0.80	2.50
C.V.		5.55	0.95

Conclusion

The result from the present investigation concluded that in terms of Growth parameters Sweet pepper volante was recorded with maximum Plant height (72.87 cm), Number of branches (3.47), Days to first flower initiation (9.67). In terms of Yield parameters Sweet pepper volante was recorded with maximum fruit weight (154.20 g), Fruit length (8.83cm), Fruit

Comment [D45]: indicated

Comment [D46]: remove

yield per plant (1.47 kg), Fruit yield (17.72(q)/200sq.m.). In terms of Yield parameters Sweet Pepper Volante was recorded with maximum TSS (8.00), Vitamin C (154.05 mg).

References

Comment [D47]: follow journal guidelines

- Agarwal, A., Gupta, S. & Ahmed, Z. (2007). Influence of plant densities on productivity of bell pepper (*Capsicum annuum* L.) under greenhouse in high altitude cold desert of Ladakh. *ISHS Acta Horticulturae*, 756, 309-314. DOI: 10.17660/ActaHortic.2007.756.32
- Ahmed, M. A., Mahmoud and Ahmed, A.S., 2015, Production and evaluation of high yielding sweet pepper hybrids under greenhouse conditions. *J. Agric. Environ. Sci.*, 15 (4): 573-580.
- Baghel BS, Gupta M, Tiwari R. 2003. Potential and prospects of protective cultivation of horticultural crops. In: Proceedings of All India on Seminar Potential and Prospects for Protective Cultivation, Institute of Engineers, Ahmednagar.;389.
- Banaras, M., Bosland, P. W. and Lownds, N. K., 2005, Effects of harvest time and growth conditions on storage and post-storage quality of fresh peppers (*Capsicum annuum* L.). *Pakistan J. Biotech.*, 37 (2): 337-344.
- Bergeford, B. R., Lewis, W., Harker, T., Miller, L., Welch, A. and Weaks, E., 2011, Bell pepper cultivar performance trial grown in southern Ohio. www.southcenters.osu.edu.
- Bhatt, R. M. and Rao, N. K. S., 1993, Response of bell pepper to photosynthesis, growth, flower and fruit setting to night temperature. *Photosynthetica*, 28: 127-132.
- Biwalkar, N., Singh, K. G., Jain, A. K., Sharda, R., Jindal, S. K., Singh, K and Chawla, N. 2015. Response of coloured sweet pepper (*Capsicum annuum* L. var. *grossum*) to fertigation and irrigation levels under naturally ventilated greenhouse. *Agricultural Research Journal*. 52(1): 19-25.
- Bukasov, S. M. (1930). The cultivated plants of Mexico, Guatemala and Colombia. *Bull. Appl. Bot. Genet. and Plant Breed*, 553.
- CABI. (2019). *Capsicum annuum* (Bell pepper). Retrieved from <https://www.cabi.org/isc>.
- Choudhary, N., Pathania, N. K., Singh, S. P., Pardeep, K. and Yudhvir, S., 2011, Evaluation of red and yellow capsicum hybrids for quality attributes in naturally ventilated polyhouse in mid hills of western Himalayas. *Veg. Sci.*, 38: 218-220.
- Hajarica, T. K. and Phookan, D. B., 2005, Performance of tomato cultivars for polyhouse cultivation during spring summer in Assam. *Indian J. Hort.*, 62 (3): 268-271.

- Kumar, V., Pathania, N. K. & Kumar, N. (2015). Evaluation of bell pepper (*Capsicum annuum* L. Var. *GrossumSendt.*) genotypes for quality traits in modified naturally ventilated polyhouse. *Asian Journal of Plant Science and Research*, 5(4), 35-37.
- Kurubetta, Y., 2008, Performance of coloured capsicum hybrids under different protected structures. M.Sc. (Agri.) Thesis, Univ. Agric. Sci., Dharwad, Karnataka (India).
- Ochigbo, A. A. and Harris, G. P., 1989, Effect of film plastic cover on the growth and yield of bush tomato grown in a bed system. *J. Hort. Sci.*, 64 (1): 61-68.
- Pevicharova, G., Todorova, V. and Todorov, J., 2007, Ascorbic acid and total sugars content of kapyra type pepper depending on cultivars and climatic conditions. *Rasteniev-dni-Nauki*, 44 (1): 52-56.
- Rajender, S. and Shukla, Y. R., 2013, Performance of coloured bell pepper in naturally-ventilated polyhouse under mid-hill conditions of Himachal Pradesh. *J. Hort. Sci.*, 8 (2): 259-261.
- Reddy, S. R. (1999). *Principles of Agronomy*. Kalyani Publishers, New Delhi, pp. 25-28.
- Sharma, A., Kumar, M., Dogra, R. K., Kumar, N., Kumari, R. & Kansal, S. (2019). Estimation of genetic variability in bell pepper (*Capsicum annuum* L. var. *grossum*). *International Journal of Chemical Studies*, 7(3), 10-13.
- Sood, S, Anil Bindal and Akhilesh Sharma. 2007. Genetical study for quality traits in bell pepper (*Capsicum annuum* L. var. *grossumSendt.*). *Indian Journal of Genetics*. 67(1):95-97.
- Sweta Rani. 2003. Variability and path coefficient studies in bell pepper (*Capsicum annuum* L.) M.Sc thesis submitted to Dr. Yashwant Singh Parmar University of Horticulture and Forestry, Nauni, Solan.(H.P).
- Yellavva and Patil, A. A. 2009. Performance of coloured capsicum hybrids under different protected structures. *Karnataka Journal of Agricultural Sciences*. 22(5): 1058-1061.