

# Characterization of indigenous *Bacillus thuringiensis* isolates and toxicity analysis against Brinjal shoot and fruit borer, *Leucinodes orbonalis* (Guenee)

## ABSTRACT

The toxins derived from *Bacillus thuringiensis* (*Bt*) exhibit significant promise in managing specific orders of harmful insects without the drawbacks in using of chemical insecticides. This study examines 20 indigenous *Bt* isolates, with a focus on their colony and crystal morphology, gene content, protein profiles, and toxicity against the larvae of *Leucinodes orbonalis*. The results revealed that all colonies were creamy white colonies in colour with fried egg type appearance, flat surface, and undulated margin with spherical shaped crystals. PCR screening detected the existence of *cry1*, *cry2*, and *vip3A* genes with varying gene combinations among the isolates. Protein profiles exhibited the presence of multiple proteins with molecular weights ranging from 20 to 135 kDa, particularly ~135 and ~65 kDa represents Cry1 and Cry2 protein. In vitro bioassays against *L. orbonalis* revealed that four isolates T193, T339, T374, and T380 having substantial toxicity (93.33 to 100% mortality), suggesting the potential for their utilization in sustainable pest management strategies.

**Keywords:** *Bacillus thuringiensis*, insecticidal toxins, indigenous isolates, *Leucinodes orbonalis*, in vitro bioassay.

## 1. INTRODUCTION

Farmers across the world mostly depend on pesticides for pest management, but chemical control agents pose serious risks, including insect resistance, rise of secondary pest outbreak, harm to non-target organisms, environmental deterioration, and residues on crops and environment. Hence, there is a critical need for exploration of eco-friendly pesticides [1, 2, 3]. Biological control offers a potential alternate solution by utilizing microorganisms to manage pest species [4, 5]. *Bacillus* spp. stands out among the microorganisms employed for biological control, particularly in integrated pest management in agriculture,

particularly, *Bacillus thuringiensis* based formulations constitute a dominant share, comprising up to 90% of the market [6, 7, 8] among bio pesticides.

*Bacillus thuringiensis* (*Bt*) is a gram-positive soil bacterium found in diverse ecosystems, including soil, water, dead insect, and various plant ecosystems like phylloplanes and rhizosphere [9, 10]. ~~It has been~~ ~~Recognized-recognized~~ for its ability to produce crystalline inclusions containing insecticidal proteins called  $\delta$ -endotoxin. *Bt* strains produce crystal (Cry) and cytolytic (Cyt) toxins during sporulation phase [11, 12]. Furthermore, at vegetative phase, *Bt* isolates have the capability to generate additional insecticidal proteins, like vegetative insecticidal proteins (Vip) [10, 13, 14, 15] and secreted insecticidal proteins (Sip) [16, 17]. These cry proteins are effective in managing a diverse array of insect orders viz Lepidoptera, Coleoptera, Diptera, Hymenoptera, Homoptera, Orthoptera, and also exhibit toxicity towards nematodes [12, 18, 19, 20, 21, 22]. These cry toxins can be categorized based on amino acid homology, and the proteins with the same primary rank in phylogenetic tree, typically exhibit toxicity to specific insect orders [23]. Over 800 *Cry* genes are categorized into 75 families (from *Cry*1 to *Cry*75), while 40 *Cyt* genes are grouped into three families (*Cyt*1, *Cyt*2, and *Cyt*3). Additionally, 146 *Vip* genes are classified under four families (*Vip*1 to *Vip*4) [24]. While *Bt* proves to be an effective alternative to synthetic insecticides for controlling important insect pests, its prolonged use has the drawback of developing resistance in target insect populations in both laboratory and field studies [25, 26]. In order to combat these challenges, in the current study, toxicity of 20 indigenous *Bt* isolates were evaluated against *Leucinodes orbonalis* under laboratory conditions.

## 2. MATERIALS AND METHOD

### 2.1 Insect culture

Laboratory culture of *L. orbonalis* was initiated from the damaged field collected brinjal fruits. The mass culturing of Brinjal Shoot and Fruit Borer (BSFB) was done by following the method outlined by Visnupriya and Muthukrishnan [27] with minor modification. These collected fruits were placed in a tray filled with double autoclaved river sand and were maintained under controlled environmental conditions, including temperature ( $25 \pm 1$  °C), humidity ( $75 \pm 5\%$ ), and a photoperiod of 16: 8 hours (L:D). The larva pupates in the soil and emerged adults were provided a 10% sugar solution enriched with vitamin E and laid eggs in the black cloth. Newly hatched ~~larva-larvae~~ were transferred to potato tubers, containing 4 – 5 longitudinal slits and allowed to grow and pupate in sand. After being reared for two generations under laboratory condition, a stable insect population was utilized for the bioassay studies.

### 2.2 *Bt* isolates

Twenty native *Bt* isolates, Standard strain HD1 (*Btk*) and acrySTALLiferous strain 4Q7 were obtained from *Bt* Laboratory, Department of Plant Biotechnology, Centre for Plant Molecular Biology and Biotechnology, TNAU, Coimbatore, India. These bacterial cultures were revived from glycerol stock and maintained in T3

medium (For a litre of media: 2 g of tryptose, 3 g of tryptone, 1.5 g yeast extract, 6.9 g sodium dihydrogen phosphate, 8.9 g disodium hydrogen phosphate, 100 µl of manganese chloride (0.05 g in 1 ml of water), 20 g Agar, pH – 6.8 to 7.0).

### 2.3 Colony and crystal morphology of *Bt* strains

Colony characteristics like colour, type, elevation and margin of bacterial colonies were studied by visually observing a single isolated colony. To access the morphology, spore crystal inclusions were heatfixed on glass slides, stained with 0.133% Coomassie Brilliant Blue stain and examined under a bright field microscope at 100X magnification (EuromexiScope).

### 2.4 PCR screening for *cry* and *vip* genes

The extraction of genomic DNA from *Bt* strains was done, following the protocol of Kalman et al. [28]. The DNA concentration was measured using a Nano-Drop Spectrometer (Genova Nano, Jenway) and 0.8% agarose gel electrophoresis was employed to assess the quality of the DNA. PCR screening was done to confirm the presence of lepidopteran toxic genes, namely *cry1*, *cry2*, *cry9*, and *vip3A*. For carrying out PCR analysis, each reaction mixture consisted of 1 µl (50 ng) of template DNA, 1 µl of each primer (10 pmol), and 10 µl of 2X PCR Master Mix (Smartprime), comprising dNTPs, Taq polymerase, and 7 µl of sterile distilled water was used. The PCR temperature profile for each gene was maintained as specified in Table 1. Subsequently, the PCR products were separated in an agarose gel with EtBr (Ethidium bromide) staining and documented using a gel documentation system (Bio-Print imaging device, Vilber, France).

Table 1. Details of primers used and its temperature profile for PCR screening *Bt* isolates

S. No	Gene	Primer sequence	Product size	Temperature profile	Reference
1	<i>cry1</i>	FP: 5'-CATGATTCATGCGGCAGATAAAC-3' RP: 5'-TTGTGACACTTCTGCTTCCCATT-3'	~ 277 bp	94 °C for 2 min 94 °C for 40 s 62 °C for 40 s 72 °C for 1 min 72 °C for 7 min	Ben-Dov et al., 1997

Formatted: Highlight

Comment [A1]: Citation style(numbering) and please be aware that it has been included in your Reference section.

---

					30 cycles
2	<i>cry2</i>	FP: 5'-GTTATTCTTAATGCAGATGAATGGG-3' RP: 5'-CGGATAAAATAATCTGGGAAATAGT-3'	~ 700 bp	94 °C for 2 min 94 °C for 40 s 60 °C for 40 s 72 °C for 40 s 72 °C for 10 min 30 cycles	
3	<i>cry9</i>	FP: 5'-CGGTGTTACTATTAGCGAGGGCGG-3' RP: 5'-GTTGAGCCGCTTCACAGCAATCC-3'	~ 345 bp	59 °C for 40 s 61.5 °C for 40 s 72 °C for 1 min 72°C for 7 min 72°C for 1 min 25 cycles	
4	<i>vip3A</i>	FP: 5'-CCTCTATGTTGAGTGATGTA-3' RP: 5'-CTATACTCCGCTTCACTTGA-3	~ 1.0 Kb	94 °C for 5 min 94 °C for 1 min 55 °C for 1 min 72 °C for 40 s 72 °C for 10 min 35 cycles	Jain et al., 2012

---

**Formatted:** Highlight

**Comment [A2]:** Citation style(numbering) and please be aware that it has been included in your Reference section.

## 2.5 Spore-crystal mixture isolation and Protein profiling of *Bt* isolates

Spore crystal mixture of *Bt* isolates and standard strains were obtained using standard procedure [29]. From each *Bt* culture, a single colony was carefully picked using a sterile loop and introduced into test tubes containing 5 ml of T3 broth. Subsequently, these cultures underwent an overnight incubation at 30 °C with continuous shaking at 200 rpm (Orbitek, Scigenics Biotech). Following this, a 1% inoculum (250 µl) from the overnight cultures was transferred into T3 broth (25 ml), and then incubated for 48 hours at 30 °C and 200 rpm. After observing 90% of the cells lysis at 48-hour period, the culture was subjected to centrifugation at 7500 rpm for 15minutes (at 4°C). The resulting pellet was then resuspended in 500 µl of

Sterile Water with 10 µl of 1mM phenylmethyl sulfonyl fluoride (PMSF) and stored at -20°C for subsequent use. Protein profiling of *Bt* isolates was done by using SDS-PAGE (Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis), involving 10% separating gel and 4% stacking gel [30].

## 2.6 Bioassay

With Bradford assay [31], to quantify the protein concentration in spore crystal mixture was quantified using Bovine Serum Albumin (BSA) as a standard. The insect bioassay was conducted against neonates of *L. orbonalis* through surface coating method at 25 µg/ml concentration [32] using potato. Ten first-instar larvae (pre-starved for 8 hours) were released onto the potato without any physical damage. The standard strain HD1 and water served as a positive and negative controls respectively. Each treatment was replicated thrice and observation on larval mortality were recorded upto 72hrs using a stereo zoom microscope (Labomed stereoscopic microscope CSM2, California, United States) and expressed in percentage.

## 2.7 Statistical analysis

The laboratory experiment followed a completely randomized design (CRD). Using Abbott's formula [33], larval mortality was corrected for control mortality, and ~~followed by conducting~~ one-way analysis of variance (ANOVA) was done using R software (~~version 4.3.1~~) [Version 4.3.1 \[34\]](#). Subsequently, significant difference between means were identified through Tukey HSD test.

## 3. RESULTS AND DISCUSSION

### 3.1 Colony and crystal morphology of *Bt* strains

The 20 screened *Bt* isolates were uniformly creamy white in color. Among them, 17 isolates exhibited fried egg type colony and 3 isolates showed mucoid colonies. In terms of colony shape and margin, 12 isolates showed circular colony with complete margin, whereas 8 isolates had irregular colony with undulated margin. Colony elevation of 15 isolates were flat while 5 isolates exhibited raised colonies (Table 2). Different types of crystal inclusions were observed among the isolates viz., spherical (10), bipyramidal (7), cuboidal (6), and rectangular (2). These findings were in accordance with the results of Navya et al. [32]. Rashki et al. [\[3435\]](#) reported varying crystal shapes viz., bipyramidal, spherical, cubical, irregular, and elliptical, of which spherical shape (73.33%) was the most abundant among the *Bt* isolates of Iran.

**Table 2. Colony morphology of *Bt* isolates**

Colour	Type	Elevation	Margin	Shape	Bt isolates	Occurrence	
						Nos.	%
Creamy	Fried egg	Flat	Complete	Circular	T52, T194, T195	3	15.0

white			Irregular	T46, T380	2	10.0
		Undulated	Circular	T193, T197, T374	3	15.0
			Irregular	T100, T159, T196, T338, T339	5	25.0
	Raised	Complete	Circular	T51, T198, T199	3	15.0
			Irregular	T54	1	5.0
Mucoid	Flat	Complete	Circular	T45, T53	2	10.0
	Raised	Complete	Circular	T158	1	5.0

### 3.2 PCR screening of *Bt* isolates

Examination of 20 *Bt* isolates indicated that five isolates possessed only *cry1* gene, three isolates harbored both *cry1* and *cry2* genes, and two isolates possessed *cry1*, *cry2*, and *vip3A* genes. No amplification was observed for *cry9* gene in any of the isolates. (Table 3). The coexistence of *cry* and *vip* genes in *Bt* isolates was formerly reported by Navya et al. [33] and Gothandaraman et al. [3536]. Lone et al. [3637] confirmed the existence of both *cry1* and *cry2* gene in 12 out of 44 isolates screened. Karuppaiyan et al. [3738] and Maheesha et al. [3839] reported the occurrence of *cry1A* (*cry1Ab*, *cry1Ac*), *cry2* (*cry2Aa*, *cry2Ab*) and *vip3A* genes in the native *Bt* isolates using PCR analysis. *Cry1* was the most common gene among the 20 isolates screened followed by *cry2*, *vip3A* genes. Previously, *cry1* type genes were reported as predominant gene, present in indigenous *Bt* strains [3940, 4041, 4442].

**Table 3. Distribution of *Cry* and *Vip* genes among indigenous *Bt* isolates**

Bt isolates	Genes			
	<i>cry1</i>	<i>cry2</i>	<i>cry9</i>	<i>vip3A</i>
T45	-	-	-	-

T46	-	-	-	-
T51	-	-	-	-
T52	-	-	-	-
T53	-	-	-	-
T54	-	-	-	-
T100	-	-	-	-
T158	-	-	-	-
T159	-	-	-	-
T193	+	-	-	-
T194	+	-	-	-
T195	+	-	-	-
T196	+	-	-	-
T197	+	-	-	-
T198	-	-	-	-
T199	-	-	-	-
T338	-	-	-	-
T339	+	+	-	-
T374	+	+	-	+
T380	+	+	-	+
HD1	+	+	-	+
4AT1	-	-	+	-
4Q7	-	-	-	-

HD1 – positive standard strain for cry1, cry2 and vip3A; 4AT1 – standard strain for cry9 only; 4Q7 – negative strain for all genes. (+) = presence and (-) = absence of respective genes

### 3.3 Sodium dodecyl sulphate–polyacrylamide gel electrophoresis (SDS-PAGE) analysis

The insecticidal crystal proteins of the *Bt* isolates varied in their molecular weight between 20 and 135 kDa (Table 4). Eight isolates showed protein bands of ~ 135 kDa size (Cry1) and three isolates with protein size of ~ 65 kDa (Cry2), respectively. Vip3 protein of size ~ 88 kDa was observed in two isolates

(Fig 1). Protein profiles were comprising of varying no. of bands: a single band in 1 isolate, two bands in 2 isolates, and more than 2 bands in 16 isolates, while one isolate did not exhibit any distinct band. Hassan et al. [26] reported multiple protein bands of 20 to 130kDa size, among *Bt* strains. Of the 70 *Bt* strains that were screened through SDS-PAGE analysis, Ramalakshmi and Udayasuriyan [29] found that 17 strains (24.2%) showed two prominent protein bands, which corresponded to Cry1 and Cry2 protein, with molecular weights ~ 135 and 65 kDa. Sujayanand et al. [4243] reported protein bands of various sizes viz., 175, 135, 65, 51, 35, and 21 kDa; including 135 and 65 kDa size bands of Cry1 and Cry2 protein. Similarly, Kaviyapriya et al. [4344], Karuppaiyan et al. [3738], Maheesha et al. [3839], Reyaz et al. [4445] reported the Cry1 and Cry2 proteins of 135 and 65 kDa size. Lone et al. [4546] revealed that Vip3A protein produce bands at approximately 88 kDa in size.

**Table 4. Various protein bands produced by *Bt* isolates**

Protein size	Occurrence (n= 20)	
	Nos.	%
~ 135 kDa	8	40.0
~ 100 kDa	3	15.0
~ 91 kDa	2	10.0
~ 88 kDa	4	20.0
~ 80 kDa	1	5.0
~ 71 kDa	2	10.0
~ 65 kDa	7	35.0
~ 54 kDa	7	35.0
~ 52 kDa	5	25.0
~ 50 kDa	10	50.0
~ 43 kDa	3	15.0
~ 40 kDa	4	20.0
~ 33 kDa	1	5.0
~ 30 kDa	6	30.0
~ 27 kDa	13	65.0
~ 25 kDa	2	10.0



**Table 5. Percent mortality of *L. orbonalis* caused by indigenous *Bt* isolates (@ 25 µg/ml)**

<b>S. No.</b>	<b>Bt isolates</b>	<b>% Mortality at 72 HAT</b>
1	T45	10.0 (18.43) <sup>fg</sup>
2	T46	3.33 (10.51) <sup>g</sup>
3	T51	6.66 (14.96) <sup>g</sup>
4	T52	0.00 (0.91) <sup>g</sup>
5	T53	10.0 (18.43) <sup>fg</sup>
6	T54	0.00 (0.91) <sup>g</sup>
7	T100	0.00 (0.91) <sup>g</sup>
8	T158	0.00 (0.91) <sup>g</sup>
9	T159	3.33 (10.51) <sup>g</sup>
10	T193	93.33 (75.03) <sup>abc</sup>
11	T194	66.66 (54.73) <sup>cd</sup>
12	T195	80.0 (63.43) <sup>bc</sup>
13	T196	80.0 (63.43) <sup>bc</sup>
14	T197	76.66 (61.11) <sup>bc</sup>
15	T198	33.33 (35.26) <sup>def</sup>
16	T199	30.0 (33.21) <sup>ef</sup>
17	T338	40.0 (39.23) <sup>de</sup>
18	T339	93.33 (75.03) <sup>abc</sup>
19	T374	100 (89.09) <sup>a</sup>
20	T380	100 (89.09) <sup>a</sup>
21	HD1	100 (89.09) <sup>a</sup>
22	Control	0.00 (0.91) <sup>g</sup>

\*HAT – Hours After Treatment

Figures in parentheses are arcsine transformed values of percentages. Values followed by the same letters in a column are not significantly different by the Tukey Honestly Significant Difference (HSD) test with  $\alpha = 0.05$  ( $F = 79.04$ ;  $df = 21, 44$ ;  $P < 0.001$ )

#### 4. CONCLUSION

Four out of 20 *Bt* isolates exhibited higher toxicity (93.33 – 100%) against *L. orbonalis*. Identification of toxic genes cry1 and cry2 in these *Bt* isolates can be utilized in development of transgenic plants. These effective isolates can ~~additionally exploited~~ also be used for formulation studies. Overall, these findings underscore the multifaceted nature of *Bt* isolates and their promising role as an effective biopesticides for insect control.

#### REFERENCES

1. Shelton AM, Zhao JZ, Roush RT. Economic, ecological, food safety, and social consequences of the deployment of *Bt* transgenic plants. *Annu Rev Entomol.* 2002;47(1):845-81.
2. Muhammad W, Ahmad I, Bhatti HT, Zubair M. Linking pesticide exposure with covid-19 among agricultural professionals in perspectives of immunity and safety: a review. *Plant Prot.* 2020;4(2).
3. Sunitha P, Kalia VK. Insecticidal activity of native *Bacillus* species against brinjal shoot and fruit Borer, *Leucinodes orbonalis* (Lepidoptera: Crambidae). *Pest manage. hortic. ecsyst.* 2021;27(2):240-5.
4. Sarkar D, Rakshit A, Al-Turki AI, Sayyed RZ, Datta R. Connecting bio-priming approach with integrated nutrient management for improved nutrient use efficiency in crop species. *Agriculture.* 2021;11(4):372.
5. Abdullah S, Zahoor I. Biopesticides: A Green Substitute to Chemical Pesticide. *Int. j. chem. biochem. Sci.* 2023;24 (4):141-156.
6. Ortiz A, Sansinenea E. Genetically Modified Plants Based on *Bacillus* Genes and Commercial *Bacillus*-Based Biopesticides for Sustainable Agriculture. *Horticultrae.* 2023;9(9):963.
7. Kaur S. Molecular approaches towards development of novel *Bacillus thuringiensis* biopesticides. *World J. Microbiol. Biotechnol.* 2000;16: 781-93.
8. Lacey LA, Frutos R, Kaya HK, Vail P. Insect pathogens as biological control agents: do they have a future?. *Biol. control.* 2001;21(3):230-48.
9. Roh JY, Choi JY, Li MS, Jin BR, Je YH. *Bacillus thuringiensis* as a specific, safe, and effective tool for insect pest control. *J. Microbiol. Biotechnol.* 2007;17(4):547-59.
10. Gupta M, Kumar H, Kaur S. Vegetative insecticidal protein (Vip): A potential contender from *Bacillus thuringiensis* for efficient management of various

- detrimental agricultural pests. *Front. Microbiol.* 2021;12:659736. <https://doi.org/10.3389/fmicb.2021.659736>
11. Palma L, Muñoz D, Berry C, Murillo J, Caballero P. *Bacillus thuringiensis* toxins: an overview of their biocidal activity. *Toxins.* 2014;6(12):3296-3325. <https://doi.org/10.3390/toxins6123296>
  12. Schnepf E, Crickmore N, Van Rie J, Lereclus D, Baum J, Feitelson J, Zeigler DR, Dean D. *Bacillus thuringiensis* and its pesticidal crystal proteins. *Microbiol. Mol. Biol. Rev.* 1998;62(3):775-806. <https://doi.org/10.1128/mmlr.62.3.775-806.1998>
  13. Estruch JJ, Warren GW, Mullins MA, Nye GJ, Craig JA, Koziel MG. Vip3A, a novel *Bacillus thuringiensis* vegetative insecticidal protein with a wide spectrum of activities against lepidopteran insects. *Proceedings of the National Academy of Sciences.* 1996;93(11):5389-94.
  14. Warren G. Vegetative insecticidal proteins: novel proteins for control of corn pests. In: Carozzi N, Koziel M, editors. *Advances in Insect Control: The Role of Transgenic Plants.* 1<sup>st</sup> ed. London: Taylor & Francis; 1997:109-21.
  15. Chakroun M, Banyuls N, Bel Y, Escriche B, Ferré J. Bacterial vegetative insecticidal proteins (Vip) from entomopathogenic bacteria. *Microbiol. Mol. Biol. Rev.* 2016;80(2):329-50.
  16. De Maagd RA, Bravo A, Berry C, Crickmore N, Schnepf HE. Structure, diversity, and evolution of protein toxins from spore-forming entomopathogenic bacteria. *Annu. Rev. Genet.* 2003;37(1):409-33.
  17. Donovan WP, Engleman JT, Donovan JC, Baum JA, Bunkers GJ, Chi DJ, Clinton WP, English L, Heck GR, Ilagan OM, Krasomil-Osterfeld KC. Discovery and characterization of Sip1A: A novel secreted protein from *Bacillus thuringiensis* with activity against coleopteran larvae. *Appl. Microbiol. Biotechnol.* 2006;72:713-9. <https://doi.org/10.1007/s00253-006-0332-7>
  18. Crickmore N, Zeigler DR, Feitelson J, Schnepf ES, Van Rie J, Lereclus D, Baum J, Dean D. Revision of the nomenclature for the *Bacillus thuringiensis* pesticidal crystal proteins. *Microbiol. Mol. Biol. Rev.* 1998;62(3):807-13. <https://doi.org/10.1128/mmlr.62.3.807-813.1998>
  19. De Maagd RA, Bravo A, Crickmore N. How *Bacillus thuringiensis* has evolved specific toxins to colonize the insect world. *Trends Genet.* 2001;17(4):193-9.
  20. Wei JZ, Hale K, Carta L, Platzer E, Wong C, Fang SC, Aroian RV. *Bacillus thuringiensis* crystal proteins that target nematodes. *Proceedings of the National Academy of Sciences.* 2003;100(5):2760-5. <https://doi.org/10.1073/pnas.0538072100>
  21. Ye W, Zhu L, Liu Y, Crickmore N, Peng D, Ruan L, Sun M. Mining new crystal protein genes from *Bacillus thuringiensis* on the basis of mixed plasmid-enriched genome sequencing and a computational pipeline. *Appl. Environ. Microbiol.* 2012;78(14):4795-801. <https://doi.org/10.1128/AEM.00340-12>

22. van Frankenhuyzen K. Cross-order and cross-phylum activity of *Bacillus thuringiensis* pesticidal proteins. *J. Invertebr. Pathol.* 2013;114(1):76-85. <https://doi.org/10.1016/j.jip.2013.05.010>
23. George Z, Crickmore N. *Bacillus thuringiensis* applications in agriculture. In: Estibaliz Sansinenea editor. *Bacillus thuringiensis* biotechnology. Dordrecht, Heidelberg, London: Springer;2012:19-39. [https://doi.org/10.1007/978-94-007-3021-2\\_2](https://doi.org/10.1007/978-94-007-3021-2_2)
24. Crickmore N, Berry C, Panneerselvam S, Mishra R, Connor TR, Bonning BC. A structure-based nomenclature for *Bacillus thuringiensis* and other bacteria-derived pesticidal proteins. *J. Invertebr. Pathol.* 2021;186:107438. <https://doi.org/10.1016/j.jip.2020.107438>
25. Jurat-Fuentes JL, Heckel DG, Ferré J. Mechanisms of resistance to insecticidal proteins from *Bacillus thuringiensis*. *Annu. Rev. Entomol.* 2021;66:121-40. <https://doi.org/10.1146/annurev-ento-052620-073348>
26. Hassan AA, Youssef MA, Elashtokhy MM, Ismail IM, Aldayel M, Afkar E. Isolation and identification of *Bacillus thuringiensis* strains native of the Eastern Province of Saudi Arabia. *Egypt. J. Biol. Pest Control.* 2021;31(1):1-1. <https://doi.org/10.1186/s41938-021-00369-7>
27. Visnupriya M, Muthukrishnan N. Negative Cross Resistance of *Leucinodesorbonalis* Population of Brinjal to Newer Molecule Spinetoram 12 SC W/W (11.7 W/W). *Int. J. Curr. Microbiol. App. Sci.* 2017;6(12):1790-6. <https://doi.org/10.20546/ijcmas.2017.612.202>
28. Kalman S, Kiehne KL, Libs JL, Yamamoto T. Cloning of a novel cryIC-type gene from a strain of *Bacillus thuringiensis* subsp. *galleriae*. *Appl. Environ. Microbiol.* 1993;59(4):1131-7. <https://doi.org/10.1128/aem.59.4.1131-1137.1993>
29. Ramalakshmi A, Udayasuriyan V. Diversity of *Bacillus thuringiensis* isolated from western ghats of Tamil Nadu state, India. *Curr. Microbiol.* 2010;61:13-18. <https://doi.org/10.1007/s00284-009-9569-6>
30. Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature.* 1970;227(5259):680-5.
31. Bradford, M.M., A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* 1976;72:248-54.
32. Navya RN, Balasubramani V, Raveendran M, Murugan M, Lakshmanan A. Diversity of indigenous *Bacillus thuringiensis* isolates toxic to the diamondback moth, *Plutellaxylostella* (L.) (Plutellidae: Lepidoptera). *Egypt. J. Biol. Pest Control.* 2021;31:1-7. <https://doi.org/10.1186/s41938-021-00495-2>
33. Abbott, WS. A method of computing the effectiveness of an insecticide. *J. Econ. Entomol.* 1925;18(2):265-267. <https://doi.org/10.3109/13880209.2012.674950>

- 33-34. [R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing \(2023\), Vienna, Austria. URL: <https://www.R-project.org/>.](#)
- 34-35. Rashki M, Maleki M, Torkzadeh-Mahani M, Shakeri S, Nezhad PS. Isolation of Iranian *Bacillus thuringiensis* strains and characterization of lepidopteran-active cry genes. Egypt. J. Biol. Pest Control. 2021;31:1-0. <https://doi.org/10.1186/s41938-021-00432-3>
- 35-36. Gothandaraman R, Venkatasamy B, Thangavel T, Eswaran K, Subbarayalu M. Molecular characterization and toxicity evaluation of indigenous *Bacillus thuringiensis* isolates against key lepidopteran insect pests. Egypt. J. Biol. Pest Control. 2022;32(1):1-1. <https://doi.org/10.1186/s41938-022-00639-y>
- 36-37. Lone SA, Malik A, Padaria JC. Characterization of lepidopteran-specific cry1 and cry2 gene harbouring native *Bacillus thuringiensis* isolates toxic against *Helicoverpa armigera*. Biotechnol. Rep. 2017;15:27-32. <https://doi.org/10.1016/j.btre.2017.05.001>
- 37-38. Karuppaiyan T, Balasubramani V, Murugan M, Raveendran M, Rajadurai G, Kokiladevi E. Characterization and evaluation of indigenous *Bacillus thuringiensis* isolate T352 against fall armyworm, *Spodoptera frugiperda* (JE Smith). Int. J. Plant Soil Sci. 2022;34(21):729-36. <https://doi.org/10.9734/IJPSS/2022/v34i2131325>
- 38-39. Maheesha M, Balasubramani V, Murugan M, Raveendran M, Rajadurai G, Tamilnayagan T, Kokiladevi E, Sathiah N. Characterisation of native *Bacillus thuringiensis* isolates toxicity to fall armyworm, *Spodoptera frugiperda* (JE Smith). J. Biol. Control. 2021;35(3):171-80. <https://doi.org/10.18311/jbc/2021/28812>
- 39-40. Jain D, Sunda SD, Sanadhya S, Nath DJ, Khandelwal SK. Molecular characterization and PCR-based screening of cry genes from *Bacillus thuringiensis* strains. 3 Biotech. 2017;7:1-8. <https://doi.org/10.1007/s13205-016-0583-7>
- 40-41. Salama HS, Abd El-Ghany NM, Saker MM. Diversity of *Bacillus thuringiensis* isolates from Egyptian soils as shown by molecular characterization. J. Genet. Eng. Biotechnol. 2015;13(2):101-9. <https://doi.org/10.1016/j.jgeb.2015.10.001>
- 41-42. Ramalakshmi A, Annakodi P, Udayasurian V, Balasubramani V. Toxicity Analysis and cry Gene Profiling of *Bacillus thuringiensis* Isolated from Western Ghats of Tamil Nadu State, India. In Proc Indian Natn Sci Acad. 2018;84(3):723-9. <https://doi.org/10.16943/ptinsa/2018/49400>
- 42-43. Sujayanand GK, Akram M, Konda A, Nigam A, Bhat S, Dubey J, Kumar K, Muthusamy SK. Distribution and toxicity of *Bacillus thuringiensis* (Berliner) strains from different crop rhizosphere in Indo-Gangetic plains against

polyphagous lepidopteran pests. Int. J. Trop. Insect Sci. 2021;1-9.  
<https://doi.org/10.1007/s42690-021-00451-5>

- 43-44. Kaviyapriya M, Lone R, Balakrishnan N, Rajesh S, Ramalakshmi A. Cloning and characterization of insecticidal cry/vip genes from an indigenous *Bacillus thuringiensis* isolate T29 and evaluation of its toxicity to maize fall armyworm *Spodoptera frugiperda*. J. Entomol. Zool. Stud. 2019;7(3):1314-21.
- 44-45. Reyaz AL, Gunapriya L, Indra Arulselvi P. Molecular characterization of indigenous *Bacillus thuringiensis* strains isolated from Kashmir valley. 3 Biotech. 2017;7:1-1. <https://doi.org/10.1007/s13205-017-0756-z>
- 45-46. Lone SA, Yadav R, Malik A, Padaria JC. Molecular and insecticidal characterization of Vip3A protein producing *Bacillus thuringiensis* strains toxic against *Helicoverpa armigera* (Lepidoptera: Noctuidae). Can. J. Microbiol. 2016;62(2):179-90. <https://doi.org/10.1139/cjm-2015-032>
- 46-47. Satpathy S, Rai AB, Singh M, Kumar A, Sharma BK, Rai M. Susceptibility of Two Populations of *Leucinodes orbonalis* Guen. to Cry1 AC Toxin. Veg. Sci. 2009;36(3s):349-52.
- 47-48. Reyaz AL, Balakrishnan N, Udayasuriyan V. Genome sequencing of *Bacillus thuringiensis* isolate T414 toxic to pink bollworm (*Pectinophora gossypiella* Saunders) and its insecticidal genes. Microb. Pathog. 2019;134:103553. <https://doi.org/10.1016/j.micpath.2019.103553>