

PRANAM-Ca: An Eggshell derived Calcium fertilizer and its impact on blossom end rot in Tomato cultivation

Abstract:

Eggshells, amounting to a staggering 76.7 million tons annually, are often inadequately disposed of in landfills, leading to significant nutrient loss for both plants and humans, while contributing to environmental pollution and emitting unpleasant odors. In response to these challenges, recent research has been dedicated to repurposing eggshells, exploring their applications in diverse fields such as biochemistry, biomedicine, environmental technology, and agriculture. Recognized as an exceptional source of calcium, eggshells offer notable benefits for calcium-demanding crops, including tomatoes, chilli and cotton. In a comprehensive study conducted, we investigated the application of organic foliar calcium derived from eggshells on tomato plants. The results demonstrated substantial improvements in addressing physiological disorders, promoting overall growth, and enhancing yield, signifies the importance of calcium fertilizer for growing healthy crop. With this background, Multiplex R&D team has made an effort to develop an in-house technology to derive the calcium from waste eggshell. Afterwards, how to scale-up and commercialise it as a sprayable form of Organic calcium fertilizer, PRANAM-CA. Present paper aims to describe the technique to derive calcium from eggshell and its impact on managing the blossom end rot in Tomato cultivation under protected glass-house condition.

Keywords: Egg shell derive Calcium (ESDC), Blossom end rot (BER), Tomato cultivation, Foliar spray, Calcium chloride, Calcium carbonate

Introduction:

Eggs are one of the most widely consumed foods worldwide, making the egg production business a critical component of the global food market (Anon, 2023). “The volume of global egg production exceeded 86.3 million metric tons in 2021, down from about 87.07 million metric tons in 2020. Since 1990, the volume of global egg production has increased by over 100 per cent” (Shahbandeh, 2023). “Worldwide, China is the highest producer of eggs (462 billion eggs annually) followed by United states (99 billion eggs annually) and India (93 Billion eggs)” (Anon, 2023). “An eggshell is the outer covering of a hard-shelled egg and of some forms of eggs with soft outer coats” (Hunton, P, 2005). “The composition of good quality

eggshell contains approximately 2.2 grams of calcium in the form of calcium carbonate. About 95 per cent of the dry eggshell is calcium carbonate weighing 5.5 gram. Along with this, it also contains 0.3 per cent of phosphorus and magnesium. The traces of sodium, potassium, zinc, manganese, iron and copper are also present in egg shell” (Butcher and miles, 2019). It is mostly discarded as a landfill or burned that contributes to different types of pollution like soil, water, air (Kattimani *et al.*, 2022) and also incur considerable disposal costs in the world (Verma *et al.*, 2012). “The disposal of eggshells not only provide site for flies and abrasiveness, but also contributes to the loss of many materials” (Hamester *et al.*, 2012). “Eggshell waste has been ranked as the 15th major food industry pollution problem by the Environmental Protection Agency” (Ajala *et al.*, 2018; Faridi and Arabhosseini, 2018).

Hence eggshell waste from industrial processes can be used for agriculture, as an agent to increase soil pH (Vandeginste, 2021; Faridi and Arabhosseini, 2018), as a fertilizer and it can also be processed into more useful products such as animal feeds, medicines, and building materials (Ajala *et al.*, 2018). “The use of these shells as an alternative source of CaCO_3 (calcium carbonate) may reduce the impact on the natural reserves of limestone, a non-renewable natural source” (Neves 1998; Boron 2004 cited in Faridi and Arabhosseini, 2018).

Calcium (Ca) deficiency in plants can be caused by insufficient level of available calcium in the growing medium (Khan *et al.*, 2017), which indirectly leads to a major physiological disorder like BER in tomatoes etc.

Tomato (*Solanum lycopersicum* L.) is a key member of the Solanaceae family, renowned as one of the world's most extensively cultivated and economically significant vegetables. Calcium plays a crucial role in the normal growth and development of plants by maintaining membrane structures, enhancing nutrient uptake, and activating metabolic processes. (Tuna *et al.*, 2007; Sarwat *et al.*, 2013). In addition, the Ca is needed to maintain cell wall integrity and to ensure bindings between cells (Marschner, 1995). Calcium also reduces detrimental effects of stress by regulating antioxidant metabolism (Zorrig *et al.*, 2012; Ahmad *et al.*, 2015). Several factors are involved in the incidence of blossom-end rot (BER) in tomato fruit, but the main one is insufficient Ca uptake and transport through the plant, resulting in Ca deficiency in the fruits. Sprays of Ca-containing products are considered to be a possible measure to overcome the local Ca deficiency in tomato fruit.

Blossom-end rot (BER) is a common nutritional disorder of tomato, pepper, eggplant, pumpkin, squash and watermelon that is caused by a shortage of calcium in enlarging fruits.

This disorder typically occurs when plants are growing rapidly and the first fruits are developing. As fruit cells breakdown due to a lack of calcium, dark blemishes appear on the blossom-end of affected fruits. These may enlarge until the entire bottom of the fruit becomes dark, shrunken and leathery. Factors that encourage blossom-end rot include: low soil pH and low levels of calcium, inconsistent watering, shallow watering or droughty conditions, and excessive use of nitrogen fertilizers (Khan *et al.*, 2017).

Many studies have demonstrated that Ca deficiency have initiated BER incidence in the tomato fruits (Shear 1975; Ho *et al.*, 2015; Taylor and Locascio 2004). Calcium is an important nutrient that plays a key role in the structure of cell walls and cell membranes, fruit growth and development, as well as general fruit quality (Usten *et al.*, 2006). Foliar application of fertilizer stands out as the most effective method for enhancing the nutritional status of plants (Shabbir *et al.*, 2015). The aim of this greenhouse study was to investigate the effects of organic foliar calcium application in compared to inorganic calcium chloride (CaCl₂) and calcium nitrate (Ca(NO₃)₂) form of application on yield and quality of tomato.

Material and Methods

A pot experiment was conducted in commercial greenhouse, during the season of 2021/2022 at Research & Development unit, Peenya, Bangalore, Karnataka. Greenhouse temperature was maintained at 26 °C ± 4 with 80 per cent of moisture. Soil mixture was prepared by mixing 25 per cent compost in 75 per cent sandy loam soil and 8 kg of soil mixture at 18 per cent moisture in a bags of size 12 x 12-inch. The nature of the compost was coco peat.

The pot mixture of the experiment was sterilised before use for the study. The experiment was laid out in completely randomized design with four treatments and five replications. Treatment includes 0.5 per cent concentration of calcium chloride, calcium nitrate and egg shell derived calcium which were sprayed three times at an interval 20 days. One-month old tomato (Cherry tomato variety sun gold) seedlings were transplanted to the pots. The concentrations of calcium chloride, calcium nitrate and egg shell derived calcium foliar solutions were fixed based on their pH value in water, to avoid adverse effect of acidic pH on plants, 0.5 per cent was chosen. Industrial grade nitric acid of 70% concentration and Calcium carbonate derived from eggshell of 95% concentration were used for the study. The calcium content in eggshell is 38-40%.

Table 1: Details of preparation

Sl.No.	Treatment details	Dosage (%)
1	T1- Calcium Chloride	0.5
2	T2- Calcium Nitrate	0.5
3	T3- Egg shell derived Calcium	0.5
4	T4-Control	0

2.1 Ca Preparations and Plant Treatments:

To test the possibilities to increase Ca content in tomato fruits, treatments with 3 different sources of Ca along with **PRANAM Ca** (A commercially available Product by Multiplex, used for this study) were applied. The method of preparations, application rates and intervals were as follows.

Specification of PRANAM-Ca: Calcium percentage:11-15%, pH :5.0-7.0, Density at 28-30 °C:1.55-1.66 kg/m², Specific gravity:1.73-1.82, Physical appearance: reddish brown. The percentage of Ca in Ca(NO₃)₂ is 18% and N is 15% and Ca in CaCl₂ is 27% and Cl₂ is 50% since nitrogen is having growth impact on plant it has showed good results compared to chloride which has no significant role in plant growth. In PRANAM Ca the concentration of Ca is 15% and 10% N.

Preparation of PRANAM Ca:

The chicken and duck egg shells were selected due to their inherent variations in calcium concentrations. The shells, procured from diverse restaurants and hatcheries, underwent a systematic processing regimen. Initial cleaning involved washing the shells in water to eliminate impurities, followed by a meticulous drying process in a hot air oven. This step aimed to remove excess moisture and pre-emptively address concerns related to rotting and unpleasant odors associated with the yolky part. Drying was carried out at 105 °C for 48 hr. It's a mixture of 90% of chicken egg shell and remaining 10% from unknown bird egg shells.

Acid hydrolysis:

Given that calcium carbonate constitutes a substantial portion of egg shells and possesses a weak molecular bond, a strategic approach involving chemical treatment was employed. Nitric acid was introduced to react with the carbonate bond, facilitating the

conversion of calcium carbonate into calcium nitrate, a water-soluble compound. The acid-induced reaction was allowed to transpire over a known duration, resulting in the generation of a dense, dark reddish-brown liquid. Subsequently, filtration using Watt Mann filter paper was employed to isolate the liquid from residual particulate matter.

To quantify the concentration of the water-soluble calcium solution, the specific gravity of the liquid was determined using the formula $\text{Density} = \text{Mass}/\text{Volume}$, yielding an observed value of 1.62 g/ml. Notably, the water-soluble calcium content in the liquid was established at 15 per cent. This scientifically rigorous methodology ensures the production of a potent and consistent source of calcium suitable for application in agricultural settings, particularly for augmenting the calcium uptake by tomato plants. The flow chart for the same is displayed below (Fig.1).




Fig.1: Preparation of PRANAM Ca

3. Results and Discussion:

The foliar spray treatments exhibited a positive impact on the fruit yield, height and physiological parameters of the tomato. Analysis revealed a noticeable increase in the overall mass of fruits, indicating a positive correlation between calcium supplementation and enhanced fruit development. The treatment T2- Calcium Nitrate 0.5% recorded maximum fruit yield (402 g/plant) followed by T3- Egg shell derived Calcium 0.5% (389 g/plant), whereas, the lowest fruit yield was observed in the plants which were left untreated (252.4 g/plant) as referred in Table 2. The incidence of blossom end rot, a common symptom associated with calcium deficiency, showed a considerable reduction in plants treated with calcium foliar sprays. The treatment T3- Egg shell derived Calcium 0.5% showed less incidence of fruits infected with BER (1.2 fruits) and more incidence was observed in the plants which were left untreated (6.2) as mentioned in the Table 2. This suggests a potential protective effect of the treatment against blossom end rot. Calcium is the one of the important nutrient that maintains the cell integrity and cell structure which enhanced the fruit quality by increasing the fruit firmness. Foliar application of calcium significantly reduced the blossom end rot. Peyvast *et al.* (2009), observed that calcium application affected yield and quality parameter, further calcium deficiency reduces yield while less fruit firmness, high fruit cracking and more blossom end rot was recorded in calcium deficient plants. The results are also in close proximity with findings of Gaonkar and Chakraborty (2016) that regular irrigations and calcium along with phosphorus fertilizer controlled the blossom end rot infestation in tomato and the number of fruits exhibiting cracking symptoms were lower in plants subjected to calcium foliar spray treatments. This observation implies a role for calcium supplementation in mitigating fruit cracking issues and the occurrence of fruit dropping, a phenomenon often linked to calcium imbalance, was notably reduced in Ca treated plants suggesting the foliar spray treatments positively influenced fruit retention and the lowest number of fruit cracking was observed in the plants treated with T2- Calcium Nitrate 0.5% (0.4), followed by T3- Egg shell derived Calcium 0.5% showed less incidence of fruit cracking (0.6) as against (4.2) in the control plants as mentioned in Table 2 and Table 3. Besides, increase in the biomass yield, the supplementation of Ca to the plants at the vegetative growth has also showed increased plant height (5.7-6.0 cm) when compared to untreated control (4.1 cm). The increasing tendency in

tomato plant height was similar with the findings of (Chaudhry *et al.*, 1999; Jaha and Krishi 2001) observed differences in plant height among cultivars of tomato pot under evaluation and screening trial. Our findings were in accordance with Jadhav *et al.* (2020) who reported that, dry fruit yield of chilli was significantly influenced by foliar spray of $\text{Ca}(\text{NO}_3)_2$. Foliar spray of $\text{Ca}(\text{NO}_3)_2$ resulted in increased absorption of Ca and $\text{NO}_3\text{-N}$ by plant canopy, this $\text{NO}_3\text{-N}$ might have enhanced fruit yield by enhancing cell division, that led to increased flowering and fruiting synergized by Ca present in $\text{Ca}(\text{NO}_3)_2$ and also due to timely and spot availability of calcium and nitrogen in readily available forms to the chilli plants which resulted in better absorption of nutrients by plant leaves leading to more flowers and fruits. Similar observations on increased fruit yield in chilli crop due to foliar spray of mono potassium phosphate and KNO_3 were reported by Veerendra Patel (2014) and Somimol (2012). The results of the present investigation were in accordance with the findings of Sarkar *et al.* (1999) and Chaity and Sarkar (2009).

Conclusion

By utilizing eggshells in this innovative manner, we not only address the challenge of waste disposal but also harness a valuable resource to enhance agricultural productivity in an environmentally sustainable manner. These findings underscore the potential of integrating organic calcium derived from eggshells as a viable and effective solution in modern agricultural practices.





Fig.2: Physiological parameters of tomato fruits. A-C: Symptoms of blossom end rot; B-D: PRANAM Ca sprayed healthy fruits

UNDER PEER REVIEW

References:

- Adams P, Ho LC. Effects of environment on the uptake and distribution of calcium in tomato and on the incidence of blossom-end rot. *Plant and soil*. 1993 Jul;154:127-32.
- Ahmad P, Sarwat M, Bhat NA, Wani MR, Kazi AG, Tran LS. Alleviation of cadmium toxicity in *Brassica juncea* L.(Czern. & Coss.) by calcium application involves various physiological and biochemical strategies. *PloS one*. 2015 Jan 28;10(1):e0114571.
- Ajala EO, Eletta OA, Ajala MA, Oyeniya SK. Characterization and evaluation of chicken eggshell for use as a bio-resource. *Arid Zone Journal of Engineering, Technology and Environment*. 2018 Mar 1;14(1):26-40.
- Amor FD, Marcelis LF. Regulation of nutrient uptake, water uptake and growth under calcium starvation and recovery. *The Journal of Horticultural Science and Biotechnology*. 2003 Jan 1;78(3):343-9.
- Amu OO, Fajobi AB, Oke BO. Effect of eggshell powder on the stabilizing potential of lime on an expansive clay soil. *Research journal of Agriculture and biological sciences*. 2005;1(1):80-4.
- Anonymous, 2020, Egg shell – an important poultry by-product. <https://www.pashudhanpraharee.com/egg-shell-an-important-poultry-byproduct/#:~:text=Egg%20shells%20are%20used%20to,without%20using%20eco%20Dunfriendly%20pesticides>
- Anonymous, 2023, Top Five egg producing countries & companies, Essential Protein Trade and shipping news. <https://essfeed.com/top-5-egg-producing-countries-companies/> (Accessed on 8th December, 2023)
- Boron L. Citrato de cálcio obtido da casca de ovo: biodisponibilidade e uso como suplemento alimentar.
- Butcher, G. D. and Miles, R., 2019, Concepts of eggshell quality. *Veterinary Medicine-Large Animal Clinical Sciences Department, UF/IFAS extension*. University of Florida. VM69 series. <https://edis.ifas.ufl.edu>. (Accessed on 8th December 2023)

- Chaudhry MF, Khokhar KM, Hussain SI, Mahmood T, Iqbal SM. Comparative performance of some local and exotic tomato cultivars during spring and autumn seasons. *Pakistan Journal of Arid Agriculture (Pakistan)*. 1999;2(2).
- Daengprok W, Garnjanagoonchorn W, Mine Y. Fermented pork sausage fortified with commercial or hen eggshell calcium lactate. *Meat science*. 2002 Oct 1;62(2):199-204.
- Das DK. *Micronutrients: Their behaviour in soils and plants*. Bangladesh Forest Research Institute; 2001.
- De Freitas ST, do Amarante CV, Labavitch JM, Mitcham EJ. Cellular approach to understand bitter pit development in apple fruit. *Postharvest Biology and Technology*. 2010 Jul 1;57(1):6-13.
- Douglas, S, M., Blossom-end rot of tomato. The Connecticut Agricultural Experiment Station. [www.ct.gov/caes.https://portal.ct.gov//media/CAES/DOCUMENTS/Publications/Fact_Sheets/Plant_Pathology_and_Ecology/BLOSSOMENDROTOFTOMATO110410Rpdf.pdf](https://portal.ct.gov//media/CAES/DOCUMENTS/Publications/Fact_Sheets/Plant_Pathology_and_Ecology/BLOSSOMENDROTOFTOMATO110410Rpdf.pdf) (Accessed on 8th December 2023)
- Dublin, 2023, Global Egg Industry Analysis 2017-2022 & 2023-2027: Forecast Data, Trends, Growth Drivers, and Challenges. Research and Markets. <https://www.globenewswire.com/en/newsrelease/2023/08/17/2727026/28124/en/Global-Egg-Industry-Analysis-2017-2022-2023-2027-Forecast-Data-Trends-Growth-Drivers-and-Challenges.html> (Accessed on 8th December 2023)
- Faridi H, Arabhosseini A. Application of eggshell wastes as valuable and utilizable products: A review. *Research in Agricultural Engineering*. 2018 Apr 1;64(2).
- Gaonkar M, Chakraborty AP. Application of eggshell as fertilizer and calcium supplement tablet. *International journal of innovative research in science, engineering and technology*. 2016 Mar;5(3):3520-5.
- Gebhardt SE. *Nutritive value of foods*. US Department of Agriculture, Agricultural Research Service; 2002.
- Hamster MR, Balzer PS, Becker D. Characterization of calcium carbonate obtained from oyster and mussel shells and incorporation in polypropylene. *Materials Research*. 2012;15:204-8.

- Hecht J. Eggshells break into collagen market.
https://scholar.google.com/scholar?hl=en&as_sdt=0%2C5&q=Hecht+J.+Eggshells+break+into+collagen+market.&btnG=
- Ho LC, Adams P, Li XZ, Shen H, Andrews J, Xu ZH. Responses of Ca-efficient and Ca-inefficient tomato cultivars to salinity in plant growth, calcium accumulation and blossom-end rot. *Journal of Horticultural Science*. 1995 Jan 1;70(6):909-18.
- Horst Marschner. Mineral nutrition of higher plants. Academic press; 1995.
- Huntun P, 2005, *Brazilian Journal of Poultry Science*, Jun 2005 7 (2): 67 – 71.
- Jadhav KP, Bidari BI, Shashidara GB, Venkatesh MS. Effect of calcium nitrate foliar application on quality and yield of Byadgi Chilli. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(6):1628-31.
- Jadhav KP, Bidari BI, Shashidara GB, Venkatesh MS. Effect of calcium nitrate foliar application on quality and yield of Byadgi Chilli. *Journal of Pharmacognosy and Phytochemistry*. 2020;9(6):1628-31.
- Jaha JC, Krishi B. Studies on performance of different tomato hybrids in offseason under different planting method in Terai agro-climatic zones of west Bangal. *J. Interacademia*. 2001;5(2):186-9.
- Johnson, G., 2022, calcium disorders in vegetables and fruits.
<https://sites.udel.edu/weeklycropupdate/?p=20474>
- Kattimani V, Panga GS, Girija EK. Eggshell membrane separation methods-waste to wealth-a scoping review. *Journal of Poultry Research*. 2022 Jun 6;19(1):11-8.
- Khan A, Anwar Y, Hasan MM, Iqbal A, Ali M, Alharby HF, Hakeem KR, Hasanuzzaman M. Attenuation of drought stress in Brassica seedlings with exogenous application of Ca²⁺ and H₂O₂. *Plants*. 2017 May 13;6(2):20.
- Kundu C, Sarkar RK. Effect of foliar application of potassium nitrate and calcium nitrate on performance of rainfed lowland rice (*Oryza sativa*). *Indian Journal of Agronomy*. 2009;54(4):428-32.
- Lahoz I, Perez-de-Castro A, Valcárcel M, Macua JI, Beltran J, Roselló S, Cebolla-Cornejo J. Effect of water deficit on the agronomical performance and quality of processing tomato. *Scientia Horticulturae*. 2016 Mar 8;200:55-65.

- Naeem MU, Naeem MS, Ahmad R, Ahmad R. Foliar-applied calcium induces drought stress tolerance in maize by manipulating osmolyte accumulation and antioxidative responses. *Pak. J. Bot.* 2017 Apr 1;49(2):427-34.
- Neves MA. Alternativas para a valorização da cacao de ovo como complemento alimentar e em implantes ósseos.
- Peyvast G, Olfati JA, Ramezani-Kharazi P, Kamari-Shahmaleki S. Uptake of calcium nitrate and potassium phosphate from foliar fertilization by tomato. *Journal of Horticulture and Forestry.* 2009;1(1):7-13.
- Sajid M, Ullah I, Rab A, Shah ST, Basit A, Bibi F, Ahmad M. 2. Foliar application of calcium improves growth, yield and quality of tomato cultivars. *Pure and Applied Biology (PAB).* 2020 Feb 7;9(1):10-9.
- Sajid M, Ullah I, Rab A, Shah ST, Basit A, Bibi F, Ahmad M. 2. Foliar application of calcium improves growth, yield and quality of tomato cultivars. *Pure and Applied Biology (PAB).* 2020 Feb 7;9(1):10-9.
- Sarkar RK, Chakraborty A, Saha A. Effect of foliar application of potassium nitrate and calcium nitrate on groundnut (*Arachis hypogaea*). *Indian Journal of Agronomy.* 1999;44(4):809-12.
- Sarwat M, Ahmad P, Nabi G, Hu X. Ca²⁺ signals: the versatile decoders of environmental cues. *Critical reviews in biotechnology.* 2013 Mar 1;33(1):97-109.
- Shabbir RN, Ashraf MY, Waraich EA, Ahmad R, Shahbaz M. Combined effects of drought stress and NPK foliar spray on growth, physiological processes and nutrient uptake in wheat. *Pak. J. Bot.* 2015 Aug 1;47(4):1207-16.
- Shahbandeh. M., 2023, Egg production worldwide 1990-2021. <https://www.statista.com/statistics/263972/egg-production-worldwide-since-1990/#statisticContainer> (Accessed on 8th December 2023)
- Shear CB. Calcium-related Disorders of Fruits and Vegetables1. *HortScience.* 1975 Aug 1;10(4):361-5.
- Somimol PV. Effect of foliar feeding of 19: 19: 19 and potassium nitrate (KNO₃) water soluble fertilizers on yield and quality of Byadgi chillies in Vertisols, M. Sc (Agri.) Thesis, Univ. Agric. Sci., Dharwad (India). 2012.
- Taylor MD, Locascio SJ. Blossom-end rot: a calcium deficiency. *Journal of Plant nutrition.* 2004 Dec 27;27(1):123-39.

- Tuna AL, Kaya C, Ashraf M, Altunlu H, Yokas I, Yagmur B. The effects of calcium sulphate on growth, membrane stability and nutrient uptake of tomato plants grown under salt stress. *Environmental and Experimental Botany*. 2007 Mar 1;59(2):173-8.
- Ustun N, Altunlu H, Yokaş I, Saygili H. Influence of potassium and calcium levels on severity of tomato pith necrosis and yield of greenhouse tomatoes. In II International Symposium on Tomato Diseases 808 2007 Oct 8 (pp. 347-350).
- Vandeginste V. Food waste eggshell valorization through development of new composites: A review. *Sustainable Materials and Technologies*. 2021 Sep 1;29: 00317.
- Verma N, Kumar V and Bansal M. C. Utilization of egg shell waste in cellulase production by *neurospora crassa* under wheat bran-based solid state fermentation. *Polish journal of environmental studies*.2012; 21(2):491-497.
- Watanabe T, Tomizaki R, Watanabe R, Maruyama H, Shinano T, Urayama M, Kanayama Y. Ionomic differences between tomato introgression line IL8–3 and its parent cultivar M82 with different trends to the incidence of blossom-end rot. *Scientia horticulturae*. 2021 Sep 20; 287:110266.
- Zorrig W, Shahzad Z, Abdelly C, Berthomieu P. Calcium enhances cadmium tolerance and decreases cadmium accumulation in lettuce (*Lactuca sativa*). *African Journal of Biotechnology*. 2012;11(34):8441-8.

Table 2: Effect of spraying different sources of Calcium on plant height and fruit physiological characters of tomato (local variety)

Treatment details	Height (Feet)	Blossom end rot	Fruit Cracking	Fruit dropping
T1- Calcium Chloride 0.5%	5.70 ^a (39.02)	1.4 ^b (77.42)	0.8 ^b (80.95)	1 ^b (84.38)
T2- Calcium Nitrate 0.5%	6.00 ^a (46.34)	1.2 ^b (80.65)	0.4^b (90.48)	1.2 ^b (81.25)
T3- Egg shell derived Calcium 0.5%	5.84^a (42.44)	1.2^b (80.65)	0.6 ^b (85.71)	1.4 ^b (78.13)
T4-Control	4.1 ^b (0.00)	6.2 ^a (0.00)	4.2 ^a (0.00)	6.4 ^a (0.00)
CD@1%	0.96	1.538	1.116	1.597

* Values are mean of three replications. Mean ratings a column that are followed by the same letter are not significantly different according to LSD at $P \leq 0.05$

Table 3: Effect of spraying different sources of Calcium on height and yield parameters of tomato (local variety)

Treatment details	Yield (g/plant) at 1 st picking
T1- Calcium Chloride 0.5%	319 ^b (26.38)
T2- Calcium Nitrate 0.5%	402 ^a (59.27)
T3- Egg shell derived Calcium 0.5%	389^a(54.12)
T4-Control	252.4 ^c (0.00)
CD@1%	90.37

* Values are mean of three replications. Mean ratings a column that are followed by the same letter are not significantly different according to LSD at $P \leq 0.05$




Fig.3: Effect of PRANAM Ca on Growth and Yield of tomato




Fig.4: Effect of PRANAM Ca on Physiological parameters of tomato fruits