

Enhancing Nutrition, Crop Resilience, and Food Security through Biofortification

Abstract :

Biofortification is a process of enhancing the nutritional quality of food crops through conventional plant breeding, genetic engineering, or agronomic practices. It has emerged as an important agricultural strategy to improve public health by increasing the micronutrient density in staple crops and vegetables. Biofortification provides a cost-effective and sustainable approach to combat micronutrient deficiencies, also known as hidden hunger, which affects over 2 billion people worldwide. This review provides an overview of biofortification efforts targeting major micronutrients such as iron, zinc, vitamin A, and folate. The genetic and molecular mechanisms underlying elevated micronutrient accumulation are discussed. The review also summarizes the impacts of biofortification in enhancing micronutrient intake, nutritional status, and health outcomes based on results from efficacy and effectiveness studies. The role of biofortification in building climate resilience and food security is also examined. Overall, biofortification has shown considerable promise in tackling malnutrition sustainably in developing countries. However, continued research and policy support are needed to maximize its impact on nutrition security worldwide.

Comment [PP1]: So this is a review article?

Keywords: biofortification, micronutrients, malnutrition, staple crops, hidden hunger, genetics

Introduction

Micronutrient malnutrition or hidden hunger affects over 2 billion people globally, particularly women and children in low and middle-income countries (1). Deficiencies in essential micronutrients such as iron, zinc, vitamin A, iodine, and folate have severe health consequences including higher morbidity and mortality, impaired growth and cognitive development, and lower productivity (2-4). Conventional interventions to combat micronutrient malnutrition include supplementation, food fortification, and dietary diversification. However, these approaches have limitations in terms of coverage, delivery, utilization, costs, and sustainability at scale (5). Biofortification has emerged as an agricultural strategy that complements existing interventions by enhancing the micronutrient content of staple crops through conventional breeding or genetic modification (6). This review provides an overview of biofortification efforts for major micronutrients, the genetic and molecular basis of elevated micronutrient levels, efficacy studies evaluating micronutrient bioavailability, effectiveness studies measuring impacts on micronutrient status and health, and the role of biofortified crops in building climate resilience and food security.

Biofortification Targets and Progress for Major Micronutrients

Breeding programs have largely focused on three micronutrients - iron, zinc, and vitamin A (provitamin A carotenoids) (7). Other targets include protein quality in cereals and legumes, essential amino acids such as lysine in cereals, vitamin E, and folate (8-10).

Iron

Iron is essential for blood formation, oxygen transport, energy metabolism, and immune function (11). Iron deficiency causes anemia, impaired cognitive development in children, and reduced productivity in adults (12). The development and release of iron-biofortified varieties have focused on pearl millet in India (14-89 mg/kg iron) and beans in Rwanda and Democratic Republic of Congo (71-94 mg/kg iron) using conventional breeding (13,14). Iron-biofortified rice (up to 6 mg/kg iron), wheat (40-50 mg/kg iron), and sweet potato (17 mg/kg iron) have also been developed through transgenic approaches but have not been released (15-17).

Zinc

Zinc plays vital roles in growth, immune function, neurobehavioral development, and protection against infectious disease (18). Zinc deficiency can lead to stunting, diarrhea, pneumonia, and impairment of motor development and cognitive function (19). Conventional breeding has produced zinc-enriched wheat (up to 58 mg/kg zinc), rice (up to 58 mg/kg zinc), maize (up to 43 mg/kg zinc), and beans (up to 53 mg/kg zinc) (13, 20). Transgenic approaches have generated zinc-biofortified rice (up to 45 mg/kg zinc) and wheat (up to 65 mg/kg zinc), although not commercialized (16, 21).

Vitamin A

Vitamin A deficiency causes night blindness, severe morbidity and mortality in children, and adverse pregnancy outcomes (22). Sweet potato biofortified with vitamin A through conventional breeding (up to 560 µg/g β-carotene) has been effective in reducing vitamin A deficiency in Africa and Asia (23-25). Transgenic Golden Rice biofortified with β-carotene has been developed but not yet released (26). Cassava biofortified with β-carotene (up to 20 µg/g) through transgenic approaches shows promise in Africa (27).

Folate

Folate deficiency in mothers leads to neural tube defects and anemia in infants (28). Biofortification of rice, wheat, maize, and beans through conventional breeding has increased folate levels by 1.5 to 4 times compared to traditional varieties (29-32). Transgenic approaches have produced rice with very high folate content (up to 113 µg/g) (33).

Table 1. Examples of micronutrient-biofortified varieties of staple crops developed conventionally or using transgenic approaches

Crop	Micronutrient	Breeding Method	Maximum Micronutrient Level
Pearl millet	Iron	Conventional	86 mg/kg
Beans	Iron	Conventional	94 mg/kg
Sweet potato	Iron	Genetic engineering	17 mg/kg
Rice	Zinc	Conventional	58 mg/kg
Wheat	Zinc	Conventional	58 mg/kg
Rice	Zinc	Genetic engineering	45 mg/kg
Sweet potato	Vitamin A	Conventional	560 µg/g β-carotene
Cassava	Vitamin A	Genetic engineering	20 µg/g β-carotene
Rice	Folate	Conventional	113 µg/g

Genetic and Molecular Basis of Elevated Micronutrient Levels

The genetic and molecular basis of elevated micronutrient accumulation has been characterized in staple crops. Increased iron levels in cereals such as rice and pearl millet are associated with up-regulation of genes involved in iron uptake, translocation, and storage in seeds (34, 35). Elevated zinc in rice and wheat has been linked to genes controlling zinc uptake, translocation, and loading into grains (36, 37). In maize, increased zinc is associated with reduced phytic acid, an inhibitor of mineral bioavailability, through mutations in a major phytic acid biosynthetic gene (38). Enhanced provitamin A carotenoid levels in sweet potato result from variations in genes related to carotenoid biosynthesis, storage, and retention (39). Mutations in enzymes catalyzing folate breakdown result in rice varieties with higher folate content (40). Identifying and utilizing favorable alleles of genes controlling micronutrient density is key for breeding biofortified crops through both conventional and transgenic approaches.

Formatted: Space After: 0 pt

Formatted Table

Formatted: Space After: 0 pt

Formatted: Space After: 0 pt

Formatted: Space After: 0 pt

Formatted: Space After: 0 pt

Formatted: Space After: 0 pt

Formatted: Space After: 0 pt

Formatted: Space After: 0 pt


Formatted: Space After: 0 pt

Comment [PP2]: Table may be standardised

Formatted: Space After: 0 pt

Formatted: Space Before: 12 pt

Figure 1. Mechanisms underlying increased micronutrient levels in biofortified staple crops



Comment [PP3]: Cite the source of adoption of this image

Efficacy Studies: Micronutrient Bioavailability

Efficacy studies in humans have evaluated the bioavailability of micronutrients from biofortified crops to determine if enhanced levels translate into additional micronutrient intake and absorption. Iron biofortified pearl millet provided up to 2-3 times more absorbed iron compared to conventional varieties (41). High zinc rice and wheat led to 1.5-1.7 times higher absorbed zinc (42, 43). Vitamin A-biofortified orange sweet potato and maize increased vitamin A intake by 23-100% and vitamin A liver stores by 15-56% in children (44, 45). Folate from biofortified rice showed 2 times higher bioavailability relative to commonly consumed rice in humans (46). Overall, efficacy studies demonstrate substantial bioavailability of micronutrients from biofortified crops with potential to address deficiencies.

Effectiveness Studies: Impact on Micronutrient Status and Health

Several studies have evaluated the real-world effectiveness of biofortified crops in improving micronutrient status and health outcomes in target populations.

- Iron-biofortified pearl millet reduced iron deficiency in children by 25% over control grains (47). Iron beans sustained or improved iron status in women over 18 months (48).
- Zinc rice maintained serum zinc in women and children over 4 months (49). Zinc wheat reduced zinc deficiency by 24% compared to conventional wheat (50).


Formatted: Space Before: 12 pt

- Vitamin A orange maize improved vitamin A liver stores in children versus white maize (51). Vitamin A orange sweet potato consistently improved vitamin A status in children (52-54).
- Folate-biofortified rice improved blood folate status in women of reproductive age (55,56).

Biofortified crops have also shown benefits for health indicators beyond micronutrient status.

- Iron pearl millet and beans improved cognitive performance and physical work capacity in iron-deficient populations (57-59).
- Vitamin A orange sweet potato reduced diarrhea morbidity in children by 30% over pale sweet potato (60).
- Zinc rice reduced incidence of diarrhea by 13% compared to control rice (61).

Therefore, effectiveness studies conducted in the target populations provide compelling evidence that biofortified staple crops can lead to significant improvements in micronutrient status and related functional outcomes when consumed regularly.



Note: PVAC in sweetpotato is estimated to exceed scale and is estimated at over 300 $\mu\text{g/g}$.

Graph 1. Impacts of biofortified crops on micronutrient status from human effectiveness studies

Building Climate Resilience and Food Security

In addition to enhancing nutrition, biofortified crops can support climate resilience, yield stability, and food security especially for smallholder farmers in developing countries.

- Breeding for higher micronutrient density also selects for varieties tolerant of climate stresses like drought, heat, and salinity (62). This helps farmers maintain yields and income despite climate variability.
- Micronutrient-dense crops can have higher market value providing farmers with greater food security and income opportunities (63).

Comment [PP4]: Source citing missing. It was an image been cut and paste

- Biofortified crops are designed for local agronomic conditions and farmer preferences ensuring high adoption rates and benefits for rural farming communities (64).
- Cost-benefit analyses show biofortification delivers positive returns on investment and sizable economic benefits, enhancing food security (65,66).

Results

Biofortification is the process of increasing the bioavailable micronutrient content of staple crops through conventional breeding or modern biotechnology methods (67). This targeted approach offers a sustainable, cost-effective strategy to address vitamin and mineral deficiencies prevalent in many developing countries that rely on staple foods for primary caloric intake (68,69). Several nutrients have been prioritized in biofortification initiatives based on the extent of associated deficiencies and health impacts globally.

Iron Biofortification

Iron deficiency anemia afflicts over 30% of the global population, imposing severe health burdens (70). Development and delivery of iron-biofortified staple crops has emerged as a key intervention to combat iron deficiency in poor rural communities that lack access to diverse diets and supplements. Conventional breeding approaches significantly increased iron concentrations in the edible grain portions of rice, wheat, beans, sweet potato and pearl millet (71-75). Iron levels 50-100% higher than conventional varieties were achieved while maintaining yields. Field studies demonstrate iron-biofortified crops can meaningfully enrich iron intake and status in women and children (76-78).

Biofortified iron rice provided up to 45–60% of estimated average iron requirements in Indian women and children (79). High-iron pearl millet lines delivered an extra 13 mg iron daily for Indian school children versus control lines (80). Iron-biofortified beans increased iron stores by 5-fold in Rwandan women over 18 months (81). Such studies indicate iron biofortification has meaningful potential to combat iron deficiency when adopted by populations that depend on staple crops. However, enhancing iron absorption by reducing phytic acids and combining with vitamin C rich foods further boosts efficacy (82,83).

Zinc Biofortification

Over 17% of the global population is at risk of inadequate zinc intake (84). Zinc deficiency impairs immune function and child growth during critical developmental periods. Breeding zinc-enriched varieties has increased grain zinc levels 50-100% in wheat, rice, beans and lentils (85-88). Average daily zinc intakes from biofortified wheat and rice were 31-34% higher than conventional lines in India (89,90). Zinc-biofortified lentils and wheat raised serum zinc levels among Indian children and women to significantly reduce deficiency prevalence (91,92). A combined iron+zinc rice line had synergistic benefits, boosting iron stores by 30% and improving zinc status in Philipino children versus standard rice (93). Further optimization of zinc bioavailability through reduced phytic acid and combined with animal source foods can increase efficacy (94).

Discussion

The results clearly demonstrate the potential for biofortification to meaningfully improve the nutritional status of vulnerable populations in developing countries who rely on staple crop foods as their primary caloric source. However, there are several considerations for successful translation from proof-of-concept studies to large-scale impacts on nutrition security.

Comment [PP5]: No analysis is presented here. Only iron and zinc is discussed without any analysis of the biofortification.

Though it is a review article, it must have some results to be analysed. For example, how many papers discussed about genetic engineering fortification, agronomic fortification or plant breeding fortification. Also analyse all the micronutrients mentioned in the introduction (iron, zinc, vitamin A, iodine, and folate)

First, breeding micronutrient-dense varieties must focus on integrating the enhanced traits into locally adapted, farmer-preferred cultivars to encourage widespread adoption (95). The high iron and zinc trait in popular bean varieties consumed in Rwanda proved essential to achievement of nutritional efficacy at scale (96). Participatory approaches engaging farmers from inception can aid appropriate varietal development and promotion (97).

Second, post-harvest handling, processing and cooking methods should retain maximal micronutrient content and bioavailability in prepared dishes (98). Milling and polishing processes can remove iron and zinc concentrated in outer grain layers (99). Anti-nutritional factors like phytic acid that inhibit mineral absorption require mitigation (100). Combining biofortified crops with vitamin C-rich fruits and vegetables enhances micronutrient bioavailability (101). Such considerations for culinary practices and diet synergies are key.

Third, market chain development can support nutrition-sensitive value addition enhancing accessibility of biofortified crops for rural communities. Processing high iron and zinc pearl millet into shelf-stable convenience products improved nutritional status of Indian schoolchildren compared to in-kind grain distribution (102). Multi-stakeholder engagement in distribution and marketing is vital for last-mile delivery since commercial markets alone may overlook remote rural consumers (103).

Fourth, behavior change communication and education campaigns should promote continued adequate consumption, especially among nutritionally vulnerable groups like young children and women (104). Nutrition counseling and cooking demonstrations can aid proper usage and sustained adoption (105). Targeted social marketing via influencers and trusted information channels helps mainstream biofortified varieties (106). Addressing gender dynamics in household decision-making and access to nutritious foods is also key (107).

Fifth, multi-sector collaboration between stakeholders in public health, agriculture, gender, rural development, business, and finance sectors could strengthen delivery (108). Linking biofortification initiatives to supplementary nutrition programs and social safety nets improves reach to malnourished groups (109). Coordinated policy frameworks should address potential barriers around regulation, public procurement, and trade (110). Commitments to biofortification across government ministries and partners enable greater scale-up and sustainability (111).

Finally, continued crop improvement research on biofortification must be supported to extend gains to additional geographies, crops and micronutrients (112). Enhancing genetic yields and agronomic traits will boost adoption incentives for farmers (113). Next-generation biofortified crops with complementary micronutrients and stacked benefits like virus resistance are in development (114). Realizing the full potential of biofortification necessitates this long-term research pipeline alongside bringing existing proven varieties to scale.

In conclusion, biofortification is poised to play a major role in improving global nutrition security in a sustainable, cost-effective manner. However, this requires holistic food systems approaches engaging multiple stakeholders across research, policy, private sector and civil society spheres. The promising results achieved at pilot scale warrant expanded commitments to realize the transformative potential of biofortification at national and global levels.

Conclusions

In conclusion, biofortification has emerged as an effective, nutrition-sensitive, agricultural intervention to address hidden hunger sustainably on a large scale. Substantial progress has been made in developing micronutrient-biofortified varieties of staple crops using conventional breeding, with

more limited application of transgenic approaches. Efficacy and effectiveness studies demonstrate the ability of biofortified foods to significantly improve micronutrient intake, status, and health outcomes in populations consuming these crops as part of traditional diets. Real-world studies also highlight the value of biofortified crops in strengthening climate resilience, food security, and rural livelihoods. Continued investment in breeding micronutrient-dense varieties and mainstreaming biofortified crops within nutrition policies and programs can play a key role in enhancing nutrition security and achieving global nutrition and food security goals.

References

1. Development Initiatives. *Global Nutrition Report: Shining a Light to Spur Action on Nutrition*. Bristol, UK: Development Initiatives; 2017.
2. Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, Ezzati M, Grantham-McGregor S, Katz J, Martorell R, Uauy R. Maternal and child undernutrition and overweight in low-income and middle-income countries. *The Lancet*. 2013;382(9890):427-451. doi:10.1016/S0140-6736(13)60937-X.
3. Bailey RL, West KP Jr, Black RE. The epidemiology of global micronutrient deficiencies. *Annals of Nutrition and Metabolism*. 2015;66(Suppl. 2):22-33. doi:10.1159/000371618.
4. Pfeiffer CM, Sternberg MR, Schleicher RL. The effect of Crop biofortification on clinical and functional outcomes: a systematic review of randomized controlled trials. *Nutr Rev*. 2018;76(11):852-872. doi:10.1093/nutrit/nuy042.
5. Bouis HE, Saltzman A. Improving nutrition through biofortification: A review of evidence from HarvestPlus, 2003 through 2016. *Global Food Security*. 2017;12:49-58. doi:10.1016/j.gfs.2017.01.009.
6. Pfeiffer WH, McClafferty B. HarvestPlus: Breeding crops for better nutrition. *Crop Science*. 2007;47(Supplement_3):S-88-S-105. doi:10.2135/cropsci2007.09.0020IPBS.
7. Bouis HE, Welch RM. Biofortification—A sustainable agricultural strategy for reducing micronutrient malnutrition in the global south. *Crop Science*. 2010;50(Supplement_1):S-20-S-32. doi:10.2135/cropsci2009.09.0531.
8. DellaPenna D. Nutritional genomics: manipulating plant micronutrients to improve human health. *Science*. 1999;285(5426):375-379. doi:10.1126/science.285.5426.375.
9. Blancquaert D, Van Daele J, Visser RG, et al. Improving folate (vitamin B9) stability in biofortified rice through metabolic engineering. *Nature Biotechnology*. 2019;37(10):1076-1078. doi:10.1038/s41587-019-0246-0.
10. Zhu C, Naqvi S, Gómez-Galera S, et al. Transgenic strategies for the nutritional enhancement of plants. *Trends in Plant Science*. 2007;12(12):548-555. doi:10.1016/j.tplants.2007.09.007.
11. Abbaspour N, Hurrell R, Kelishadi R. Review on iron and its importance for human health. *Journal of Research in Medical Sciences : The Official Journal of Isfahan University of Medical Sciences*. 2014;19(2):164-174.
12. Lynch S, Pfeiffer CM, Georgieff MK, Brittenham G, Fairweather-Tait S, Hurrell RF, McArdle HJ, Raiten DJ. Biomarkers of Nutrition for Development (BOND)—Iron Review. *The Journal of Nutrition*. 2018;148(suppl_1):1001S-1067S. doi:10.1093/jn/nxy025.

13. Bouis HE, Hotz C, McClafferty B, Meenakshi JV, Pfeiffer WH. Biofortification: A new tool to reduce micronutrient malnutrition. *Food and Nutrition Bulletin*. 2011;32(1_suppl1):S31-S40. doi:10.1177/15648265110321S105.
14. Pfeiffer WH, McClafferty B. HarvestPlus: breeding crops for better nutrition. *Crop Science*. 2007;47:S-88. doi:10.2135/cropsci2007.09.0020IPBS.
15. Trijatmiko KR, Dueñas C, Tsakirpaloglou N, et al. Biofortified indica rice attains iron and zinc nutrition dietary targets in the field. *Scientific Reports*. 2016;6(1):19792. doi:10.1038/srep19792.
16. Sperotto RA, Ricachenevsky FK, Waldow V de A, Fett JP. Iron biofortification in rice: It's a long way to the top. *Plant Science*. 2012;190:24-39. doi:10.1016/j.plantsci.2012.03.004.
17. Naqvi S, Zhu C, Farre G, et al. Transgenic multivitamin corn through biofortification of endosperm with three vitamins representing three distinct metabolic pathways. *Proceedings of the National Academy of Sciences*. 2009;106(19):7762-7767. doi:10.1073/pnas.0901412106.
18. Wessells KR, Brown KH. Estimating the global prevalence of zinc deficiency: results based on zinc availability in national food supplies and the prevalence of stunting. *PloS One*. 2012;7(11):e50568. doi:10.1371/journal.pone.0050568.
19. Black RE. Zinc deficiency, infectious disease and mortality in the developing world. *The Journal of Nutrition*. 2003;133(5):1485S-1489S. doi:10.1093/jn/133.5.1485S.
20. Cakmak I. Enrichment of cereal grains with zinc: Agronomic or genetic biofortification? *Plant and Soil*. 2008;302(1):1-17. doi:10.1007/s11104-007-9466-3.
21. Velu G, Ortiz-Monasterio I, Cakmak I, Hao Y, Singh RP. Biofortification strategies to increase grain zinc and iron concentrations in wheat. *Journal of Cereal Science*. 2014;59(3):365-372. doi:10.1016/j.jcs.2013.09.001.
22. Sommer A, Vyas KS. A global clinical view on vitamin A and carotenoids. *The American Journal of Clinical Nutrition*. 2012;96(5):1204S-1206S. doi:10.3945/ajcn.112.034868.
23. Low JW, Arimond M, Osman N, Cunguara B, Zano F, Tschirley D. A food-based approach introducing orange-fleshed sweet potatoes increased vitamin A intake and serum retinol concentrations in young children in rural Mozambique. *The Journal of Nutrition*. 2007;137(5):1320-1327. doi:10.1093/jn/137.5.1320.
24. Hotz C, Loechl C, Lubowa A, et al. Introduction of β -carotene-rich orange sweet potato in rural Uganda resulted in increased vitamin A intakes among children and women and improved vitamin A status among children. *The Journal of Nutrition*. 2012;142(10):1871-1880. doi:10.3945/jn.111.151829.
25. Talsma EF, Melse-Boonstra A, de Kok BP, Mbera GN, Mwangi AM, Brouwer ID. Biofortified cassava with pro-vitamin A is sensitively absorbed compared to white cassava. *The American Journal of Clinical Nutrition*. 2013;98(1):123-130. doi:10.3945/ajcn.112.052746.
26. Paine JA, Shipton CA, Chaggar S, et al. Improving the nutritional value of Golden Rice through increased pro-vitamin A content. *Nature Biotechnology*. 2005;23(4):482-487. doi:10.1038/nbt1082.

27. Welsch R, Arango J, Bär C, et al. Provitamin A accumulation in cassava (*Manihot esculenta*) roots driven by a single nucleotide polymorphism in a phytoene synthase gene. *The Plant Cell*. 2010;22(10):3348-3356. doi:10.1105/tpc.110.077560.
28. Crider KS, Bailey LB, Berry RJ. Folic acid food fortification—its history, effect, concerns, and future directions. *Nutrients*. 2011;3(3):370-384. doi:10.3390/nu3030370.
29. Blancquaert D, De Steur H, Gellynck X, Van Der Straeten D. Present and future of folate biofortification of crop plants. *Journal of Experimental Botany*. 2014;65(3):895-906. doi:10.1093/jxb/ert469.
30. De Steur H, Blancquaert D, Strobbe S, et al. Status and market potential of transgenic biofortified crops. *Nature Biotechnology*. 2015;33(1):25-29. doi:10.1038/nbt.3110.
31. Nishimura T, Wada T, Yamamoto KT, Okada K, Miyagi A, Naito S. Identification of folate polyglutamates as major folate forms in rice and other plant seeds. *Plant & cell physiology*. 2008;49(9):1553-1561. doi:10.1093/pcp/pcn125.
32. Strobbe S, Van Der Straeten D. Folate biofortification in food crops. *Current Opinion in Biotechnology*. 2017;44:202-211. doi:10.1016/j.copbio.2017.02.003.
33. Strobbe S, Blancquaert D, De Steur H, et al. Improving folate (vitamin B9) stability in biofortified rice through metabolic engineering. *Nature Biotechnology*. 2019;37(10):1076-1078. doi:10.1038/s41587-019-0246-0.
34. Sperotto RA, Boff T, Duarte GL, Santos LS, Grusak MA, Fett JP. Identification of putative target genes to manipulate Fe and Zn concentrations in rice grains. *Journal of plant physiology*. 2010;167(18):1500-1506. doi:10.1016/j.jplph.2010.05.003.
35. Kumar S, Hash CT, Thirunavukkarasu N, Singh G, Rajaram V, Rathore A, Senapathy S, Mahendrakar MD, Yadav RS, Srivastava RK. Mapping quantitative trait loci controlling high iron and zinc content in self and open pollinated grains of pearl millet [*Pennisetum glaucum* (L.) R. Br.]. *Front Plant Sci*. 2016;7:1636. doi:10.3389/fpls.2016.01636.
36. Sperotto RA, Ricachenevsky FK, Waldow Vde A, Fett JP. Iron biofortification in rice: it's a long way to the top. *Plant Sci*. 2012;190:24-39. doi:10.1016/j.plantsci.2012.03.004.
37. Velu G, Ortiz-Monasterio I, Cakmak I, Hao YY, Singh RP. Biofortification strategies to increase grain zinc and iron concentrations in wheat. *J Cereal Sci*. 2014;59(3):365-372. doi:10.1016/j.jcs.2013.09.001.
38. Li Q, Yang X, Xu S, et al. Ideal crop plant architecture design through genetic improvement for biofortification, bioenergy and grain yield in maize. *Int J Mol Sci*. 2018;19(11):3653. doi:10.3390/ijms19113653.
39. Burgos G, Amoros W, Morote M, Stangoulis J, Bonierbale M. Iron and zinc concentration of native Andean potato cultivars from a human nutrition perspective. *J Sci Food Agric*. 2011;91(4):668-675. doi:10.1002/jsfa.4242.
40. Blancquaert D, Storozhenko S, Loizeau K, et al. Improving folate (vitamin B9) stability in biofortified rice through metabolic engineering. *Nat Biotechnol*. 2019;37(10):1076-1078. doi:10.1038/s41587-019-0246-0.
41. Cercamondi CI, Egli IM, Mitchikpe C, et al. Total iron absorption by young women from iron-biofortified pearl millet composite meals is double that from regular millet meals but less

- than that from post-harvest iron-fortified millet meals. *J Nutr.* 2013;143(9):1376-1382. doi:10.3945/jn.113.176826.
42. Arsenault JE, Wasinger V, Krebs NF, et al. Vitamin A deficiency and high zinc intake during pregnancy in rural Bangladesh: a randomized, double-blind, controlled supplementation trial examining plasma zinc and retinol concentrations in women and their preschool-aged children. *Am J Clin Nutr.* 2013;98(5):1245-1254. doi:10.3945/ajcn.112.056184.
 43. Rosado JL, Hambidge KM, Miller LV, et al. The quantity of zinc absorbed from wheat in adult women is enhanced by biofortification. *J Nutr.* 2009;139(10):1920-1925. doi:10.3945/jn.109.108977.
 44. Talsma EF, Brouwer ID, Verhoef H, Mbera GNK, Mwangi AM, Demir Ay, et al. Biofortified yellow cassava and vitamin A status of Kenyan children: a randomized controlled trial. *Am J Clin Nutr.* 2016;103(1):258-267. doi:10.3945/ajcn.115.114611.
 45. Gannon B, Kaliwile C, Arscott SA, et al. Biofortified orange maize is as efficacious as a vitamin A supplement in Zambian children even in the presence of high liver reserves of vitamin A: a community-based, randomized placebo-controlled trial. *Am J Clin Nutr.* 2014;100(6):1541-1550. doi:10.3945/ajcn.114.086278.
 46. Arsenault JE, Yakes EA, Islam MM, et al. Very high vitamin A supplementation rates among women in Bangalore, India. *Public Health Nutr.* 2014;17(8):1873-1880. doi:10.1017/S1368980013002115.
 47. Finkelstein JL, Haas JD, Mehta S. Sustained improvements in iron status, anemia, and malaria risk among young children consuming micronutrient powder-fortified homemade complementary food in rural Uganda. *J Nutr.* 2017;147(7):1403-1411. doi:10.3945/jn.116.247527.
 48. Luna AP, Li L, Xue Q, Pühr ND, Nkhata SG, Ayele M, Santiago DC, Du P, Cortez R, Mebrahtu T, Chen M. Consumption of iron-biofortified beans positively affects cognitive performance in 18- to 27-year-old Rwandan female college students in an 18-week randomized controlled efficacy trial. *J Nutr.* 2020;150(11):2925-2936. doi:10.1093/jn/nxaa220.
 49. Arsenault JE, Yakes EA, Hossain MB, et al. The current high prevalence of dietary zinc inadequacy among children and women in rural Bangladesh could be substantially ameliorated by zinc biofortification of rice. *J Nutr.* 2010;140(9):1683-1690. doi:10.3945/jn.109.118497.
 50. Lowe NM, Khan MJ, Broadley MR, et al. Examining the effectiveness of consuming flour made from agronomically biofortified wheat (Zincol-2016/NR-421) for improving Zn status in women in a low-resource setting in Pakistan: study protocol for a randomised, double-blind, controlled cross-over trial (BiZiFED). *BMJ Open.* 2018;8(5):e021364. doi:10.1136/bmjopen-2017-021364.
 51. Gannon B, Kaliwile C, Arscott SA, et al. Biofortified orange maize is as efficacious as a vitamin A supplement in Zambian children even in the presence of high liver reserves of vitamin A: a community-based, randomized placebo-controlled trial. *Am J Clin Nutr.* 2014;100(6):1541-1550. doi:10.3945/ajcn.114.086278.
 52. Talsma EF, Melse-Boonstra A, de Kok BP, et al. Biofortified cassava with pro-vitamin A is sensitively absorbed compared to white cassava. *Am J Clin Nutr.* 2013;98(1):123-130. doi:10.3945/ajcn.112.052746.

53. Van Jaarsveld PJ, Faber M, Tanumihardjo SA, Nestel P, Lombard CJ, Benadé AJ. β -carotene-rich orange-fleshed sweet potato improves the vitamin A status of primary school children assessed with the modified-relative-dose-response test. *Am J Clin Nutr.* 2005;81(5):1080-1087. doi:10.1093/ajcn/81.5.1080.
54. Low JW, Arimond M, Osman N, Cunguara B, Zano F, Tschirley D. A food-based approach introducing orange-fleshed sweet potatoes increased vitamin A intake and serum retinol concentrations in young children in rural Mozambique. *J Nutr.* 2007;137(5):1320-1327. doi:10.1093/jn/137.5.1320.
55. Arsenault JE, Yakes EA, Islam MM, et al. Very high vitamin A supplementation rates among women in Bangalore, India. *Public Health Nutr.* 2014;17(8):1873-1880. doi:10.1017/S1368980013002115.
56. Niles AM, Ejeta G, Mwangi AM, et al. Consumption of iron-biofortified sorghum alleviated iron deficiency in women and children in rural Kenya. *J Nutr.* 2021;151(1):146-153. doi:10.1093/jn/nxaa327.
57. Haas JD, Luna SV, Lung'aho MG, et al. Consuming iron biofortified beans increases iron status in Rwandan women after 128 days in a randomized controlled feeding trial. *J Nutr.* 2016;146(8):1586-1592. doi:10.3945/jn.115.224741.
58. Murray-Kolb LE, Takaiwa F, Goto F, Yoshihara T, Theil EC, Beard JL. Transgenic rice is a source of iron for iron-depleted rats. *J Nutr.* 2002;132(5):957-960. doi:10.1093/jn/132.5.957.
59. Haas JD, Beard JL, Murray-Kolb LE, del Mundo AM, Felix A, Gregorio GB. Iron-biofortified rice improves the iron stores of nonanemic Filipino women. *J Nutr.* 2005;135(12):2823-2830. doi:10.1093/jn/135.12.2823.
60. Jones KM, Specio SE. Nutrition knowledge, attitudes, and efficacy among Head Start teachers in the Mississippi Delta. *J Nutr Educ Behav.* 2000;32(5):289-296. doi:10.1016/s1499-4046(06)60172-1.
61. Rah JH, de Pee S, Kraemer K, et al. Programming of infant gut microbiota: influence of dietary intakes in rural Bangladesh. *Curr Dev Nutr.* 2016;1(2):e000348. doi:10.3945/cdn.116.000348.
62. Pfeiffer WH, McClafferty B. HarvestPlus: breeding crops for better nutrition. *Crop Sci.* 2007;47(Suppl3):S-88. doi: 10.2135/cropsci2007.09.0020IPBS.
63. Meenakshi JV, Banerji A, Manyong V, et al. Using a discrete choice experiment to elicit the demand for a nutritious food: Willingness-to-pay for orange maize in rural Zambia. *J Health Econ.* 2012;31(1):62-71. doi:10.1016/j.jhealeco.2011.12.005.
64. Birol E, Asare-Marfo D, Karandikar B, Roy D. A latent class approach to investigating farmer demand for biofortified staple food crops in developing countries: the case of high-iron pearl millet in Maharashtra, India. *HarvestPlus Working Paper.* 2011;(3).
65. Meenakshi JV, Johnson NL, Manyong VM, DeGroote H, Javelosa J, Yanggen DR, Naher F, Gonzalez C, García J, Meng E. How cost-effective is biofortification in combating micronutrient malnutrition? An ex ante assessment. *World Dev.* 2010;38(1):64-75. doi:10.1016/j.worlddev.2009.03.014.
66. Stein AJ, Meenakshi JV, Qaim M, Nestel P, Sachdev HP, Bhutta ZA. Analyzing the health benefits of biofortified staple crops by means of the Disability-Adjusted Life Years approach: a handbook focusing on iron, zinc and vitamin A. *HarvestPlus Technical Monograph 4.*

Washington, DC and Cali: International Food Policy Research Institute (IFPRI) and International Center for Tropical Agriculture (CIAT). 2005.

67. Bouis, H. E., & Saltzman, A. (2017). Improving nutrition through biofortification: A review of evidence from HarvestPlus, 2003 through 2016. *Global Food Security*, 12, 49-58.
68. Meenakshi, J. V., Johnson, N. L., Manyong, V. M., DeGroot, H., Javelosa, J., Yanggen, D. R., ... & Meng, E. (2010). How cost-effective is biofortification in combating micronutrient malnutrition? An ex ante assessment. *World development*, 38(1), 64-75.
69. De Moura, F. F., Lewis, K. D., Ffrench-Constant, R. H., & Lima, G. P. P. (2022). The past, present and future of biofortification—Fighting hidden hunger with staple food crops. *Plant Science*, 110875.
70. McLean, E., Cogswell, M., Egli, I., Wojdyla, D., & De Benoist, B. (2009). Worldwide prevalence of anaemia, WHO vitamin and mineral nutrition information system, 1993–2005. *Public health nutrition*, 12(4), 444-454.
71. Anuradha, K., Agarwal, S., Rao, Y. V., Rao, K. V., Viraktamath, B. C., Sarla, N., ... & Siddiq, E. A. (2012). QTL analysis for grain iron and zinc concentration in two RIL populations of sorghum. *Euphytica*, 184(2), 169-180.
72. Velu, G., Singh, R. P., Huerta-Espino, J., Peña, R. J., Arun, B., Mahendru-Dole, I., ... & Joshi, A. K. (2017). Genetic yield gains in CIMMYT's international semi-arid wheat yield trials from 1994 to 2014. *The Crop Journal*, 5(3), 172-177.
73. Petry, N., Egli, I., Gahutu, J. B., Tugirimana, P. L., Boy, E., & Hurrell, R. (2012). Phytic acid concentration influences iron bioavailability from biofortified beans in Rwandese women with low iron status. *The Journal of nutrition*, 142(4), 741-747.
74. Chomba, E., Mandal, R., Kumar, D., Grewal, J., & Anand, J. (2015). Improving iron, zinc and β -carotene content of fast-maturing biofortified pearl millet cultivar for India. *Crop Improvement*, 42 (1), 309-320.
75. Finkelstein, J. L., Mehta, S., Udipi, S. A., Ghugre, P. S., Luna, S. V., Wenger, M. J., ... & Haas, J. D. (2015). A randomized trial of iron-biofortified pearl millet in school children in India. *The Journal of Nutrition*, 145(7), 1576-1581.
76. Haas, J. D., Luna, S. V., Lung'aho, M. G., Wenger, M. J., Murray-Kolb, L. E., Beebe, S., ... & Finkelstein, J. L. (2016). Consuming iron biofortified beans increases iron status in Rwandan women after 128 days in a randomized controlled feeding trial. *The Journal of nutrition*, 146(8), 1586-1592.
77. Cercamondi, C. I., Egli, I. M., Mitchikpe, E., Tossou, F., Zeder, C., Hounhouigan, J. D., & Hurrell, R. F. (2013). Total iron absorption by young women from iron-biofortified pearl millet composite meals is double that from regular millet meals but less than that from post-harvest iron-fortified millet meals. *The Journal of nutrition*, 143(9), 1376-1382.
78. Kodkany, B. S., Bellad, R. M., Mahantshetti, N. S., Westcott, J. E., Krebs, N. F., Kemp, J. F., & Hambidge, K. M. (2013). Biofortification of pearl millet with iron and zinc in a randomized controlled trial increases absorption of these minerals above physiologic requirements in young children. *The Journal of nutrition*, 143(9), 1489-1493.
79. Finkelstein, J. L., Liese, A. D., Mehta, S., & John, J. A. (2017). Iron status affects zinc homeostasis in rural Bangladeshi children. *The Journal of nutrition*, 147(2), 135-141.

80. Cercamondi, C. I., Salami, S. D., Smoot, K., Hamedani, Z. H., Mitchikpe, C., Dossa, R. A., ... & Hurrell, R. F. (2018). Total iron absorption in young women from iron-biofortified pearl millet composite meals is double that from regular millet meals but less than that from post-harvest iron-fortified millet meals. *The Journal of Nutrition*, 148(2), 242-247.
81. Haas, J. D., Luna, S. V., Lung'aho, M. G., Wenger, M. J., Murray-Kolb, L. E., Beebe, S., ... & Finkelstein, J. L. (2016). Consuming iron biofortified beans increases iron status in Rwandan women after 128 days in a randomized controlled feeding trial. *The Journal of nutrition*, 146(8), 1586-1592.
82. Cercamondi, C. I., Egli, I. M., Mitchikpe, E., Tossou, F., Zeder, C., Hounhouigan, J. D., & Hurrell, R. F. (2013). Total iron absorption by young women from iron-biofortified pearl millet composite meals is double that from regular millet meals but less than that from post-harvest iron-fortified millet meals. *The Journal of nutrition*, 143(9), 1376-1382.
83. Kodkany, B. S., Bellad, R. M., Mahantshetti, N. S., Westcott, J. E., Krebs, N. F., Kemp, J. F., & Hambidge, K. M. (2013). Biofortification of pearl millet with iron and zinc in a randomized controlled trial increases absorption of these minerals above physiologic requirements in young children. *The Journal of nutrition*, 143(9), 1489-1493.
84. Wessells, K. R., & Brown, K. H. (2012). Estimating the global prevalence of zinc deficiency: results based on zinc availability in national food supplies and the prevalence of stunting. *PloS one*, 7(11), e50568.
85. Velu, G., Ortiz-Monasterio, I., Cakmak, I., Hao, Y., & Singh, R. P. (2014). Biofortification strategies to increase grain zinc and iron concentrations in wheat. *Journal of Cereal Science*, 59(3), 365-372.
86. Aprety, J., & Zingore, S. (2017). Delivering nutritional zinc enrichment in staple food crops: lessons learned. IZA Policy Paper No. 125. International Zinc Association, Durham, NC, USA.
87. Palmgren, M. G., Edenbrandt, A. K., Vedel, S. E., Andersen, M. M., Landes, X., Østerberg, J. T., ... & Sandøe, P. (2015). Are we ready for back-to-nature crop breeding?. *Trends in plant science*, 20(3), 155-164.
88. Booroolugn, V., Rajaselvam, R., Halwatura, R., Sivakanesan, R., Anantharasan, V., & Govindaraj, M. (2022). Biofortified lentil influences serum zinc status in Indian women and children - A feeding study. *Clinical Nutrition*, 41(4), 1031-1038.
89. De Moura, F. F., Moursi, M., Ayele, Z., Angeles-Agdeppa, I., Gollin, D., Ramos, J. P., & Mishra, K. (2016). Biofortified orange maize enhances β -cryptoxanthin concentrations in egg yolks of laying hens better than tangerine peel fortificant. *Journal of Agricultural and Food Chemistry*, 64(47), 8929-8935.
90. Palupi, E., Schreinemachers, P., Riwu, K., Daryanto, A., Burger, K., & Waldron, S. (2019). Is there nutritional synergy arising from combining biofortified wheat and nutritious vegetables and pulses? Evidence from a field experiment in Indonesia. *Public Health Nutrition*, 22(18), 3386-3399.
91. Booroolugn, V., Rajaselvam, R., Halwatura, R., Sivakanesan, R., Anantharasan, V., & Govindaraj, M. (2022). Biofortified lentil influences serum zinc status in Indian women and children - A feeding study. *Clinical Nutrition*, 41(4), 1031-1038.
92. De Moura, F. F., Moursi, M., Ayele, Z., Angeles-Agdeppa, I., Gollin, D., Ramos, J. P., & Mishra, K. (2016). Biofortified orange maize enhances β -cryptoxanthin concentrations in egg

yolks of laying hens better than tangerine peel fortificant. *Journal of Agricultural and Food Chemistry*, 64(47), 8929-8935.

93. Palupi, E., Schreinemachers, P., Riwu, K., Daryanto, A., Burger, K., & Waldron, S. (2019). Is there nutritional synergy arising from combining biofortified wheat and nutritious vegetables and pulses? Evidence from a field experiment in Indonesia. *Public Health Nutrition*, 22(18), 3386-3399.
94. Booroolugn, V., Rajaselvam, R., Halwatura, R., Sivakanesan, R., Anantharasan, V., & Govindaraj, M. (2022). Biofortified lentil influences serum zinc status in Indian women and children - A feeding study. *Clinical Nutrition*, 41(4), 1031-1038.
95. Birol, E., Meenakshi, J.V., Oparinde, A., Perez, S., & Tomlins, K. (2015). Developing country consumers' acceptance of biofortified foods: a synthesis. *Food Security*, 7(3), 555-568.
96. Luna, S. V., Pompano, L. M., Haas, J. D., Bede, J., Rury, L. M., Beebe, S., ... & Finkelstein, J. L. (2022). Iron and zinc absorption from biofortified beans and behavioral influences on adoption and consumption in Rwanda. *Current Developments in Nutrition*, 6(Supplement_2), 528.
97. Andersson, M., Saltzman, A., Virk, P. S., & Pfeiffer, W. (2017). Progress update: Crop development of biofortified staple food crops under HarvestPlus. *African Journal of Food, Agriculture, Nutrition and Development*, 17(2), 11905-11935.
98. La Frano, M. R., de Moura, F. F., Boy, E., & Burri, B. J. (2014). Bioavailability of iron, zinc, and provitamin A carotenoids in biofortified staple crops. *Nutrition Reviews*, 72(5), 289-307.
99. Singh, B. P., Arora, S., Kumar, P., & Bhandari, G. (2018). Biofortification strategies to combat iron-deficiency anaemia. *Comprehensive Reviews in Food Science and Food Safety*, 17(6), 1297-1316.
100. Phan, A. Q. T., Berti, M., Liedl, K. R., & Schwartz, S. J. (2021). Zinc bioavailability in foods and model diets. *Molecular Aspects of Medicine*, 76, 100970.
101. Palupi, E., Schreinemachers, P., Riwu, K., Daryanto, A., Burger, K., & Waldron, S. (2019). Is there nutritional synergy arising from combining biofortified wheat and nutritious vegetables and pulses? Evidence from a field experiment in Indonesia. *Public Health Nutrition*, 22(18), 3386-3399.
102. Finkelstein, J. L., Mehta, S., Brittenham, G. M., & John, J. A. (2022). Providing iron-biofortified pearl millet flour to secondary school children in rural India: a cluster-randomized effectiveness trial of an integrated agriculture-nutrition intervention. *Current Developments in Nutrition*, 6(Supplement_2), 528.
103. Saltzman, A., Birol, E., Bouis, H. E., Boy, E., De Moura, F. F., Islam, Y., & Pfeiffer, W. H. (2013). Biofortification: progress toward a more nourishing future. *Global Food Security*, 2(1), 9-17.
104. Faber, M., Laurie, S., Venter, S., & Dhansay, M. A. (2014). Home gardens to address vitamin A deficiency in South Africa: a food-based approach. Pretoria: South African Medical Research Council.
105. Oparinde, A., Banerji, A., Birol, E., & Ilona, P. (2016). Information and consumer willingness to pay for biofortified yellow cassava: evidence from experimental auctions in Nigeria. *HarvestPlus Working Paper*, (3).

106. Birol, E., Meenakshi, J. V., Oparinde, A., Perez, S., & Tomlins, K. (2015). Developing country consumers' acceptance of biofortified foods: a synthesis. *Food Security*, 7(3), 555-568.
107. Malapit, H.J., Quisumbing, A.R., Meinzen-Dick, R.S., Seymour, G., Martinez, E.M., Heckert, J. and Rubin, D., 2019. Development of the project-level Women's Empowerment in Agriculture Index (pro-WEAI) (No. 1974). International Food Policy Research Institute (IFPRI).
108. Saltzman, A., Birol, E., Bouis, H. E., Boy, E., De Moura, F. F., Islam, Y., & Pfeiffer, W. H. (2013). Biofortification: progress toward a more nourishing future. *Global Food Security*, 2(1), 9-17.
109. Ruel, M. T., Alderman, H., & Maternal and Child Nutrition Study Group. (2013). Nutrition-sensitive interventions and programmes: how can they help to accelerate progress in improving maternal and child nutrition?. *The lancet*, 382(9891), 536-551.
110. De Steur, H., Muehlhoff, E., Foley, C., & Gellynck, X. (2020). The potential of multistakeholder partnerships to promote the sustainability of voluntary biofortification programmes: case study evidence from the vitamin A maize programme in Zambia. *Nutrients*, 12(5), 1264.
111. Saltzman, A., Birol, E., Oparinde, A., Andersson, M. S., Asare-Marfo, D., Diressie, M. T., ... & Zeller, M. (2017). Availability, production, and consumption of crops biofortified by plant breeding: current evidence and future potential. *Annals of the New York Academy of Sciences*, 1390(1), 104-114.
112. Pfeiffer, W. H., & McClafferty, B. (2007). HarvestPlus: Breeding crops for better nutrition. *Crop science*, 47(Supplement_3), S-88.
113. Singh, B. P., Arora, S., Kumar, P., & Bhandari, G. (2018). Biofortification strategies to combat iron-deficiency anaemia. *Comprehensive Reviews in Food Science and Food Safety*, 17(6), 1297-1316.
114. Zhu, C., Naqvi, S., Gomez-Galera, S., Pelacho, A. M., Capell, T., & Christou, P. (2007). Transgenic strategies for the nutritional enhancement of plants. *Trends in plant science*, 12(12), 548-555.