

Original Research Article
Correlation studies in Ber (*Ziziphus mauritiana* Lamk.) in Eastern region of Uttar Pradesh

Abstract

The present investigation was carried out in thirty genotype of ber at department of Fruit Science, college of Horticulture, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur (UP) during 2021-22 to 2022-23. Phenotypic correlations were worked out among eighteen characters of ber to know the nature of association existing among the characters. Correlation among the quantitative characters was worked out. Length of fruit was found significantly and positively correlated with fruit weight (0.617**), the width of fruit analysis exhibited significant positive correlation with fruit weight (0.853**). Leaf length showed significant positive correlations with length of fruit(0.443*) and non-significant but positive correlation with weight of fruit and width of fruit, Leafwidth was showed significant positive correlations with fruit weight(0.794**), length of fruit (0.563**) leaf length (0.487*). Stone weight has showed significant positive correlations with fruit weight(0.626**) width of fruit (0.874**), Stone length has showed significant positive correlations with fruit weight (0.440*) leaf width (0.409*) Stone weight(0.584**). Stone width has showed significant positive correlations with leaf length (0.417*) leaf width (0.701**) stone weight(0.487*). Hight of tree has showed significant positive correlations with length of fruit (0.576**) width of fruit (0.977**) stone length (0.803**) stone width (0.628**). The stem girth also exhibited significant positive correlation with fruit weight (0.966**), width of fruit (0.496*) leaf length (0.650**) stone weight (0.397*) hight of tree (0.406*). Specific gravity has showed significant positive correlations with fruitweight (0.817**) leaf length (0.641**) leaf width (0.424*) stone weight (0.455*), stone length (0.417*) stone width (0.734**).

TSS content was also found significantly and positively correlated with fruit weight (0.774**) leaf length (0.645**) leaf width (0.970**) stone weight (0.539**), stone length (0.425*) stone width (0.765**) stem girth (0.855**) specific gravity (0.547**). Total acidity content in fruit was found significantly and positively correlated with fruit weight (0.725**) length of fruit (0.882**) width of fruit (0.948**) leaf length (0.809**) leaf width (0.989**) stone weight (0.772**), stone length (0.605**) stone width (0.892**) hight of tree (0.792**).

Commented [WU1]: Abstract should be brief and simple and should be less than 250 words

Commented [WU2]: genotypes

Commented [WU3]: follow similar terminology fruit weight

Commented [WU4]: check the grammatical mistakes, full stops, capital letters and spacings in the entire document.

specific gravity (0.702**) .TSS: acid ratio has showed significant positive correlations with fruit weight (0.981**) length of fruit (0.455*) width of fruit (0.963**) leaf length (0.929**) leaf width (0.854**) stone length (0.813**) stone width (0.727**) height of tree (0.832**) stem girth (0.453*) specific gravity (0.728**). Reducing sugar has showed significant positive correlations with fruit weight (0.680**) leaf width (0.821**) stone length (0.835**) stone width (0.503*) height of tree (0.747**) stem girth (0.812**) TSS acidity ratio (0.410*). Non-reducing sugar has showed significant positive correlations with fruit weight (0.782**) length of fruit (0.999**) width of fruit (0.499*) leaf width (0.482*) stone weight (0.662**) stone length (0.399*) stone width (0.453*) specific gravity (0.488*) tss (0.787**). Total sugar has showed significant positive correlations with fruit weight (0.960**) length of fruit (0.412*) leaf length (0.539**) leaf width (0.710**) stone length (0.745**) stone width (0.897**) height of tree (0.658**) stem girth (0.841**) acidity (0.584**). Ascorbic acid has showed significant positive correlations with length of fruit (0.938**) leaf width (0.642**) stone length (0.664**) stone width (0.685**) Height of tree (0.400*) specific gravity (0.942**).

Keywords: Ber, Correlation Coefficient, Variability, Character Association.

Introduction

The expression of a particular character is an aggregate of complex contribution of so many other characters. In tree improvement programme, a clear understanding of the nature and degree of association among different traits is of great importance because the choice of one character can favour the appearance or disappearance of the other. Correlation, an important statistical tool, helps in determining such associations among different factors under consideration.

Ber (*Zizyphus mauritiana* Lam.) an indigenous important fruit crop for arid and semi-arid regions of India belong to the family Rhamnaceae. It is mainly grown in India and other countries in central Asia, China and Taiwan and is more associated with the Indian culture since ancient times (Balodaei *et al.*, 2012). It is cultivated widely for its resistance to grow in drought and other diversified soil and climatic conditions. It is a hardy tree that tolerates extreme temperature and thrives under dry conditions. Fruit quality is best under hot, sunny and dry conditions, but there should be a rainy season to support growth and flowering, leaving enough soil moisture to carry the fruit to maturity (Awasthi and More, 2009). Along with the ability to withstand drought, ber can also tolerate adverse conditions, such as salinity and water logging. In view of the recent development in production technology of

Commented [WU5]: Write capital letters

Commented [WU6]: Check spelling and spacing

Commented [WU7]: spacing

this crop, the cultivation of ber is becoming increasingly popular in many parts of country. It is an ideal fruit tree for tropical and subtropical regions of the country viz., Rajasthan, Haryana, Punjab, U.P, Gujarat, M.P., Bihar, Maharashtra, Andhra Pradesh and Tamil Nadu. Ber fruit have both sour and sweet tastes and are rich in nutritive value. It is rich in vitamin 'B' complex and vitamin 'C' (Pareek, 2013). However, ber is richer than apple in protein, phosphorus, calcium, carotene and vitamin C and than oranges in phosphorus, iron, vitamin C and carbohydrates and exceeds them in calorific value (Nissi and Vinayak, 2016).

Commented [WU8]: write in full form

Materials and Method

Commented [WU9]: check

Commented [WU10]: nissi et al

Commented [WU11]: methods

The present investigation was carried out at Department of Fruit Science, college of Horticulture, Chandra Shekhar Azad University of Agriculture & Technology, Kanpur (UP) during 2021-2022 and 2022 - 2023. Fruit of forty genotypes were collected from eastern region (Balrampur, Bahraich, Sidarthnagar, Shravasti, Gonda, Basti, Ayodhya, SultanpurBanaras and Allahabad district) of Utter Pradesh. Twenty fruits of ber were randomly selected from all the direction of trees from each site collected then kept into bags and tagged by the number and subjected to physico-chemical analysis in laboratory. Physical parameters like stone weight was measure by electronic weighting machine, height of tree and stem girth were estimated with the help of measuring tape data on fruit shape, fruit apex, fruit base, mature fruit colour, immature fruit colour, tree shape, leaf shape, leaf apex, leaf margin, Stone shape, stone apex, stone base, leaf colour, thorn shape, pulp colour and , pulp texture were observed using standard and recommended techniques.

Commented [WU12]: in the abstract you have mentioned thirty genotypes and here you mentioned forty genotypes

Commented [WU13]: spacing

Results and Discussion

Phenotypic correlations were worked out among 40 characters of ber to know the nature of association existing among the characters. The results are presented and pertaining to correlation at phenotypic levels between fruit weight and other related characters are presented here.

Commented [WU14]: mentioned only 18 characters

Length of fruit was found significantly and positively correlated with fruit weight (0.617**), the width of fruit analysis exhibited significant positive correlation with fruit weight (0.853**), Leaf length showed significant positive correlations with length of fruit (0.443*) and non-significant but positive correlation with weight of fruit and width of fruit, Leafwidth was showed significant positive correlations with fruit weight (0.794**), length of fruit (0.563**) leaf length (0.487*), Verma *et al.* (2002) Kevadiya *et al.* (2006) Rathod *et al.* (2007) Patel *et al.* (2015) Singh *et al.* (2015) Patel *et al.* (2016) Tatari *et al.* (2016) Dolkar *et*

al. (2017) Wangchu *et al.* (2017) Kumar *et al.* (2020) Kumar *et al.* (2021) Siddique *et al.* (2022) Panthorat. *al.* (2023) Dong *et al.* (2023) Ahmad *et al.* (2023)

Commented [WU15]: check mentioned as 2013 in the references

Commented [WU16]: not found in the references

Stone weight has showed significant positive correlations with fruit weight (0.626**), width of fruit (0.874**), Stone length has showed significant positive correlations with fruit weight (0.440*) leaf width (0.409*) Stone weight (0.584**), Stonewidth has showed significant positive correlations with leaf length (0.417*) leaf width (0.701**) stone weight (0.487*). Verma *et al.* (2002) Kevadiya *et al.* (2006) Rathod *et al.* (2007) Patel *et al.* (2015) Tatari *et al.* (2016) Dolkar *et al.* (2017) Wangchu *et al.* (2017) Gaikwad *et al.* (2017) Arivazhagan *et al.* (2019) Kumar *et al.* (2021) Kumar *et al.* (2021) Siddique *et al.* (2022) Sheikh *et al.* (2023) Panthorat. *al.* (2023) Dong *et al.* (2023) Ahmad *et al.* (2023)

Commented [WU17]: mention 2021a and 2021b

Hight of tree has showed significant positive correlations with length of fruit (0.576**) width of fruit (0.977**) stone length (0.803**) stone width (0.628**). The stem girth also exhibited significant positive correlation with fruit weight (0.966**), width of fruit (0.496*) leaf length (0.650**) stone weight (0.397*) hight of tree (0.406*). Specific gravity has showed significant positive correlations with fruit weight (0.817**) leaf length (0.641**) leaf width (0.424*) stone weight (0.455*), stone length (0.417*) stone width (0.734**). Verma *et al.* (2002) Kevadiya *et al.* (2006) Rathod *et al.* (2007) Patel *et al.* (2015) Singh *et al.* (2015) Patel *et al.* (2016) Tatari *et al.* (2016) Dolkar *et al.* (2017) Wangchu *et al.* (2017) Gaikwad *et al.* (2017) Kumar *et al.* (2021) Siddique *et al.* (2022) Sheikh *et al.* (2023) Panthorat. *al.* (2023) Dong *et al.* (2023)

Commented [WU18]: what is the need of writing the references. Describe briefly

Commented [WU19]: not found in references

TSS content was also found significantly and positively correlated with fruit weight (0.774**) leaf length (0.645**) leaf width (0.970**) stone weight (0.539**), stone length (0.425*) stone width (0.765**) stem girth (0.855**) specific gravity (0.547**). Total acidity content in fruit was found significantly and positively correlated with fruit weight (0.725**) length of fruit (0.882**) width of fruit (0.948**) leaf length (0.809**) leaf width (0.989**) stone weight (0.772**), stone length (0.605**) stone width (0.892**) hight of tree (0.792**) specific gravity (0.702**). TSS: acid ratio has showed significant positive correlations with fruit weight (0.981**) length of fruit (0.455*) width of fruit (0.963**) leaf length (0.929**) leaf width (0.854**) stone length (0.813**) stone width (0.727**) hight of tree (0.832**) stem girth (0.453*) specific gravity (0.728**). Verma *et al.* (2002) Kevadiya *et al.* (2006) Rathod *et al.* (2007) Patel *et al.* (2015) Singh *et al.* (2015) Patel *et al.* (2016) Tatari *et al.* (2016) Dolkar *et al.* (2017) Wangchu *et al.* (2017) Gaikwad *et al.* (2017) Arivazhagan *et al.* (2019) Kumar *et al.* (2021) Kumar *et al.* (2021) Siddique *et al.* (2022) Sheikh *et al.* (2023) Panthorat. *al.* (2023) Dong *et al.* (2023) Ahmad *et al.* (2023)

Commented [WU20]: not found in references

Commented [WU21]: describe briefly

Reducing sugar has showed significant positive correlations with fruit weight (0.680**) leaf width (0.821**) stone length (0.835**) stone width (0.503*) hight of tree (0.747**) stem girth (0.812**) TSS acidity ratio (0.410*), Non-reducing sugar has showed significant positive correlations with fruit weight (0.782**) length of fruit (0.999**) width of fruit (0.499*) leaf width (0.482*) stone weight (0.662**) stone length (0.399*) stone width (0.453*) specific gravity (0.488*) tss (0.787**), Total sugar has showed significant positive correlations with fruit weight (0.960**) length of fruit (0.412*) leaf length (0.539**) leaf width (0.710**) stone length (0.745**) stone width (0.897**) hight of tree (0.658**) stem girth (0.841**) acidity (0.584**) and Ascorbic acid has showed significant positive correlations with length of fruit (0.938**) leaf width (0.642**) stone length (0.664**) stone width (0.685**) Hight of tree (0.400*) specific gravity (0.942**). Verma *et al.* (2002) Kevadiya *et al.* (2006) Rathod *et al.* (2007) Patel *et al.* (2015) Singh *et al.* (2015) Patel *et al.* (2016) Tatari *et al.* (2016) Dolkar *et al.* (2017) Wangchu *et al.* (2013) Gaikwad *et al.* (2017) Arivazhagan *et al.* (2019) Kumar *et al.* (2021) Kumar *et al.* (2021) Siddique *et al.* (2022) Sheikh *et al.* (2023) Panthorat. *al.* (2023) Dong *et al.* (2023) Ahmad *et al.* (2023)

Commented [WU22]: check the spelling

CONCLUSION

Length of fruit was found significantly and positively correlated with fruit weight. The width of fruit analysis exhibited significant positive correlation with fruit weight. Leaf length showed significant positive correlations with length of fruit and non-significant but positive correlation with weight of fruit and width of fruit. Leaf width was showed significant positive correlations with fruit weight, length of fruit and leaf length. Stone weight has showed significant positive correlations with fruit weight and width of fruit. Stone length has showed significant positive correlations with fruit weight, leaf width and Stone weight. Stone width has showed significant positive correlations with leaf length, leaf width and stone weight. Hight of tree has showed significant positive correlations with length of fruit, width of fruit, stone length and stone width. The stem girth also exhibited significant positive correlation with fruit weight, width of fruit, leaf length, stone weight and hight of tree. Specific gravity has showed significant positive correlations with fruit weight, leaf length, leaf width, stone weight, stone length and stone width.

TSS content was also found significantly and positively correlated with fruit weight, leaf length, leaf width, stone weight, stone length, stone width, stem girth and specific gravity.

Total acidity content in fruit was found significantly and positively correlated with fruit weight, length of fruit, width of fruit, leaf length, leaf width, stone weight, stone length, stone width, height of tree and specific gravity. TSS: acid ratio has showed significant positive correlations with fruit weight, length of fruit, width of fruit, leaf length, leaf width, stone length, stone width, height of tree, stem girth and specific gravity. Reducing sugar has showed significant positive correlations with fruit weight, leaf width, stone length, stone width, height of tree, stem girth and TSS acidity ratio. Non-reducing sugar has showed significant positive correlations with fruit weight, length of fruit, width of fruit, leaf width, stone weight, stone length, stone width, specific gravity and TSS. Total sugar has showed significant positive correlations with fruit weight, length of fruit, leaf length, leaf width, stone length, stone width, height of tree, stem girth and acidity. Ascorbic acid has showed significant positive correlations with length of fruit, leaf width, stone length, stone width, Height of tree, specific gravity.

Commented [WU23]: spelling

Reference:

Ahmad R., Ali H. M., Lisek A., Mosa W. F. A., Ercisli S. and Anjum M. A. (2023). Correlation among some phenological and biochemical traits in date palm (*Phoenix dactylifera* L.) germplasm. *Frontiers in Plant Science*. **35**, 1–16.

Arivazhagan, E., Kandasamy, R., and Ramadoss, N. 2019. Variability and Correlation Analysis in Sapota (*Manilkara Sapota*) Under Coastal Ecosystem. *Plant Archives Vol. 19 No. 1*, pp. 652-654.

Awasthi, O.P. and More, T.A (2009). Genetic Diversity and Status of *Ziziphus* in India. *Acta Hort.*; 840:33-40

Baloda S., Sehrawat S.K., Yadav B.S., Ahlawat V.P., and Singh (2012). Present status of ber production and future thrusts in India - a review. *Agri. Reviews*, 33 (3): 256- 264.

Baloda S., Sehrawat S.K., Yadav B.S., Ahlawat V.P., and Singh (2012). Present status of ber production and future thrusts in India - a review. *Agri. Reviews*; 33 (3): 256- 264.

Commented [WU24]: Repeated above

Debbarma, Pand Hazarika, T, K: (2023). Genetic diversity of Bael (*Aegle marmelos* (L.) Corr.) accessions from north-east India based on principal component and cluster analysis. *Genet Resour Crop Evol.* **71**:253–277.

Commented [WU25]: not found in the article

Dolkar, T., Sharma, M.K., Kumar A., Mir M.S., Hussain, S. 2017. Genetic variability and correlation studies in grapes (*Vitis vinifera* L.) in Leh District of Jammu and Kashmir. *Adv. Hort. Sci.* **31**(4): 241-247

Dong Yi, Sun W., Yue Z., Gong B., Yang Xu, Wu K., and Xu Y. (2023). Phenotypic

Diversity and Relationships of Fruit Traits in Persimmon (*Diospyros kaki* Thunb.). *Germplasm Resources Agriculture*. 13: 1804

Gaikwad, K.A., Patil, S.R., Nagre, P.K., Potdukhe, N.R., Paithankar, D.H., Gahukar, S.J. and Umbarkar, P.S. (2017). Correlation studies of different physic morphological characters in citrus rootstock genotypes with fruit yield under Vidarbha region. *International Journal of Chemical Studies*. 5(4): 1421-1425

Kevadiya, P. K. (2006). Variability, correlation and path analysis for yield and fruit characters in mango. M.Sc. (Agri.) thesis, *Anand Agricultural University, Anand*.

Kumar P., Prakash Om, Srivastava A.K., Thakur N., Chugh V and Singh R. S., (2021). Correlation Studies in Wood Apple (*Feronia limonia* L.) in Bundelkhand Region of Uttar Pradesh. *Ind. J. Pure App. Biosci.* 9(1), 316-321

Kumar, P., Srivastava A.K., Prakash Om, Thakur, N. and Singh P (2021). Germplasm Survey, Collection and Evaluation of ber (*Ziziphus mauritiana* Lamk) under Bundelkhand Region of Uttar Pradesh. *Ind. J. Pure App. Biosci.* 9(1), 259-266.

Nissi F Godi, Vinayak R Joshi, Vinay S Supe (2016). Physical fruit characteristics assessment of selected Ber (*Zizyphus mauritiana* Lamk.) Genotypes. *Int J Appl Res*; 2(2): 757-761.

Panase VG, Sukhatme PV. Statistical Methods for Agricultural workers, 2nd ed. New Delhi, ICAR. 1967, 99-116.

Commented [WU26]: Not found in article

Pareek, O.P. (2013) Fruit of the Future Ber. International centre for Underutilization crops, University of Southampton, Southampton UK.

Patel, M.C., Patel, D.A., Patel, K.V., Soni, N.V., Satodiya, B.N. and Jadav R.G. (2016). Genetic variability and correlation studies for fruit yield and quality parameters in mango (L.) *Green Farming* 3: 706-709.

Patel, R.K., Maiti, C.S., Deka, C.B., Deshmukh, N.A., Verma, V.K. and Nath, A. (2015). Physical and biochemical change in guava during various stage of fruit growth and development. *Int.J. Agri. Environ. Biotech.* 8(1):63-70.

Rathod, B. P. (2007). Genetic variability, correlation, path coefficient and D analysis for morphological and biochemical parameters of mango fruit. M.Sc. (Agri.) thesis, Navsari Agricultural University, Navsari.

Sheikh, A, A., Kumar, R., Bakshi, P., Sharma, R., Reena, Kumar, V., Sharma, N., Raina, V and Sinha, B, K. (2023): Genetic diversity analysis of loquat (*Eriobotrya japonica* Lind L.) Germplasm through morphological traits. *The Pharma Innovation*

Commented [WU27]: Not found in article

Journal.,12,5020-5026.

Siddique I. M., Sharif N., Naseer M., Khan M. N, Noreen A. Asif M., Faiz H., Aslam K. And Mehreen N. (2022). A morpo-biochemical perspective on sixteen promising sweet orange (*Citrus sinensis* (L.) Osbeck) cultivars *Plant Cell Biotechnology and Molecular Biology* 23(13&14):69-87.

Singh S. K., Singh A., Nath V., Parthasarathy VA., Sthapit B., Rajan .S and Vinoth.,(2015). Genetic Diversity in Seedling Populations of Mango. *Indian J. Plant Genet. Resour.* 28(1): 123-131.

Tatari, M., Ghasemi, A. and Mousavi, A. (2016). Genetic diversity in Jujube germplasm (*Ziziphus jujuba* Mill.) based on morphological and pomological traits in Isfahan province, Iran. *Crop Breeding Journal.* 4, 5 and 6 (2; 1 and 2): 79-85.

Verma, S.K., Singh, R.K. and Arya, P.R. (2002). Variability and correlation studies in strawberry germplasm for quantitative traits. *Indian J. Hort.* 59(1):39-43.

Wangchu LD. Singh, Mitra S.K., (2013). Studies on the diversity and selection of superior types in jackfruit (*Artocarpus heterophyllus* Lam.). *Genet. Resour. Crop Evol.*; 60:1749-1762.

UNDER PEER REVIEW

table 1 : Result of Correlation analysis

Commented [WU28]: Not mentioned in the results

	Fruit weight (g.)	Length of Fruit (cm.)	Width of Fruit (cm.)	Leaf Length (cm.)	Leaf Width (cm.)	Stone Weight (g.)	Stone length (cm.)	Stone Width (cm.)	Height of Tree (m.)	Stem girth (cm.)	Specific gravity (g/cm ³)	TSS (Brix)	Acidity (%)	TSS/ Acidity ratio	Reducing sugar (%)	Non reducing sugar (%)	Total Sugar (%)	Ascorbic Acid (mg/100g)
Fruit weight(g.)		0.743**	0.844**	0.109	0.818**	0.446*	0.519**	0.164	0.110	0.967**	0.856*	0.821**	0.466*	0.877**	0.623**	0.974*	0.628*	0.285
Length of Fruit (cm.)			0.149	0.324	0.846**	0.294	0.014	0.140	0.788**	0.185	0.095	0.138	0.606**	0.459*	0.334	0.679*	0.940*	0.767**
Width of Fruit (cm.)				0.318	0.473*	0.859*	0.015	0.171	0.291	0.526**	0.064	0.135	0.765**	0.827**	0.117	0.430*	0.048	0.336
Leaf Length (cm.)					0.341	0.140	0.301	0.524**	0.348	0.631**	0.589*	0.572**	0.683**	0.974**	0.036	0.340	0.404*	0.048
Leaf Width (cm.)						0.363	0.420*	0.575**	0.259	0.089	0.593*	0.947**	0.897**	0.753**	0.914**	0.649*	0.859*	0.653**
Stone Weight (g.)							0.600**	0.710**	0.442*	0.624**	0.586*	0.568**	0.513**	0.168	0.208	0.675*	0.291	0.104
Stone length (cm.)								0.141	0.649**	0.043	0.810*	0.255	0.755**	0.794**	0.808**	0.218	0.296	0.676**
Stone Width (cm.)									0.471*	0.103	0.664*	0.961**	0.599**	0.976**	0.878**	0.146	0.551*	0.716**
Height of Tree (m.)										0.384	0.251	0.322	0.978**	0.920**	0.924**	0.478*	0.582*	0.300
Stem girth (cm.)											0.214	0.821**	0.261	0.420*	0.832**	0.322	0.521*	0.171
Specific gravity (g/cm ³)												0.787**	0.951**	0.576**	0.282	0.533*	0.759*	0.992
TSS (Brix)													0.045	0.009	0.049	0.923*	0.177	0.018
Acidity (%)														0.000	0.231	0.270	0.251	0.550**
TSS/ Acidity ratio															0.338	0.188	0.284	0.014
Reducing sugar (%)																0.002	0.000	0.243
Non reducing sugar (%)																	0.000	0.077
Total Sugar (%)																		0.076

Significance 0.05- (0.396) * and Highly significance 0.01 (0.505) **

	Fruit weight(g.)	Length of Fruit (cm.)	Width of Fruit (cm.)	Leaf Length (cm.)	Leaf Width (cm.)	Stone Weight (g.)	Stone length (cm.)	Stone Width (cm.)	Hight of Tree (m.)	Stem girth (cm.)	Specific gravity	TSS (Brix)	Acidity (%)	TSS/ Acidity ratio	Reducing sugar (%)	Non reducing sugar (%)	Total Sugar (%)	Ascorbic Acid (mg/100 g)
Fruit weight		0.617*	0.853**	0.092	0.794**	0.626*	0.440*	0.179	0.092	0.966**	0.817*	0.774**	0.725*	0.981*	0.680*	0.782*	0.960**	0.279
Length of Fruit			0.240	0.443*	0.563**	0.296	0.000	0.138	0.576**	0.160	0.075	0.167	0.882*	0.455*	0.173	0.999*	0.412*	0.938**
Width of Fruit (cm.)				0.264	0.233	0.874*	0.026	0.217	0.977**	0.496*	0.060	0.037	0.948*	0.963*	0.062	0.499*	0.094	0.173
Leaf Length (cm.)					0.487*	0.155	0.353	0.417*	0.378	0.650**	0.641*	0.645**	0.809*	0.929*	0.048	0.273	0.539**	0.062
Leaf Width (cm.)						0.352	0.409*	0.701**	0.286	0.072	0.424*	0.970**	0.989*	0.854*	0.821*	0.482*	0.710**	0.642**
Stone Weight (g.)							0.584**	0.487*	0.286	0.397*	0.455*	0.539**	0.772*	0.122	0.217	0.662*	0.225	0.115
Stone length (cm.)								0.076	0.803**	0.042	0.417*	0.425*	0.605*	0.813*	0.835*	0.399*	0.745**	0.664**
Stone Width (cm.)								0.628**	0.079	0.734*	0.765**	0.892*	0.727*	0.503*	0.453*	0.897**	0.685**	
Hight of Tree (m.)									0.406*	0.154	0.364	0.792*	0.832*	0.747*	0.230	0.352	0.400*	
Stem girth (cm.)										0.261	0.855**	0.310	0.453*	0.812*	0.350	0.658**	0.161	
Specific gravity											0.547**	0.702*	0.728*	0.222	0.488*	0.841**	0.942**	
TSS (Brix)												0.051	0.000	0.048	0.787*	0.249	0.017	
Acidity (%)													0.000	0.268	0.170	0.584**	0.254	
TSS/ Acidity ratio														0.410*	0.221	0.387	0.009	
Reducing sugar															0.002	0.00	0.245	
Non reducing sugar (%)																0.00	0.064	
Total Sugar (%)																	0.071	

Significance 0.05- (0.396) * and Highly significance 0.01 (0.505) **

table 3 : Pooled Corrilation analysis

UNDER PEER REVIEW

