

“A Review on Effectsof Biofertilizers and Bio Capsules on Vegetative growth, Flowering and Fruiting attributes of Strawberry”

Abstract:-

This survey paper analyzes the impact of biofertilizers and bio containers on the key improvement phases of strawberry plants, zeroing in on vegetative development, blossoming, and fruiting qualities. The paper gives an extensive investigation of the current writing, examining the systems through which biofertilizers and bio containers influence strawberry plants. Critical evaluation is given to pertinent studies that investigate the application strategies, microbial strains involved, and their effects on various physiological and biochemical processes. Today, organic foods are renowned for their universal health benefits. Natural product culture in a natural manner is likewise picking up speed. Delivering natural organic products has extended rapidly throughout recent years because of the appeal it in both the homegrown and worldwide business sectors, which has forestalled medical conditions welcomed on by pesticides and perilous synthetics. This survey sums up the various techniques for the development of strawberries utilizing bio-composts and their effect on plant development, yield, and quality.

Keywords: Bio-fertilizer, Bio-Capsule, Vegetative Growth, Flowering, Fruiting.

Introduction

Strawberry (*Fragaria × ananassa* Duch.), a total product of Rosaceae family, possesses a huge spot in organic product industry, since it is developed in both in fields as well as in slopes. It is herbaceous yield with prostate development propensity, which acts as a yearly in sub-tropical area and enduring in temperature locale and has acquired the situation with being one of the main delicate product of the world after grape [31]. Strawberries are one of the main berry organic products. They are generally filled on the planet and are affectionately consumed by everybody. Strawberries are crucial for well-being as they contain anticancer parts, for example, ellagic

corrosive, and are wealthy in normal cell reinforcements. Strawberries have more huge degrees of L-ascorbic corrosive, phenolics, and flavonoids than other berry natural items.[8,11].Strawberry is one of the significant delectable and delicate organic product among the berries. Different strawberry species fill wild in everywhere, except the developed strawberry depends on two American species *Fragaria chiloensis* and *Fragaria virginiana*. Cross breed between these two species was the progenitor of all the cutting edge strawberry cultivar [10]. It is the least demanding natural product to fill in the kitchen garden, beautifying pots, and hanging crate and is likewise filled in level or raised beds as well as safeguarded structure. It is taken care of somewhere around a half year after planting.[1].It is industrially developed overall for its exceptionally valued sweet, fragrant, and delicious natural product. The US is the world's driving maker of strawberries delivering north of 36 billion pounds in 2012 and representing 29% of the complete world creation.[2].

Albeit synthetic composts assume a critical part in gathering the supplement prerequisites, their predictable, over-the-top, and lopsided use might bring about well-being and ecological dangers and the consumption of the physical and substance properties of the dirt. Bio-fertilizers have been considered to be an unassuming and eco-obliging way to deal with additional creating soil readiness status.[24].Thus, there is an earnest need to contemplate the utilization of elective wellspring of safe manures which might upgrade economic harvest yield without affecting soil properties and natural dangers and diminish the cost of development. Bio-manures are considered monetarily appealing and naturally sound courses for diminishing outside inputs expanding supplement supply and working on the quality and amount of inward assets with the greatest results. These bio-manures go about as transporters containing gainful miniature organic entities in a reasonable state planned for seed or soil application, intended to further develop soil richness status and help in plant development by expanding the number and natural movement of wanted microorganisms in root climate. Bio-compost viz. *Azotobacter*, PSB, and *Azospirillum* fix climatic nitrogen and solubilize phosphorus to build ripeness of soil and increment the number and natural exercises [7].

ICAR (Indian Chamber of Horticultural Exploration) researchers have fostered the innovation to pach bio-manures in small containers comprised of a transporter medium rich in live microorganisms uniquely *bacillus*, and *pseudomonas*. Each container weighed 1gm and

Comment [MAB1]: It's better to start the introduction section with this paragraph instead of the strawberry introduction. Because the title has been started with biofertilizers and bio capsules.

contained a convergence of 1000 ppm. Before application, the cases are broken up in 1 liter of tepid water for 12 hours. After that, one liter of a 500 ppm solution is dissolved in fifty liters of water, or as needed. [21].

Role of bio-fertilizers

Synthetic manures are fundamental for meeting supplement necessities, however, they ought not to be utilized too much of the time, unreasonably, or in an unequal sum. This can hurt individuals' well-being and the climate, exhaust the synthetic and actual properties of the dirt, and ultimately bring about low harvest yields. Table 1: It is necessary to take into consideration alternative sources of safe fertilizers that can boost agricultural output without harming the quality of the soil. Hence, the usage of bio manures gives some expectation around here. Bio-manures have for some time been seen as a conservative and moral method for further developing soil fruitfulness. N-fixing bacteria can either fix nitrogen from the atmosphere on their own, in a symbiotic or nonsymbiotic way, or they can use a biological process to turn useless nutrients in the soil into forms that can be used. They don't hurt plants, the dirt, or the general climate in any capacity. Notwithstanding their part in fixing nitrogen and phosphorus, these plants likewise add to the feeling of plant development chemicals. A kind of compost known as a "bio-manure" (like Azotobacter, PSB, and Azospirillum, among others) can fix nitrogen from the air and solubilize phosphorus in the dirt, subsequently expanding the natural movement and richness of the dirt. Free-living, gram-negative, nitrogen-fixing organism of the family Azotobacteriaceae, Azotobacter To increment organic nitrogen obsession and phosphorus accessibility, the two of which are fundamental for sound vegetative turn of events, biofertilizers like PSB are used[29].

The utilization of a bio-manure containing Azotobacter, chroococcum and Azospirillumbrasilense worked on the development of strawberries. The plants developed with the bio-manure had a higher root weight, stem weight, and leaf region than those developed with substance composts. Furthermore, the investigation discovered that the utilization of bio-manure additionally worked on the yield of strawberries. The plants developed with bio-composts delivered more natural product than those developed with substance manures[19].

Strawberry quality was also improved by using bio-fertilizers. The natural products developed with bio-manures had a higher L-ascorbic acid substance and a lower pH than those developed

Comment [MAB2]: No need to add this paragraph in the introduction section. Instead of this, you can add the sources of acquisition of biofertilizers and bio capsules.

Comment [MAB3]: Role of biofertilizers should be reduced to a single page, which is enough

with compound composts. Moreover, the investigation discovered that the utilization of bio-manures likewise expanded the TSS (all-out dissolvable solids) content of strawberries, which is a mark of natural product pleasantness. Bio-composts are wealthy in microorganisms that can further develop soil structure, increment the number of inhabitants in advantageous microorganisms, and lessen the requirement for synthetic pesticides [20]. Treating strawberries with a blend of chicken compost, Azotobacter, wood debris, phosphorus-debasing microorganisms, and suitable NPK manure levels brought about better plant level, leaf development, and apical width. It has been shown that involving bio-manures notwithstanding natural and inorganic composts essentially further develops strawberry plant development. These microbial products aid in nitrogen fixation and organic nutrient mineralization, thereby improving fertilizer utilization efficiency and nutrient uptake. [17].

It has been shown that involving bio-manures notwithstanding natural and inorganic composts essentially further develops strawberry plant development.[12]. Vermicompost application works on the physio-substance attributes of the dirt, which thusly increment compound movement, microbial populace, and plant development chemicals. Nitrogen and phosphate in composts encourage vegetative growth and the emergence of floral buds. [13].Azotobacter application might be ideal which brings about additional blossoms and organic products set per plant, more crowns and inflorescences per plant, and quicker N₂ obsession are undeniably influenced by the utilization of extra bio-compost. TSS and all out sugar items in berries essentially increment with various medicines of Azotobacter, Azospirillum, and PSB[25]. The plants ingest more supplements and minerals, which expands the quantity of leaves and plant advancement. A plant with additional leaves creates more carbs through expanded photosynthetic movement, which are then used for natural product improvement.[26].

Impact of Bio-fertilizers on:

I. Vegetative Growth

Organic fruit is in high demand, making it very profitable per unit area. The large-scale and miniature components are viewed as exceptionally fundamental due to the reasonable development and advancement of plants. The utilization of stable fertilizer expanded vegetative development which was seen by the creation of additional sprinters and their spreading had a direct connection to the measures of excrement applied. Strawberry's vegetative growth was also

enhanced by stable manure, whereas chemical fertilizers appear to inhibit it. An adequate measure of FYM 80-85 tons/ha has been suggested per establishing portion during land readiness [27,3].

II. Flowering

Desai (1963) reported that strawberry's flowering season varies from place to place and year to year. Based on photoperiodic reactions, strawberry cultivars are essentially arranged as June carriers, ever conveyors or day neutrals. June conveyor cultivars are facultative as brief day (SD) plant [4]. Drawn out day (LD) plants expected underneath 16°C temperature for blossom bud separation so evergreen plants delegated Drawn out day plants [9]. (Durner et al. 1984). tested day-neutral strawberry cultivars In enthusiastic plants, a few additional parts are normal while as under horrible circumstances, some bloom plants might be smothered. According to Darrow (1966), the extent to which sepals, petals, and epicalyxes open in is partly influenced by both genetics and the environment. After applying food and paper waste vermi-composts to the cv, the growth, flowering, and yield of field strawberries increased significantly. Chandler [18]. Increasing gibberellic acid in roots, which breaks bud dormancy and increases flowering buds and fruiting sites, is largely facilitated by the quantity of nutrients like N, P, and K and hormones provided by vermin-composts. [28].

III Fruiting Attributes

Knight and Wallace noticed enormous berries by the utilization of FYM than control. Over time, the acid content of strawberry fruits was significantly reduced when organic manures were applied. [15]. the piece of strawberry changes from one phase to another during development. Weight, dissolvable solids, and sharpness all ascent with development, while chlorophyll, pH, and carotenoids fall with it. [30]. The plants treated with organic manures have higher percentages of juice, total sugars, and TSS. [6].

Impact of Bio Capsules on:

I. Vegetative Growth

Bio containers could have likewise contributed to expanding the development and blooming ascribes of strawberries. Shown in Table No. 1, they contain helpful microorganisms that are used as bio-composts or bio-energizers and can colonize the rhizosphere, plant roots, or both which upgrades the plant development. Bio Containers likewise help in expanding the no. of sprinters in plants [16].

II. Flowering

These nano particles upsurge the creation and movement of photosynthates to various plant parts bringing about expanded pace of photosynthesis and chlorophyll content, also, nanoparticles are consumed all the more productively because of their bigger surface region contrasted with composts[22]. Bio Capsules and ZnO and FeO applied topically have been found to be the most effective, with significantly fewer days required for first flowering. Utilizing of various composts with Bio Cases builds number of blossoms per plant as well[21].

III Fruiting Attributes

Nano particles like ZnO and FeO are additionally answered to build the development and fruiting traits in strawberry as well as in nut, soybean, mungbean, wheat, onion, spinach, tomato, potato and mustard[5].Bio Containers additionally helps in expanding TSS of the organic product, organic product size and upgrades the nature of the fruit [21].

Comment [MAB4]: In a review paper, it is better to start the discussion with the author's name. like "ABC et al (2023)" concluded in their research article regarding the effects of biofertilizers... and then the research findings must be discussed. It is better to reorder the discussion portion, if possible.

Table 1: List of bio-fertilizer used in cultivation of strawberry

S. No.	Name of bio-fertilizer	Effect	Source
1.	Pseudomonas	Effective root colonizing bacteria.	Fikki <i>et al.</i> (2008)
2.	Azotobacter	Increased vegetative growth.	Singh <i>et al.</i> (2009) ^[23]
3.	Azospirillum	Fix atmospheric nitrogen, solubilize phosphorus, and promote soil fertility, biological activity and plant height. more	Singh <i>et al.</i> (2009) ^[23]

		vegetative growth.	
4.	Azotobacter	TSS increased, fruit size, weight and volume.	Mishra <i>et al.</i> (2010)
5.	Phosphorus Solublizing Bacteria	Plant height, development of inflorescence, increased berry size.	Mishra <i>et al.</i> (2010)
6.	Phosphorus Solubilizing Bacteria	Increase biological nitrogen fixation and phosphorus availability, which are necessary for robust vegetative development.	Deshmukh <i>et al.</i> (2014)
7.	Phosphorus Solublizing Bacteria	Promote root growth.	Prasad <i>et al.</i> (2016)
8.	Azotobacter	Enhanced yield and growth of strawberry.	Soni <i>et al.</i> (2018)
9.	Azospirillum	Fix atmospheric nitrogen, solubilize phosphorus and promote soil fertility, biological activity, and plant height. more vegetative growth.	Kumar <i>et al.</i> (2020) ^[14]
10.	Azotobacter	Increased number of leaves, numbers of runners and plant height.	Kumar <i>et al.</i> (2020) ^[14]
11.	Pseudomonas	Increased plant height and induced early flowering.	Negi <i>et al.</i> (2021)

Conclusion

From the examination of bio-composts and bio-containers on strawberries, it can securely be presumed that fundamentally increment the number of leaves per plant, length of leaves, plant spread, width of leaves, length of petioles, level of plants, absolute number of natural products, quantities of sprinters, weight of organic product per plant, weight of organic product per berry, T.S.S. content of organic products, though created most extreme Ascorbic corrosive limit of leafy foods corrosive substance.

References

- [1] Asad. Strawberry production and marketing potentials. Advisory leaflet of MFVDP, 1997; 30:1-2.
- [2] Chauhan S. S.; Mishra S.; Tarafdar M. Studies on Strawberry Cultivation Using Bio-Fertilizer (Bio-Capsule). *Agri Articles*, ISSN: 2582-9882, Volume: 03, Issue: 01 (JAN-FEB, 2023), Page 413-414.
- [3] Cooke GW. Fertilizers for maximum yield. Richard caly (the Chaucer press) Ltd. Bungay suffok, great Britain. 1972, 457.
- [4] Darrow GM. *proc. Amer. Soc. Hort Sci.*, 1936; 34:360- 363.
- [5] Dubey, A., Mailapalli, D. R. 2016. Nano fertilizers, nano pesticides, nano sensors of pest and nanotoxicity in agriculture. In *Sustainable Agriculture Reviews*; Lichtfouse, E., Ed.; Springer: Cham, Switzerland, Volume 19, pp. 307-330.
- [6] El-Hamid Aza AS, Abbou AA, Mansour SAA, El-Sayed AAA. Effect of some biofertilizers on yield and fruit quality of strawberry. *Ann Agr Sci.* 2006; 44(10):251-64.
- [7] Gabr, S.M., Ghoneim, I.M. and Hassan, H.M.F. (2001), Effects of bio-and nitrogen fertilization on growth, flowering, chemical contents, yield and quality of sweet pepper. *J.Adv.Agril. Res.*, 6: 939-955.
- [8] Giampieri, F.; Forbes-Hernandez, T.Y.; Gasparini, M.; Alvarez-Suarez, J.M.; Afrin, S.; Bompadre, S.; Battino, M. Strawberry as a health promoter: An evidence based review. *Food Funct.* 2015, 6, 1386–1398.
- [9] Ito Hand Saito T. Studies on the flower formation in the strawberry plants. *Tohoku J Agric. Res.* 1962; 13:191- 203.
- [10] John. Fruit size and general characteristics of strawberry varieties. *Infos-Paris*, No. 1994; 117:19-23.
- [11] Kumra, R.R.; Saravanan, S.; Bakshi, P.; Kumar, A.; Singh, M.; Kumar, V. Influence of plant growth regulators on Strawberry: A review. *Int. J. Chem. Stud.* 2018, 6, 1236–1239.

Comment [MAB5]: 1.The authors and titles of some references are not showing up on Google Scholar.
2.Avoid very old references.
3.Avoid references that are not available on Google Scholar.

- [12] Kumar MS, Reddy GC, Phogat M, & Korav S . Role of bio-fertilizers towards sustainable agricultural development: A review Res. Rev. J. Pharmacogn. Phytochem., 2018;7(6), 1915-1921.
- [13] Kumar N, Singh NK, Mishra P. Impact of Organic Manures and Biofertilizers on Growth and Quality Parameters of Strawberry cv. Chandler. Indian Journal of Science and Technology. 2015;8(15):51-107.
- [14] Kumar L, Kumar S, Singh R, Singh V, Yadav S, Maurya S. A Review on Effect of Organic Manure and Biofertilizers on Growth, Yield and Quality of Strawberry. Ind. J Pure App. Bio Sci. 2020;8(2):227-132.
- [15] Knight LOM, Wallace T. Effect of various manorial treatment on chemical composition of strawberries. Journal of Pomology and Horticultural Sciences. 1932; 10:147-80.
- [16] Lingua, G., Bona, E., Manassero, P., Marsano, F., Todeschini, V., Cantamessa, S. 2013. Arbuscular mycorrhizal fungi and plant growth-promoting pseudomonads increases anthocyanin concentration in strawberry fruits (*Fragaria × ananassa* var. Selva) in conditions of reduced fertilization. International Journal of Molecular Sciences. 14(8):16207-16225.
- [17] Nazir N, Singh SR, Aroosa K, Masarat J, Shabeena M. Yield and growth of strawberry cultivar Senga Sengana as influenced by the integrated organic nutrient management system. Envi. Ecol. 2006; 243:651-54.
- [18] Odongo T, Isutsa DK, Aguyoh JN. Effects of integrated nutrient sources on growth and yield of strawberry grown under tropical high altitude conditions. Afr. J Hort. Sci. 2008; 1:53-69.
- [19] Rana, R.K. and Chandel, J.S. (2003) Effect of biofertilizers and nitrogen on growth, yield and fruit quality of strawberry. Prog Horti. 2003;35(1): 25-30.
- [20] Rueda D, Valencia G, Soria N, Rueda BB, Manjunatha B, Kundapur RR, Selvanayagam M. Effect of Azospirillum spp. and Azotobacter spp. on the growth and yield of strawberry (*Fragaria vesca*) in the hydroponic system under different nitrogen levels. J Appl Pharm Sci. 2016;6(1):48-54.

- [21] Singh R. K.; Mishra S.; Bahadur V. Effect of Nano-chitosan, Nano-micronutrients and Bio Capsules on yield and Quality of Strawberry (*Fragaria ananassa*) cv. Winter Dawn. Biological Forum- An International Journal. ISSN No. (Print): 0975-1130, ISSN No. (Online): 22249-3239. 15(5): 289-295(2023).
- [22] Sabir, S., Arshad, M. and Chaudhari, S. K. 2014. Oxide Nanoparticles for Revolutionizing Agriculture: Synthesis and Application. The Scientific World Journal, Vol-14, Article ID-925494.
- [23] Singh A, Singh JN. Effect of biofertilizers and bioregulators on growth, yield and nutrient status of strawberry cv. Sweet Charlie. Indian J Hort. 2009;66(2):220-224.
- [24] Singh A., Singh J.N. Effect of biofertilizers and bioregulators on growth, yield and nutrient status of strawberry cv. Sweet Charlie. June 2009. Indian Journal of Horticulture 66(2): 220-224.
- [25] Tripathi VK, Kumar S, Kumar K, Kumar S, Dubey V. Influence of Azotobacter, Azospirillum and PSB on vegetative growth, flowering, yield and quality of strawberry cv. Chandler. Progressive Horticulture, 2016, 48. Doi: 10.5958/2249-5258.2016.00009.9
- [26] Thoudam J, Kotiyal A. Effect of Biofertilizers and Organic Fertilizers on Growth and Yield of Strawberry cv. Camarosa. The Pharma Innovation Journal. 2022;11(5):2324-2327.
- [27] Tucker LR. Observation on the growth habits of the strawberry as affected by fertilizers. In: Proceeding of American Society for Horticultural Science. 1926; 23:149-53.
- [28] Tagliavini ME, Baldi E, Lucchi P, Antonelli M, Sorrenti G, Baruzzi G, et al. Dynamics of nutrient uptake by strawberry plants (*Fragaria x Ananassa* Duch.) grown in soil and soilless culture. Eur. J Agro. 2005; 23:15-25.
- [29] Willer H, Lernoud J. eds. The World of Organic Agriculture. Statistics and Emerging. Frick and IFOAM Organics International, 2019.
- [30] Woodward JR. Physical and chemical changes in developing strawberry fruits. J. Sci. Food Agric. 1972; 23:465-473.

[31] Zargar, M.Y., Baba, Z.A. and Sofi, P.A. (2008), Effect of N, P and biofertilizers on yield and physiochemical attributes of strawberry (*Fragaria × ananassa*). Agro Thesis, 6: 3-8.

UNDER PEER REVIEW