

Review Article

L- ~~Thianine~~Theaninean Astounding Amino Acid in Tea, its Synthesis and Health Benefits: A Review

ABSTRACT

Tea is a traditional beverage that has been consumed for thousands of years, and it is also a daily habit that makes healthful living easier. Both black and green tea drinking have historically and generally been linked to feelings of refreshment and relaxation. Tea's L-Theanine (L-R-glutamylethylamide), a non-protein amino acid primarily found in tea leaves, is thought to be responsible for this consequence. Besides enhancement of the ~~flavor~~flavour of tea, L-Theanine has a number of health advantages. Diseases related to lifestyle, including heart problems, high blood pressure, stress reduction, and tumour suppression, are significantly impacted by theanine. This amino acid can mitigate the negative effects of neurotoxins, promote normal sleep patterns, and enhance memory.

This review paper aims to explain the chemical and physical properties of L-Theanine, as well as current methods for its synthesis and extraction. It also discusses the factors that influence the theanine content of tea plants and potential health benefits that it could have for humans.

(i) INTRODUCTION

Tea (*Camellia sinensis L.*), one of the three most consumed beverages worldwide, has its origins in China. It has a complex ~~flavor~~flavour and health benefits because of the diversity of bioactive substances it contains, including theanine, catechins, and caffeine. Theanine is the most prevalent free amino acid in tea leaves, accounting for 40% to 70% of the total amount (12). It was identified in 1949 as a component of green tea, and it was isolated from gyokuro leaves in 1950 (36). It makes up between one and two percent of the dry weight of green tea leaves (40). Tea contains L-configuration theanine naturally, but synthesized theanines are a combination of D- and L-configuration (15). It is also one of the most essential qualities of tea since it gives tea infusions their "umami" ~~flavor~~flavour, which promotes relaxation (14 & 35).

In ~~addition~~addition, it also offers numerous health advantages, including impacts on the immune system, urogenital protection, cardiovascular protection, cognition, memory improvement, anti-cancer and anti-anxiety properties (1, 26 & 52). This study summarizes the information on tea's L-Theanine content and how it affects the scent of the beverage and its possible health advantages on human.

(ii) Chemical, Physical, structural, and Flavour, Properties of L-Theanine-

L-Theanine is a non-proteinaceous transparent, colourless, odourless, water-soluble amino acid with a glutamine backbone in its core as well as an ethylamide derivative of glutamate. Although it is stable in acidic environments, glutamic acid and ethylamine are produced when the base is hydrolysed. L-Theanine can be easily separated from caffeine, catechins, and other lipophilic tea components because it is insoluble in organic solvents like methanol and chloroform (51). Additionally, it is designated by the systematic nomenclature (2S) 2-amino-5-(ethylamino)-5-oxopentanoic acid (C₇H₁₄N₂O₃, M.W. = 174.2 g/mol) ([NCBI, 2023 36](#)).

Theanine is a chiral molecule that primarily exists as the L(S) enantiomer in nature, similar to other amino acids, as opposed to synthesized theanine, which is often a racemic mixture of L and D enantiomers. Because of this, theanine obtained synthetically might not always have the same physiological effects as theanine found "naturally" occurring in food (37).

L-Theanine has been reported to cause a distinct "umami" ~~flavor~~flavour as well as sweetness in the mouth which was explained by its ability to bind to the T1R1 + T1R3 umami taste receptors (35).




Figure: 1. Chemical structure of L-Theanine (47).

(iii) Extraction and Synthesis of L-Theanine

Various techniques to obtain L-Theanine have been developed like: Tea leaf extraction, chemical synthesis, Enzymatic & Microbiological production.

a. Chemical Synthesis

A more practical and affordable alternative to direct separation of the amino acid from the *Camellia sinensis* plant for large-scale production might be chemical synthesis of L-Theanine using biosynthetic techniques (57). For the first time L-Theanine was chemically created by Lichenstein in 1942 by heating an aqueous solution of pyrrolidone-5-carboxylic acid to 37 degree C (51). Since then, a number of additional methods for generating L-Theanine on a large scale has been devised, such as using an L-glutamine-Zn (II) complex with ethylamine was more successful (due to its high yield and reagent availability) with respect to reducing the transfer of amino acids from one peptide chain to another during the three-hour incubation at 37 degree C and subsequently leading to an increase in the L-Theanine yield. This method might appear to be the most appropriate for large-scale production of L-Theanine (49).

The fact that the synthetic theanine was a racemic mixture of L- and D-enantiomers rather than the pure L form found in plants was one of the main obstacles to its production and the way was less eco-friendly (48).

b. Enzymatic Synthesis of Theanine

It has been proposed that the theanine synthetase synthesizes the compound (L-Theanine) from glutamic acid and ethylamine in the roots of tea plants and then transport to the

Comment [sh1]: Please refer to this figure in the above section

Comment [sh2]: according to the previous sentence, it's better to title it **L-Theanine Isolation from Tea** moved to "a." (here).

shoots. Due to high degradability of the enzyme commercial use of this technique in industry might not be plausible (48).




Figure 2. Theanine synthesis from glutamic acid and ethylamine. Ethylamine formation from alanine is also shown (34).

Formatted: Pattern: Clear (Background 1)

Comment [sh3]: According to the fig. 2, the role of alanine in theanine synthesis should be explain.

Comment [sh4]: Please refer to this figure in text "b."

c. L-Theanine Isolation from Tea

An alternate method to chemical and enzymatic synthesis for the generation of L-Theanine on an industrial scale might be the extraction of L-Theanine from tea leaves. This included ethyl acetate-based Green Tea leaf extraction, followed by preparative high performance liquid chromatography (HPLC)-based L-Theanine separation, producing an extract with about 500 g/kg L-Theanine (48). Even though the final L-Theanine product from this process had a relatively high purity, but the higher manufacturing costs and lower overall yield made this technology less desirable to the industry (51). Utilizing molecularly imprinted polymer (MIP) technology, L-Theanine could also be extracted. Phase inversion methods were used to create MIP compositions. After that, polymers were cleaned of any impurities using an acetic acid solution. Due to the finished product's subpar purity, the procedure was discovered to be less than viable (23 & 37).

d. Microbial ~~synthesis~~ Synthesis of L-Theanine

Microbial production pathways for L-theanine could be broadly classified into two categories: glutamine-mediated and glutamate-mediated pathways. ATP was required to generate energy simultaneously in the glutamate-mediated process, which uses glutamate and ethylamine as precursors (4). Yang (2020) evaluated a *Methylovorus* may *γ*-Glutamylmethylamidesynthetase (GMAS) and ~~Succeeded~~ succeeded in producing 34.49 g/L of L-theanine required for optimizing both the protein expression and the reaction conditions (5554). By using fed-batch fermentation to introduce GMAS from *Paracoccus aminovorans* heterologously into a 5-liter bioreactor, scientists were able to reach a high level of L-Theanine synthesis of 70.6 g/L (4).

Formatted: Font: Italic

γ -glutamyl transfer reactions with glutamine and ethylamine as precursors were used by the glutamine-mediated pathway (39). Glutamine synthetase (GLS) and γ -glutamyltransferase (GGT) were the main catalysts in the glutamine-mediated synthesis pathway of L-Theanine, as they ~~eatalyze~~ catalyze the conversion of glutamine and ethylamine into L-Theanine. In contrast to the glutamate-mediated synthesis pathway, ATP was not needed for this pathway. Cloning and expressing the γ -glutamyltranspeptidase from *Bacillus amyloliquefaciens* has been done in *B. subtilis* in order to perform a biosynthetic reaction using live cells (59). Ultimately, 190mM of L-Theanine was produced by a combination of promoter screening and ~~mutagenes~~ mutagens. The yield of L-Theanine increased significantly from 58% to 83% by using a cell-free reaction with purified enzymes and salt-tolerant mutants of the γ -

Formatted: Font: Italic

Formatted: Font: Italic

glutamyltranspeptidase variant (V319A/S437G) to increase the enzyme's catalytic activity (26).

(iv) Factors Affecting L-Theanine Content in Tea

The content of L-theanine varies among different tea cultivars. Within the structures of the *Camellia sinensis* plant, throughout maturity, and during the growth phase, the relative concentration of L-Theanine fluctuates. According to research, L-Theanine was found throughout the entire plant at amounts ranging from 1.2 to 6.2 mg/g fresh weight, with the highest concentrations being found in the roots (6.2-13.7 mg/g). The production of L-Theanine might take place in the plant's roots and then transferred to the leaves (10).

Formatted: Font: Not Italic

First of all, the levels of L-Theanine vary depending on the type of tea. The average L-theanine concentration of infusions prepared from commercial tea samples quantified by high-performance liquid chromatography-diode array detector was 5.13 mg/g, 6.56 mg/g, 6.09 mg/g, and 6.26 mg/g, for oolong, black, green, and white teas respectively (3). The accumulation mechanism of albino yellow tea was linked to the delayed catabolism of L-Theanine, and the quantity of L-Theanine in albino yellow tea was higher than that in regular green tea (7).

Secondly, the expression of its metabolism-related genes was correlated with L-Theanine content. The transcription levels of CsTS2 (*Camellia sinensis* L-Theanine Synthetase 2), CsGS1 (*Camellia sinensis* glutamate synthetase 1), and CsGDH2 (*Camellia sinensis* glutamate dehydrogenase 2) were positively connected with L-Theanine content among the 17 discovered genes related to L-Theanine metabolism, while the majority of the other genes were negatively correlated (28).

Thirdly, seasonality and temperature also influence the L-Theanine content in Tea Plant. Melatonin was discovered to boost the production of L-Theanine in tea leaves at sub-high temperatures (35/30 °C) and speed up the photosynthesis of tea plants (25). L-Theanine content in spring was shown to be substantially higher than that in summer and autumn, according to a combination of transcriptomics and metabolomics study (16).

Lastly, Tea leaves from various stages such as bud, first leaf, second leaf, third leaf, and old leaf were used for research and it was discovered that the bud and first leaf had the highest concentrations of L-Theanine than matured leaves as its concentration steadily dropped (28).

In matcha and gyokuro green tea L-Theanine content rises when tea plants were purposefully shaded from sunlight (6). Additionally, the ration of tea polyphenol to amino acid was reduced by application of nano-Se through significantly decreasing the total amounts of catechin, caffeine, and tea polyphenols while increasing the total amounts of amino acids and theanine (20).

Comment [sh5]: Please explain nano-Se in a ()




Figure: 3. Influencing factors of L-theanine Theanine content in tea leaves (24).

Comment [sh6]: 1-Please refer to this figure in text.
2- this fig. has not a good resolution

(V) Health Benefits of L-Theanine

(a) Anti-Tumour & Anti –Cancer ~~property-Property~~

One of the biggest public health concerns in many countries of the world is the occurrence of cancer. Numerous studies have demonstrated theanine's anti-tumour properties both in vivo and invitro(27).Theanine derivatives such as ethyl 6-nitrocoumarin-3-carboxylyl L-Theanine (TNC) and ethyl 6-fluorocoumarin-3-carboxylyl L-Theanine (TFC) showed efficient suppression of lung cancer cell proliferation in vitro, ex vivo, and in vivo by focusing on the epidermal growth factor receptor/vascular endothelial growth factor receptor-Akt/nuclear factor-kappa B (EGFR/VEGFR-Akt/NF-kappa B) signaling pathways, which involve some of the most significant pathways in regulating the survival and proliferation of cells (58).

Table: 1 Proposed Effect of different ex vivo and in vitro study

Study Type	Proposed Effect	References
Ex Vivo/In Vitro		
Cancer Suppression	L-Theanine (Methyl coumarin-3-carboxylyl L-Theanine, ethyl coumarin-3-carboxylyl L-theanine, ethyl 6-fluorocoumarin-3-carboxylyl L-Theanine, and ethyl 6-nitrocoumarin-3-carboxylyl L-Theanine) showed significant inhibition of lung cancer cell migration, growth, and leukaemia in human and mouse.	58
Tumour Suppressor	Both separately and in combination, theanine and theobromine reduced the growth of tumours by down regulating the Akt/mTOR and JAK2/STAT3 pathways and raising the Smad2 tumour suppressor.	41

Comment [sh7]: Please refer to this table in text

(b) Improved Cognitive Learning ~~ability-Ability -~~

Cognitive ability was improved by ~~thianine-theanine~~ through increasing the concentration of brain neurotransmitters like dopamine, 5-hydroxytryptamine (5-TH), glycine, and GABA (γ-aminobutyric acid) (53& 44).According to a different study, consuming 47.5 mg of L-Theanine prevented extracellular glutamine from being incorporated into neurons, indicating that L-Theanine might help elderly people with cognitive impairment. Furthermore, it has been documented that L-Theanine could control the brain's levels of serotonin and dopamine by releasing the inhibitory neurotransmitter -aminobutyric acid (51).

Table: 2 Improved Cognitive Learning ability

Impact	Pure Theanine Treatment	References
Cognition	100.6 mg of L-Theanine showed increase in cognition among 12 men and 14 women (average age, 57.7 – 4.8 years).	2
Cognition	Sensorimotor gating was enhanced by the administration of 200–400 mg of L-Theanine.	38
Learning ability	Co-administration of 40 mg of caffeine and 97 mg of L-Theanine enhanced focus on an intersensory attention	13

Comment [sh8]: Please refer to this table in text

switching task.

(c) Improved Sleeping Quality & Relaxation

L-Theanine showed a positive effect on betterment of mental and cognitive health by supporting healthy sleep patterns, which in turn promote better brain growth. After being tested on a specific population, the impact of theanine was found to be helpful in maintaining regular sleep patterns (38). Studies showed that by enhancing slow-brain waves, controlling brain electrical activity, and raising neurotransmitter and GABA receptor levels, Mg-L-Theanine compounds enhanced the effects of L-Theanine on sleep (9).

Table: 3 Improved Sleeping Quality & Relaxation

Comment [sh9]: Please refer to this table in text

Impact	Pure Theanine Treatment	References
Improved sleep quality	The eight-week L-Theanine (250 mg/day) treatment of patients (17) with diagnosed schizophrenia improved the quality of their sleep.	38
Relaxation	L-Theanine (200 mg) administration promoted relaxation when the body was at rest.	30
Relaxation	Following an acute stress task, administration of L-Theanine (200 mg) led to a decrease in both heart rate and salivary immunoglobulin A (s-IgA).	38

(d) Anti-Depressant and Stress Relief

It is commonly believed that tea (*Camellia sinensis*) encourages feelings of tranquillity and comfort. Tea's L-Theanine (L-R-glutamylethylamide), a non-protein amino acid primarily found in tea leaves, is thought to be responsible for this consequence (50).

Formatted: Normal, Indent: Left: 0.69", No bullets or numbering

Formatted: Font: (Default) Times New Roman, 12 pt, Bold, Font color: Custom Color(RGB(35,31,32))

Table: 4.a Anti-Depressant and Stress Relief

Comment [sh10]: Please refer to this table in text

Impact	Pure Theanine Treatment	References
Stress related Symptoms Reduced	Four weeks of L-Theanine administration (250mg daily) on 30 individuals (nine men and 21 women; age: 48.3 ± 11.9 years) was found to be associated with reduced stress related symptoms.	17
Depressive symptoms Reduced	Eight weeks of L-Theanine administration (250mg/day) on 20 people with major depressive disorder (open-label study) showed lower depression.	18

(a) (f) Improve Immune System

Older patients have weakened immune systems, which makes them more susceptible to influenza virus infection. Serum IgG (Immunoglobulin G) and antigen-specific IgM (Immunoglobulin M) levels were increased when L-Theanine and L-Cystine were administered together. According to clinical and epidemiological research, L-theanine

Formatted: Font: (Default) Times New Roman, 12 pt, Bold, Font color: Custom Color(RGB(35,31,32))

administration was found to prevent colds and influenza by boosting immunity and minimize the immunosuppression induced by hard activity (5).L-Theanine might mitigate heat-induced immune dysfunction by modulating the P38 signaling pathway, suppressing the rise in p-P65/P65 resulting from HSP27 (Heat Shock Protein 27) overexpression, and regulating the levels of PPAR- γ (peroxisome proliferator-activated receptor- γ) and Foxp3 (forkhead box P3) (19).

Table: 4.b5: Improve Immune System

Impact	Pure Theanine Treatment	References
Immune function	Co-administration of L-Theanine (70 mg) and cysteine (175 mg) among 176 participants was found to be associated with a decreased risk of developing the common cold.	22
Post-operative recovery	In patients with distal gastrectomy for cancer, co-administration of L-Theanine (280 mg) and cysteine (700 mg) during a randomised, single blind, parallel-group trial reduced post-gastrectomy inflammation.	33

Comment [sh11]: Please refer to this table in text

(g) Anti-Oxidant Activity

L-Theanine demonstrated strong in vitro and in vivo antioxidant activity, according to recent studies. Due to its neuroprotective and antioxidant characteristics, L-Theanine could significantly reduce the risk of cadmium-induced brain damage in rats (43). In another study, the administration of L-Theanine improved the striatum's antioxidant capacity by lowering the levels of lipid peroxide and nitric oxide (NO) in orofacial dyskinesia (OD) rat models induced by haloperidol (HAL) (45).

Formatted: Font: (Default) Times New Roman, 12 pt, Bold, Font color: Custom Color(RGB(35,31,32))

Formatted: Font: (Default) Times New Roman, 12 pt, Font color: Custom Color(RGB(35,31,32))

Formatted: Normal, Indent: Left: 0.69", No bullets or numbering

(h) Cardiovascular Protection

The risk of cardiovascular disease was lowered and vascular function was enhanced by tea drinking through lowering blood cholesterol, increasing nitric oxide production and arterial vasodilation and shielding the brain from cerebral ischemia injury (42). L-Theanine was found to reduce the risk of cerebrovascular disorders by lowering the serum cholesterol levels (27).

Formatted: Normal, Indent: Left: 0.69", No bullets or numbering

Formatted: Font: (Default) Times New Roman, 12 pt, Bold

(i) Improvement of Intestinal Immunity

Intestinal mucositis and diarrhea could be prevented by combining L-Theanine and cysteine (5:2, w/w) as they could also prevent the reduction of glutathione (GSH) levels, inhibit ROS (Reactive Oxygen Species) production and oxidative stress, and shorten intestinal villi and destroy crypts caused by 5-FU (5-Fluorouracil) (56).

Formatted: Font: (Default) Times New Roman, 12 pt, Bold

Formatted: Normal, Indent: Left: 0.69", No bullets or numbering

In addition to causing severe intestinal damage, radiation therapy also suppresses bone marrow. The crypt cells found in small intestines are especially vulnerable to radiation and prone to dying. L-Theanine and cysteine pre-treatment could lengthen the villus, deepen the crypt, and decrease the amount of apoptotic cells in the jejunal crypt following radiation exposure (32).

(j) Urogenital Protection-

L-Theanine also demonstrated both in vitro and in vivo protection of the urogenital system. L-Theanine inhibited pro-inflammatory protein kinase C (PKC)/ERK (extracellular signal-

Formatted: Normal, Indent: Left: 0.69", No bullets or numbering

Formatted: Font: (Default) Times New Roman, 12 pt, Bold, Font color: Custom Color(RGB(35,31,32))

regulated kinase)/NF- κ B(Nuclear factor kappa B)/intercellular adhesion molecule 1 (ICAM-1)/IL-33 (Interleukin-33) signaling, oxidative stress, apoptosis, and autophagy in female Wistar rats given urethane anesthesia, thereby mitigating substance P-induced bladder hyperfunction (45). Furthermore, research on stem cells revealed that 50 mM L-theanine might stimulate human stem cell proliferation and glucose metabolism to sustain the Krebs cycle, which was critical to avoiding interruption of spermatogenesis (11). According to the findings, oral L-Theanine might, at least in part, prevent bladder dysfunctions by preserving bladder contractility and preventing chronic sympathetic hyperactivity (31).

(vii) Possible Side Effects of L-Theanine

There is little information on L-Theanine used alone, and its adverse effects are poorly understood. Nonetheless, L-Theanine was categorized by the U.S. Food and Drug Administration (FDA) as a food ingredient that is generally acknowledged to be safe up to 250 milligrams per serving (40). It was found that tea could upset the stomach, especially when taken in large amounts, therefore those who drink it or take L-Theanine as green tea extract should be mindful of this. Furthermore, a small number of reports of liver issues had involved users of green tea extract (55).

Pregnant or nursing women are not yet safe to take L-Theanine, according to medical experts. Before taking L-Theanine, green tea extract, or drinking more tea, pregnant or nursing woman should consult with their doctors (8).

(viii)Conclusions

A major amino acid called L-Theanine is present in tea leaves and is responsible for the "umami" flavour. Climate and growing period have an impact on the amount of L-theanine found in different types of tea. The yields and commercial viability of the L-Theanine synthesis have varied; the majority were labour- and time-intensive with relatively low L-Theanine yields. The commercial production of purified L-Theanine appears to lack a technique that could produce the product in an environmentally and financially sustainable manner. To meet the high market demand in recent time eco-friendly microbiological techniques have been developed for production of L-Theanine (29).

L-Theanine has also generated a great deal of interest as a functional food ingredient and supplement due to its therapeutic benefits, which include effects on immunological function, learning capacity, memory improvement, urogenital protection and cancer suppression. Additionally, L-Theanine stimulates the brain's production of α -waves, which relaxes the body without making it feel sleepy. However, it is also important to acknowledge that additional research is required to support any clinically meaningful claims regarding L-Theanine as a potential functional food ingredient when compared to its pure encapsulated form (51).

COMPETING INTERESTS DISCLAIMER:

Authors have declared that they have no known competing financial interests OR non-financial interests OR personal relationships that could have appeared to influence the work reported in this paper.

References

1. Adhikary R & Mandal V. L-theanine- a potential multifaceted natural bioactive amide as health supplement. *Asian Pac J Trop Bio.* 2017; 7(9), 842–848. <https://doi.org/10.1016/j.apjtb.2017.08.005>
2. Baba Y, Inagaki S, Nakagawa S, Kaneko T, Kobayashi M, Takihara T. Effects of L-Theanine on Cognitive Function in Middle-Aged and Older Subjects: A Randomized Placebo-Controlled Study. *Journal of medicinal food.* 2021; 24(4), 333–341. <https://doi.org/10.1089/jmf.2020.4803>
3. Boros K, Jedlinszki N, Csupor D. Theanine and Caffeine Content of Infusions Prepared from Commercial Tea Samples. *Pharmacognosy magazine.* 2016; 12(45), 75–79. <https://doi.org/10.4103/0973-1296.176061>
4. Cao R, Hu S, Lu Y, Wang W, Fu Z, Cheng J. Fermentative Production of L-Theanine in *Escherichia coli* via the Construction of an Adenosine Triphosphate Regeneration System. *Fermentation.* 2023; 9(10), 875. <https://doi.org/10.3390/fermentation9100875>
5. Chen S, Kang J, Zhu H, Wang K, Han Z, Wang L, Liu J, Wu Y, He P, Tu Y, Li B. L-Theanine and Immunity: A Review. *Molecules (Basel, Switzerland).* 2023; 28(9), 3846. <https://doi.org/10.3390/molecules28093846>
6. Chen X, Ye K, Xu Y, Zhao Y, & Zhao D. Effect of Shading on the Morphological, Physiological, and Biochemical Characteristics as Well as the Transcriptome of Matcha Green Tea. *International journal of molecular sciences.* 2022; 23(22), 14169. <https://doi.org/10.3390/ijms232214169>
7. Cheng S, Fu X, Liao Y, Xu X, Zeng L, Tang J, Li J, Lai J, Yang Z. Differential accumulation of specialized metabolite l-theanine in green and albino-induced yellow tea (*Camellia sinensis*) leaves. *Food chemistry.* 2019; 276, 93–100. <https://doi.org/10.1016/j.foodchem.2018.10.010>
8. Choi JH, Ryan LM, Cramer DW, Hornstein MD, & Missmer SA. Effects of Caffeine Consumption by Women and Men on the Outcome of In Vitro Fertilization. *Journal of caffeine research.* 2011; 1(1), 29–34. <https://doi.org/10.1089/jcr.2011.0001>
9. Dasdelen MF, Er S, Kaplan B, Celik S, Beker MC, Orhan C, Tuzcu M, Sahin N, Mamedova H, Sylla S, Komorowski J, Ojalvo SP, Sahin K, Kilic E. A Novel Theanine Complex, Mg-L-Theanine Improves Sleep Quality *via* Regulating Brain Electrochemical Activity. *Frontiers in nutrition.* 2022; 9, 874254. <https://doi.org/10.3389/fnut.2022.874254>
10. Deng WW, Ashihara H. Occurrence and de novo biosynthesis of caffeine and theanine in seedlings of tea (*Camellia sinensis*). *Natural product communications.* 2015; 10(5), 703–706.
11. Dias TR, Bernardino RL, Alves MG, Silva J, Barros A, Sousa M, Casal S, Silva BM, & Oliveira PF. L-Theanine promotes cultured human Sertoli cells proliferation and modulates glucose metabolism. *European journal of nutrition.* 2019; 58(7), 2961–2970. <https://doi.org/10.1007/s00394-019-01999-2>
12. Dong C, Li F, Yang T, Feng L, Zhang S, Li F, Li W, Xu G, Bao S, Wan X, Lucas WJ, Zhang Z. Theanine transporters identified in tea plants (*Camellia sinensis* L.). *The Plant journal: for cell and molecular biology.* 2020;101(1), 57–70. <https://doi.org/10.1111/tpj.14517>

13. Einöther SJ, Martens VE, Rycroft JA, De Bruin EA. L-theanine and caffeine improve task switching but not intersensory attention or subjective alertness. *Appetite*. 2010; 54(2), 406–409. <https://doi.org/10.1016/j.appet.2010.01.003>
14. Feng L, Gao MJ, Hou RY, Hu XY, Zhang L, Wan XC, Wei S. Determination of quality constituents in the young leaves of albino tea cultivars. *Food chemistry*. 2014; 155, 98–104. <https://doi.org/10.1016/j.foodchem.2014.01.044>
15. Fiori J, Pasquini B, Caprini C, Orlandini S, Furlanetto S, Gotti R. Chiral analysis of theanine and catechin in characterization of green tea by cyclodextrin-modified micellar electrokinetic chromatography and high performance liquid chromatography. *Journal of chromatography. A*, 2018; 1562, 115–122. <https://doi.org/10.1016/j.chroma.2018.05>.
16. Gong AD, Lian SB, Wu NN, Zhou YJ, Zhao SQ, Zhang LM, Cheng L, & Yuan HY. Integrated transcriptomics and metabolomics analysis of catechins, caffeine and theanine biosynthesis in tea plant (*Camellia sinensis*) over the course of seasons. *BMC plant biology*, 2020; 20(1), 294. <https://doi.org/10.1186/s12870-020-02443-y>
17. Hidese S, Ogawa S, Ota M, Ishida I, Yasukawa Z, Ozeki M., Kunugi H. Effects of L-Theanine Administration on Stress-Related Symptoms and Cognitive Functions in Healthy Adults: A Randomized Controlled Trial. *Nutrients*. 2019; 11(10), 2362. <https://doi.org/10.3390/nu11102362>
18. Hidese S, Ota M, Wakabayashi C, Noda T, Ozawa H, Okubo T, Kunugi H. Effects of chronic l-theanine administration in patients with major depressive disorder: an open-label study. *Actaneuropsychiatrica*. 2017; 29(2), 72–79. <https://doi.org/10.1017/neu.2016.33>
19. Hu Y, Lin L, Liu K, Liu e, Han S, Gong Z, Xiao W. l-Theanine alleviates heat stress-induced impairment of immune function by regulating the p38 MAPK signalling pathway in mice†. *Food Funct*. 2023; 14(1), 335-343. <https://doi.org/10.1039/D2FO02775E>
20. Huang X, Tang Q, Chen C, Li Q, Lin H, Bai S, Zhao J, Li J, Wang K, Zhu M. Combined analysis of transcriptome and metabolome provides insights into nano-selenium foliar applications to improve summer tea quality (*Camellia sinensis*). *Science Direct*. 2023; 175(8), 114496
21. Jiang HB. Diversity of tea landraces based on agronomic and quality traits in yunnan province. *J. Plant Genet. Res.* 2013; 14, 634–640
22. Kurihara S, Hiraoka T, Akutsu M, Sukegawa E, Bannai M, Shibahara S. Effects of (L)-cystine and (L)-theanine supplementation on the common cold: a randomized, double-blind, and placebo-controlled trial. *Journal of amino acids*. 2010; 307475. <https://doi.org/10.4061/2010/307475>
23. Lachova M, Lehotay J, Karasova G, Skacani I. Isolation of L-Theanine from Plant Material using a Molecularly Imprinted Polymer. *Journal of Liquid Chromatography & Related Technologies*. 2007; 30(13-16), 2045-2058. <https://doi.org/10.1080/10826070701435053>
24. Li MY, Liu HY, Wu DT, Kanaan A, Geng F, Li HB, Gunaratne A, Li H, Gan RY. L-Theanine: A Unique Functional Amino Acid in Tea (*Camellia sinensis* L.) With Multiple Health Benefits and Food Applications. *Frontiers in nutrition*. 2022; 9, 853846. <https://doi.org/10.3389/fnut.2022.853846>

Comment [sh12]: This reference is not used in the text.

25. Li X, Li MH, Deng WW, Ahammed GJ, Wei JP, Yan P, Zhang L P, Fu J Y, Han WY. Exogenous melatonin improves tea quality under moderate high temperatures by increasing epigallocatechin-3-gallate and theanine biosynthesis in *Camellia sinensis* L. *Journal of plant physiology*. 2020; 253, 153273. <https://doi.org/10.1016/j.jplph.2020.153273>
26. Li Z, Zhu R, Liu Y, Li J, Gao H, Hu N. γ -Glutamyltranspeptidase from *Bacillus amyloliquefaciens*: transpeptidation activity enhancement and L-theanine production. *Enzyme and Microbial Technology*. 2020; 140, 109644.
27. Liang Y R, Liu C, Xiang LP and Zheng XQ. Health Benefits of Theanine in Green Tea: A Review. *Tropical Journal of Pharmaceutical Research*. 2015; 14(10), 1943-1949
28. Liu ZW, Wu ZJ, Li H, Wang YX, Zhuang J. L-theanine content and related gene expression: novel insights into theanine biosynthesis and hydrolysis among different tea plant (*Camellia sinensis* L.) tissues and cultivars. *Front Plant Sci*. 2017; 8, 498. doi: 10.3389/fpls.2017.00498
29. Liu S, Li J, Huang J, Liu Z, Xiong L. New advances in genetic engineering for l-theanine biosynthesis, *Trends in Food Science & Technology*. 2021; 114, 540-551
30. Lyon MR, Kapoor MP, Juneja LR. The effects of L-theanine (Suntheanine®) on objective sleep quality in boys with attention deficit hyperactivity disorder (ADHD): a randomized, double-blind, placebo-controlled clinical trial. *Alternative medicine review : a journal of clinical therapeutic*. 2011; 16(4), 348–354.
31. Matsuoka K, Akaihata H, Hata J, Tanji R, Honda-Takinami R, Onagi A, Hoshi S, Koguchi T, Sato Y, Kataoka M, Ogawa S, Kojima Y. l-Theanine Protects Bladder Function by Suppressing Chronic Sympathetic Hyperactivity in Spontaneously Hypertensive Rat. *Metabolites*. 2021; 11(11), 778. <https://doi.org/10.3390/metabo11110778>
32. Matsuu-Matsuyama M, Shichijo K, Tsuchiya T, Kondo H, Miura S, Matsuda K, Sekine I, Nakashima M. Protective effects of a cystine and theanine mixture against acute radiation injury in rats. *Environmental toxicology and pharmacology*. 2020; 78, 103395. <https://doi.org/10.1016/j.etap.2020.103395>
33. Miyachi T, Tsuchiya T, Oyama A, Tsuchiya T, Abe N, Sato A, Chiba Y, Kurihara S, Shibakusa T, Mikami T. Perioperative oral administration of cystine and theanine enhances recovery after distal gastrectomy: a prospective randomized trial. *Journal of parenteral and enteral nutrition*. 2013; 37(3), 384–391. <https://doi.org/10.1177/0148607112458798>
34. Mu W, Zhang T, Jiang B. An overview of biological production of L-theanine. *Biotechnology Advances*. 2015; 33(3–4), 335-342
35. Narukawa M, Toda Y, Nakagita T, Hayashi Y, Misaka T. L-Theanine elicits umami taste via the T1R1 + T1R3 umami taste receptor. *Amino acids*. 2014; 46(6), 1583–1587. <https://doi.org/10.1007/s00726-014-1713-3>
36. National Center for Biotechnology Information (2023). PubChem Compound Summary for CID 439378, Theanine. Retrieved November 2, 2023 from <https://pubchem.ncbi.nlm.nih.gov/compound/Theanine>.
37. Nobre AC, Rao A, Owen GN. L-theanine, a natural constituent in tea, and its effect on mental state. *Asia Pacific journal of clinical nutrition*. 2008; 17, 167–168.
38. Ota M, Wakabayashi C, Sato N, Hori H, Hattori K, Teraishi T, Ozawa H, Okubo T, Kunugi H. Effect of L-theanine on glutamatergic function in patients with

schizophrenia. *Actaneuropsychiatrica*. 2015; 27(5), 291–296.
<https://doi.org/10.1017/neu.2015.22>

39. Pan X, Yu J, Du Q, Zeng S, Liu J, Jiao Q, Zhang H. Efficient synthesis of γ -glutamyl compounds by co-expression of γ -glutamylmethylamidesynthetase and polyphosphate kinase in engineered *Escherichia coli*. *Journal of Industrial Microbiology & Biotechnology: Official Journal of the Society for Industrial Microbiology and Biotechnology*. 2020; 47(8), 573-583.
40. Saeed M, Khan MS, Kamboh AA, Alagawany M, Khafaga AF, Noreldin AE, Kumar M, Safdar M, Hussain M, Abd El-Hack ME, Chao S. L-theanine: an astounding sui generis amino acid in poultry nutrition. *Poultry science*. 2020; 99(11), 5625–5636. <https://doi.org/10.1016/j.psj.2020.07.016>
41. Shojaei-Zarghani S, YariKhosroushahi A, Rafraf M. Oncopreventive effects of theanine and theobromine on dimethylhydrazine-induced colon cancer model. *Biomedicine & pharmacotherapy = Biomedecine&pharmacotherapie*. 2021; 134, 111140. <https://doi.org/10.1016/j.biopha.2020.111140>
42. Siamwala JH, Dias PM, Majumder S, Joshi MK, Sinkar VP, Banerjee G, Chatterjee S. L-theanine promotes nitric oxide production in endothelial cells through eNOS phosphorylation. *The Journal of nutritional biochemistry*. 2013; 24(3), 595–605. <https://doi.org/10.1016/j.jnutbio.2012.02.016>
43. Souparnika V, Karthik Ganesh Mohanraj, Vidya S. Antioxidant Activity Of L - Theanine On Cadmium Induced Oxidative Stress Mediated Neurodegeneration -An In Vivo Analysis. *Therapeutics and Clinical Pharmacology*. 2023; 29(02), 123–130.
44. Tian X, Sun L, Gou L, Ling X, Feng Y, Wang L, Yin X, Liu Y. Protective effect of l-theanine on chronic restraint stress-induced cognitive impairments in mice. *Brain research*. 2013; 1503, 24–32. <https://doi.org/10.1016/j.brainres.2013.01.048>
45. Tsai CC, Wang MH, Chang KC, Soung HS, Yang CC, Tseng HC. Possible nitric oxide mechanism involved in the protective effect of L-theanine on haloperidol-induced orofacial dyskinesia. *The Chinese journal of physiology*. 2019; 62(1), 17–26. https://doi.org/10.4103/CJP.CJP_8_19
46. Tsai WH, Wu CH, Yu HJ, Chien CT. l-Theanine inhibits proinflammatory PKC/ERK/ICAM-1/IL-33 signaling, apoptosis, and autophagy formation in substance P-induced hyperactive bladder in rats. *Neurourology and urodynamics*. 2017; 36(2), 297–307. <https://doi.org/10.1002/nau.22965>
47. Unno K. Prevention of brain aging by green tea components: Role of catechins and theanine, *The Journal of Physical Fitness and Sports Medicine*. 2016; 5(2), 117-122. <https://doi.org/10.7600/jpfs.5.117>
48. Vuong QV, Bowyer MC, Roach PD. L-Theanine: properties, synthesis and isolation from tea. *Journal of the science of food and agriculture*. 2011; 91(11), 1931–1939. <https://doi.org/10.1002/jsfa.4373>
49. Wang HQ, Yao Z, Zhou Z, Sun Y, Wei P, Ouyang P. Enzymatic synthesis of theanine with L-glutamine-Zn (II) complexes. *Biotechnology and Bioprocess Engineering*. 2012; 17, 1135–1139. <https://doi.org/10.1007/s12257-012-0205-0>
50. Wanga L, Brennanb M, Lic S, Zhaod H, Langeek, Brennanb C. How does the tea L-theanine buffer stress and anxiety. *Food Science and Human Wellness*. 2022; 11(3) 467-475

Comment [sh13]: This reference is not used in the text.

51. Williams, J., Kellett, J., Roach, P., McKune, A., Mellor, D., Thomas, J., & Naumovski, N. (2016). L-Theanine as a Functional Food Additive: Its Role in Disease Prevention and Health Promotion. *Beverages*. 2(2), 13. <https://doi.org/10.3390/beverages2020013>
52. Xu XY, Zhao CN, Cao SY, Tang GY, Gan R. Y, Li HB. Effects and mechanisms of tea for the prevention and management of cancers: An updated review. *Critical reviews in food science and nutrition*. 2020; 60(10), 1693–1705. <https://doi.org/10.1080/10408398.2019.1588223>
53. Yamada T, Terashima T, Wada K, Ueda S, Ito M., Okubo T, Juneja LR., Yokogoshi H. Theanine, r-glutamylethylamide, increases neurotransmission concentrations and neurotrophin mRNA levels in the brain during lactation. *Life sciences*. 2007; 81(16), 1247–1255. <https://doi.org/10.1016/j.lfs.2007.08.023>
54. Yang SY, Han YH, Park YL, Park JY, No SY, Jeong D, Park S, Park HY, Kim W, Seo SO, Yang YH. Production of L-Theanine Using *Escherichia coli* Whole-Cell Overexpressing γ -Glutamylmethylamide Synthetase with Baker's Yeast. *J. Microbiol. Biotechnol.* 2020; 30(5), 785–792. <https://doi.org/10.4014/jmb.1910.10044>
55. Yin X, Yang J, Li T, Song L, Han T, Yang M, Liao H, He J, Zhong X. The effect of green tea intake on risk of liver disease: a meta analysis. *International journal of clinical and experimental medicine*. 2015; 8(6), 8339–8346.
56. Yoneda J, Nishikawa S, Kurihara S. Oral administration of cystine and theanine attenuates 5-fluorouracil-induced intestinal mucositis and diarrhea by suppressing both glutathione level decrease and ROS production in the small intestine of mucositis mouse model. *BMC cancer*. 2021; 21(1), 1343. <https://doi.org/10.1186/s12885-021-09057-z>
57. Zhang F, Zheng QZ, Jiao QC, Liu JZ, Zhao GH. Enzymatic synthesis of theanine from glutamic acid γ -methyl ester and ethylamine by immobilized *Escherichia coli* cells with γ -glutamyltranspeptidase activity. *Amino acids*. 2010; 39(5), 1177–1182. <https://doi.org/10.1007/s00726-010-0553-z>
58. Zhang G, Ye X, Ji D, Zhang H, Sun F, Shang C, Zhang Y, Wu E, Wang F, Wu F, Tian H, Liu X, Chen L, Liu K, Wang Y, Liu H, Zhang W, Guan Y, Wang Q, Zhao X, Wan X. Inhibition of lung tumor growth by targeting EGFR/VEGFR-Akt/NF- κ B pathways with novel theanine derivatives. *Oncotarget*. 2014; 5(18), 8528–8543. <https://doi.org/10.18632/oncotarget.2336>
59. Zhang R, Zheng L, Zhou L, Xiang L, Jiang B, Zhang T, Chen J. Characterization of alkaline *Bacillus amyloliquefaciens* γ -glutamyltranspeptidase expressed in *Bacillus subtilis* and its application in enzymatic synthesis of L-Theanine. *Process Biochem*. 2023; 131, 125–132.

Comment [sh14]: This reference is not used in the text.