

Original Research Article

**Effect of seed biopriming on germination_&_seedling attributes in chickpea
(*CicerarietinumL.*)**

ABSTRACT

The Study was conducted during *Zaid* season of 2022-2023 with IPC-05-59 genotype of chickpea (*Cicer arietinum L.*) Find out the Effect of seed biopriming on germination & seedling attributes in chickpea (*Cicer arietinum L.*). The design of experiment was Completely Randomized Design comprising treatments [viz.T0-uninoculatedT1-*Bacillus* sp, T2- *Mesorhizobium* sp., T3- *Pseudomonas* sp, T4- *Bacillus* sp(0.5)+ *Mesorhizobium* sp.(0.5),T5- *Pseudomonas* sp.(0.5)+ *Mesorhizobium* sp.(0.5), T6- *Bacillus* sp.(0.5)+*Pseudomonas* sp(0.5).+ *Mesorhizobium* sp.(0.5), T7- *Bacillus* sp,T8- *Mesorhizobium*sp, T9- *Pseudomonas* sp, T10- *Bacillus* sp.(0.5) + *Mesorhizobium*sp(0.5),T11-*Pseudomonas* sp(0.5)+*Mesorhizobium*sp(0.5) T12- *Bacillus* sp.(0.5) + *Pseudomonas* sp. (0.5) + *Mesorhizobium*sp. (0.5) . The treated seeds along with control were evaluated for their morpho-physiological, growth, yield, root parameters and biochemical parameters under laboratory conditions, The study revealed that seeds biopriming techniques with *Bacillus* sp.(0.5)+*Pseudomonas* sp(0.5).+ *Mesorhizobium*sp.(0.5) concentration @ 1.0ml for 1 hours T6 recorded significantly higher germination percent (93%), rate of germination (5.38), mean germination time (11.16), root length (11.73cm),shoot length(13.27cm),seedling length(25.00cm),freshweight (7.91gm),dry weight (1.62gm), vigor index-I (2324.91), vigor index – II (135.82), seed density (1.27), electrical conductivity (0.817), seed metabolic efficiency (1.018). which was followed by seedprimingwithT12- *Bacillus*sp.(0.5)+*Pseudomonas*sp.(0.5)+*Mesorhizobium*sp.(0.5) @ 1.5ml for 2 hours and minimum was recorded in T0 (control).

Keywords:~~Chickpea, Biopriming treatment, CRBD~~: *Chickpea, Biopriming treatment, +CRBD* (Completely Randomized Block Design), germination percent, speed of germination, seedling length, vigor index.

Commented [A1]: Abstract should contain the following points: introduction, objectives, materials and methods, results, conclusions and recommendations

Please add, introduction, conclusions and recommendations in your abstract

Commented [A2]: Which design you used, in your key words you put, CRBD. (CRD or CRBD please re visit)

Formatted: Font: Italic

Formatted: Font: Italic

Commented [A3]: Scientific name should be written in italic form

Formatted: Font: Italic

Formatted: Font: 12 pt, Italic

Formatted: Font: 12 pt, Italic

Formatted: Font: 12 pt, Italic

Formatted: Font: 12 pt, Italic

Formatted: Font: 12 pt, Italic

Formatted: Font: 12 pt, Italic

Formatted: Font: 12 pt, Italic

Formatted: Font: 12 pt, Italic

Formatted: Font: 12 pt, Italic

Commented [A4]: Please put correct design, in laboratory CRD is appropriate design

Commented [A5]: Please remove your design from key word

INTRODUCTION

Commented [A6]: There is spelling and Grammatical error (check again)

Chickpea(Chickpea) (*Cicer arietinum*.L)belongstothe familyFabaceae,subfamilyFaboideae.It is commonly known by other names like gramor Bengalgram, Egyptian pea, garbanzobean or garbanzo. Chickpea have protein (23.3- 28.9%), carbohydrates (61.5%), fats (4.5%) and minerals (phosphorus, calcium, magnesium, iron, zinc). Seed priming is a physiological strategy that involves soaking of seeds in a solution of a specific priming agent followed by drying of seeds that initiates germination related process. This crop was cultivated in about 99Lha. The country harvested a record production of 107 Lt at a highest productivity level of 1086kg/ha.Asusual,MadhyaPradeshhascontributedasignificant 28% of the total gram area and 34% of total gram production in the country, there by ranking first both in area and production followed by Maharashtra (20%and 18%),Rajasthan(19%&18%)and Karnataka (10% & 6%).About 97 percent of gram production of the country during the period under report has beenrealized by10states ofMadhya Pradesh,Maharashtra, Rajasthan, Karnataka, Uttar Pradesh, Andhra Pradesh, Gujarat, Chhattisgarh and Jharkhand (DAC&FW, GOI, 2022).

Among all the different priming methods, the recent modern techniques biopriming which is a combinationofbeneficialmicroorganisms'applicationonsurface ofseed and seed hydration Upadhyay *et al.*, (2016) that integrates both biological and physiological methods to improve growth of plant, along with controlling diseases. Beneficial microorganisms like Pseudomonas and Rhizobium improve plant growth by colonizing on root surface, aiding in enhanced growthofplant. Chickpea is mostly consumed in the formofprocessed whole seed (boiled,roasted,parched,fried,steamed,sproutedetc.)ordalorasdalflour(besan).Itisused in preparing a variety of snacks, sweets and condiments. It is mixed with wheat flour for "chapati" making. Fresh green seeds are consumed as green vegetable. Green leaves are used as vegetable. Grains are also used as vegetable (chole). Husk and bits of dal are used as nutritiousfeedforanimals.Chickpeacanalso beusedasgreenfodderforanimals.Itismainly growninmorethan50countries includingIndia, Pakistan, Turkey, Iran, Myanmar, Australia, Ethiopia, Canada, Mexico and Iraq Gaur *et al.*, (2010).

Biofertilizers may colonizes the rhizosphere and promotes growth by increasing the availability and supply of nutrients and/or growth stimulus to crop. Nitrogen fixer and phosphate solubilizing microorganisms play an important role in supplementing nitrogen and phosphorus to the plant, allowing a sustainable use of nitrogen and phosphate fertilizers.

Some important strains are mentioned as plant growth promoting rhizobacteria (PGPR) and that can be used as biofertilizers **Kennedy et al., (2004)** i.e. Rhizobium, Azospirillum, Azotobacter, Bacillus, Burkholderia, Erwinia, Mycobacterium, Flavobacterium etc. The bio-priming techniques changes physiological behavior of the host under stress moreover significantly plays vital role to improve seeds tolerance level under environmental stress conditions (**Entesari et al., 2013**). Priming of seeds with helpful microorganisms and bio control means has been testified more proficiently for the management of diseases and pests as equated to other available methodologies (**Prabha et al., 2019**). The seed bio-priming is a recently adapted method of seed priming. Seed priming is a pre-sowing treatment which leads to a physiological state that allows seed to germinate more proficiently. The preponderance of seed treatments are based on seed imbibition allowing the seeds to go through the first reversible stage of germination but does not allow radical protrusion through the seed coat. Seeds keeping their desiccation tolerance are then dehydrated and can be stored until final sowing (**Stanley et al., 2016**).

Physiological and biochemical changes in seeds on priming:

During seed germination an initial rapid uptake of water is followed by a visible lag phase. Only after the germination is completed a further increase in water uptake occurs, as the embryonic axis elongates. Movement of water across cell membranes is essential for the initiation of metabolism and the water movement is mediated by aquaporins. Transmembrane proteins are aquaporins (AQPs) which plays a vital role in plant water relations. High expression of AQPs during priming increases the water transport across the plasma membrane which facilitate water supply to the expanding tissues and enhance the germination potential of primed seeds. This may be the reason for the faster imbibition of water by primed seeds in comparison with non-primed ones. Priming leads to different biochemical and physiological changes in seeds. It synchronizes germination after breaking dormancy **Nascimento et al., (2004)**, diminishes the lag time required for imbibition, hydrolyses or metabolises inhibitors, activates enzymes, mobilises reserved food and enhances embryonic tissue outgrowth **Rafi et al., (2015)**. Starch metabolism is of great importance during seed metabolism which influences seedling vigour under stress. This metabolism is brought about by α -amylases which hydrolyse the starch reserves into metabolisable sugars providing energy to the developing embryo. Seed priming enhances α -amylase and dehydrogenase activity that could hydrolyse the starch macromolecules into

smaller and simple sugars with increased ATP production and respiration. Phytase, amylase, and protease also increase during this process. ~~Likewise~~ ~~Likewise~~ Kamithiet *et al.*, (2016) have shown increased activity of α -amylase in primed barley seeds by 2.8 times, whereas in primed wheat seeds activity was increased by 2.7 times compared to unprimed seeds. This might have led to enhanced germination events in primed seeds.

Bio-Priming of Seeds in Seed Production:

Seed bio-priming offers significant advantages to the seed manufacturing business. In the same way that genetic traits are passed down through generations, seed bacteria and other promising bio-gents are now being employed to increase seed vigour. In many crops, the addition of these helpful microorganisms to the seed and soil interface is a crucial strategy for producing high-quality seeds. Therefore, the seed producing sectors will be the first to gain from seed bio-priming techniques. By undergoing a number of structural, physiological, and biochemical changes within the plants, seed priming with advantageous microorganisms or biocontrol agents increases the availability of nutrients to the plants and induces systemic resistance against biotic and abiotic stresses in various ecological conditions. These microorganisms encompass a variety of bacterial and fungal agents that encourage plant development.

Keeping in view, the importance of crop and benefits of seed bio priming with microbes, the aim of the study is to identify the most effective agents for bio priming of chickpea seeds and their ~~and their~~ effect on emergence, growth and seed ~~quality parameters~~ quality parameters.

OBJECTIVES:

- To produce the inoculum of the indigenous bacterial isolates.
- To evaluate the effect of selected indigenous potential isolates on germination and seedling parameters of chickpea.

Commented [A7]: Please is there any why the genetic traits are passed down

Commented [A8]: I didn't see any seed quality parameters in your methodology please refer again

Commented [A9]: Is there any germination problem of chickpea in your area, please justify the gap of your study under introduction part

JUSTIFICATION:

There is to be proper study regarding impact of seed biopriming treatment for improvement of plant growth attributes in chickpea. So this study helps to investigate effect of biopriming seed treatment. Biofertilizers may colonize the rhizosphere and promote growth by increasing the availability and supply of nutrients and/or growth stimulus to crop chickpea. Selecting appropriate seeds and treatments is key in crop production. Biofertilizers has shown better improvement of physiological and morphological growth attributes of crop growth.

MATERIALS AND METHODS

The materials, methodology and techniques adopted in the present experiment entitled, "Effect of seed biopriming on germination & seedling attributes in chickpea (*Cicer arietinum* L.)" with a brief description regarding the site of the experiment, temperature conditions, sampling techniques and statistical analysis are presented in this chapter with following headings

Experimental design:

The experiment consists of 13 treatments including control which are replicated four times and allocated in Completely Randomized Block Design in each replication.

List 1: Details of layout

Crop	Chickpea (<i>Cicer arietinum</i> L.)
Genotype	IPC-05-59
Season	Zaid 2022-23
Number of Replications	4
Number of Treatments	13(12+1)
Number of seeds per replication	100
Method of germination	Between the paper method
First count	4th DAS
Final Count	8th DAS

Commented [A10]: Please include under introduction part. All justification should be completed under introduction part

Commented [A11]: Re write your Materials and methods. -description of the study area including temperature, altitude, rain fall, temperature...
> There is no any reference in materials and methods
> How to combine your treatments, Please refer any previous work related this title
> What is your base??
> Data collected (germination rate...) please include under methodology

Commented [A12]: Please remove this title from methodology part

Commented [A13]: Where is your sampling technique

Commented [A14]: Please justify how to analyze your data? Which soft ware you used R or SAS soft ware or any other please justify???

Commented [A15]: Please what does it means between paper method? Clearly write your method of germination, for other readers not clear

Commented [A16]: DAS represent what? Please write under table

Commented [A17]: DAS?

List 2: Treatment Combinations

Treatments	Organisms	Concentration	Duration (hours)
T0	Uninoculated	NIL	-
T1	<i>Bacillus</i> sp.	1ml	1
T2	<i>Mesorhizobium</i> sp.	1ml	1
T3	<i>Pseudomonas</i> sp.	1ml	1
T4	<i>Bacillus</i> sp.(0.5)+ <i>Mesorhizobium</i> sp.(0.5)	1ml	1
T5	<i>Pseudomonas</i> sp.(0.5)+ <i>Mesorhizobium</i> sp.(0.5)	1ml	1
T6	<i>Bacillus</i> sp.(0.5)+ <i>Pseudomonas</i> sp.(0.5)+ <i>Mesorhizobium</i> sp.(0.5)	1ml	1
T7	<i>Bacillus</i> sp.	1.5ml	2
T8	<i>Mesorhizobium</i> sp.	1.5ml	2
T9	<i>Pseudomonas</i> sp.	1.5ml	2
T10	<i>Bacillus</i> sp.(0.5)+ <i>Mesorhizobium</i> sp.(0.5)	1.5ml	2
T11	<i>Pseudomonas</i> sp.(0.5)+ <i>Mesorhizobium</i> sp.(0.5)	1.5ml	2
T12	<i>Bacillus</i> sp.(0.5)+ <i>Pseudomonas</i> sp.(0.5)+ <i>Mesorhizobium</i> sp.(0.5)	1.5ml	2

Formatted: Font: Italic, (Intl) Calibri

Formatted: Font: Italic

Formatted: Font: Italic, (Intl) Calibri

Formatted: Font: Italic

Formatted: Font: Italic, (Intl) Calibri

Formatted: Font: Italic

Commented [A18]: Write all scientific name in italic form

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic, (Intl) Calibri

Formatted: Font: Italic

Formatted: Font: Italic, (Intl) Calibri

Formatted: Font: Italic

Formatted: Font: Italic, (Intl) Calibri

Formatted: Font: Italic

Formatted: Font: Italic

Formatted: Font: Italic, (Intl) Calibri

Commented [A19]: What is your base to use these level of concentration

Formatted: Font: Italic, (Intl) Calibri

Formatted: Font: Italic, (Intl) Calibri

Formatted: Font: Italic, (Intl) Calibri

Commented [A20]: Please put Reference (e.g. according to this outour....)??

Preparation of solutions:

Bacterial cultures:

Preparation of media:

For *Bacillus* sp. Nutrient broth media will be used which is a composition of

Manitol agar - 10 gm/litre

K₂HPO₄ -0.5gm/litre
 MgSO₄ - 0.2gm/litre
 NaCl -0.1 gm/litre
 CaCO₃ -
 4gm/litre Yeast extract-
 0.1gm/litre

For *Rhizobium sp* Yamabrothmedia will be used which is a composition of

Peptone - 5 gm/litre Beef
 extract - 3gm/litre NaCl
 -5gm/litre

For *Pseudomonas sp*: pseudomonas agar is used which is a composition of

Casine hydrolysate -
 10gm/litre Protease, peptone -
 10gm/litre
 K₂HPO₄ -1.5gm/litre
 MgSo₄ .7H₂O -
 1.5gm/litre pH - 7

- Autoclave media at 120° for 15 min.
- After media gets cooled down fresh culture of *Bacillus* spp, *Pseudomonas* spp, *Rhizobium* spp. are separately inoculated in specific broth.
- Incubate at 28°c for 2-3 days.
- Seeds were surface disinfected with ethanol
- The sterilized seeds were soaked in bacterial suspension (1×10^7 cells/ml) for 1, 2 hours respectively, in case of control, the seeds are soaked in distilled water.
- Seeds were air dried for 48 hours
- Seeds were placed on each germination paper for germination tests
- Observations were recorded on first count and last count i.e. on 4th and 8th day after kept for germination.

Commented [A21]: Please describe all of your parameters in materials and methods (germination percentage...

RESULTS AND DISCUSSIONS

An experiment entitled “Effect of seed biopriming on germination & seedling attributes in chickpea (*Cicer arietinum* L)” was conducted at Seed Testing Laboratory, Department of Genetics and Plant Breeding, Naini Agricultural Institute, SHUATS, Prayagraj, (U.P.) during Zaid season of 2022-2023. The results of this experiment ~~are~~ were presented in the following pages under appropriate sub headings. Experimental findings are discussed in the light of scientific reasoning and their conformity with the previous researchers.

Analysis of Variance: The analysis of variance for Germination and Seedling Attributes in chickpea was given below table. Analysis of variance revealed that the differences among thirteen treatments (Table 1) were significant for germination and seedling attributes viz., Germination percentage (%), Rate of germination, Mean germination time, Root length (cm), Shoot length (cm), Seedling length (cm), Fresh weight (g), Dry weight (g), Vigour Index-I, Vigour Index-II, Seed density, Electrical conductivity (dSm⁻¹), Seed metabolic efficiency (Table 2).

Commented [A22]: No any discussion, please re write your discussion parts
➤ No any reference in your results and discussion. Please discuss your paper with previous work

Commented [A23]: Which researcher?? Please cite the researcher

Table 1: Analysis of variance for seed germination and seedling parameters of bio-priming by *Rhizobium*, *Bacillus*, *Pseudomonas* in chickpea (*Cicer arietinum* L.).

S.no	Characters	Meansumofsquares	
		Treatments (df=12)	Error (df=39)
1	Germinationpercentage(%)	125.4231*	4.0000
2	Rateofgermination	0.5409*	0.4288
3	Meangerminationtime	1.8061*	0.0576
4	Rootlength(cm)	7.8748*	0.0112
5	Shootlength(cm)	7.4281*	0.0617
6	Seedlinglength(cm)	29.8297*	0.0921
7	freshweight(g)	4.8217*	0.0999
8	dryweight(g)	0.0296*	0.0051
9	vigourindex-I	418188.5837*	283.6725
10	vigourindex-II	527.8566*	42.1658
11	Seed density	0.0032*	0.0013
12	Electricalconductivity(dSm-1)	0.5077*	0.0338
13	Seedmetabolicefficiency	0.0545*	0.0192

*Indicates 5% Level of Significance

Commented [A24]: Add CV

Formatted: Space Before: 0 pt

Formatted Table

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Indent: Left: 0"

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Table 2

UNDER PEER REVIEW

Treatments	Germination percentage (%)	Rate of germination	Mean germination time	Root length (cm)	Shoot Length (cm)	Seedling length (cm)	Fresh weight (g)	Dry weight (g)	Vigour index I	Vigour index II	Seed density	Electrical conductivity	Seed metabolic efficiency
T0	68	3.77	8.16	6.64	7.67	14.32	3.07	1.25	973.18	84.71	1.15	1.905	0.556
T1	78	4.66	9.36	9.31	10.95	20.27	4.54	1.35	1580.45	105.72	1.21	1.766	0.823
T2	79	4.76	9.48	10.4	10.65	21.05	3.81	1.36	1662.81	128.26	1.21	1.686	0.690
T3	81	4.95	9.72	7.49	9.15	16.65	4.21	1.39	1348.03	112.68	1.19	1.625	0.789
T4	82	4.95	9.84	10.14	10.77	20.91	4.88	1.46	1714.33	126.28	1.20	1.078	0.722
T5	86	5.10	10.32	10.62	12.41	23.04	5.46	1.51	1980.81	130.21	1.23	0.902	0.591
T6	93	5.38	11.16	11.73	13.27	25.00	7.91	1.62	2324.91	135.82	1.27	0.817	1.018
T7	73	4.23	8.76	6.85	8.59	15.45	3.26	1.34	1127.26	106.24	1.17	1.848	0.861
T8	79	5.09	9.48	9.77	10.51	20.29	3.8	1.36	1602.14	109.57	1.24	1.734	0.862
T9	80	4.95	9.60	9.21	10.77	19.99	5.53	1.39	1598.59	114.2	1.20	1.686	0.841
T10	82	4.81	9.84	8.85	10.43	19.29	4.04	1.43	1581.16	111.48	1.22	1.331	0.650
T11	85	4.81	10.20	9.41	11.19	20.61	4.59	1.46	1751.22	114.94	1.19	1.002	0.772
T12	89	5.23	10.68	11.60	12.63	24.23	5.51	1.54	2156.37	119.02	1.25	0.891	0.948
Grand mean	81	4.82	9.74	9.39	10.69	20.08	4.66	1.42	1646.25	115.66	1.21	1.406	0.779

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted Table

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

S.Em	1.155	0.378	0.139	0.06 1	0.143	0.175	0.182	0.041	9.724	3.749	0.021	0.106	0.080
SE(d)	1.633	0.535	0.196	0.08 6	0.203	0.248	0.258	0.058	13.752	5.302	0.029	0.150	0.113
C.D(0.05)	3.357	1.099	0.403	0.17 8	0.417	0.509	0.530	0.120	28.267	10.8 98	0.060	0.309	0.233

Table no: 2 Mean Performance of germination and seedling parameters of bio-primed seeds by *Rhizobium, Bacillus, Pseudomonas* in Chickpea (*Cicer arietinum L.*) Genotype: *IPC-05-59*

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Formatted: Space Before: 0 pt

Commented [A25]: Please summarize your table, put in clear form. Refer published research papers

Commented [A26]: Write the table name on top of your table

Commented [A27]:

UNDER PEER REVIEW

Germination percentage:

A perusal of data on germination percentage was recorded during the experimental period revealing a significant difference in germination percent, which was recorded As per the data tabulated germination percent was recorded significantly higher in treatment (T6 - *Bacillus* sp.(0.5)+ *Pseudomonas* sp(0.5)+ *Mesorhizobium* sp.(0.5) @ 1ml) with 93% and treatment (T12- *Bacillus* sp.(0.5)+*Pseudomonas*sp.(0.5)+*Mesorhizobium*sp.(0.5)@1.5ml)with89%wasstatisticallyat par with treatment T6 and the lowest germination percent was recorded in treatment T0 (control) with 68%. *Pseudomonas* strains are known to produce plant growth-promoting hormones such as auxins and ~~cytokinins~~cytokines. These hormonescanstimulate seed germinationand earlyseedling growthbypromoting cellelongation,rootdevelopment,andnutrientuptake.*Pseudomonas*treatmentscaninducesystemic resistance in plants. This means that chickpea seeds treated with *Pseudomonas* may become more resilienttostressorsandpathogens,therebyimprovingtheirchancesofgerminationandearlygrowth. there by improving their chances of germination and early growth (Reference).

Commented [A28]: Please briefly describe this parameters in materials and method parts

Formatted: Font: 12 pt, Italic

Formatted: Font: 12 pt, Italic

Formatted: Font: 12 pt, Italic

Commented [A29]: See the spelling again

Rate of germination:

From the data recorded it is obvious that treatment (T6 - *Bacillus* sp.(0.5)+ *Pseudomonas* sp(0.5)+ *Mesorhizobium* sp.(0.5) @ 1ml) gave significantly higher rate of germination with 5.38% over all other treatments. However, treatment (T12- *Bacillus* sp.(0.5)+ *Pseudomonas* sp.(0.5)+ *Mesorhizobium*sp.(0.5)@1.5ml)recordedarateofgerminationof5.23%andwasstatisticallyatpar with T3 and the lowest rate of germination was obtained in T0 (control) with 3.77%.

Commented [A30]: Please briefly describe this parameters in materials and method parts

Formatted: Font: 12 pt, Italic

Formatted: Font: 12 pt, Italic

Formatted: Font: 12 pt, Italic

Mean germination time:

As per the data recorded the treatment (T6 - *Bacillus* sp.(0.5)+ *Pseudomonas* sp(0.5)+ *Mesorhizobium* sp.(0.5) @ 1ml) had significantly highest mean germination time of 11.16 over all other treatments and treatment T12- *Bacillus* sp.(0.5)+ *Pseudomonas* sp.(0.5)+ *Mesorhizobium* sp.(0.5) @ 1.5ml with 10.68 of mean germination time was statistically at par with T3. The lowest mean germination time was recorded in T0 (control) with 8.16.

Commented [A31]: Please briefly describe this parameters in materials and method parts

Commented [A32]: Why?? Please refer any previous work

Commented [A33]: Please briefly describe this parameters in materials and method parts

Formatted: Font: 12 pt, Italic

Root length:

From the data recorded it is obvious that treatment (T6 - *Bacillus* sp.(0.5)+ *Pseudomonas* sp(0.5)+ *Mesorhizobium* sp.(0.5) @ 1ml) gave significantly higher root length with 11.73cm over all other treatments considered. Treatment (T12- *Bacillus* sp.(0.5)+ *Pseudomonas* sp.(0.5)+ *Mesorhizobium* sp.(0.5)@1.5ml)recordedarootlengthof11.60cmdandwasstatisticallyatparwithT6andthelowest root length was obtained in T0 (control) with 6.64cm.

Pseudomonas treatments can stimulate the development of root hairs, which are extensions of root epidermalcells.Roothairsincreasethesurfaceareafor nutrientandwaterabsorption,enhancing

overall root growth. *Pseudomonas* bacteria can trigger the production of antioxidants in chickpea

UNDER PEER REVIEW

seedlings. Antioxidants protect plant tissues from oxidative stress, allowing roots to grow without hindrance.

Shootlength:

As per the data recorded the treatment (T6 - Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1ml) had significantly highest shoot length of 13.27cm over all other treatments and treatment T12- Bacillus sp.(0.5)+Pseudomonas sp.(0.5)+Mesorhizobiumsp.(0.5)@ 1.5ml with 13.27cm of shoot length was statistically at par with T6. The lowest shoot length was recorded in T0 (control) with 7.67cm.

Commented [A34]: Please briefly describe this parameters in materials and method parts

Seedlinglength:

The significantly higher seedling length (25.00cm) was recorded with treatment (T6 - Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1ml) over the rest of the treatments taken. In contrast a significantly lower seedling length of 14.32cm was recorded with treatment (T0-control). However, treatment (T12- Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1.5ml) with 24.23cm was statistically at par with T6.

Commented [A35]: Please briefly describe this parameters in materials and method parts

Pseudomonas treatments can trigger the production of antioxidants in chickpea seedlings. Antioxidants protect plant tissues from oxidative stress, which can hinder growth. With lower oxidative stress, seedlings can focus on elongating their stems and leaves. Pseudomonas treatments can induce systemic resistance in chickpea seedlings. When the seedlings are primed for defense responses, they allocate more resources to growth, resulting in increased seedling length.

Freshweight:

The significantly higher seedling fresh weight was obtained in treatment (T6 - Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+Mesorhizobiumsp.(0.5) @ 1ml) with 7.91gm among all the other treatments and the lowest seedling fresh weight was recorded in T0 (control) with 3.07gm respectively. The treatment (T12- Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1.5ml) was statistically at par with T6 with 5.51gm.

Commented [A36]: Please briefly describe this parameters in materials and method parts

Pseudomonas treatments can promote more efficient nutrient uptake by these seedlings. This increased nutrient absorption can support the development of larger and heavier seedlings. Pseudomonas can produce plant growth-promoting hormones like auxins and cytokinins. These hormones stimulate cell division and elongation, leading to increased biomass and fresh weight in seedlings. Pseudomonas treatments can induce systemic resistance in chickpea seedlings. As a result, seedlings are better protected against pathogens and environmental stresses, allowing them to allocate more energy to growth and fresh weight gain.

Dryweight:

The significantly higher seedling dry weight (1.62gm) was recorded with treatment (T6 - Bacillus sp.(0.5)+ Pseudomonas sp(0.5)+ Mesorhizobium sp.(0.5) @ 1ml). In contrast a significantly lower dry weight of 1.25 was recorded with treatment (T0-control). However, treatment (T12- Bacillus sp.(0.5)+Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @1.5ml) with 1.54gm was statistically at par with treatment T2.

Pseudomonas treatments often lead to the development of more extensive and well- branched root systems. A healthy root system allows for better nutrient and water absorption, ultimately contributing to increased seedling dry weight. Pseudomonas treatments may enhance the efficiency of photosynthesis in chickpea seedlings, leading to increased production of carbohydrates and dry biomass.

Vigour index I:

Significantly higher vigour index I was recorded in treatment T6 - Bacillus sp.(0.5)+ Pseudomonas sp(0.5)+ Mesorhizobium sp.(0.5) @ 1ml with 2324.91 and the treatment T12- Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1.5ml with vigour index I of 2156.37 was statistically at par with T6. The lowest vigour index I was recorded in treatment T0-control with vigour index I of 973.18.

Pseudomonas treatments often lead to the development of a more extensive and well- branched root system. A healthy root system enhances nutrient and water uptake, supporting overall seedling vigor.

Pseudomonas treatment promotes germination, seedling length, and dry weight through better cell division increasing the seedling vigour.

Vigour index II:

As per the data recorded during laboratory study the vigour index II was recorded significantly higher in treatment T6 - Bacillus sp.(0.5)+ Pseudomonas sp(0.5)+ Mesorhizobium sp.(0.5) @ 1ml with vigour index II of 135.82 and statistically at par value was recorded in treatment T12- Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1.5ml with 130.21. The lowest data on vigour index II was recorded in treatment (T0-control) with 84.71.

Seed density:

The treatments varied significantly and the highest seed density was recorded in treatment T6 - Bacillus sp.(0.5)+ Pseudomonas sp(0.5)+ Mesorhizobium sp.(0.5) @ 1ml with 1.27 dsm^{-1} . The treatment (T12- Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1.5ml) with seed density of 1.25 dsm^{-1} was statistically at par with T6 and the lowest seed density was recorded in treatment T0 - control with 1.15 dsm^{-1} .

Commented [A37]: Please briefly describe this parameters in materials and method parts

Commented [A38]: Please briefly describe this parameters in materials and methods parts

Commented [A39]: Please briefly describe this parameters in materials and method parts

Commented [A40]: Please briefly describe this parameters in materials and methods parts

Commented [A41]: It is your control, why you compare with other treatments

Pseudomonas treatments can stimulate the development of a more extensive root system in chickpea seedlings. These well-developed root systems enable seedlings to explore a larger soil volume for resources, indirectly contributing to their overall vigor and potential seedling density.

Electrical conductivity:

As per the data tabulated electrical conductivity was recorded significantly lower in treatment (T6 - Bacillus sp.(0.5)+Pseudomonas sp.(0.5)+Mesorhizobium sp.(0.5) @ 1ml) with 0.817 and treatment (T12-Bacillus sp.(0.5)+Pseudomonas sp.(0.5)+Mesorhizobium sp.(0.5) @ 1.5ml) with 0.891 was at par with T6 and the highest electrical conductivity was recorded in treatment (T0-control) with 1.905.

Pseudomonas seed treatment may influence the ion-selective transport systems in chickpea root cells. These bacteria can modulate ion channels and transporters to favor the uptake of essential nutrients (e.g., potassium and calcium) while limiting the uptake of potentially detrimental ions (e.g., sodium and chloride). This selective ion regulation can lead to an altered ion composition within the plant and, consequently, a lower EC of the plant sap.

Seed metabolic efficiency:

A perusal of data on seed mobilization efficiency was recorded during the experimental period on different treatments revealing a significant difference in seed mobilization efficiency. As per the data tabulated seed mobilization efficiency was recorded significantly higher in treatment (T6 - Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1ml) with 1.018 and treatment (T12- Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1.5ml) with 0.948 was statistically at par with T6 and the lowest seed mobilization efficiency was recorded in treatment (T0- control) with 0.556.

No discussion part

Summary:

The present lab experiment entitled “Effect of seed biopriming on germination & seedling attributes in chickpea (*Cicer arietinum* L)” was conducted at Seed Testing Laboratory, Department of Genetics and Plant Breeding, Naini Agricultural Institute, SHUATS, Prayagraj, (U.P.) during Zaid season of 2022-2023.

The study conducted in the seed testing laboratory revealed that germination percent (93%), rate of germination (5.38), mean germination time (11.16), root length (11.73 cm), shoot length (13.27 cm), seedling length (25.00 cm), fresh weight (7.91 gm), dry weight (1.62 gm), vigor index-I (2324.91), vigor index – II (135.82), seed density (1.27), electrical conductivity (0.817), seed metabolic efficiency (1.018) were recorded significantly higher in T6 - Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1ml over other treatments and treatment T12- Bacillus sp.(0.5)+ Pseudomonas sp.(0.5)+ Mesorhizobium sp.(0.5) @ 1.5ml was at par with T6.

Commented [A42]: Please briefly describe this parameters in materials and method parts

Commented [A43]: Please briefly describe this parameters in materials and method parts

Commented [A44]: See your summary again

➤ Analysis of variance showed significant difference among the treatments for the characters studied of the

UNDER PEER REVIEW

Chickpea which indicates significant effect of the treatments on seedling parameters of the crop.

The salient finding of the experiment is summarized and conclusion is drawn. The study was undertaken to know the effect of seed biopriming treatments on Chick pea seeds have been presented and discussed in the previous chapter. The data was recorded for seedling parameters of Chick pea.

Commented [A45]: No any discussion please

Conclusions and Recommendations:

Studying indigenous bacteria can improve plant health and plant growth and inoculating seed with beneficial indigenous bacteria can enhance nutrient availability and protect plants from harmful pathogens. Indigenous bacteria can provide insights into the local microbial community and its role in various ecosystems. They contain unique enzymes or metabolic pathways that can be harnessed for biotechnological applications in crop.

It is concluded from the present study that (T6) treatment in which seeds treated with *Bacillus sp.*(0.5)+ *Pseudomonas sp.*(0.5)+ *Mesorhizobium sp.*(0.5) @ 1ml had out performed all treatments in laboratory conditions with highest germination percentage, seedling length, fresh weight, vigor indices, electrical conductivity followed by (T12) treatment in which seeds treated with *Bacillus sp.*(0.5)+ *Pseudomonas sp.*(0.5)+ *Mesorhizobium sp.*(0.5) @ 1.5ml and the lowest were recorded in (T0- control).

Future prospects:

The bio priming technique can be recommended for priming seeds prior to field planting as an environmentally friendly strategy to improve seed germination and initial seedling growth. Extensive field trials under different environmental conditions, using several seed lots of cultivars, will be necessary in order to establish the efficiency of these strains as potential bio-priming seed treatments for improving Crop productivity.

Commented [A46]: Remove

Commented [A47]: Include under conclusions and Recommendations

References:

Abdul-Baki, A., and Anderson, J. D. (1973). Vigor Determination in Soybean Seed by Multiple Criteria. *Crop Science*, 13 (6):630-633.

Baliyan, N., Dheeman, S., Maheshwari, D. K., Dubey, R. C., & Vishnoi, V. K. (2018). Rhizobacteria isolated under field first strategy improved chickpea growth and productivity. *Environmental Sustainability*. 1: 461-469.

Chaurasia, A. K., Naik, B. P., and Bara, B. M. (2018). Influence of priming on seed yield parameters of chickpea (*Cicer arietinum* L.). *Journal of Pharmacognosy and Phytochemistry*. 7(2): 753-755.

Ditta, A., Imtiaz, M., Mehmood, S., Rizwan, M. S., Mubeen, F., Aziz, O., Qian, Z., Ijaz, R., Tu, S., (2018). Rock phosphate-enriched organic fertilizer with phosphate-solubilizing microorganisms improves nodulation, growth, and yield of legumes. *Commun. Soil Sci. Plant Anal.* 49(21): 2715-2725.

Entesari, M., Sharifzadeh, F., Ahmadzadeh, M., & Farhangfar, M. (2013). Seed bio-priming with *Trichoderma* species and *Pseudomonas* fluorescent on growth parameters, enzyme activity and nutritional status of soybean. *International Journal of Agronomy and Plant Production*. 4(4): 610-619.

Commented [A48]: Please re write your references
Some of your references you write is not cited in your paper
>Some your reference you write under introduction parts not available in reference parts
>Please cross check all reference parts

Commented [A49]: Some of your reference is very old please update your reference

Commented [A50]: Put your reference in alphabetical order

Commented [A51]: Number your references

Commented [A52]: Very old references

Commented [A53]: Not available (not cited in your paper, please check

Commented [A54]: Not cited in your paper please

Fahsi, N., Mahdi, I., Mesfioui, A., Biskri, L., and Allaoui, A. (2021). Plant growth promoting rhizobacteria isolated from the jujube (*Ziziphus lotus*) plant enhance wheat growth, Zn uptake, and heavy metal tolerance. *Agriculture* 11:316.

Commented [A55]: Not cited in your paper

Gupta, S.C. (2006) Effect of combined inoculation on nodulation, nutrient uptake and yield of chickpea in Vertisol. *Journal of the Indian Society of Soil Science* 54: 251-254

Hayat, R., Ali, S., Amara, U., Khalid, R., & Ahmed, I. (2010). Soil beneficial bacteria and their role in plant growth promotion: a review. *Annals of microbiology*. 60: 579-598.

Jeffries, P., Gianinazzi, S., Perotto, S., Turnau, K., & Barea, J.M. (2003). The contribution of arbuscular mycorrhizal fungi in sustainable maintenance of plant health and soil fertility. *Biology and fertility of soils*. 37: 1-16.

Kamithi, K. D., Wachira, F., & Kibe, A. M. (2016). Effects of different priming methods and priming durations on enzyme activities in germinating chickpea (*Cicer arietinum* L.). *American Journal of Natural and Applied Sciences*. 1(1): 1-9.

Khan, M. S., Zaidi, A. and Wani, P.A. (2007). Role of phosphate solubilizing microorganisms in sustainable agriculture. A Review. *Agronomy for sustainable Development*. 27: 29-43.

Lohitha, B., Dutta, A. (2022) Biopriming in chickpea (*Cicer arietinum* L.): An ecofriendly approach for quality seed production. *Pharma Innovation* ;11(2):375-379

Moeinzadeh, A., Sharif-Zadeh, F., Ahmadzadeh, M., & Tajabadi, F. H. (2010). Biopriming of Sunflower (*Helianthus annuus* L.) Seed with *Pseudomonas fluorescens* for Improvement of Seed Invigoration and Seedling Growth. *Australian Journal of Crop Science*. 4(7): 564-570.

Nascimento, W. M., Cantliffe, D. J., & Huber, D. J. (2005). Seed aging affects ethylene production and endo- β -mannanase activity during lettuce seed germination at high temperature. *Seed Science and Technology*. 33(1): 11-17.

Negi, S., Bharat, N. K., Kaushal, R., & Rohiwala, P. (2020). Screening of bioagents for seed biopriming in French bean (*Phaseolus vulgaris* L.) under Laboratory condition. *International Journal of Communication Systems*. 8(3): 790-793.

Pandey, S., and Gupta, S. (2020). Evaluation of *Pseudomonas* sp. for its multifarious plant growth promoting potential and its ability to alleviate biotic and abiotic stress in tomato (*Solanum lycopersicum*) plants. *Scientific Reports* 10:20951.

Panase, V.G., and Sukhatme, P.V. (1967). Statistical Methods for Agricultural workers. Indian council of agricultural research, New Delhi. pp 381

Commented [A56]: Not cited in your paper

Pramanik, K., and Bera, A.K. (2012). Response of biofertilizer and phytohormone on growth and yield of chickpea (*Cicer arietinum* L.). *Journal of crop and weed*. 8(2): 45- 49.

Rafi, H., Dawar, S., & Zaki, M.J. (2015). Seed priming with the extracts of *Acacia nilotica* (L.) Willd. ex Delile and *Sapindus mukorossi* (L.) plant parts in the control of root rot fungi and growth of plants. *Pak. J. Bot.* 47(3): 1129-1135.

Rao, D. G. and Sinha, S. K. (1993). Efficiency of mobilization of seed reserves in sorghum hybrids and their parents as influenced by temperature regimes. *Seed Research*. 2(2): 97-100.

Commented [A57]: Not available check all references

Reddy, A. S. R., Madhavi, G. B., Reddy, K. G., Yellareddygari, S. K., & Reddy, M. S. (2011). Effect of seed biopriming with *Trichoderma viride* and *Pseudomonas fluorescens* in chickpea (*Cicer arietinum*) in Andhra Pradesh, India. In Plant growth-promoting rhizobacteria (PGPR) for sustainable agriculture, Proceedings of the 2nd Asian PGPR conference, Beijing, China (pp. 324-429).

Singh, V., Upadhyay, R. S., Sarma, B. K., & Singh, H. B. (2016). Seed bio-priming with *Trichoderma asperellum* effectively modulate plant growth promotion in pea. *International Journal of Agriculture, Environment and Biotechnology*. 9(3): 361- 365.

Tambekar, D.H., Gulhane, S.R., Somkuwar, D.O., Ingle, K.B., Kanchalwar, S.P., Upadhye, M. A., & Bidwai, U. A. (2009). Potential Rhizobium and phosphate solubilizers as a biofertilizers from saline belt of Akola and Buldhana district, India. *Res. J. Agric. Biol. Sci.* 5(4): 578-582.

Tanwar, A., Aggarwal, A., Kaushish, S., & Chauhan, S. (2013). Interactive effect of AM fungi with *Trichoderma viride* and *Pseudomonas fluorescens* on growth and yield of broccoli. *Plant Protection Science*. **49**(3), 137-145.

Valverde, A., Burgos, A., Fiscella, T., Rivas, R. and Igual, J.M. (2006). Differential effects of coinoculations with *Pseudomonas jessenii* PS06 (aphosphate-solubilizing bacterium) and *Mesorhizobium ciceri* C-2/2 strains on the growth and seed yield of chickpea under green house and field conditions. *Plant and Soil*. **287**: 43-50.

Van der Maesen L.J.G. (1987). Origin, history and taxonomy of chickpea. In *The chickpea*, 11-34

Commented [A58]: Not cited please

Veenashri Jainapur, Shalini N. Huilgol, S.M. Vastrad and R.B. Jolli (2018). Biopriming of chickpea seeds with biocontrol agents for enhanced seedling vigour and reduced seed borne diseases. *International Journal of Current Microbiology and Applied Sciences*. **7**(2): 2746-2750.

Vishwas, S., Chaurasia, A.K., Bara, B.M., Debnath, A., Parihar, N.N., Brunda, K., & Saxena, R. (2017). Effect of priming on germination and seedling establishment of chickpea (*Cicer arietinum* L.) seeds. *Journal of pharmacognosy and phytochemistry*. **6**(4): 72- 74.

Vishwas, S., Chaurasia, A.K., Bara, B.M., Debnath, A., Parihar, N.N., Brunda, K., & Saxena, R. (2017). Effect of priming on germination and seedling establishment of chickpea (*Cicer arietinum* L.) seeds. *Journal of pharmacognosy and phytochemistry*. **6**(4): 72- 74.

Yadav, S.K., Dave, A., Sarkar, A., Singh, H.B. and Sarma, B.K. (2013). Co- inoculated biopriming with *Trichoderma*, *Pseudomonas* and *Rhizobium* improves crop growth in *Cicer arietinum* and *Phaseolus vulgaris*. *International Journal of Agriculture, Environment and Biotechnology*. **6**: 255-259.