

Case study Necrotising Soft Tissue Infection (NSTI), a Unique, Uncommon Complication of Pressure Ulcer – A Case Report.

ABSTRACT

Introduction: Infected pressure sore, also known as bedsore or pressure ulcer is a very common condition presented to hospitals especially in elderly and bedbound patients with reported incidence rate of 12%^[1]. According to Malaysian Registry of Intensive Care Report 2017, the rate of pressure ulcer following the admission to Intensive Care Unit (ICU) was 4.9 per 1000 ICU days^[2]. Managing infected bedsore is challenging and involve multidisciplinary care. Unfortunately, despite best possible efforts, infected pressure sore may progress into necrotising soft tissue infection (NSTI).

Case Report: We present a case of 73-year-old lady with infected pressure sore which turned out to be NSTI at gluteal region. She was treated aggressively with broad spectrum antibiotic and underwent multiple surgical debridement.

Conclusion: NSTI arising from pressure sores is rare but has high morbidity and mortality. Early diagnosis and treatment is crucial in handling such cases.

Keywords: pressure, ulcer, sore, necrotising, fasciitis

1. INTRODUCTION

Pressure sores are wounds over the skin and underlying tissues resulting from prolonged pressure over the affected area which usually affect those who are immobile for long periods or bedridden patients. This prolonged pressure, which are higher than capillary pressure leads to localized soft tissue ischemic necrosis^[3]. In time, if left untreated, the sores may grow large and lead to infections. Pressure sore are classified into four grades based on National Pressure Ulcer Advisory Panel (NPUAP) system as per Table 1.

Table 1 : Gradation based on National Pressure Ulcer Advisory Panel (NPUAP) system

Grade	Features
Grade I	Pressure sore with intact epidermis and non-blanchable redness over localised area, commonly

Comment [91]: Abstract has never contained references. Please correct.

Comment [92]: Abstract has never contained references. Please correct.

Comment [93]: Correct the sentence grammatically

Comment [94]: Rewrite the proper keywords

Comment [95]: Follow the uniformity in referencing

Comment [96]: Give the reference

	over bony prominent
Grade II	Pressure sore involving partial thickness loss of dermis with red wound bed and without slough
Grade III	Pressure sore involving full thickness tissue loss. Bone, tendon or muscle are not exposed. Slough may present.
Grade IV	Pressure sore involving full thickness tissue loss with exposed bone, tendon or muscle. Exposed bone or muscle is visible and can be palpable. Slough may present.

Necrotizing soft tissue infection (NSTI) previously known as necrotising fasciitis, is a type of life threatening, aggressive skin and soft tissue infection that causes necrosis of the muscle fascia and subcutaneous tissues. The rapid infection spreads initially along fascia plane which may then cause secondary infection of the overlying and underlying skin, soft tissue and muscle^[4].

NSTI rarely developed from pressure sore. Whether it is really rare or under-reported is not known. Herein we present such case in which pressure bed sore from immobilisation leads to infected bed sore and progress into NSTI.

2. CASE REPORT

A 73 year-old woman with underlying diabetes mellitus, hypertension and Parkinson's disease presented to local hospital with painful gluteal sore with foul smelling pus discharge which progressively worsening for the past 1 month. She had history of bilateral lower limbs cellulitis about 2 month ago which was treated with antibiotic. Since then, she was bedridden due to bilateral lower limb weakness. Physical examination revealed gluteal sore measuring 10 x 15 cm with necrotic patch (Figure 1A).

Blood investigations shows leucocytosis with white cell count (WCC) of 27×10^6 with high C-reactive protein (CRP) of 109.17. She was started on broad spectrum antibiotics and early surgical wound debridement was performed. Intraoperatively, unhealthy and necrotic tissue extend beyond the gluteal fascia with presence of fishy-odour smell. Gluteal maximus muscle over left side noted to be non-viable and gluteal medius appeared flaccid. Extensive wound debridement was done (Figure 1B). Tissue culture and sensitivity (C+S) intraoperatively showed presence of *Escherichia coli* (*E. coli*) and *Proteus mirabilis*. Antibiotic was changed and tailored according to sensitivity. Second wound debridement was performed on Day 12 admission due to persistent slough with necrotic tissue (Figure 1C and 1D). On Day 22 admission, third wound debridement was done due to persistent foul smelling discharge with necrotic tissue over superior medial and superior lateral quadrant of the wound despite daily modern dressing (Figure 1E, 1F and 1G). Post third wound debridement, wound appeared healthy with no pus discharge (Figure 1H). She was subsequently able to be discharged home with daily dressing at nearest clinic. Wound inspection at surgical outpatient clinic showed healthy granulation tissue with no slough or discharge (Figure 1I).




Figure 1: A case of gluteal sore progression to NSTI. (A) An infected gluteal sore with necrotic patch. (B) Finding intraoperatively showed necrosis tissue extending to fascia. (C) The patient's wound after a week of daily wound dressing. (D) Wound before second wound debridement shows slough with pocket. (E) The wound shown after second re-wound debridement, prior to third wound debridement. (F) Wound before third surgical debridement with necrotic and unhealthy tissue. (G) Wound prior third wound debridement. (H) Wound prior discharge shows clean, pink tissue. (I) Wound during follow-up.

3. DISCUSSION

Population with conditions such as diabetes, malignancies, obesity, alcoholics and post-surgical infections are at higher risk of developing NSTI. Most common risk factor is diabetes mellitus which account for 40 to 60%^[5]. Patient with diabetes are more susceptible to infections as higher glucose level gives an optimum medium for bacterial growth especially in a low oxygen tension environment.

NSTI itself is a rare condition with an incidence rate ranges from 0.3-15 cases per 100,000 population^[6]. A report has concluded that NSTI secondary to pressure sore is even uncommon.^[7] Patient with grade III and IV pressure sore are at higher risk to develop

NSTI.^[8] However, the aetiology of pressure sores evolve to necrotising fasciitis is yet, unclear.

Pressure ulcers only limits to pressured area which cause localized soft tissue ischemic necrosis meanwhile NSTI is differentiated by the extension of the infection. In pressure sores that develop into NSTI, the extension of infection is beyond the area of bony prominence and does not limited to the primary necrotic tissue caused by pressure induced ischemia.

Diagnostic criteria for infected pressure sore progressing to NSTI are not well established. The diagnosis can be determined based on presence of necrotic fascia and fat tissue. In addition, diagnosis can be made based on evidence found through direct observation during surgical debridement or by pathological findings or through evidence of presence of gas within soft tissue adjacent to pressure ulcer in computed tomography^[9]. Intraoperatively, fascia and muscle of a NSTI appear grey-blackish while infection pressure ulcers only display yellowish debris known as slough^[7].

The Laboratory Risk Indicator for Necrotizing Fasciitis (LRINEC) scoring system was developed to differentiate NSTI from the other severe soft tissue infections. Yet, the sensitivity of this scoring system ranged from 43% to 80% in different studies which may limit its use in clinical setting.

The principle of management of NSTI and a high grade infected pressure sore are the same. Prompt, aggressive wound debridement with early initiation of broad spectrum antibiotic are crucial. Surgical debridement is required to remove dead tissue and control the infection. In most of the cases, multiple surgeries are needed to fully control the wide extension of infection and to remove all of the dead tissue^[9].

In our study, diagnosis of NSTI was made through direct observation through surgical wound debridement as evidence by extension of necrotic tissue beyond the fascia. Our case was managed with a total of 3 extensive surgical wound debridement and 21 days course of intravenous antibiotics before being discharged home. At the same time, the patient was also reviewed by dietician, physiotherapy, occupational therapist, general medicine specialist and other relevant team to optimise healing and prevent secondary infection.

4. CONCLUSION

In conclusion, NSTI develop from pressure sore is a rare condition with high morbidity and mortality. Therefore, early diagnosis and treatment is of paramount importance. Aggressive surgical debridement, with early initiation of antibiotic are the gold standard in managing NSTI. As NSTI from pressure sores are more common to develop in a higher grade sores, care need to be taken seriously to prevent development of pressure sore in the first place. As the saying goes, prevention is always better than cure.

REFERENCES

- [1] Borojeny, L. A., Albatineh, A. N., Dehkordi, A. H., & Gheshlagh, R. G. (2020). The incidence of pressure ulcers and its associations in different wards of the hospital: a systematic review and meta-analysis. *International Journal of Preventive Medicine*, 11.
- [2] Tai Li Ling, Lim Chew Har, Mohd Ridhwan Mohd Nor, Nahla Irtiza Ismail, Wan Nasrudin Wan Ismail. (2017). *Malaysian Registry of Intensive Care 2017 report*.
- [3] Agrawal, K., & Chauhan, N. (2012). Pressure ulcers: Back to the basics. *Indian Journal of Plastic Surgery*, 45(02), 244-254.
- [4] Wallace, H. A., & Perera, T. B. (2017). Necrotizing fasciitis.
- [5] Misiakos, E.P. *et al.* (2014) *Current concepts in the management of necrotizing fasciitis*, *Frontiers in surgery*. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4286984/> (Accessed: 02 August 2023).
- [6] Tantirat, P., Rattanathumsakul, T., Praekunatham, H., Pachanee, K., & Suphanchaimat, R. (2020). Epidemiological situation of necrotizing fasciitis and factors in Thailand and factors associated with its morbidity and mortality, 2014–2018. *Risk Management and Healthcare Policy*, 1613-1624.
- [7] Esayag, Y., Brautbar, A., Popov, A., & Wiener-Well, Y. (2011). Necrotizing soft tissue infection: an unusual and devastating complication of pressure sores. *The Israel Medical Association journal: IMAJ*, 13(7), 442-443.
- [8] Citak, M., Backhaus, M., Tilkorn, D. J., O'loughlin, P. F., Meindl, R., Muhr, G., & Fehmer, T. (2011). Falcone, M., & Tiseo, G. (2023). Skin and soft tissue infections in the elderly. *Current Opinion in Infectious Diseases*, 36(2), 102-108.
- [9] Mizokami, F., Furuta, K., & Isogai, Z. (2014). Necrotizing soft tissue infections developing from pressure ulcers. *Journal of Tissue Viability*, 23(1), 1-6.